

# Spatial Dynamics of Malaria Transmission

## Authors

- Sean L. Wu<sup>†</sup> (seanwu89@uw.edu), Institute for Health Metrics and Evaluation, University of Washington
- John M. Henry<sup>†</sup> (henry529@uw.edu), Quantitative Ecology and Resource Management and Institute for Health Metrics and Evaluation, University of Washington
- Daniel T Citron<sup>†</sup> (daniel.citron@nyulangone.org), Department of Population Health, Grossman School of Medicine, New York University
- Doreen Mbabazi Ssebuliba (doreenresty@gmail.com), Department of Mathematics and Statistics, Kyambogo University.
- Juliet Nakakawa Nsumba (juliet.nakakawa@mak.ac.ug), Department of Mathematics, Makerere University Department of Mathematics, School of Physical Sciences, College of Natural Science, Makerere University, P.O. Box 7062, Kampala, Uganda.
- Héctor M. Sánchez C. (sanchez.hmsc@berkeley.edu), Divisions of Epidemiology and Biostatistics, School of Public Health, University of California Berkeley
- Oliver J. Brady (Oliver.Brady@lshtm.ac.uk) Centre for Mathematical Modelling of Infectious Diseases, London School of Hygiene & Tropical Medicine, London, UK; Department of Infectious Disease Epidemiology, Faculty of Epidemiology and Population Health, London School of Hygiene & Tropical Medicine, London, UK;
- Carlos A. Guerra (cguerra@mcd.org), MCD Global Health
- Guillermo A. García (ggarcia@mcd.org), MCD Global Health
- Austin R. Carter, aucarter@uw.edu, Institute for Health Metrics and Evaluation, University of Washington
- Heather M. Ferguson (Heather.Ferguson@glasgow.ac.uk), Faculty of Biomedical and Life Sciences, University of Glasgow
- Bakare Emmanuel Afolabi (emmanuel.bakare@fuoye.edu.ng); International Centre for Applied Mathematical Modelling and Data Analytics, Federal University Oye Ekiti, Ekiti State, Nigeria; Department of Mathematics, Federal University Oye Ekiti, Ekiti State, Nigeria
- Simon I. Hay (sihay@uw.edu), Department of Health Metrics Science and Institute for Health Metrics and Evaluation, University of Washington

It is made available under a [CC-BY 4.0 International license](#) .

- Robert C. Reiner Jr. (bcreiner@uw.edu), Department of Health Metrics Science and Institute for Health Metrics and Evaluation, University of Washington
- Samson Kiware (skiware@ihi.or.tz), Ifakara Health Institute
- David L Smith<sup>‡</sup> (smitdave@uw.edu), Department of Health Metrics Science and Institute for Health Metrics and Evaluation, University of Washington

<sup>†</sup> authors contributed equally; <sup>‡</sup> corresponding author

**Target Journal** PLoS Computational Biology

## Author summary

The Ross-Macdonald model, a simple mathematical model of malaria transmission based on the parasite life-cycle, established basic theory and a set of metrics to describe and measure transmission. Here, we extend the Ross-Macdonald model so it has the skill to study, simulate, and analyze parasite dispersal and heterogeneous malaria spatial transmission dynamics in a defined geographical area with malaria importation. This extended framework was designed to build models with complexity that scales to suit the needs of a study, including models with enough realism to support monitoring, evaluation, and national strategic planning. Heterogeneity in human epidemiology or behaviors – differences in age, immunity, travel, mobility, care seeking, vaccine status, bed net use, or any trait affecting transmission – can be handled by stratifying populations. Mosquito spatial ecology and behaviors are responding to heterogeneous resource availability and weather, which affects adult mosquito dispersal, blood feeding, and egg laying in a structured set of aquatic habitats. We propose new formulas for human biting rates and entomological inoculation rates that integrate exposure as humans move around. We rigorously define parasite dispersal, and we develop matrices describing the spatial dimensions of vectorial capacity and parasite dispersal in mobile humans. We relate these to the parasite’s overall reproductive success, local reproductive numbers and thresholds for endemic transmission. [185/200 words]

## Abstract

The Ross-Macdonald model has exerted enormous influence over the study of malaria transmission dynamics and control, but it lacked features to describe parasite dispersal, travel, and other important aspects of heterogeneous transmission. Here, we present a patch-based differential equation modeling framework that extends the Ross-Macdonald model with sufficient skill and complexity to support planning, monitoring and evaluation for *Plasmodium falciparum* malaria control. We designed a generic interface for building structured, spatial models of malaria transmission based on a new algorithm for mosquito blood feeding. We developed new algorithms to simulate adult mosquito demography, dispersal, and egg laying in response to resource availability. The core dynamical components describing mosquito ecology and malaria transmission were decomposed, redesigned and reassembled into a modular framework. Structural elements in the framework – human population strata, patches, and aquatic habitats – interact through a flexible design that facilitates construction of ensembles of models with scalable complexity to support robust analytics for malaria policy and adaptive malaria control. We propose updated definitions for the human biting rate and entomological inoculation rates. We present new formulas to describe parasite dispersal and spatial dynamics under steady state conditions, including the human biting rates, parasite dispersal, the “vectorial capacity matrix,” a human transmitting capacity distribution matrix, and threshold conditions. An **R** package that implements the framework, solves the differential equations, and computes spatial metrics for models developed in this framework has been developed. Development of the model and metrics have focused on malaria, but since the framework is modular, the same ideas and software can be applied to other mosquito-borne pathogen systems.

## Introduction

*Plasmodium falciparum* transmission dynamics are complex: they involve multiple-agents, non-linear dynamics, localized spatial interactions, spatial, temporal and behavioral heterogeneity, stochasticity, and exogenous forcing by weather, hydrology, and malaria control. Over time, these processes can be modified by economic development; by changing socioeconomic status, human incentives and social norms; and by the evolution of resistance. Every one of these features of malaria transmission dynamics and control presents its own set of challenges to the quantitative study of malaria for scientific research and for analytics to support policy. An important practical problem is how to quantify and synthesize all of the factors affecting transmission at some particular place and time to support malaria control programs in various ways, including monitoring and evaluation of malaria control. The study of complex spatial processes are best addressed using some sort of mathematical model. Here, to fill a need to give robust policy advice, we have developed a modular framework with accompanying software to build and analyze suites of models with scalable complexity for malaria spatial transmission dynamics and control.

A starting point for the quantitative study of malaria transmission dynamics has been the Ross-Macdonald model, which played a central role in developing basic theory and metrics for malaria [1, 2]. That model is simple, general, and conceptually useful, but it is not realistic enough to describe many important features of transmission [3]. The model's lack of realism has also limited its applicability: simple models support generic policy advice, but specific advice – tailored to context – must be based on models that can quantify and weigh the effects of locally relevant details [4]. A basic limitation of the Ross-Macdonald model was that it lacked features required to describe spatial transmission dynamics and control. Mathematical models for spatial dynamics of mosquito-borne pathogens have been developed [5–18], but there is a need for a generalized synthetic framework to develop and use spatial dynamic models, to extend the Ross-Macdonald model to define and analyze parasite dispersal, to define and measure malaria connectivity [19], and to link spatial dynamics to spatial data. The Ross-Macdonald model is also missing other features that are relevant for malaria dynamics and control, which can be identified from a survey of studies that have modeled mosquito-borne diseases (see Box #1) [2]. Modeling and analyzing real systems can become overwhelming because of computational, parametric, or conceptual challenges that arise from combining all the factors, dimensions, interactions, features, and processes. Individual-based models (IBMs) have been developed around algorithms that make it possible to deal with the complexity by simulating individual states and transitions *in silico* [20], but these high-dimensional computational approaches have some limitations that limit their use and applicability. IBMs require intensive computation, are challenging to parameterize, are difficult to critically evaluate, and their output that is often as difficult to analyze and understand as malaria itself. Using a modular framework, we present an alternative way of dealing with the complexity that is analytically tractable, including some new algorithms to understand mosquito ecology, parasite transmission by mosquitoes, and parasite dispersal on spatial landscapes.

In most places, malaria transmission has been modified by control. The extent of effect modification by malaria control is occasionally revealed when health systems are

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

disrupted (e.g., [21,22]), when malaria control is relaxed or abandoned [23], or when 46  
resistance evolves to drugs or insecticides (e.g., [24,25]). Programs must weigh evidence 47  
and make decisions through analysis of counterfactuals, rather than through direct 48  
estimation of control effect sizes, since there would be drastic consequences to 49  
experimentally disrupting control. A predominant need in most contexts is thus a set of 50  
methods to quantify transmission in its local context as a baseline that has been 51  
modified by control. A challenge to achieving this has been that the responses to 52  
control efforts are context dependent and have been highly variable across settings. 53  
Relevant factors affecting responses to control include details about blood feeding, 54  
mosquito ecology, and mosquito behaviors that affect contact with interventions (*e.g.*, 55  
resting indoors and IRS). To reconstruct the counterfactual baseline, transmission must 56  
be understood in terms of innate mosquito behaviors responding to local resources, 57  
vector control, and other contextual factors that have been modified by control. All 58  
these have been characterized as being notoriously context dependent and 59  
heterogeneous [26–28]. What are the local factors that determine baseline malaria 60  
transmission, effect modification, and differences in effect modification at some 61  
particular place and time? Basic concerns about the heterogeneous impacts of vector 62  
control raise a larger set of questions about how to study and quantify transmission in a 63  
way that is relevant for planning malaria control. 64

This new framework is thus an attempt to bridge two well-established but somewhat 65  
contradictory views of malaria. One view is that human malaria transmission dynamics 66  
and control are so moulded by local ecology and other conditions that the factors 67  
driving transmission or responses to control at one time and place are unlikely to hold 68  
elsewhere [27]. Another view – encouraged by the rigorous analysis of the 69  
Ross-Macdonald model and extensions of it – is that malaria transmission intensity can 70  
be quantified using a small set of bionomic parameters to compute basic reproductive 71  
numbers, which also provide a basis for computing threshold conditions for endemic 72  
malaria. To build a bridge, the contextual factors affecting basic bionomic parameters 73  
must be identified and integrated with new theory describing spatial extensions of the 74  
basic metrics, including rigorous, quantitative description of parasite dispersal, and 75  
some estimates of the appropriate spatial scales to measure malaria transmission [3]. 76

Context-dependency is an uncomfortable but unavoidable fact of malaria ecology. 77  
The heterogeneous nature of transmission and the causes and consequences of variable 78  
responses to control have been a difficult and sometimes contentious problem for 79  
scientists studying malaria, for national malaria programs and funding agencies making 80  
malaria policy, and for malaria advocates. Historical trends in malaria and the outcomes 81  
of malaria control have been so variable that case studies can be found to support rosy 82  
projections, alarmist warnings, or contradictory claims about the underlying causes of 83  
trends or patterns. To be useful, studies of malaria and programmatic evaluations must 84  
acknowledge the important role of context, the multi-factorial nature of causation in 85  
these complex systems, non-linear responses to control, the difficulty of measuring 86  
heterogeneous systems, and the resulting uncertainty. A consequence of context 87  
dependency is the difficulty in drawing conclusions that generalize across systems. 88

The framework is designed to support development of robust malaria policy advice 89  
and to find practical ways of dealing with uncertainty. While scientific research and 90  
policy analytics grapple with the same issues and use similar methods, they often put 91

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

very different weights on uncertainty. Uncertainty affects the ability to do effective inference for scientific research *versus* policy analytics – questions about what is known *versus* what should be done. To address these concerns and give policy advice despite uncertainty, an integrated inferential framework is needed to weigh evidence, integrate the effects of multiple exogenous factors (often involving experts from distinct specialties), estimate their effect sizes, quantify uncertainty, and identify critical gaps. Statistical theory and inferential methods have been developed around the principle of parsimony for scientific inference, but substantially less attention has been given to appropriate designs for analyses that can give advice that is robust to uncertainty. Are the conclusions of an analysis robust to reasonable alternative formulations of a model, and how well are policy recommendations really supported by the evidence? Concerns about robustness could lead to study designs that make different tradeoffs between realism and abstraction. For example, compared with parsimonious models, models with a high degree of realism might be more useful for identifying critical missing data and prioritizing studies to collect it. Robust analytics requires having a modeling framework to build suites of models that are realistic enough to weigh the importance of the major drivers of transmission despite major knowledge gaps.

To address these needs, we have developed a new, modular framework designed to support development of models for robust, simulation-based analytics and adaptive malaria control with scalable complexity. With scalable complexity in model building, members of a model ensemble could range from very simple to very complex, and that models along that spectrum are related to one another through a logical sequence of structural or parametric changes. At one extreme, this framework includes the Ross-Macdonald model, a simple system of differential equations describing the parasite life-cycle in mosquito and vertebrate host populations linked by transmission during blood feeding [1, 29, 30]. By extending the Ross-Macdonald model, simple models can be extended step by step to add complexity or heterogeneity that could be important – based on *a priori* considerations – yet difficult to quantify or poorly informed by existing data (Box #1). With modularity, it is possible to develop new dynamical systems models describing some parts of the system, add or modify components, or add a set of exogenous factors that force a system. It is also relatively straightforward to modify functional responses, or to modify some basic parameters affecting the outcome. Swarms of models can thus be developed to analyze data and to test the robustness of any conclusions. To demonstrate scalable complexity, we here present a complicated family of models that has terms and variables anticipating modification by weather or malaria control. For practical reasons, the model family we present here was scaled back to include a limited set of elements describing transmission, but leaving in place the elements that facilitate modeling control (Box #1). The resulting extensible framework that is capable of describing and analyzing malaria spatial transmission dynamics and control with a high degree of realism in any particular setting. An **R** package which implements the modular differential equations and spatial metrics presented in the article is available with documentation (Supplement 1 - <https://dd-harp.github.io/exDE/>). Despite being programmed in R, the implementation of the mathematical framework into code should be easily adapted to any high-level programming language.

In **Framework**, we first present the modular concepts and structural elements,

It is made available under a [CC-BY 4.0 International license](#) .

including a new blood feeding model. Next we present one exemplar model family for 138  
each dynamical component. In **Spatial Metrics**, we develop a set of metrics that 139  
describe various aspects of parasite spatial dynamics, including metrics for parasite 140  
dispersal, connectivity, and the parasite's reproductive success. Finally, in 141  
**Quantifying Transmission in a Place**, we discuss the application of these models to 142  
the investigation of malaria transmission dynamics and control in a particular place. 143



**Box 1: Features** This generalized, modular framework presents equations integrating multiple agents and interacting processes. Many of these innovations appeared first elsewhere, but here they are integrated into a single framework:

- Immature mosquito population dynamics structured in distinct aquatic habitats linked to adult populations through egg laying and emergence [31,32];
- Spatially heterogeneous blood feeding and parasite mixing on vertebrate populations (*i.e.*, blood hosts) with dynamically changing availability, such that feeding rates and the human fraction change adjust to changing conditions [33–36];
- Heterogeneous adult mosquito behaviors, including dispersal, survival, blood feeding, egg laying, mating, and sugar feeding on landscapes in response to spatially heterogeneous resource availability (*e.g.*, mating sites, sugar sources, blood hosts, aquatic habitats) [37–39];
- Multiple vector species or types with different host preferences, daily activity patterns, habitats, *etc.* [40], and potentially with inter-specific resource-based competition in habitats;
- Human mobility based on a concept of time at risk, which combines time spent by humans in places where they are at risk with mosquito blood feeding activity, preferences and other factors [9,18];
- The capability to model indoor and outdoor spaces for blood feeding, exposure, and vector control;
- A non-linear relationship between the daily entomological inoculation rate (EIR) and the daily force of infection (FoI) due to heterogeneous exposure [41].
- Malaria importation through multiple routes [42];
- An exogenously forced, time-varying extrinsic incubation period (EIP) to model effects of temperature on parasite development;

The model has flexible structural elements to stratify an area into patches, to model any distribution of aquatic habitats, and to stratify a human population into sub-populations by age, immunity, or any heterogeneous, epidemiologically relevant trait. The software also includes time-dependent terms and structures to model exogenous forcing by weather, modification of exposure or transmission by vector control in relation to coverage, including effects of spatial repellents and mosquito behaviors that result in heterogeneous local contact patterns with vector-based interventions.

## Framework

To describe malaria spatial dynamics with scalable complexity, we designed a modular framework for model building around four core dynamical components, each one a (potentially non-linear) state-space model. An interface rigidly defines interactions among those components, based on passing terms we call *dynamical quantities*. All state variables are vectors of arbitrary length, to accommodate models with different structure or spatial granularity.

To model mosquito ecology, we consider immature mosquitoes in a set of aquatic habitats, and adult mosquitoes in a set of patches. A state space model describes aquatic immature mosquito populations ( $\mathcal{L}$ ) with dynamics  $d\mathcal{L}/dt$  requiring an input term from adult mosquito populations: the daily rate eggs are laid in each habitat ( $\eta$ ). A coupled state space model describes mature adult female mosquito populations ( $\mathcal{M}$ ) with dynamics  $d\mathcal{M}/dt$  requiring an input term from the aquatic mosquito populations: the rate adults emerge from all the habitats in each patch ( $\Lambda$ ). A state space model for parasite infection dynamics in mosquitoes ( $\mathcal{Y}$ , which extends  $\mathcal{M}$ ) with dynamics  $d\mathcal{Y}/dt$ , requires an input term from human malaria epidemiology: the net infectiousness of humans (NI), the probability a mosquito becomes infected after blood feeding on a human (denoted  $\kappa$ ). A state space model describing parasite infection dynamics in humans, immunity, and disease ( $\mathcal{X}$ ) with dynamics  $d\mathcal{X}/dt$ , requires an input term from adult mosquito infection dynamics: the daily EIR ( $E$ ). The inputs to one component can be passed as trace functions or as the outputs of another coupled component, which is called the *interface* of each dynamical component; a generic interface is coded for each term and if needed specialized methods can be written for particular models. Models in the framework have the following form:


$$\begin{aligned} d\mathcal{L}/dt &= F_{\mathcal{L}}(\eta, \mathcal{L}) \\ d\mathcal{M}/dt &= F_{\mathcal{M}}(\Lambda, \mathcal{M}) \\ d\mathcal{Y}/dt &= F_{\mathcal{Y}}(\kappa, \mathcal{M}, \mathcal{Y}) \\ d\mathcal{X}/dt &= F_{\mathcal{X}}(E, \mathcal{X}) \end{aligned} \quad (1)$$

The interactions among these dynamical components are thus defined by four input terms ( $\eta$ ,  $\Lambda$ ,  $\kappa$ , and  $E$ ), which may be computed as outputs of another component or provided as an external forcing term (Fig. 1). Because these terms can be computed from the state of the model and are used to couple different model components together, we call these dynamical quantities. These terms are rates which determine how components interact (*e.g.*, flows between components). Because construction of these dynamical quantities can be done in a generic way, computation of these quantities in code can be done for any model which fulfills the interface of its dynamical component.

The dynamical quantities responsible for transfer of pathogens between hosts and vectors are  $E$  and  $\kappa$ , the EIR and NI of humans, respectively. These quantities couple the dynamics between the human  $\mathcal{X}$  and mosquito  $\mathcal{Y}$  dynamical components. To allow computation of  $E$  and  $\kappa$  to be highly generic across various types of models of human and mosquito infection, we developed a new model of blood feeding which produces  $\beta$ , the biting distribution matrix describing how bites arising from mosquitoes at patches are taken on human population strata.

Similarly, the adult  $\mathcal{M}$  and aquatic  $\mathcal{L}$  mosquito components are coupled via egg laying from adults in aquatic habitats, and emergence of new adults from those aquatic habitats. Because the patches where adult mosquitoes are found may contain many (or no) aquatic habitats, another matrix translates the rate of egg production from adults into egg deposition in each aquatic habitat  $\eta$ . Likewise, each aquatic habitat produces newly emerging adult mosquitoes at some rate  $\alpha$ , which in general depends on the current aquatic population, and therefore on lagged adult densities. Another matrix maps this into the rate at which new adults are added to each mosquito population,  $\Lambda$ .

In addition to reformulating blood feeding and egg laying, the framework includes mathematical descriptions of survival, search for blood hosts or habitats, and dispersal. These new models of adult mosquito behaviors have all been reformulated around the concept of heterogeneous resource availability and functional responses to available resources.



**Fig 1.** Models for malaria transmission dynamics are naturally modular (see Eq. 1). The dynamic modules describe a stratified human population (purple) that interacts through blood feeding (red) with adult mosquito populations in a discrete spatial domain; each patch could contain a set of aquatic habitats. Two components,  $\mathcal{L}$  and  $\mathcal{M}$ , describe mosquito ecology: dynamics of immature mosquitoes (blue) in aquatic habitats are described by a system of equations  $d\mathcal{L}/dt$ ; and dynamics of adult mosquitoes (green) are described by  $d\mathcal{M}/dt$ . Habitat locations within patches are described by a membership matrix,  $\mathcal{N}$ . Eggs hatch into larval mosquitoes, that develop, pupate, and later emerge from habitats as mature adults ( $\alpha$ ) and added to the adult populations in each patch ( $\Lambda$ ). Adults lay eggs ( $\nu$ ), which are distributed spatially according to which patch habitats belong ( $\mathcal{N}$ ). Egg deposition rates at the habitats are ( $\eta$ ). Two additional components,  $\mathcal{Y}$  and  $\mathcal{X}$ , describe parasite infection dynamics and transmission: that for mosquitoes, described by  $d\mathcal{Y}/dt$  and in humans, described by  $d\mathcal{X}/dt$ , are linked through parasite transmission. A new model for blood feeding describes how blood meals are allocated among humans ( $\beta$ ) and associated parasite transmission rates: the density of infectious humans by strata ( $X$ ) is used to compute net infectiousness (NI) of humans to mosquitoes in patches ( $\kappa$ ); and the density of infectious blood feeding mosquitoes ( $Z$ ) is used to compute the entomological inoculation rate (EIR) on each strata ( $E$ ).

The modular framework was implemented as a software package in **R** [43] (Supplement 1 is the website <https://dd-harp.github.io/exDE/>). The software builds dynamical models of malaria in a modular way using method dispatch to define generic code which implements the framework described here. The dynamical models are functions which return arrays of derivatives of state variables, and can be solved using the integrators available in *deSolve*, or other tools in **R** [43, 44]. The software also

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

includes routines that compute steady state conditions and spatial metrics (see Spatial Metrics, below). Because each component has an interface – the generic functions that compute and pass of dynamical quantities between components – any new model can be implemented which fulfills a specific interface, independent of the rest of the framework. In this way, building and testing new models of particular components is straightforward, and the framework is flexible and extensible. As new models are required, they will be added to the package, increasing its applicability and scope over time.

We have developed a glossary of terms (Supplement 2). In the equations that follow, for each dynamical component, we describe the model structure in detail, and we present one family of models describing transmission dynamics in a single vector species. In Supplement 3, we formulate a model using both conventional notation and the modular notation of this framework. In Supplement 1, we have implemented a previously published model of malaria transmission on Bioko Island [45]. In Supplement 4, we extend the discussion of vector dynamics, including a discussion of models with multiple vector species. All the terms and parameters may be time dependent to accommodate seasonality or modification by exogenous factors: seasonal travel, exogenous forcing by weather, and parameter modification by vector control. Analysis of temporal heterogeneity in this same framework is outside the scope of this study, it but would be straightforward extension following approaches analogous to those shown in the supplements.

**Box 2: Notation** Equations describing spatial processes include terms describing scalar quantities, vectors of scalars, vectors of functions, and matrices. We have avoided using any notation to designate a vector or indicate it could be time-dependent, in part, because it would be ubiquitous; most parameters could vary by space and time. The most general form of a term or parameter is usually described when it is first presented, but most terms describing a vector or matrix should be assumed to be modifiable. In writing out the equations, we consistently use the center dot, “.”, in equations to denote the dot product of two matrices, or a matrix and a vector. The juxtaposition of two vectors denotes element-wise product, and  $1/*$  denotes the vector of the inverses of each element. The symbol  $\odot$  denotes the Hadamard product (*i.e.*, element-wise multiplication) of two matrices. When  $x$  is a vector,  $\text{diag}(x)$  is a matrix with the elements of  $x$  on the main diagonal. The identity matrix is denoted  $I$ , and  $\mathbf{1}$  denotes a row or column vector with each element equal to 1. When  $F$  is a functional response, we assume it accepts vector arguments and returns a vector of the same length, *i.e.*,  $|F(X)| = |X|$ . The glossary (Supplement 1) discusses the dimensions of each term.

## Model Structure

The following describes, in detail, the structural elements and the algorithms that connect them. Adult mosquito and human population strata are connected through blood feeding and transmission, and adult and aquatic mosquito populations are connected through egg laying and emergence.

**Structural Elements** The framework has been designed to build model ensembles with the goal of studying the spatial transmission dynamics of malaria in a defined geographical area, called the spatial domain. An important part of this framework is having flexibility in defining the model structure to describe spatial and population heterogeneity at the appropriate level of detail, depending on the needs of a study and the available data. The structural elements – the patches, the aquatic habitats, and the population strata – were designed to handle arbitrary patch definitions, arbitrary human population residency patterns and stratification, and arbitrary numbers and locations of aquatic habitats.

To deal with spatial heterogeneity in transmission, we subdivide the spatial domain and identify a set of  $p$  patches that includes all locations relevant for studying and quantifying mosquito ecology or transmission: places where people live; places where mosquitoes blood feed; or places with aquatic habitats where mosquitoes lay eggs. We assume that there are  $l$  aquatic habitats with actual physical locations that are nested within the patches. To deal with heterogeneity in the human population, the model accommodates stratification. The human population is sub-divided into a set of  $n$  population strata by residency, immunity, behaviors affecting risk, or any other epidemiologically relevant factors (Supplement 5). Human populations are assigned a single residency patch, where they live and spend most of their nights. Other subdivisions of the human population could take into account age, sex, travel patterns, ITN usage, or any trait that is heterogeneous and epidemiologically relevant. The total census population size, the number of people who reside in each patch in the spatial domain, is given by a vector denoted  $P$  (of length  $p$ ). The number of people in each stratum is given by a vector  $H$  (of length  $n$ ). In this model, it is not necessary for every patch to have some residents.

To manage terms for interactions among structural elements, we create two mathematical objects called membership matrices that aggregate quantities to patches (Supplement 3). Since the  $l$  aquatic habitats are scattered among the patches, we define the habitat membership matrix  $\mathcal{N}$ , a  $p \times l$  matrix, that aggregates quantities from the  $l$  aquatic habitats to  $p$  patches where they are found. Similarly, we define the strata membership matrix  $\mathcal{J}$ , a  $p \times n$  matrix, that aggregates the  $n$  human population strata to the  $p$  patches where they reside. The census population size, for example, is  $P = \mathcal{J} \cdot H$ . If a human population were stratified by other traits, such as frequent travel or age, a membership matrix could be created to aggregate model output by trait.

The framework has also been designed to accommodate models with multiple mosquito vector species or types (see Supplement 4). Most of the following discussion assumes there is just one vector species, but we point out where the framework has can generalize to multiple vector species.

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

**Human Mobility** After defining the model structure (*i.e.*, the patches and population strata), the next challenge is to construct the algorithms describing local human mobility and travel. Local mobility determines where and when humans are available and exposed to blood feeding mosquitoes within the spatial domain. We define *travel* in this model by time spent outside the spatial domain; travel and mobility are thus different modalities and handled with different constructs.

To model local human mobility patterns within the patches, we develop a model describing the fraction of time spent by humans in each stratum among the patches [9, 18]. The information is summarized in a time-dependent  $p \times n$  matrix  $\Theta(t)$ , called the *Time Spent* (TiSp) matrix (Supplement 5). Each column in a TiSp matrix describes the fraction of time spent in each patch by an individual from a single stratum. In formulating the TiSp matrix, we account for time spent by time of day in the patches where mosquitoes are blood feeding. Total time spent should subtract time spent traveling and time spent in the spatial domain in places where there is no risk (*e.g.*, in office buildings).

Blood feeding combines human and mosquito behaviors. Since mosquito blood feeding has a daily rhythm [46], time at risk modifies time spent to account for differences in mosquito daily blood feeding activity rates. We let  $\xi(t)$  denote a species-specific *circadian weighting function* for blood feeding rates, constrained such that  $\int_0^1 \xi(t) dt = 1$ , which appropriately assigns a *weight* to time spent by time of day (Supplement 5). Using  $\xi$ , we compute the *Time At Risk* (TaR) matrix as time spent weighted by mosquito activity:  $\Psi(t) = \text{diag}(\xi(t)) \cdot \Theta(t)$ .

This distinction between TiSp and TaR matrices makes it possible to study human mosquito contact in detail, to quantify differential transmission by multiple vectors with the same human mobility patterns, and to quantify other aspects of mosquito-human contact [47, 48]. A model could have two or more vector species, each with different blood feeding patterns ( $\xi_1$  and  $\xi_2$ ), so that one TiSp matrix would be transformed into two different TaR matrices ( $\Psi_1 = \xi_1 \Theta$  and  $\Psi_2 = \xi_2 \Theta$ ).


**Denominators and Availability** After defining host population movement, it is necessary to compute appropriate denominators to model blood feeding, based on the models for time spent and time at risk. Because of mobility, mosquito preferences, and human behaviors, the denominators for blood feeding are different from the resident population size – the number that would be used by most studies (Fig. 2).

An important intermediate quantity is ambient population density, which describes the population present in patches at a point in time. In a mobile population, the ambient population density will tend to be different from resident population density. From the time spent matrix, the ambient population density is a vector of length  $p$  given by:

$$A(t) = \Theta(t) \cdot H. \tag{2}$$

Similarly, ambient population density at risk is given by:  $\Psi(t) \cdot H$ . One way to understand what the TiSp matrix means is by taking ratios of ambient to resident populations. The ambient density of residents is  $A_r = (\mathcal{J} \odot \Theta) \cdot H$ , where  $\odot$  denotes the Hadamard (element-wise) product. The non-resident, non-visitor, ambient population is  $A - A_r$ . The ratios of various census and ambient population densities (*e.g.*, the ratio of

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).



**Fig 2. Denominators and Mixing** A schematic diagram relating various concepts of population density under a model of human mobility, resulting in a biting distribution matrix,  $\beta$ . Here, and in Figures 3-6, rounded rectangles denote endogenous state variables, sharp rectangles denote endogenous dynamical quantities, and parallelograms represent exogenous data or factors. Purple indicates the element is related to human populations, green for mosquitoes, and red for biting and transmission. Population strata ( $H$ ) describe how persons are allocated across demographic characteristics. The matrix  $\mathcal{J}$  distributes these strata across space (patch), according to place of residency. By combining information on how people spend their time across space ( $\Theta(t)$ ) and mosquito activity ( $\xi(t)$ ) a time at risk (TaR) matrix  $\Psi$  is generated describing how person-time at risk is distributed across space. Because blood feeding can be modified by human and mosquito factors (*e.g.*, net use and biting preferences), search weights ( $w_f(t)$ ) may further weight person-time at risk. The final result is a biting distribution matrix  $\beta$ , which is the fraction of each bite in each patch that would arise on an individual in each stratum, so  $\text{diag}(H) \cdot \beta = 1$ .

residents to ambient population  $P/A$ , defined wherever  $A > 0$ ), can be used to understand and diagnose unrealistic terms in a TiSp or TaR matrix. The ambient population thus provides one easy statistic to understand TiSp or TaR matrices.

To model the denominators for blood feeding, we also consider other factors – mosquito preferences or human behaviors or traits such as ITN usage – that affect host availability to mosquitoes and relative biting rates on the strata [33]. We assign biting weights,  $w_f$ , to each strata, where we think of  $w_f = 1$  as the value that would be assigned to an average person under baseline conditions (*e.g.*, without a net). These weights affect both the total biting rates and the relative biting rates on the ambient population. We define the *availability* of the host populations to mosquitoes for blood feeding as:

$$W = \Psi \cdot w_f H. \tag{3}$$

Availability is thus defined in units of weighted person-days at risk, and  $W$  is a vector of length  $p$  describing total human availability in each patch.

We also consider the presence of a population of visitors, a non-resident population spending time in the spatial domain (Supplement 5). We assume that some visitors could be present, and that some of them could be infectious. We can let  $A_\delta$  denote the ambient density of visitors, but we let  $W_\delta$  denote their availability by patch. The *resident fraction* or fraction of human blood meals taken on a resident in each patch, a vector of length  $p$  denoted  $v$ , is:


$$v = \frac{W}{W + W_\delta}. \tag{4}$$



It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

The total availability of humans for blood feeding, in each patch, is thus  $W + W_\delta$ .

330



**Fig 3. Blood Feeding and Human Biting Rates** The daily human biting rates (HBR) for the resident population strata are defined as the expected number of bites by vectors, per person, per day. To compute the HBR, we count up exposure over all the patches where residents spend time. We also consider the presence of visitors and other bloodhosts (yellow input), which increases the total available hosts.

**Blood Feeding** With a well-defined population denominator, we can compute the frequency of blood feeding rates and the human fraction (*i.e.*, the fraction of human blood meals among all blood meals) in each patch in response to the availability of humans and other available vertebrate hosts. To do so, we use functional responses to model blood feeding rates and habits [33–36].

331  
332  
333  
334  
335

Human availability,  $W$ , is often highly variable among patches and over time, which could affect the rate mosquitoes blood feed (Fig. 3). Mosquitoes could also feed on other vertebrate hosts. To model blood feeding, we supply a vector of functions describing the availability of non-human vertebrate hosts in each patch over time, denoted  $O(t)$ . We assume that mosquito preferences could scale with host densities, so we assign a shape parameter,  $\zeta$ , that modifies how preferences scale with host densities. Total availability of all vertebrate hosts for blood feeding is  $B = W + W_\delta + O^\zeta$  (Supplement 5).

336  
337  
338  
339  
340  
341  
342

Let  $f(t)$  denote the blood feeding rate, the number of blood meals, per mosquito, per day. To guarantee mathematical consistency in computing blood feeding rates (*e.g.*, if  $B = 0$ , then it should be true that  $f = 0$ ), we can model time-dependent blood feeding rates, where  $f(t)$  is a vector of length  $p$ , as:

343  
344  
345  
346

$$f(t) = F_f(B) = f_x \frac{s_f B}{1 + s_f B}. \quad (5)$$

Depending on a shape parameter(s),  $s_f$ , blood feeding rates increase with host availability up to a maximum (or maxima)  $f_x$ , which is limited by the time it takes to search, process the blood meal, lay eggs, and perhaps to sugar feed. The fraction of blood meals taken on humans at a point in time, a vector of length  $p$  denoted  $q(t)$ , is called the human blood feeding fraction or human fraction:

347  
348  
349  
350  
351

$$q(t) = \frac{W + W_\delta}{B}. \quad (6)$$

The local human fraction, the fraction feeding on resident humans, is thus  $vq = W/B$ . The functional forms guarantee that when no humans are present, it must be true that  $f q = 0$ ; and when only humans are available, it must be true that  $q = 1$ .

352  
353  
354



It is made available under a [CC-BY 4.0 International license](#) .

**Mixing and Parasite Transmission** The model for mixing is an answer to the question: How are blood meals in a patch allocated among humans in the strata? The time at risk matrix and the factors affecting blood feeding rates and habits in each patch must be consistent with the algorithm that computes the distribution of biting and parasite mixing.

To allocate mosquito bites in patches among the resident strata, we let  $\beta$  denote a  $n \times p$  biting distribution matrix:

$$\beta(t) = \text{diag}(w_f) \cdot \Psi^T \cdot \text{diag}\left(\frac{1}{W(t)}\right). \quad (7)$$

Each column of  $\beta$  describes the fraction of a bite in a patch that lands on an individual in each strata, so the matrix  $\text{diag}(H) \cdot \beta$  gives the fraction of bites that land on each stratum, and its columns sum to unity.

In the models for mosquito ecology and infection dynamics, we define variables (vectors of length  $p$ ) for the density of mosquitoes ( $M$ ) and infectious mosquitoes ( $Z$ ). From these, we derive an expression for the daily human biting rate (HBR) and entomological inoculation rate (EIR) for all the strata. The sporozoite rate (SR) in each patch is given by:

$$z = \frac{Z}{M}. \quad (8)$$

The net per-capita human blood feeding rates in each patch, or  $f q M / W$ , are hereafter called the patch HBR (pHBR), and  $f q Z / W$  is hereafter called the patch EIR (pEIR) for infectious mosquitoes. By way of contrast, exposure risk for the strata – the HBR and EIR – are defined as the number of bites / infectious bites by vectors, per person, per day. The HBR is  $\beta \cdot f q v M$ , and the EIR is the product of the HBR and the SR, or

$$E = \beta \cdot f q v Z. \quad (9)$$

To draw a sharp contrast between the terms, the pHBR and pEIR describe the number of bites / infectious bites, per person, in patches. They are stratified by location, so they are vectors of length  $p$ . The HBR and the EIR are stratified quantities that sum exposure over all locations for the strata, so they are vectors of length  $n$ .


Each model for parasite infection dynamics in humans defines a quantity,  $x$ , the probability a mosquito becomes infected after biting a human in each stratum. The quantity  $X = xH$ , a vector of length  $n$ , is herein called the *infective density* of infectious human residents. We can also specify the infective density of visitors,  $X_\delta$  where  $X_\delta = x_\delta W_\delta$  is intrinsically using the availability of visitors. The net infectiousness (NI) for the mosquito populations in all the patches, denoted  $\kappa$ , is:

$$\kappa = v\beta^T \cdot X + (1 - v)X_\delta \quad (10)$$

The force of infection for the mosquito population is thus  $f q \kappa$ .

**Egg Laying** To compute quantities affecting mosquito ecology and population dynamics, we need to formulate algorithms to compute egg laying rates and egg laying distributions: how many eggs are laid by adult mosquitoes in a patch, and how are they

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .



**Fig 4. Egg Laying and Egg Deposition** The availability of aquatic habitats ( $Q$ ) the patch sum of habitat search weights ( $Q = \mathcal{N} \cdot w_\nu$ ), and the egg distribution matrix ( $\mathcal{U}$ ) describes the locally normalized search weights. Available habitat determines per-capita oviposition rates ( $\nu$ ) by the population of gravid mosquitoes ( $G$ ) in a patch through a functional response to availability,  $F_\nu(Q)$ . The net egg laying rate, per-patch, is  $\Gamma = \chi\nu G$ . The eggs are distributed among the aquatic habitats ( $\mathcal{U}$ ) so that the egg deposition rates in habitats is  $\eta = \mathcal{U} \cdot \Gamma$ .

distributed among the aquatic habitats in that patch? To do so, we develop the concept 389  
of habitat availability. We assign a search weight to each aquatic habitat,  $w_\nu$ . Using the 390  
patch membership matrix,  $\mathcal{N}$ , we define aquatic habitat availability as: 391

$$Q(t) = \mathcal{N} \cdot w_\nu(t) \quad (11)$$

For each patch, total habitat availability is the sum of the search weights for habitats in 392  
that patch. 393

Daily, per-capita oviposition rates of gravid mosquitoes are computed using a 394  
functional response to habitat availability, such as: 395

$$\nu = F_\nu(Q) = \nu_x \frac{s_\nu Q}{1 + s_\nu Q}. \quad (12)$$

where  $\nu_x$  is the highest possible egg-laying rate for a gravid female, and  $s_\nu$  is a shape 396  
parameter. We note that if  $Q = 0$ , then  $\nu = F_\nu(0) = 0$ . We let  $G = F_G(\mathcal{M})$  denote the 397  
density of gravid mosquitoes, and we let  $\chi$  denote the number of eggs laid, per batch. 398  
The net egg laying rate, per patch, per day, is: 399

$$\Gamma = \chi\nu G \quad (13)$$

To model egg distribution among habitats, we formulate an egg distribution matrix ( $\mathcal{U}$ ) 400  
that allocates eggs to habitats in proportion to local habitat availability. To compute  $\mathcal{U}$ , 401  
for computational reasons we first create  $Q^*$  by setting any zero entries to an arbitrary 402  
positive value (if  $Q = 0$ , then  $\nu = 0$ , so associated products will later be multiplied by 403  
zero), and the egg deposition rate,  $\eta$ , is computed by: 404

$$\mathcal{U}(\mathcal{N}, w_\nu) = \text{diag}(w_\nu) \cdot \mathcal{N}^T \cdot \text{diag}\left(\frac{1}{Q^*}\right). \quad (14)$$

Finally, we can compute egg deposition rates in the habitats: 405

$$\eta = \mathcal{U} \cdot \Gamma \quad (15)$$

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

While  $\Gamma$  (a vector of length  $p$ ) describes the net egg-laying rate of the adult mosquito population in each patch, per day  $\eta$  (a vector of length  $l$ ) describes the number of eggs laid, in each habitat, per day.

## Core Dynamical Components

The dynamical quantities whose computation was described above, are configurable elements that connect the four dynamical components: aquatic mosquito ecology; adult mosquito ecology and infection dynamics; and infection and immunity, including human demography. In the following, we describe one model family for each component, including functions that compute terms required for the dynamical quantities; in code these are the generic interface of each dynamical component. In Supplement 1 (<https://dd-harp.github.io/exDE/>), we have formulated alternative model families for some of the components.

**Aquatic Mosquito Ecology** The first core dynamical component describes aquatic mosquito population dynamics; the algorithm computes mosquito survival and development from eggs laid through adults emerging. For aquatic population dynamics, we here adapt a previously published model [31, 32].

Let  $L(t)$  denote the total density of immature mosquitoes. We let  $\psi(t)$  denote maturation rates,  $\phi(t)$  the density independent mortality rate, and  $\theta(t)L(t)$  describes increased per-capita mortality due to mean crowding. The aquatic dynamics are thus:

$$\frac{dL}{dt} = \eta - (\psi + \phi + \theta L)L \quad (16)$$

The total emergence rate of female mosquitoes in this model, per aquatic habitat, is:

$$\alpha(t) = F_{\alpha}(L(t)) = \frac{\psi(t)L(t)}{2}. \quad (17)$$

These are recruited into the adult population in the patch, so that the net emergence rate per patch is:


$$\Lambda(t) = \mathcal{N} \cdot \alpha \quad (18)$$

While  $\alpha$  is a vector of length  $l$ ,  $\Lambda$  is a vector of length  $p$ . This is passed as input to the equations describing adult populations (below).

Given uncertainty about the factors affecting immature mosquito populations, we assume studies might choose to formulate and analyze alternative dynamics. Other dynamical systems models for aquatic ecology in the framework are defined by state variables,  $\mathcal{L}$ , with dynamics defined by a system of equations  $d\mathcal{L}/dt = \eta - F_{\mathcal{L}}(\mathcal{L})$ , and a function such that  $\alpha = F_{\Lambda}(\mathcal{L})$ , such that  $\Lambda = \mathcal{N} \cdot \alpha$  (Supplement 4).

**Adult Mosquito Ecology** The second core dynamical component describes adult mosquito ecology. Given all the functions, terms and parameters above, we have formulated a set of algorithms describing adult mosquito mortality and dispersal that are internally consistent. All this is embodied in the mosquito demographic matrix, called  $\Omega(t)$ .

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).



**Fig 5. Adult mosquito demography** is defined by survival and dispersal. Mobility rates and dispersal are determined by the available of resources: aquatic habitats ( $Q$ ), available humans ( $W + W_\delta$ ) and other blood hosts ( $O^\delta$ ), and sugar ( $S$ ). The emigration rate is a functional response ( $F_\sigma$ ) that increases if any one of the resources is missing. Resource availability and distance also play a role in computing the dispersal kernel,  $\mathcal{K}$ , that determines where mosquitoes land if they leave a patch. When combined with mortality, a matrix  $\Omega$  is produced which describes the behavior of adult mosquitoes after emergence.

We assume mosquito mobility is driven by a search for resources. We have already defined total blood host availability  $B$ , and aquatic habitat availability  $Q$ . We also consider sugar availability,  $S(t)$ , which is passed to the model as a function vector of length  $p$ . We assume mosquitoes leave a patch while searching for resources, and that they leave a patch more frequently if the resources are less available. Patch-specific emigration rates,  $\sigma(t)$ , are a functional response to resource availability:

$$\sigma = F_\sigma(B, Q, S) = \sigma_x \left( \frac{\sigma_B}{1 + s_B B} + \frac{\sigma_Q}{1 + s_Q Q} + \frac{\sigma_S}{1 + s_S S} \right) \quad (19)$$

The parameters  $\sigma_B$ ,  $\sigma_Q$ , and  $\sigma_S$  determine the rate that mosquitoes leave a patch if no resources are available, and the shape parameters  $s_B$ ,  $s_Q$ , and  $s_S$  determine how the rate of patch leaving is reduced by the availability of resources. The shape parameter  $\sigma_x$  is a scaling parameter that can be used to adjust models with differing patch sizes. Similarly, we formulate a mosquito dispersal matrix,  $\mathcal{K}(t)$  that describes where mosquitoes land after they leave each patch (the diagonal elements of  $\mathcal{K}$  are constrained to be equal to zero, Supplement 4).

We let  $g(t)$  denote the local per-capita mortality rate of mosquitoes in each patch. The matrix  $\Omega(t)$  describes adult mosquito survival and dispersal:

$$\Omega = \text{diag}(g) + (I - \mathcal{K}) \cdot \text{diag}(\sigma) \quad (20)$$

where  $I$  is the identity matrix.

We let  $\Lambda(t)$  be the net emergence rate of mosquitoes into the patches from aquatic habitats (see Eq. 18, above). The dynamics of adult mosquitoes are described by the equation:

$$\frac{dM}{dt} = \Lambda - \Omega \cdot M \quad (21)$$

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Under the assumptions of this model, the density of gravid mosquitoes,  $G$ , is: 459

$$\frac{dG}{dt} = f(M - G) - \nu G - \Omega \cdot G \quad (22)$$

This model thus assumes that only gravid mosquitoes can lay eggs (Eq. 13), but that all mosquitoes (including gravid mosquitoes) can blood feed. 460  
461

Other models for adult mosquito ecology, denoted  $dM/dt$ , could be formulated that describe separate functions for mosquito survival and dispersal, depending on their behavioral states (possibly including sugar feeding, mating and maturation), or that describe a mosquito's reproductive states, or its chronological age or reproductive age. All models developed in this framework must accept the adult emergence rates,  $\Lambda$ , and they must be formulated in enough detail to specify a population of egg-laying mosquitoes,  $G$ , to compute  $\nu$  (see Eq. 13). 462  
463  
464  
465  
466  
467  
468

**Parasite Infection Dynamics in Mosquitoes** The third core dynamical component describes parasite infection dynamics in adult mosquito populations. Here, we extend a previously published delay differential equation for the density of infectious mosquitoes to include space and a time-varying extrinsic incubation period (EIP) [49]. 469  
470  
471  
472

Let  $Y(t)$  denote the density of infected mosquitoes. Using  $\kappa$  from Eq. 10, the dynamics of infection in mosquitoes are described by: 473  
474

$$\frac{dY}{dt} = fq\kappa(M - Y) - \Omega \cdot Y \quad (23)$$

We include a time-dependent EIP so that parasite development can be modulated by temperature or other factors exogenous to the system: let  $\tau(t)$  denote the EIP for a mosquito that becomes infected at time  $t$  (*i.e.*, it becomes infectious at time  $t + \tau(t)$ , Supplement 4). We must also define the inverse  $\tau^{-1}(t)$ , the delay for a mosquito that became infectious at time,  $t$ . Let  $\Upsilon_\tau(t)$  denote a matrix describing survival and dispersal of a cohort from time  $t - \tau^{-1}(t)$  through the EIP to become infectious at time  $t$ : 475  
476  
477  
478  
479  
480

$$-\ln \Upsilon_\tau(t) = \int_{t-\tau^{-1}(t)}^t \Omega(s) ds. \quad (24)$$

When  $\Omega$  and  $\tau$  are constant, survival and dispersal through the EIP is  $\Upsilon_\tau = e^{-\Omega\tau}$ . Otherwise, let the  $\tau$ -subscript denote the value of a variable or parameter at time  $t - \tau^{-1}(t)$ . 481  
482  
483

To model the density of infectious mosquitoes, let  $Z(t)$  denote the density of infectious mosquitoes. The dynamics of infectious mosquitoes are: 484  
485

$$\frac{dZ}{dt} = \Upsilon_\tau \cdot f_\tau q_\tau \kappa_\tau (M_\tau - Y_\tau) - \Omega \cdot Z \quad (25)$$

The number of human blood meals per patch, called the net infectious biting rate, is  $fqZ$ . 486  
487

Models for infection dynamics, generically denoted  $dY/dt$  are nested within the model for adult mosquito population dynamics  $dM/dt$  (for example, see [50]). These models accept the net infectiousness ( $\kappa$ ), they must define a variable describing the 488  
489  
490

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

density of infectious, blood feeding mosquitoes,  $Z$ , in order to compute the EIR (see Eq. 9).

**Epidemiology** The fourth core dynamical component describes parasite infection dynamics in human populations. Models for malaria infection, immunity, disease, and infectiousness in humans, denoted  $d\mathcal{X}/dt$ , can become quite complicated, depending on the needs of a study. Studies of malaria epidemiology could consider the complex time course of infections, superinfection, disease, detection, infectiousness, and immunity. The state space describing malaria infection and immunity  $\mathcal{X}$  can be modified to suit the needs of a study, and the framework also has enormous flexibility to model heterogeneity in populations through stratification. The following is one model family that is complex enough to illustrate the generic features of the framework.

Let  $h = f_h(E)$  denote the local daily force of infection (FoI) and  $\delta(t)$  the FoI during travel. In general,  $f_h(E)$  could be modified to include heterogeneous biting [41], but in this model, we assume  $h = bE$ . Both terms are defined for each sub-population. In these models, we stratify on variables relevant for the epidemiology, including immunity, and we model the effects by assigning different parameter values to each stratum.

To model infection dynamics, we modify a hybrid model for the multiplicity of infection (MoI). The dynamics are based on a queuing model, in which new infections occur at the rate  $h$ , and each parasite clears at the rate  $r$ , where we track apparent and actual clearance as linked but distinct processes. The variables  $m_1$  and  $m_2$  track the mean MoI for present and detectable parasites in each strata, which fully describe the epidemiological state space in a simple model with superinfection [51]. We assume that parasites clear at the per-capita rate,  $r_1$ , so that:

$$\frac{dm_1}{dt} = h + \delta - r_1 m_1 \quad (26)$$

In this model, the true prevalence is:

$$x_1 = 1 - e^{-m_1} \quad (27)$$

We also formulate a model for the MoI of apparent infections. We assume parasite infections are detectable for a shorter time so they appear to clear at a higher rate,  $r_2$ , and

$$\frac{dm_2}{dt} = h + \delta - r_2 m_2 \quad (28)$$

Similarly, we let  $x_2$  denote the apparent prevalence

$$x_2 = 1 - e^{-m_2} \quad (29)$$

We assume that if the infection is patent, a bite infects a mosquito with a higher probability,  $c_2$ , and  $c_1$  if it is not. A bite on a person in each stratum infects a mosquito with probability:

$$x = c_2 x_2 + c_1 (x_1 - x_2) \quad (30)$$

To compute  $\kappa$ , the infective density of infectious resident hosts by strata is  $X = xH$ . The vector  $X$  is passed to Eq. 10 to compute a vector of patch-specific net

It is made available under a [CC-BY 4.0 International license](#) .

infectiousness,  $\kappa$ .

To compute some of the spatial transmission metrics, including basic reproductive numbers (see below), a model must compute the human transmitting capacity (HTC) [52]. In this model, the number of days infecting mosquitoes at the higher probability,  $c_2$  is  $1/r_2$ . The remaining days, are spent infecting mosquitoes at the lower probability. Expressed as the equivalent number of perfectly infectious days, the HTC is:

$$D = \frac{c_2}{r_2} + c_1 \left( \frac{1}{r_1} - \frac{1}{r_2} \right) \quad (31)$$

This framework can accommodate other systems of equations describing parasite infection and immune dynamics in humans. This particular model was designed to illustrate some basic features of the modular design. These particular equations were designed to incorporate the effects of immunity on transmission through stratification, allowing parameters describing the duration of infections or detection and the infectiousness to vary among strata (*e.g.*,  $r_1$ ,  $r_2$ ,  $c_1$  and  $c_2$ ). New models for human epidemiology can use any epidemiological state space,  $\mathcal{X}$ , and any system of equations,  $d\mathcal{X}/dt$ , including models with dynamical changes in the host population size. While the travel FoI is recommended, it is not required. The modules should accept the EIR, and to interact with other components, they must provide a function to compute the infective density of infectious hosts,  $X$ .

## Spatial Metrics

The Ross-Macdonald model defined a set of concepts and metrics that have formed a basis for measuring and understanding malaria transmission, including vectorial capacity and the basic reproductive number  $R_0$ , but that model and associated metrics did not include metrics for spatial dynamics, parasite dispersal, or malaria importation [3].

Here, we define parasite dispersal by the set of locations (*i.e.* patches) where infecting bites occurred in continuous chains of transmission stretching back in time. Dispersal for any parasite transmission chain is thus defined by locations of the bites that caused each infection, and dispersal alternating between moving humans and mosquitoes between bites. We acknowledge that, due to an observational process, there is an important difference between where an infection occurred and where an infectious person or mosquito is found. There is also an important difference between the formulas defining dispersal and those used to compute reproductive numbers, which count from after a host becomes infectious. Using this definition of parasite dispersal in the context of a model, we have developed formulas and metrics to compute and study parasite dispersal and reproductive success.


To develop these metrics, we assume steady state conditions. This is done for convenience to avoid discussing the complications of understanding spatial dispersal under dynamically changing conditions, and it is a necessary first step to understanding such models. Analysis of malaria transmission dynamics under temporally varying conditions are being developed in a subsequent manuscript.

The formulation of this static model helps to clarify the role of some of the intermediate terms – if all parameters in a model were constant, the transmission model

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

could be fully defined by a much smaller set of parameters, but it may not be clear why the parameters take on those values. Some of the terms that appear in the static analysis correspond to parameters or variables in some Ross-Macdonald models, while others are new: net emergence rates ( $\Lambda$ ) or adult mosquito density ( $M$ ), scaled to the appropriate human population density denominator of host availability ( $W$ ), mosquito bionomics ( $f$ ,  $q$ , and  $\Omega$ ), and epidemiological parameters ( $r_1$ ,  $r_2$ ,  $c_1$ , and  $c_2$ ). New terms describe the spatial biting distribution matrix ( $\beta$ ) and parameters describing malaria importation ( $\delta$ ,  $v$ , and  $x_\delta$ ).

In models where the context is changing dynamically – due possibly to weather, land use changes, or vector control – exogenous forcing functions can be passed to the model that change resource availability or that perturb the dynamics; the functional forms and intermediate terms (*e.g.* availability) are used to describe changes in the local parameter values and guarantee mathematical consistency. In these static models, the functions and terms are used to set up the model, but after setting parameter values, they need not be called again.



**Fig 6.** To model malaria importation, we define a travel FoI for each stratum,  $\delta(t)$ , and two set of terms to model the role of visitors in mosquito blood feeding and parasite transmission: the available visitor population  $W_\delta$  and the NI for the visitor population, by patch  $x_\delta$ . To model blood feeding and transmission, we compute a patch-specific resident fraction for blood feeding,  $v$ , the fraction of all biting that occurs on a resident of the spatial domain. From this, we can compute the visitor reservoir fraction,  $\gamma$ , the travel fraction for incidence, and other measures of malaria importation.

## Net Malaria Importation and Travel Fractions

Terms describing the travel FoI ( $\delta$ ) and visitor populations were defined above and integrated into the models for blood feeding and human epidemiology. We define an imported malaria case as a human infection that traces back to a location outside of the spatial domain in the parasite's previous generation, *i.e.*, the mosquito and human host preceding this one in a chain of infections [42]. Net malaria importation rates describe the number of imported malaria cases, per day.

The fraction of all cases that were imported called the *travel fraction* can be defined as either: 1) the fraction of incident infections that were imported; or 2) the fraction of prevalent infections that were imported [45, 53]. To compute these travel fractions, we



It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

let  $\gamma = (1 - v)x_\delta/\kappa$  denote the *visitor fraction*, the fraction of infectious mosquitoes that were infected by visitors. We let  $h$  denote the FoI. The travel fraction for incidence is:

$$\frac{h\gamma + \delta}{h + \delta} \tag{32}$$

The travel fraction for true prevalence is:

$$\frac{1 - e^{-(\delta+h\gamma)/r_1}}{1 - e^{-(\delta+h)/r_1}} \tag{33}$$

We note that these are per-capita terms defined for the strata. The net malaria importation rate, the number of imported malaria incidence per day for each patch is:

$$\mathcal{J} \cdot (h\gamma + \delta) H \tag{34}$$

so the travel fraction for incidence for the patches would be:

$$\frac{\mathcal{J} \cdot (h\gamma + \delta) H}{\mathcal{J} \cdot (h + \delta) H} \tag{35}$$

Formulas for the travel fraction for prevalence are formulated in the same way.

## Parasite Dispersal

To compute quantities related to parasite dispersal, from bite to bite, we focus on local transmission, and we need some formulas that describe how mosquitoes move around in humans and in mosquitoes.

**Mosquito Dispersal and Steady States** In these models, we can compute steady state mosquito population density, assuming  $\Lambda$  is constant over time. At the steady state of Eq. 21,

$$M = \Omega^{-1} \cdot \Lambda \tag{36}$$

Here, the inverse  $\Omega^{-1}$  can be understood as a measure of time spent alive in each patch by mosquitoes emerging habitats in each patch. In other Markov chain models with finite state space, it has also been shown that the elements of the matrix inverse can be interpreted as residence times [54, 55]. In the simpler Ross-Macdonald model, the inverse of a mortality rate,  $g$ , is a measure of time spent alive or the average mosquito lifespan [56, 57]. The *time spent alive* interpretation of  $\Omega^{-1}$  is more apparent if there is no movement: if we set  $\sigma = 0$ , then  $\Omega^{-1} = \text{diag}(1/g)$ .

In spatial models, the matrix  $\Omega$  accounts for both survival and movement. To illustrate – and to demonstrate that if  $\Omega$  is a sensible description of mosquito demography, then the matrix inverse must exist – we construct a tracking matrix. Let  $\Xi(t)$  denote a matrix that tracks cohorts of mosquitoes:

$$\Xi(t, M_0) = e^{-\Omega t} \cdot \text{diag}(M_0) \tag{37}$$

It describes the density of mosquitoes left from an initial cohort in each patch  $M_0$  that is found in each location at each point in time. There is a duality between the

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

equilibrium population density from Eq. 21 and time spent alive by a cohort, computed by integrating Eq. 37 (*i.e.* orbits of the related equation  $dM/dt = -\Omega \cdot M$ ). Just as we can compute  $g^{-1} = \int_0^\infty e^{-gt} dt$ , we can compute:

$$M = \Omega^{-1} \cdot \Lambda = \int_0^\infty e^{-\Omega t} dt \cdot \Lambda \quad (38)$$

so that the steady state can be found by simply adding up the time spent alive in each patch by a cohort emerging from every other patch. Under generalized static conditions (*i.e.*  $\sigma > 0$ ),  $\Omega^{-1}$  can thus be interpreted as the average time spent alive in every patch by cohorts of mosquitoes initially found in each patch.

**Parasite Dispersal in Mosquitoes** Using mosquito tracking matrices, we can also track parasite dispersal in mosquitoes to derive a matrix that has the same interpretation as the formula for vectorial capacity [57, 58].

To transmit, mosquitoes must blood feed on a human to become infected: the net infection rate in each patch, per available human, is  $f q \kappa M / W$ . After becoming infected, a mosquito must survive while dispersing through the EIP ( $\Upsilon = e^{-\Omega \tau}$ ). After becoming infectious, a mosquito must blood feed to transmit parasites, so we use the matrix inverse  $\Omega^{-1}$  which describes where the mosquitoes are for each infectious human blood meal as long as they remain alive; after becoming infectious, the distribution of infectious bites is given by  $f q \Omega^{-1}$ . We can describe parasite transmission by mosquitoes by following the story of infection in mosquitoes: after emerging ( $\text{diag}(\Lambda)$ ), a mosquito must blood feed on a human to become infected ( $f q \Omega^{-1} / W$ ); then survive the EIP ( $e^{-\Omega \tau}$ ); and then blood feed to transmit ( $f q \Omega^{-1}$ ).

In the Ross-Macdonald model, the formula for vectorial capacity can be derived from the formula for the daily EIR as a limit [57]. In spatial models, a vectorial capacity matrix can be derived as the limit of a tracking matrix describing the number of infectious bites arising, per available person (*i.e.*, the denominator is  $W$ ), per day at the steady state (Supplement 4):

$$\mathcal{V} = \lim_{\kappa \rightarrow 0} \frac{f q Z}{W} = f q \Omega^{-1} \cdot e^{-\Omega \tau} \cdot \text{diag} \left( \frac{f q M}{W} \right) \quad (39)$$

Elements in the matrix  $\mathcal{V}$  are the expected number of infectious bites eventually arising in every patch from all the mosquitoes in a single patch blood feeding on a single human on a single day, computed as if each human were perfectly infectious. The derivation assumes that no mosquitoes are already infected, and the assumption that humans are perfectly infectious ( $\kappa = 1$ ) is made so that the formula deals only with phenomena related to mosquitoes. In models with multiple vector species, the notion of what it means to be “perfectly infectious” is not as simple because of differences among vector species in their capacity to be a host for the parasites, or vector competence (Supplement 4).

**Parasite Dispersal by Humans** To quantify parasite dispersal by humans, we compute the human transmitting capacity distribution (HTCD) matrix. We let human transmitting capacity (HTC) describe the net number of perfectly infectious days for

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

each stratum: since infectiousness varies over the time-course of infections, we sum partially infectious days and interpret the HTC as an equivalent number of days spent perfectly infectious [52]. For the population strata in this model, the HTC ( $D$ ) is defined by Eq. 31. Since transmission requires two bites, we use the TaR matrix to determine both where a human becomes infected and where it infects a mosquito. Using the transposed TaR matrix, we can describe where infectious days at risk are spent,  $\Psi^T \cdot D$ . Parasite dispersion by mosquitoes for the sub-populations also accounts for where a mosquito becomes infected, or  $b\Psi$ .

The HTCD matrix uses the biting distribution matrix,  $\beta$ , to count from the infectious bite and weight biting appropriately for subsequent blood feeding by all the population strata. The HTCD, a  $p \times p$  matrix ( $\mathcal{D}$ ), is:

$$\mathcal{D} = \text{diag}(W) \cdot \beta^T \cdot \text{diag}(bDH) \cdot \beta. \quad (40)$$

We note that  $\mathcal{D}$  in spatial models is analogous to  $bD$  in models with a single patch. (The equivalency of  $\mathcal{D}$  and  $bD$  is most apparent if no humans move, and if there is one stratum per patch, and if all search weights are 1, in which case  $H = W$  and  $\beta = \text{diag}(1/H)$ .) Like  $bD$ ,  $\mathcal{D}$  describes days spent infectious by an individual human, but in  $\mathcal{D}$ , describes both where a human got infected and where the mosquitoes were subsequently infected.

The definition of  $\mathcal{D}$  as a time-dependent matrix is substantially more complicated if local human mobility patterns change dynamically.

**Parasite Dispersal through one Parasite Generation** Parasite dispersal is defined by the locations where infecting bites occurred, alternatively moving in infected mosquitoes and humans. The equations for  $\mathcal{D}$  and  $\mathcal{V}$  describe the expected movement for a parasite among patches in humans or mosquitoes, respectively, counting from bite to bite. Notably, the formulas are defined for a parasite in either a mosquito or a human. We can also define parasite dispersal through one parasite generation (*i.e.*, from human to human, or from mosquito to mosquito) but the formula depends on where we start counting. If we started from all the mosquitoes blood feeding on a single human (averaged appropriately) on a single day in every patch, then we would get a matrix describing dispersal from every patch to every patch:

$$\mathcal{D} \cdot \mathcal{V}. \quad (41)$$

If we started counting from a typical human infected in a patch on a single day, we would get a different dispersal matrix:

$$\mathcal{V} \cdot \mathcal{D}. \quad (42)$$

Importantly, these formulas follow the same process in the same order, and thus closely resemble the reproductive numbers for malaria (described below), which measure reproductive success for a single parasite. These formulas are two among many that could be developed to count events through a parasite's life-cycle starting at different points.

Formulas that describe the parasite's *per-capita* reproductive success, such as

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

Eqs. 41-42, counting events arising from a single host. In some cases, we might wish to count the total number of events arising from a patch. To measure the contribution of a patch to overall transmission, we must have a measure of connectivity, or total parasite flows. A tracking matrix describing all of the infections arising from each patch on a day, is:

$$\text{diag}(W) \cdot \mathcal{D} \cdot \mathcal{V} \quad (43)$$

If we started counting infections occurring on humans in a patch, we would get an alternative patch-based tracking matrix. The number of infections arising from a patch is thus tracked by:

$$\text{diag}(W) \cdot \mathcal{V} \cdot \mathcal{D} \quad (44)$$

These measures emphasize the role of places with larger available populations.

The same sort of formulas can be devised to describe transmission from human strata to human strata, but the resulting formulas are only spatial insofar as the human strata are anchored to a residency. If we focused instead on parasite reproductive success starting with an infection in humans, regardless of location, we would get

$$\mathcal{R} = b\beta \cdot \mathcal{V} \cdot \text{diag}(W) \cdot \beta^T \cdot \text{diag}(DH). \quad (45)$$

or we could also count bulk transmission from humans as  $\text{diag}(H) \cdot \mathcal{R}$ . Notably, Eq. 45 is a stratum-based measure. To make it quasi-spatial, we would need to assign events to patches by stratum residency using the membership mapping operator  $\mathcal{J} \cdot \mathcal{R} \cdot \mathcal{J}^T$ .

**Distances Dispersed** To get a measure of the distribution of distances travelled by parasites, we match a measure of transmission intensity with the corresponding element in a patch distance matrix describing the distance. We take the couplet (distance and intensity) and sort by distance, then compute the cumulative distribution function (CDF). From the CDF, we derive a probability mass function [39]. These dispersal kernels provide a simple way of visualizing distances dispersed by mosquitoes, humans, or parasites.

These formulas and algorithms draw attention to the differences in metrics describing parasite transmission dynamics and dispersal. Because of spatial heterogeneity in mosquito and human population densities, there are many sensible formulas for counting dispersal, some of which correspond to describing rates, ratios, proportions, and numbers. Careful thought should be given to choosing or developing a metric that fits the analysis.

## Reproductive Numbers

Reproductive numbers are a measure of the parasite's average reproductive success. When transmission is spatially heterogeneous, reproductive success will vary for parasites, depending on where they are. As parasites spread over several generations, the expected success of its progeny will change. To calculate threshold criteria for persistence (in the absence of malaria importation), we want a reproductive number to be a measure of average success taken over the whole system, but we want to use an average that does not change across generations. Doing so requires that we compute the

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

*spectral average*, which is computed as the dominant eigenvalue of the parasite’s *next generation matrix*. 727  
728

For many reasons, it is useful to formulate *local* reproductive numbers that describe a parasite’s average reproductive success at a particular place and time – an arithmetic average. These *local* reproductive numbers could ignore differences across generations, so they would not serve as thresholds for parasite persistence. In this section, we define local reproductive numbers at the steady state, but the formulas could also serve as point estimates. 729  
730  
731  
732  
733  
734

Reproductive numbers describe malaria transmission under a range of different conditions that are relevant for understanding malaria transmission dynamics and control or for national strategic planning. Baseline conditions are described by the basic reproductive number,  $R_0$ , which is defined for a population with no acquired immunity and no malaria control. The adjusted reproductive number,  $R_C$ , describes a family of numbers defined for a population with no acquired immunity adjusted by malaria control, at a fixed level of control denoted  $C$ . In other words,  $R_0$  is defined as a special case of  $R_C$ , but in the absence of control. The total effect size of malaria control on transmission is  $R_0/R_C$ . Here, we also describe the endemic reproductive number,  $R_E$ , which describes potential transmission modified by immunity. The total effect size of immunity on transmission is  $R_C/R_E$ . In computing  $R_E$ , as with  $R_0$  and  $R_C$ , we ignore the fact that some hosts are already infected. In this way,  $R_E$  is defined differently than the effective reproductive number, denoted  $R_e$ , which is lower than  $R_E$  because it does not count infections occurring in someone who is already infected. We note that, by definition, at an endemic steady state  $R_e = 1$ . By way of contrast,  $R_E$  counts the number of infections that would occur after one generation, which is useful for planning because it helps to clarify how success in malaria control can be assisted by immunity that will eventually wane. 735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752

Both  $R_0$  and  $R_C$  are computed as if there were no acquired immunity. In this model, the effects of acquired immunity on transmission are quantified through the stratified values of  $b, r_1, r_2, c_1$  and  $c_2$ . These parameters determine the HTC for all the strata ( $D$ , see Eq. 31). If  $D$  were computed using values that have been tuned to a stratum with some level of immunity, we would be computing  $R_E$ . To compute  $R_C$ , we would need to replace  $D$  with values set to a non-immune baseline (*i.e.*,  $D_0$ ), and then recompute the next-generation matrix. Next generation matrices computed with values of  $D$  that include the effects of acquired immunity are thus describing an endemic reproductive number. Depending on how  $D$  is computed, and whether the bionomic parameters incorporate effects of vector control, we may thus be computing  $R_0, R_C$  or  $R_E$ . 753  
754  
755  
756  
757  
758  
759  
760  
761  
762

**Local Reproductive Numbers** One way to define local reproductive numbers is to modify Macdonald’s formula using the local values of parameters, as if there was no movement of mosquitoes or humans. To write the formula using some models in this framework, we may need to modify HTC (which is defined for the strata, of length  $n$ ) to take a patch average. To compute a patch average HTC,  $\check{D}$  (a vector of length  $p$ ), we take the population weighted average, 763  
764  
765  
766  
767  
768

$$\check{D} = \frac{\Psi^T \cdot w_f DH}{\Psi^T \cdot w_f H} \quad (46)$$

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

We can then describe a local reproductive number,  $\check{R}_C$  (or possibly  $\check{R}_E$ , depending on how the interpret parameters are defined in  $D$ ):

$$\check{R}_C = \frac{\Lambda}{W} \frac{f^2 q^2}{g^2} e^{-g\tau} \check{D} \quad (47)$$

This local measure is similar to Macdonald's formula [59]. While useful in some contexts, the formula should be applied with caution.

An alternative way to compute local reproductive numbers uses  $\mathcal{V}$  and  $\mathcal{D}$  (perhaps modified to remove the effects of immunity on transmission). Since the matrices count infections arising from each patch, and we add all infections arising to the patch where the bite originates. We let  $1$  be a row vector of ones of length  $p$ , and we can count infections arising starting from all the humans infected in a patch on a single day:

$$\hat{R}_C = 1 \cdot \mathcal{V} \cdot \mathcal{D} \quad (48)$$

that counts infections occurring on humans, or we can start from all the mosquitoes blood feeding on humans on a single day, and:

$$\tilde{R}_C = 1 \cdot \mathcal{D} \cdot \mathcal{V} \quad (49)$$

that counts infected mosquitoes. These patch reproductive numbers could provide valuable information about whether to target the mosquitoes or humans in some patch for enhanced interventions. We could also consider the equivalent formulas for total patch outputs:

$$W^T \cdot \mathcal{V} \cdot \mathcal{D} \text{ or } W^T \cdot \mathcal{D} \cdot \mathcal{V} \quad (50)$$

where  $W^T$  is a row vector. Alternatively, we can also weigh transmission from strata using Eq. 45:

$$1 \cdot \mathcal{R} \quad (51)$$

or the equivalent scaled by stratum size:


$$H^T \cdot \mathcal{R}, \quad (52)$$

where  $H^T$  is a row vector, which gives us valuable information about infections arising from every stratum on every strata, a way of identifying the relative importance of various population strata.

**Next Generation Matrix** In the Ross-Macdonald model, a parasite's reproductive success in the next generation is described by a single number. It is computed by counting forward from the moment a mosquito or human becomes infectious. Since parasites move in infected mosquitoes and humans, parasite reproductive success – measured as the number of infections in the next generation – varies across generations as the parasite distributions evolve across generations among strata and among patches. The matrices  $\mathcal{V}$  and  $\mathcal{D}$  describe parasite transmission and dispersal in mosquitoes and humans, respectively. While the product of these formulas does describe net reproductive success, the computation of threshold conditions has been developed

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

around the concept of a next generation matrix [60,61], which traces the same process in the same sequence but that start counting at a different point in the parasite’s life cycle (Fig. 7). A threshold condition is found by taking the spectral average of the next generation matrix.



**Fig 7. A Spatial Life-Cycle Model.** A diagram that illustrates how the parameters describing each stage in the parasite’s life-cycle translate into a parasite’s reproductive success spatially, when mosquitoes and hosts move. The right half of the circle represents mosquitoes and the left half humans. The flow of events is clockwise. Mosquitoes must blood feed to become infected ( $f q M$ ), and then survive and disperse through the EIP ( $e^{-\Omega\tau}$ ). Infectious bites are distributed as long as a mosquito survives, while it blood feeds and disperses ( $f q \Omega^{-1}$ ). The bites are distributed among humans ( $\beta$ ) and some of them cause an infection ( $b$ ). Parasites are transmitted for as long as humans remain infectious, measured in terms of the human transmitting capacity (HTC, or  $D$  days). Infectious humans are distributed wherever humans spend time at risk (affecting  $\beta$ ). These processes are summarized differently to model parasite dispersal and parasite reproductive success. Dispersal counts from bite to bite using the VC matrix ( $\mathcal{V}$ ) and the HTC matrix ( $\mathcal{D}$ ). Threshold computations count from when a host becomes infectious to measure a parasite’s reproductive success in infectious mosquitoes ( $R_Z$ ); in infectious humans ( $R_X$ ); from human to humans among strata after a human becomes infectious ( $\mathcal{R}$ ); and from mosquito to mosquitoes ( $\mathcal{Z}$ ).  $R_0$  is the lead eigenvalue of  $\mathcal{R}$  or  $\mathcal{Z}$ . Under endemic conditions, we can also consider how frequently parasites are actually transmitted by including the probability a mosquito gets infected  $\kappa$ , and the probability a mosquito is infectious, given by the sporozoite rate  $z$ .

In computing next generation matrices, we focus on transmission within a defined spatial domain. For mathematical convenience here, we thus set  $v = 1$ , though we could easily develop matrices leaving  $v$  undetermined to discount exported malaria cases.

We first compute offspring transmitted from a single infectious mosquito to humans or from a single infectious human to mosquitoes, each of which defines a stage in the parasite’s next-generation [60]. After a mosquito has become infectious, how many humans (in each stratum) would it infect? In these models, the answer to that question is  $n \times p$  matrix, denoted  $R_Z$ , describing transmission from an infectious mosquito in each patch to humans in each strata:

$$R_Z = b\beta \cdot f q \Omega^{-1}. \tag{53}$$



It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

How many infectious mosquitoes would arise from each human infection? The answer is a  $p \times n$  matrix, denoted  $R_X$ , describing transmission from a human in each stratum to mosquitoes:

$$R_X = e^{-\Omega\tau} \cdot fqM \cdot (\beta^T \cdot \text{diag}(DH)). \quad (54)$$

The next-generation matrix by type is:

$$\mathcal{G} = \begin{bmatrix} 0 & R_Z \\ R_X & 0 \end{bmatrix} \quad (55)$$

To describe reproductive success in terms of the parasite biology, we count reproductive success through one full parasite generation, either from humans back to humans, or mosquitoes back to mosquitoes. For the parasites, reproductive success through one full generation requires two events, one of each type, so we square the matrix given by Eq. 55 to get a new matrix in block form:

$$\mathcal{G}^2 = \begin{bmatrix} \mathcal{R} & 0 \\ 0 & \mathcal{Z} \end{bmatrix}. \quad (56)$$

We thus get two diagonal block sub-matrices describing reproductive success in the parasite's next generation, denoted  $\mathcal{R}$  and  $\mathcal{Z}$ . Reproductive success from human population strata back to human strata is described by an  $n \times n$  matrix  $\mathcal{R} = R_Z \cdot R_X$ :

$$\mathcal{R} = b\beta \cdot \mathcal{V} \cdot \text{diag}(W) \cdot \beta^T \cdot \text{diag}(DH). \quad (57)$$

Reproductive success from mosquito through the population strata back to mosquitoes, described patch-by-patch is described by the  $p \times p$  matrix  $\mathcal{Z} = R_X \cdot R_Z$ :

$$\mathcal{Z} = e^{-\Omega\tau} \cdot \text{diag}\left(\frac{fqM}{W}\right) \cdot \mathcal{D} \cdot fq\Omega^{-1} \quad (58)$$

We have also formulated the next-generation matrix for systems with multiple vector species (Supplement 4).

**The Spectral Average** We can also compute  $R_C$  as a spectral average through simulation, which is one useful way of illustrating what a spectral average means. To do so, we choose a vector describing the distribution of parasites in a founding generation,  $\mathcal{X}_0$  or  $\mathcal{Y}_0$ , and iterate parasite infections across  $i$  successive parasite generations:

$$\mathcal{Y}_{i+1} = \mathcal{Z}\mathcal{Y}_i \text{ or } \mathcal{X}_{i+1} = \mathcal{R}\mathcal{X}_i. \quad (59)$$

We define the vector:

$$\mathcal{E}_i = \frac{\mathcal{X}_{i+1}}{\|\mathcal{X}_i\|} \text{ or } \mathcal{E}_i = \frac{\mathcal{Y}_{i+1}}{\|\mathcal{Y}_i\|}.$$

where  $\|\mathcal{X}\|$  or  $\|\mathcal{Y}\|$  is a scalar that denotes its magnitude. Over many generations,  $\mathcal{E}_i$  converges to the lead eigenvector, a scalar value also called the spectral average or  $R_C$ :

$$R_C = \lim_{i \rightarrow \infty} \|\mathcal{E}_i\| \quad (60)$$



It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

and it is interpreted as the asymptotic average reproductive success expressed as a number of infected hosts per host, per generation. Note that it is asymptotic only for the linearized system defined by Eq. 55 or Eq. 56.

**Quasi-Thresholds for Endemic Malaria** Without malaria importation,  $R_C > 1$  is a threshold criterion. Analysis of models without malaria importation have consistently demonstrated that malaria is either absent or that there is a single globally, asymptotically stable equilibrium. When there is imported malaria, there are three sufficient criteria for some local parasite transmission to occur within the area:

1.  $\max\{\delta\} > 0$  and  $R_C > 0$ ;
2.  $\max\{(1 - \nu)X_\delta\} > 0$  and  $R_C > 0$ ;
3.  $R_C > 1$ .

If condition 1 or condition 2 is satisfied, then malaria will be present in an area, and if  $R_C > 0$  then there will be some local transmission. If  $R_C > 1$ , malaria transmission would be sustained in the absence of importation. We thus call  $R_C > 1$  a quasi-threshold for endemic transmission to occur within the spatial domain: *endemic* describes places where  $R_C > 1$ , and *pseudo-endemic* places where  $0 < R_C < 1$  with significant levels of transmission.

## Quantifying Transmission in a Place

The framework, models developed within it, and the associated spatial metrics were designed to have the skill required to describe and quantify heterogeneous spatial transmission dynamics of malaria in a specific place at a particular time. We have not explicitly defined algorithms for the observational processes that would map model states onto observable quantities, which would be required to extend this mathematical modeling framework into a state space modeling framework to rigorously fit models to data. Instead, we have focused on the mathematics of these processes: time spent by humans; other blood hosts; daily mosquito rhythms; mosquito host preferences, time at risk; and mosquito mobility. Similarly, the models for mosquito ecology and population dynamics describe the mathematics of mosquito mobility, in terms of explicit assumptions about the locations of aquatic habitats, heterogeneous distributions of resources, and mosquito mobility patterns that emerge from a search for resources. By quantifying spatial patterns in terms of the underlying processes – including malaria importation, mosquito ecology and spatial population dynamics, parasite transmission dynamics, human mobility, and malaria epidemiology – the equations point towards a general inferential framework.

Models developed within this framework involve substantially more parameters than the Ross-Macdonald model. This is an inevitable consequence of a decision to model transmission at a particular place and time. If any local features are important for transmission, then a larger set of quantities must be estimated to understand and quantify those features. This gives rise to an important but difficult practical question: *What is the relationship between the amount of local intelligence and the specificity of*

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

*the policy advice that can be offered?* With minimal local information, it is possible to offer generic policy advice, but it may not be necessary to know everything about a place to tailor advice to context. With this framework, it is possible for models to evolve as the amount information increases, and the models may be used to look ahead to prioritize missing data: *How can programs identify missing information that would most rapidly improve the effectiveness of malaria control?* These contextual factors and the related questions are addressed below.

**Malaria Landscapes** While the Ross-Macdonald model describes parasite transmission between abstractly defined mosquito and human populations, the framework we have described was developed to understand and quantify malaria importation and transmission among structured mosquito and human populations in a well-defined geographical area. (Using a model  $d\mathcal{X}/dt$  that describe the infection dynamics of other pathogens and immunity in vertebrate host populations, and making other appropriate choices, the framework could be used as a basis for modeling dengue, West Nile virus, or other mosquito-borne pathogen transmission dynamics, as well.) Since the models are developed to approximate malaria transmission in an actual place, after defining an observational process, the model outputs would be verifiable statements about real quantities over some specific period of time.

As a practical first step, model building starts by defining a set of structural elements – patches, human population strata, and aquatic habits – that are appropriate for the needs of a study (*e.g.* Fig 8 illustrates some options for simulating malaria on Bioko Island, Equatorial Guinea). A geographical study area is usually defined by projects, programs, or political boundaries. In planning interventions for a defined area, an important concern is connectivity to surrounding areas. How much malaria is imported by daily human movement or travel? Are the mosquito populations within the area strongly connected to others nearby?

Using spatial metrics to identify differences in transmission patterns and the flow of parasites across a landscape can help control programs prioritize drugs, outreach, and medical attention to populations, and vector or larval control to places. Using our differential equation framework to reconstruct the equilibrium analysis presented in [45], we have generated spatial bulk transmission matrices ( $\text{diag}(H) \cdot \mathcal{R}$ ) among areas for Bioko Island, Equatorial Guinea. In Fig. 9 different patterns of pathogen transport are readily apparent between persons who live in Malabo (left), the densely populated capitol of the island and a sink for travellers, and Luba (right), a small settlement in the Southern half of the island. The pattern of travel seen in Luba typifies most of the areas outside of Malabo, where individuals most often travel to the capitol but not to the other outlying settlements. These patterns affect transmission, where we see parasites originating in Malabo tend to stay in the city. Parasites originating in Luba either tend to stay highly local, or are transported to Malabo when those persons move. Because malarial mosquitoes tend to fare less well in urban settings, these spatial metrics can help understand how high prevalence can be sustained in otherwise unsuitable locations.

An equally important question is about heterogeneity in mosquito population densities within the area and heterogeneity in the risk of exposure, which should inform the definition of patches and the choice of a patch size. Patches, in this model, are defined around adult mosquito activities, and each “patch” has a geographical location.

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

The patch is the spatial unit that defines the algorithms for time spent, blood feeding, egg laying, adult mosquito survival and dispersal. The concept of a patch is flexible enough to model blood feeding indoors and outdoors at the same geographical locations, which may be useful to inform programmatic questions about the effectiveness of vector control measures that target indoor biting (Fig. 10). Since the patch is the basis for computing most aspects of blood feeding, the patches define the structure for human time spent at risk, including (if required) quantifying time spent indoors *vs.* outdoors, and mosquito movement rates from indoors to outdoors, from outdoors to indoors, or from outdoors to other outdoor patches.

An important basic concern is the spatial granularity of the patches used for simulation (see Fig. 8). Some questions remain unresolved about the appropriate spatial scales and ways to define patches for describing and analyzing malaria transmission for policy (*e.g.*, to compute IRS coverage). One advantage of this framework is that it is possible to build nested models with different spatial grains and compare them. Smaller patches more accurately capture heterogeneity in a landscape while increasing the number of parameters that need to be inferred during calibration to data.

Aquatic habitats are located in patches, but the model was designed to assign patches to habitats assuming the habitats had an actual location. Patches in this framework need not have any human residents or any available hosts, so that mosquito habitats in the uninhabited areas around human households are contributing to transmission. Mosquito population dynamics are coupled through related equations describing gravidity, egg laying and egg deposition. The framework thus does not impose any constraints on either the method for constructing patches, or on the number or arrangement of aquatic habitats within the spatial domain. Given the modular nature of these models, the dynamics of immature mosquito populations in each aquatic habitat depend only on its parameters and the egg deposition rates. The productivity of any one aquatic habitat in an area is, however, coupled to other habitats through egg laying by adult mosquitoes that could have emerged anywhere.

To improve the accuracy of models, human populations can be segmented into strata to reduce heterogeneity in traits that affect malaria: the first segmentation is by residency. In this framework, which is designed to quantify process affecting *transmission*, heterogeneity in any trait affecting transmission is dealt with by sub-dividing the population into homogeneous (or less heterogeneous) strata, such as by age, travel habits or patterns, ITN usage, vaccination, care seeking, or any effects of immunity affecting malaria epidemiology or transmission.

Notably, all this structural flexibility is achieved through membership matrices and through the variables describing resource availability, which links search weights, functional responses, and other functional forms to guarantee mathematical consistency (*e.g.* avoiding problems when denominators are zero) despite structural changes. Suites of models can be developed to address concerns about data gaps and uncertainty that are appropriate for studies. Model complexity can be modified by changing dynamical modules, by changing functional forms, by fixing or changing parameters, by splitting and joining patches, by splitting or joining strata, or by adding and subtracting aquatic habitats. With the ability to split and join patches or strata, any model can be mapped onto simpler, nested models in a series of simple join operations until it is collapsed onto a single-patch, single-stratum Ross-Macdonald model. This is functionally what is

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

meant by scalable complexity. 965

It is thus as easy to modify and evaluate the effects of model structure (*e.g.* the 966  
number of strata) as it is to vary parameters, to facilitate developing suites of models, 967  
including models with nested patches or nested strata, to explore tradeoffs in building 968  
and calibrating models at various levels of detail. 969

**Mosquito Blood Feeding and Ecology** Three constant parameters describing 970  
mosquito behavior are a standard part of the Ross-Macdonald model [56,57]: the daily 971  
death rate of mosquitoes ( $g$ ), the overall daily blood feeding rate ( $f$ ), and the human 972  
blood feeding fraction ( $q$ ). Incorporating the possibility of dynamical feedback between 973  
the future emergence of adults and current population size means we have added the 974  
population egg-laying rate ( $\Gamma$ ). Adding spatial complexity to the model means the daily 975  
emigration rate ( $\sigma$ ), mosquito dispersal ( $\mathcal{K}$ ), distribution of habitats ( $\mathcal{N}$ ) and the 976  
distribution of eggs among patches ( $\mathcal{U}$ ) are additional parameters which define how 977  
populations may interact in space. While our analysis has focused on steady states, the 978  
models were formulated with parameters that can vary over time in response to 979  
changing availability of resources [33–36]. 980

In this framework, the values of all these parameters are computed with functional 981  
responses based on resource availability, mosquito biology and innate preferences that 982  
constrain the parameters within sensible ranges. This formulation emphasizes how 983  
baseline mosquito bionomics for different species could respond to available resources 984  
and how those responses would be modified by control. In particular, the same human 985  
behaviors can give rise to very different blood feeding patterns for different vector 986  
species, depending on the daily rhythms, host preferences, and aquatic ecology of 987  
different vector species (Supplement 4). We thus have a basis for understanding 988  
mosquito behaviors and ecology as a baseline that may have been modified by vector 989  
control or weather. 990

Blood feeding in this model thus makes an important distinction between 991  
*anthropophily*, or innate mosquito preferences for hosts of different types, and 992  
*anthropophagy*, summarized by the human blood feeding rate ( $fq$ ). Models can also 993  
consider a difference between the time of day when mosquitoes are actively searching for 994  
blood ( $\xi$ ) and the blood feeding rates by time of day ( $f$ ), which vary with host 995  
availability. Innate, species-specific host preferences are embodied in functional forms 996  
and parameters, while the rates describing what has happened also depend on context. 997

Similarly, mosquito population dynamics are an emergent feature of a resource 998  
landscape. Since searching for resources is also associated with resource availability, 999  
adult mosquitoes will tend to aggregate in patches that have habitats and other 1000  
required resources. In these models, egg-deposition rates in habitats by volant adult 1001  
populations are spatially heterogeneous and only partially determined by the emergence 1002  
rates of adults from a single habitat. The concept of a carrying capacity is, perhaps, not 1003  
as useful as the concept of habitat productivity and the functional forms that determine 1004  
how the number of adults emerging is related to the number of eggs laid [31]. A 1005  
habitat’s carrying capacity only makes sense in the abstract – if adult mosquitoes 1006  
emerging from a single habitat only laid eggs in that natal habitat. In this framework, 1007  
the aquatic population dynamic module determines how adult mosquito emergence 1008  
rates respond to egg laying by the adult population. 1009

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

The parameters describing these processes are both habitat-specific and time-dependent: density-independent mortality, density-dependent mortality, the response to crowding, maturation rates, and search weights could vary for every habitat. A habitat can thus disappear seasonally (which occurs when  $w_\nu = 0$ ), or weather could affect immature mosquito maturation and mortality rates. If a study called for modeling resource-based competition or stage-structured mosquito populations, the equations describing aquatic populations ( $d\mathcal{L}/dt$ ) can be modified as needed (Fig 1). The framework thus facilitates the construction of realistic models of mosquito ecology, insofar as it is justified by data available and the needs of a study.

**Local Exposure, Human Biting Rates and Mixing** In defining the algorithms for blood feeding, we also developed a new model for the human biting rate (HBR) and by extension, the entomological inoculation rate (EIR), two basic metrics used to measure malaria transmission entomologically.

The model emphasizes that for any population stratum, the risk of exposure to biting mosquitoes is distributed spatially. In these models, this is determined by a biting distribution matrix ( $\beta$ ). A similar matrix has appeared in other models for the spatial dynamics of mosquito-borne diseases for which human mobility is based on a concept of "visitation" or time spent – classified as Lagrangian movement [7, 8, 10, 12–15, 17, 18, 45]. Here,  $\beta$  is based on a concept of availability, the weighted, ambient population at risk. Availability is computed from observable quantities, and it is computed dynamically for arbitrarily defined human strata and changing availability (the denominator). The formulas guarantee consistency in blood feeding: the number of human blood meals taken by mosquitoes is equal to the number of blood meals received by the humans.

In the new model, the HBR is defined as  $\beta \cdot fqM$  and the EIR is  $\beta \cdot fqZ$ , so that the number of bites received by each stratum depends on how they spend their time at risk. In studies that have reported a value for the HBR or EIR, the quantity reported is based on catch counts by a person or device in a place. In this model, the quantity that is closest to the quantities being estimated is pHBR or pEIR, the number human blood meals, or infectious human blood meals in a patch, per available person, per day ( $fqM/W$  or  $fqZ/W$ ). A person who is in a patch at a particular time of day would experience the local biting rates at that time scaled by a search weight ( $fqM\xi(t)\omega_f/W$  or  $fqZ\xi(t)\omega_f/W$ ). The quantity being estimated by human landing catches is a measure of the intensity of exposure in a place.

Since other hosts are also available, the number of mosquitoes caught also depends on the biases of the trapping method. In this model, each method for trapping mosquitoes can be thought of as having its own "availability," and it is competing for the attention of mosquitoes. Each method for catching mosquitoes is biased in some unknown way. We thus suggest that field methods designed to estimate the EIR are best interpreted as a location-specific measure of risk in a place, and that epidemiologically relevant measures of risk must acknowledge exposure occurring for a period of time, including all the places where a person spends time. The pEIR, weighted by total availability, is a good approximation of the EIR only if a person spends most of their time at risk in that place. The formulas presented here are useful to quantify how local measures of mosquito blood feeding in a place could differ from what the humans living in that place would experience. What is the difference between risk for a human

It is made available under a [CC-BY 4.0 International license](#) .

who moves around compared to their counterfactual self who never leaves home? 1055

**The Spatial Scales of Transmission** Important considerations for planning, 1056  
monitoring, and evaluating malaria control are the spatial scales that characterize 1057  
transmission, as defined by parasite dispersal in mosquitoes or humans. We have 1058  
defined parasite dispersal rigorously in terms of the locations where blood meals 1059  
occurred that transmitted parasites in dispersal chains. While these definitions are 1060  
compelling, the distribution of distances separating every pair of infectious bites in a 1061  
chain of malaria infections can only be approximated using other data. In practice, the 1062  
framework we have described makes a distinction between local transmission and 1063  
imported or exported malaria. The framework makes the most sense mathematically if 1064  
*most* transmission is local, but the framework also defines quantities for malaria 1065  
importation and exportation, making it possible to study connectivity using a frame 1066  
that shifts among spatial domains and across spatial scales. 1067

After drawing a bounding box to define a spatial domain and a set of patches, we 1068  
classify any pair of bites in a transmission chain where at least one occurred in the 1069  
patch: either both bites occurred somewhere in the spatial domain, called local 1070  
transmission; or the first bite occurred outside the spatial domain, called imported 1071  
malaria; or the second bite occurred outside the spatial domain, called exported malaria. 1072  
These measures of imported and exported malaria thus provide a basis for 1073  
understanding and quantifying dispersal within and among defined geographical areas. 1074  
These models weigh the consequences of imported malaria, but as a practical matter, 1075  
the importance of exported malaria is difficult to quantify because the expected number 1076  
of subsequent bites depends on conditions somewhere else. Importantly, the fraction 1077  
that stays local may differ depending on whether the parasite is moving in a mosquito 1078  
or a human. Similar definitions and arguments would apply to transmission through a 1079  
parasite, a full parasite generation encompassing three bites and two jumps. The 1080  
metrics we have developed describe transmission within a defined geographical domain, 1081  
but if there is a need, the models can be reformulated for a larger spatial domain. 1082

The models and metrics provide a way of characterizing the spatial scales of 1083  
transmission by computing the cumulative fraction of all transmission occurring within 1084  
a circle of a given radius. Sensible points on that curve can be compared by patch: 1085  
What distances contain 80%, 90%, 95%, or 98% of all transmission? These estimates 1086  
are, out of necessity, based on estimated quantities – models of mosquito mobility, 1087  
human mobility, and modeled mosquito population density – about which there is 1088  
substantial uncertainty. 1089

Despite the overall uncertainty, these spatial scales are constrained by limits on time 1090  
and travel. Some quantities are known from census data (*e.g.* population distributions). 1091  
Most mosquito dispersal distances are short. Mosquitoes can move large distances, but 1092  
most stay within 1 *km* of a natal habitat [62]. For humans, the fraction of time spent 1093  
declines sharply with distance away from home. A large fraction of time is spent at 1094  
home, especially at night, and a larger fraction of the time is spent within roughly 10 1095  
*km* of home. The fraction of time spent drops off sharply with  $\log_{10}$  distance. The 1096  
spatial scales also depend on transmission intensity. In places with highly heterogeneous 1097  
transmission, places with the highest transmission intensity, will have the greater the 1098  
fraction of transmission that occurs at short distances. 1099



It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

**Mosquitoes, Travel, and Transmission** Highly spatially resolved data describing the EIR is rarely available. It is often cheaper, albeit less accurate, to use cross-sectional blood survey data describing malaria prevalence (*i.e.* the parasite rate, PR) to estimate local transmission. Spatial models and spatial metrics described herein provide some guidelines about how patterns in the PR can be used to identify areas with the most mosquitoes, particularly given the enormous heterogeneity in human population density.

It is commonly assumed that local clustering of cases implies that there is local transmission. For models developed in this framework, the vectorial capacity matrix (Eq. 39) describes parasite dispersion by mosquitoes, and evidence suggests that the spatial scales describing parasite dispersal by mosquitoes could vary by context [62].

Importantly, imported malaria can confound the relationship between local transmission by mosquitoes and prevalence. Travel habits and other traits describing humans often cluster spatially, partly because human neighborhoods are organized by socio-economic status. Spatial clustering of cases could arise if travel habits and thus malaria importation rates are spatially clustered, giving the appearance of local transmission.

**Measuring Reproductive Success** The most complete measure of transmission in an area is a reproductive number – the number of malaria cases arising from each malaria case after one complete parasite generation. We have defined reproductive matrices in several ways as matrices describing reproductive success among patches within a spatial domain, which can be used to define local reproductive numbers as cases arising from a patch. These reproductive matrices form a basis for investigating the appropriate spatial scales to measure and model transmission, for estimating contamination in randomized control trials, and for understanding the spatial effect sizes of control. These can put other data into a context that is relevant for transmission. For example, mosquito counts data and measures of malaria can vary over very short distances [28, 62]. The functional relevance of local heterogeneity in mosquito catch counts or in malaria prevalence can be critically examined by examining a matrix that integrates the effects of parasite movement in both mosquitoes and humans. After fully considering the uncertainty, it may be possible to determine the relevant spatial scales of transmission and thus the relevant spatial units for estimating reproductive numbers for malaria dynamics and control.

## Discussion

The simplicity of the Ross-Macdonald model can be contrasted with Hackett’s description of the elaborate and context-dependent nature of malaria that he observed in the field [27]:

*... malaria is so moulded and altered by local conditions that it becomes a thousand different diseases and epidemiological puzzles. Like chess, it is played with a few pieces, but is capable of an infinite variety of situations.*

The Ross-Macdonald model clearly identified enough chess pieces to develop basic concepts and theory to describe and measure malaria transmission [1], such as vectorial

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

capacity, the basic reproductive numbers, daily human biting rates, sporozoite rates, 1141  
entomological inoculation rates, and malaria parasite rates (*i.e.* prevalence). These 1142  
basic metrics have formed the basis for quantitative studies of malaria transmission, but 1143  
they ignored heterogeneity and complexity. In particular, the metrics and associated 1144  
concepts describing *parasite dispersal* in infected mosquitoes and humans were missing. 1145

Parasite dispersal is defined by the locations where infecting bites occurred in chains 1146  
of transmission, tracing dispersal events backwards through alternating jumps in 1147  
moving, infected humans and mosquitoes. It is practically impossible to study 1148  
transmission directly, but this framework has established a quantitative basis for 1149  
studying transmission through a set of constructs describing closely related processes 1150  
that can be observed. We have established a basis for describing dispersal rigorously, 1151  
and for analyzing dispersal and simulating transmission. The metrics and concepts we 1152  
have proposed here are designed to quantify transmission (and uncertainty about 1153  
transmission) through the study of patterns and the processes that generated them. 1154  
The metrics provide a rigorous way of quantifying parasite dispersal and spatial 1155  
transmission intensity. 1156

In developing models of a specific place for monitoring and evaluating malaria, it is 1157  
important to understand where and when transmission occurs as well as the local 1158  
contextual factors that shape transmission. In the Ross-Macdonald model, the basic 1159  
notions of reproductive success, transmission, and community effect sizes of control were 1160  
based on the abstract notion of a *population*, but it was never clear how to define a 1161  
population for purposes of quantifying malaria transmission dynamics: “What, if 1162  
anything, is a malaria population?” *Focal* transmission has been described [63], but 1163  
without a quantitative basis for quantifying malaria spatial heterogeneity and spatial 1164  
dynamics, there was no basis for a nuanced quantitative discussion about “What, if 1165  
anything, is a focus?” Without defining explicit boundary conditions, it was easy to 1166  
ignore malaria importation: “What fraction of malaria in a defined area was 1167  
attributable to local transmission?” Without modeling structured populations, it was 1168  
impossible to understand how differences in human behaviors would affect 1169  
transmission [64]. *Who* is responsible for most local transmission or malaria 1170  
importation? In malaria control, these discussions have focused on the issue of 1171  
*stratification*, but it remains unclear whether those strata should define sub-populations, 1172  
spatial areas, or both. Without a framework for understanding malaria transmission 1173  
spatially in heterogeneous populations, it was difficult to develop a consistent 1174  
methodology for quantifying transmission in a specific place and time. 1175

We have synthesized a set of old, new, and revised models to fully develop concepts, 1176  
constrain parameters, and update basic concepts and metrics in a spatial context. New 1177  
algorithms have filled a need to connect model parameters with data and remove bias 1178  
while guaranteeing mathematical consistency. The new framework and spatial metrics 1179  
make model complexity scalable, and it provides a way to study the role of context in 1180  
mosquito ecology and malaria transmission. How and why do bionomic parameters vary 1181  
over space and time? What spatial scales characterize mosquito populations? What are 1182  
the appropriate spatial scales to measure transmission and intervention coverage as a 1183  
*spatial average*? What are some appropriate methods for dealing with population 1184  
heterogeneity, including heterogeneity arising from differences in behavior, exposure, or 1185  
immunity? 1186



It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

The framework emphasizes the way we organize our knowledge about malaria into bins of expertise. Given the complexity of the problem, this means modellers can build models that adapt over time as more information about transmission in a place accumulated. The first models can focus on components whose dynamics are better known, and use simpler, pragmatic approaches to parts of the model whose mechanistic foundations are more uncertain. Our framework can make building ensembles of plausible models to cover this uncertainty easier. Model building and model comparison makes it possible to weigh the importance of various factors in context. In asking *where* transmission is occurring, we are concerned about mosquito populations, human behaviors, and human blood feeding. In asking *who* is responsible for malaria, we are not just concerned about differences in infectiousness, but also populations who import malaria, and strata who play an out-sized role in moving malaria around an area. These are the basic quantities that play a role in spatial targeting and in tailoring interventions to context.

## Conclusion

The goal of this study was to develop and present framework that could support model building for planning, monitoring and evaluating malaria control programs. Suites of models developed in this framework can be used to synthesize data, to quantify the major factors affecting transmission in a particular place, to identify critical data gaps and prioritize new data collection, to propagate uncertainty through analyses, and to support policy. We plan to use the framework to synthesize evidence and to give robust policy advice about malaria control on Bioko Island, and elsewhere, iteratively as part of adaptive malaria control. The spatial metrics and concepts describe an important dimension of malaria transmission that can help tailor interventions and spatially target interventions. In future studies, we plan to address concerns about the temporal dimensions of transmission, including threshold conditions, forcing by weather, and the spatial dimensions of malaria control. In adaptive management, the goal is to support monitoring and evaluation by developing rigorous methods that quantify malaria transmission as a changing baseline (*e.g.*, forced by weather) that has been modified by control. In other settings, the framework can be used to enhance the design of randomized control trials or to help programs implement and interpret *ad hoc* experiments to fill local knowledge gaps. Simulation-based analytics in this framework can be updated using evidence collected by malaria programs to update models and analysis and revise policy recommendations, to target and tailor interventions, and to use evidence to adapt to changing local conditions.


## Acknowledgments

DLS and SLW were supported by a grant from the National Science Foundation, Directorate for Technology, Innovation, and Partnerships (TIP) as part of the Convergence Accelerator Program (NSF 2040688). DLS, JMH, and AC were supported by a grant from the National Institute of Allergies and Infectious Diseases (R01 AI163398). DLS was supported by grant from the National Institutes of Health

It is made available under a [CC-BY 4.0 International license](#) .


(2U19AI089674). DLS, DMS, and JNN, were supported by a grant from the Bill and Melinda Gates Foundation (INV 030600). This project is partly a product of discussions with the mosquito working groups of working groups over several years RAPIDD (Research and Policy for Infectious Disease Dynamics). Over that time, many of us benefitted from the unwavering support and inspiration of F. Ellis McKenzie. OJB was supported by a UK Medical Research Council Career Development Award (MR/V031112/1).

1228  
1229  
1230  
1231  
1232  
1233  
1234




**Fig 8.** An important practical concern is spatial granularity of patches for simulation-based studies. For Bioko Island, Equatorial Guinea, for example, we could define patches at several scales: the whole island; or approximately 240 occupied areas ( $1\text{km} \times 1\text{km}$ , the squares); approximately 4,400 occupied  $100\text{m} \times 100\text{m}$  sectors (points); or 8 distinct regions (the colors of the squares); or clusters of contiguous sectors (the colors of the points); or approximately 70,000 individual households. An important concern is that the weight of evidence – the number of observations per patch – declines sharply as granularity of the simulations increases. This framework makes it possible to define a set of nested (or partially nested) studies that modify the number and size of patches, which requires modifying the human and mosquito mobility sub-models, but that holds other aspects of the model constant.

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .



**Fig 9.** (Left): bulk transmission metric describing transmission from the most densely populated area in Malabo, the capitol city, seen as the bright cell in the Northern tip of the island, to all other populated areas. (Right): bulk transmission from the most highly populated area in the south of the island (Luba), seen as the bright cell in the small harbor on the Western coast of the island.



**Fig 10. Structural Elements** of the framework are flexible to facilitate building models that are appropriate for various settings. These diagrams illustrate two examples. left) A forest malaria model with seven patches (including 3 villages and 2 campsites), 6 population strata, and 5 aquatic habitats. The village residents are stratified into loggers and other residents. Loggers from different villages spend time at home or in campsites, which have no permanent residents. Aquatic habitats (the moons) can be in villages, in campsites, or in patches near villages. Some villages (*e.g.* village 3), could lack mosquitoes but still have populations at risk. Right) It is also possible to model indoor and outdoor blood feeding with indoor and outdoor patches that share the same place. In these models, movement indoors *vs.* outdoors in the same place is modeled differently from movement among outdoor patches.

It is made available under a [CC-BY 4.0 International license](#) .

## Supplements

1235

### List of Supplements

1236

1. Supplement 1 - a github repository <https://dd-harp.github.io/exDE/>

1237

2. Supplement 2 - Glossary

1238

3. Supplement 3 - Modular Notation

1239

4. Supplement 4 - Human Travel and Mobility

1240

5. Supplement 5 - Vector Dynamics

1241

## References

1. Smith DL, Battle KE, Hay SI, Barker CM, Scott TW, McKenzie FE. Ross, Macdonald, and a theory for the dynamics and control of mosquito-transmitted pathogens. *PLoS Pathog.* 2012;8(4):e1002588.
2. Reiner RC Jr, Perkins TA, Barker CM, Niu T, Chaves LF, Ellis AM, et al. A systematic review of mathematical models of mosquito-borne pathogen transmission: 1970-2010. *J R Soc Interface.* 2013;10(81):20120921.
3. Smith DL, Perkins TA, Reiner RC Jr, Barker CM, Niu T, Chaves LF, et al. Recasting the theory of mosquito-borne pathogen transmission dynamics and control. *Trans R Soc Trop Med Hyg.* 2014;108(4):185–197.
4. Levin S. The Problem of Relevant Detail. In: Busenberg S, Martelli M, editors. *Differential Equations Models in Biology, Epidemiology and Ecology: Proceedings of a Conference held in Claremont California, January 13–16, 1990.* Berlin, Heidelberg: Springer Berlin Heidelberg; 1991. p. 9–15.
5. Radcliffe J. The initial geographical spread of host-vector and carrier-borne epidemics. *J Appl Probab.* 1973;10(4):703–717.
6. Bailey NTJ. *The Biomathematics of Malaria.* Oxford: Charles Griffin & Company Ltd.; 1982.
7. Dye C, Hasibeder G. Population dynamics of mosquito-borne disease: effects of flies which bite some people more frequently than others. *Trans R Soc Trop Med Hyg.* 1986;80(1):69–77.
8. Hasibeder G, Dye C. Population dynamics of mosquito-borne disease: persistence in a completely heterogeneous environment. *Theor Popul Biol.* 1988;33(1):31–53.
9. Sattenspiel L, Dietz K. A structured epidemic model incorporating geographic mobility among regions. *Math Biosci.* 1995;128(1-2):71–91.
10. Torres-Sorando L, Rodriguez DJ. Models of spatio-temporal dynamics in malaria. *Ecol Modell.* 1997;104(2-3):231–240.
11. Jansen VA, Lloyd AL. Local stability analysis of spatially homogeneous solutions of multi-patch systems. *J Math Biol.* 2000;41(3):232–252.
12. Auger P, Kouokam E, Sallet G, Tchuente M, Tsanou B. The Ross–Macdonald model in a patchy environment. *Math Biosci.* 2008;216(2):123–131.
13. Adams B, Kapan DD. Man bites mosquito: understanding the contribution of human movement to vector-borne disease dynamics. *PLoS One.* 2009;4(8):e6763.
14. Cosner C, Beier JC, Cantrell RS, Impoinvil D, Kapitanski L, Potts MD, et al. The effects of human movement on the persistence of vector-borne diseases. *J Theor Biol.* 2009;258(4):550–560.
15. Gao D, Ruan S. A multi-patch malaria model with logistic growth populations. *SIAM J Appl Math.* 2012;72(3):819–841.

16. Perkins TA, Scott TW, Le Menach A, Smith DL. Heterogeneity, mixing, and the spatial scales of mosquito-borne pathogen transmission. *PLoS Comput Biol*. 2013;9(12):e1003327.
17. Ruktanonchai NW, DeLeenheer P, Tatem AJ, Alegana VA, Caughlin TT, Zu Erbach-Schoenberg E, et al. Identifying Malaria Transmission Foci for Elimination Using Human Mobility Data. *PLoS Comput Biol*. 2016;12(4):e1004846.
18. Citron DT, Guerra CA, Dolgert AJ, Wu SL, Henry JM, Sánchez C HM, et al. Comparing metapopulation dynamics of infectious diseases under different models of human movement. *Proc Natl Acad Sci U S A*. 2021;118(18).
19. Guerra CA, Citron DT, García GA, Smith DL. Characterising malaria connectivity using malaria indicator survey data. *Malar J*. 2019;18:440.
20. Smith NR, Trauer JM, Gambhir M, Richards JS, Maude RJ, Keith JM, et al. Agent-based models of malaria transmission: A systematic review. *Malar J*. 2018;17(1):299.
21. Grillet ME, Hernández-Villena JV, Llewellyn MS, Paniz-Mondolfi AE, Tami A, Vincenti-Gonzalez MF, et al. Venezuela's humanitarian crisis, resurgence of vector-borne diseases, and implications for spillover in the region. *Lancet Infect Dis*. 2019;19(5):e149–e161.
22. Jaramillo-Ochoa R, Sippy R, Farrell DF, Cueva-Aponte C, Beltrán-Ayala E, Gonzaga JL, et al. Effects of political instability in Venezuela on malaria resurgence at Ecuador-Peru border, 2018. *Emerg Infect Dis*. 2019;25(4):834–836.
23. Cohen JM, Smith DL, Cotter C, Ward A, Yamey G, Sabot OJ, et al. Malaria resurgence: a systematic review and assessment of its causes. *Malar J*. 2012;11:122.
24. Plowe CV, Kublin JG, Dzinjalama FK, Kamwendo DS, Mukadam RAG, Chimpeni P, et al. Sustained clinical efficacy of sulfadoxine-pyrimethamine for uncomplicated falciparum malaria in Malawi after 10 years as first line treatment: five year prospective study. *BMJ*. 2004;328(7439):545.
25. Epstein A, Maiteki-Sebuguzi C, Namuganga JF, Nankabirwa JI, Gonahasa S, Opigo J, et al. Resurgence of malaria in Uganda despite sustained indoor residual spraying and repeated long lasting insecticidal net distributions. *PLOS Global Public Health*. 2022;2(9):e0000676.
26. Hackett LW, Missiroli A. The natural disappearance of malaria in certain regions of Europe. *Am J Epidemiol*. 1931;13(1):57–78.
27. Hackett LW. *Malaria in Europe. An Ecological Study*. London, Oxford University Press; 1937.
28. Bejon P, Williams TN, Nyundo C, Hay SI, Benz D, Gething PW, et al. A micro-epidemiological analysis of febrile malaria in Coastal Kenya showing hotspots within hotspots. *Elife*. 2014;3(2-3):e02130.

29. Ross R. Some quantitative studies in epidemiology. *Nature*. 1911;87(2188):466–467.
30. Macdonald G. The analysis of equilibrium in malaria. *Trop Dis Bull*. 1952;49(9):813–829.
31. Smith DL, Perkins TA, Tusting LS, Scott TW, Lindsay SW. Mosquito population regulation and larval source management in heterogeneous environments. *PLoS One*. 2013;8(8):e71247.
32. Brady OJ, Godfray HCJ, Tatem AJ, Gething PW, Cohen JM, McKenzie FE, et al. Adult vector control, mosquito ecology and malaria transmission. *Int Health*. 2015;7(2):121–129.
33. Killeen GF, McKenzie FE, Foy BD, Bøgh C, Beier JC. The availability of potential hosts as a determinant of feeding behaviours and malaria transmission by African mosquito populations. *Trans R Soc Trop Med Hyg*. 2001;95(5):469–476.
34. Ma BO, Roitberg BD. The role of resource availability and state-dependence in the foraging strategy of blood-feeding mosquitoes. *Evol Ecol Res*. 2008;10(8):1111–1130.
35. Yakob L. How do biting disease vectors behaviourally respond to host availability? *Parasit Vectors*. 2016;9:468.
36. Thongsripong P, Hyman JM, Kapan DD, Bennett SN. Human-mosquito contact: A missing link in our understanding of mosquito-borne disease transmission dynamics. *Ann Entomol Soc Am*. 2021;114(4):397–414.
37. Service, M W. Mosquito (Diptera: Culicidae) dispersal—the long and short of it. *J Med Entomol*. 1997;34(6):579–588.
38. Smith DL, Dushoff J, McKenzie FE. The risk of a mosquito-borne infection in a heterogeneous environment. *PLoS Biol*. 2004;2(11):e368.
39. Wu SL, Sánchez C HM, Henry JM, Citron DT, Zhang Q, Compton K, et al. Vector bionomics and vectorial capacity as emergent properties of mosquito behaviors and ecology. *PLoS Comput Biol*. 2020;16(4):e1007446.
40. Brady OJ, Godfray HCJ, Tatem AJ, Gething PW, Cohen JM, McKenzie FE, et al. Vectorial capacity and vector control: reconsidering sensitivity to parameters for malaria elimination. *Trans R Soc Trop Med Hyg*. 2016;110(2):107–117.
41. Smith DL, Drakeley CJ, Chiyaka C, Hay SI. A quantitative analysis of transmission efficiency versus intensity for malaria. *Nat Commun*. 2010;1(8):108.
42. Le Menach A, Tatem AJ, Cohen JM, Hay SI, Randell H, Patil AP, et al. Travel risk, malaria importation and malaria transmission in Zanzibar. *Sci Rep*. 2011;1:93.



43. R Core Team. R: A Language and Environment for Statistical Computing; 2022.
44. Soetaert K, Petzoldt T, Setzer RW. Solving Differential Equations in R: Package deSolve; 2010.
45. Citron DT, Guerra CA, García GA, Wu SL, Battle KE, Gibson HS, et al. Quantifying malaria acquired during travel and its role in malaria elimination on Bioko Island. *Malar J.* 2021;20:359.
46. Rund SSC, O'Donnell AJ, Gentile JE, Reece SE. Daily rhythms in mosquitoes and their consequences for malaria transmission. *Insects.* 2016;7(2).
47. Monroe A, Moore S, Okumu F, Kiware S, Lobo NF, Koenker H, et al. Methods and indicators for measuring patterns of human exposure to malaria vectors. *Malar J.* 2020;19(1):207.
48. Monroe A, Msaky D, Kiware S, Tarimo BB, Moore S, Haji K, et al. Patterns of human exposure to malaria vectors in Zanzibar and implications for malaria elimination efforts. *Malar J.* 2020;19(1):1–14.
49. Aron JL, May RM. The population dynamics of malaria. In: Anderson RM, editor. *The Population Dynamics of Infectious Diseases: Theory and Applications*. Boston, MA: Springer US; 1982. p. 139–179.
50. Le Menach A, McKenzie FE, Flahault A, Smith DL. The unexpected importance of mosquito oviposition behaviour for malaria: Non-productive larval habitats can be sources for malaria transmission. *Malar J.* 2005;4(1):23.
51. Nåsell I. *Hybrid Models of Tropical Infections*. vol. 59 of *Lecture Notes in Biomathematics*. 1st ed. Berlin: Springer-Verlag; 1985.
52. Johnston GL, Smith DL, Fidock DA. Malaria's missing number: calculating the human component of  $R_0$  by a within-host mechanistic model of *Plasmodium falciparum* infection and transmission. *PLoS Comput Biol.* 2013;9(4):e1003025.
53. Guerra CA, Kang SY, Citron DT, Hergott DEB, Perry M, Smith J, et al. Human mobility patterns and malaria importation on Bioko Island. *Nat Commun.* 2019;10:2332.
54. Darroch JN, Seneta E. On quasi-stationary distributions in absorbing continuous-time finite Markov chains. *J Appl Probab.* 1967;4(1):192–196.
55. Wu SL, Bennett JB, Sánchez C HM, Dolgert AJ, León TM, Marshall JM. MGDive 2: A simulation framework for gene drive systems incorporating seasonality and epidemiological dynamics. *PLoS Comput Biol.* 2021;17(5):e1009030.
56. Macdonald G. The analysis of the sporozoite rate. *Trop Dis Bull.* 1952;49(6):569–586.
57. Smith DL, McKenzie FE. Statics and dynamics of malaria infection in *Anopheles* mosquitoes. *Malar J.* 2004;3:13.

It is made available under a [CC-BY 4.0 International license](#) .

58. Garrett-Jones C. Prognosis for interruption of malaria transmission through assessment of the mosquito's vectorial capacity. *Nature*. 1964;204:1173–1175.
59. Smith DL, Musiime AK, Maxwell K, Lindsay SW, Kiware S. A New Test of a Theory about Old Mosquitoes. *Trends Parasitol*. 2021;37(3):185–194.
60. van den Driessche P, Watmough J. Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. *Math Biosci*. 2002;180(1-2):29–48.
61. Diekmann O, Heesterbeek JAP, Metz JAJ. On the definition and the computation of the basic reproduction ratio  $R_0$  in models for infectious diseases in heterogeneous populations. *J Math Biol*. 1990;28(4):365–382.
62. Carter R, Mendis KN, Roberts D. Spatial targeting of interventions against malaria. *Bull World Health Organ*. 2000;78(12):1401–1411.
63. Bousema T, Griffin JT, Sauerwein RW, Smith DL, Churcher TS, Takken W, et al. Hitting hotspots: spatial targeting of malaria for control and elimination. *PLoS Med*. 2012;9(1):e1001165.
64. Sturrock HJW, Hsiang MS, Cohen JM, Smith DL, Greenhouse B, Bousema T, et al. Targeting asymptomatic malaria infections: active surveillance in control and elimination. *PLoS Med*. 2013;10(6):e1001467.