

Monitoring Body Composition Change for Intervention Studies with Advancing 3D Optical Imaging Technology in Comparison to Dual-Energy X-Ray Absorptiometry

Michael C. Wong^{1,2}, Jonathan P. Bennett^{1,2}, Lambert T. Leong¹, Isaac Y. Tian³, Yong E. Liu¹, Nisa N. Kelly¹, Cassidy McCarthy⁴, Julia MW Wong⁵, Cara B. Ebbeling⁵, David S. Ludwig⁵, Brian A. Irving⁶, Matthew C. Scott^{4,6}, James Stampley⁶, Brett Davis⁶, Neil Johannsen^{4,6}, Rachel Matthews⁶, Cullen Vincelle⁶, Andrea K. Garber⁷, Gertraud Maskarinec¹, Ethan Weiss⁷, Jennifer Rood⁴, Alyssa N. Varanoske^{8,9}, Stefan M. Pasiakos⁸, Steven B. Heymsfield⁴, John A. Shepherd^{1,2}

¹ Department of Epidemiology, University of Hawaii Cancer Center, Honolulu, HI;

² Department of Human Nutrition, Food and Animal Sciences, University of Hawaii at Manoa, Honolulu, HI;

³ Paul G. Allen School of Computer Science and Engineering, University of Washington, Seattle, WA;

⁴ Pennington Biomedical Research Center, Baton Rouge, LA;

⁵ New Balance Foundation Obesity Prevention Center, Boston Children's Hospital, Boston, MA;

⁶ Louisiana State University, School of Kinesiology, Baton Rouge, LA;

⁷ Department of Pediatrics, University of California San Francisco, San Francisco, CA;

⁸ Military Nutrition Division, U.S. Army Research Institute of Environmental Medicine, Natick, MA;

⁹ Oak Ridge Institute for Science and Education, Oak Ridge, TN

Funding

Phases of this study were funded by the National Institute of Diabetes and Digestive and Kidney Diseases (NIH R01 DK R01DK109008), National Institute of Aging (R21AG058181), the US Army Medical Research and Development Command, Military Operational Medicine Research Program Grant W81XWH-19-C-0162. Supported in part by an appointment to the U.S. Army Research Institute of Environmental Medicine administered by the Oak Ridge Institute for Science and Education through an interagency agreement between the U.S. Department of Energy and the U.S. Army Medical Research and Development Command.

Clinical Trials Number

Shape Up! Adults (NCT03637855, <https://clinicaltrials.gov/ct2/show/NCT03637855>)

Macronutrients and Body Fat Accumulation: A Mechanistic Feeding Study (NCT03394664, <https://clinicaltrials.gov/ct2/show/NCT03394664>)

Resistance Exercise and Low-Intensity Physical Activity Breaks in Sedentary Time to Improve Muscle and Cardiometabolic Health (NCT03771417, <https://clinicaltrials.gov/ct2/show/NCT03771417>)

Time Restricted Eating on Weight Loss (NCT03393195, <https://clinicaltrials.gov/ct2/show/NCT03393195>)

Trial of Testosterone Undecanoate for Optimizing Performance During Military Operations (NCT04120363, <https://clinicaltrials.gov/ct2/show/NCT04120363>)

Conflicts of Interest

SBH reports his role on the Medical Advisory Boards of Tanita Corporation, Amgen, and Medifast; he is also an Amazon Scholar.

The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Army or the Department of Defense. Any citations of commercial organizations and trade names in this report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.

All other authors have no disclosures.

Corresponding Author

John A. Shepherd

701 Ilalo Street

Honolulu Hawaii 96813

Email: johnshep@hawaii.edu

Running Title: Monitoring Body Composition with 3D Optical Shape

Abbreviations: 3DO (three-dimensional optical), ALM (appendicular lean mass), ALMI (appendicular lean mass index), CV (coefficient of variation), DXA (dual-energy X-ray absorptiometry), FM (fat mass), FFM (fat-free mass), LSC (least significant change), PCA (principal component analysis), PCs (principal components), VAT (visceral adipose tissue)

1 **ABSTRACT**

2 **Background**

3 Recent three-dimensional optical (3DO) imaging advancements have provided more accessible,
4 affordable, and self-operating opportunities for assessing body composition. 3DO is accurate
5 and precise with respect to clinical measures made by dual-energy X-ray absorptiometry (DXA).
6 However, the sensitivity for monitoring body composition change over time with 3DO body
7 shape is unknown.

8 **Objective**

9 To evaluate 3DO's ability to monitor body composition changes across multiple intervention
10 studies.

11 **Methods**

12 A retrospective analysis was performed using intervention studies on healthy adults that were
13 complimentary to the cross-sectional study, Shape Up! Adults. Each participant received a DXA
14 (Hologic Discovery/A system) and 3DO (Fit3D ProScanner) scan at baseline and follow-up. 3DO
15 meshes were digitally registered and reposed using Meshcapade to standardize the vertices
16 and pose. Using an established statistical shape model, each 3DO mesh was transformed into
17 principal components (PCs), which were used to predict whole-body and regional body
18 composition values using published equations. Body composition changes (follow-up minus
19 baseline) were compared to DXA with linear regression.

20 **Results**

21 The analysis included 133 participants (45 females) in six studies. The mean (SD) length of
22 follow-up was 13 (5) weeks, range 3-23 weeks. Agreement between 3DO and DXA (R^2) for
23 changes in total fat mass (FM), total fat-free mass (FFM), and appendicular lean mass,
24 respectively, were 0.86, 0.73, and 0.70 with RMSEs of 1.98 kg, 1.58 kg, and 0.37 kg in females,
25 and 0.75, 0.75, and 0.52 with RMSEs of 2.31 kg, 1.77 kg, and 0.52 kg in males. Further
26 adjustment with demographic descriptors improved the 3DO change agreement to changes
27 observed with DXA.

28 **Conclusions**

29 As compared to DXA, 3DO was highly sensitive in detecting body shape changes over time. The
30 3DO method was sensitive enough to detect even small changes in body composition during
31 intervention studies. The safety and accessibility of 3DO allows users to self-monitor on a
32 frequent basis throughout interventions.

33 **Keywords:** Body Composition, Three-dimensional Optical Imaging, DXA, Interventions, Weight-
34 loss, Monitoring

35 INTRODUCTION

36 Obesity remains an area of concern as global prevalence continues to rise (1-3).
37 According to the Center for Disease Control and Prevention (CDC), more than 42% of adults in
38 the United States were considered obese in 2018, while the prevalence was approximately 30%
39 two decades prior (4). To counter the obesity epidemic, diet and physical activity interventions
40 have been studied extensively to target weight loss (5, 6). However, metaanalyses have shown
41 that weight is only loosely associated with metabolic health (7) and initial changes in response
42 to intervention are small and quickly undone long-term (6, 8). On the other hand, a range of
43 changes in body composition (reduced total body, abdominal and visceral fat, and increased
44 muscle mass) can be produced through diet or exercise intervention and are consistently
45 associated with decreased cardiovascular disease risk (9, 10). Further, decrements in skeletal
46 muscle, particularly in the elderly, can lead to losses in strength and endurance, reductions in
47 energy expenditure, and an increased risk of insulin resistance (11). Nevertheless, measures of
48 body composition have been relegated to research and specialized facilities while clinical care
49 continues to rely on weight as a flawed marker of health.

50 Another pressing reason to limit our reliance on weight is the differing relative weight of
51 the compartments and tissues. Garrow (12) suggested, and Prentice et al. (13) concurred, that
52 weight loss is typically 25% fat-free mass (FFM, i.e. lean soft tissue + bone mineral content) and
53 75% fat loss. However, recent research suggested this “25/75 rule of thumb” may not
54 accurately describe various weight loss interventions. The amount FFM lost depends on energy
55 intake, diet composition, sex, baseline adiposity, inactivity or type and physical activity level,
56 and potentially metabolic and hormonal responses (14, 15). As such, it may not be appropriate

57 to only monitor weight in interventions as FFM might be lost at a greater proportion. By
58 monitoring body composition, investigators receive a more accurate assessment of their
59 intervention's efficacy.

60 Body composition by dual-energy X-ray absorptiometry (DXA) has been used extensively
61 in clinical settings for its accuracy and precision of whole-body and regional measurements
62 (16). Although DXA provides clinically useful measurements (e.g., bone mineral density, VAT,
63 fat, and lean mass), it requires expensive radiological equipment, qualified technicians, and may
64 not be feasible or accessible for routine clinical practice or frequent monitoring. The ideal body
65 composition method should be affordable, accessible, free of ionizing radiation, and not require
66 radiological-qualified technicians.

67 Three-dimensional optical (3DO) imaging has become increasingly accessible, made
68 large advancements in recent years, and is safe to use repeatedly (17, 18). 3DO scanners
69 provide accurate and precise digital anthropometry in comparison to criterion methods and
70 output a 3D mesh that represents a person's entire shape (19, 20). 3DO shape has shown to be
71 highly predictive of DXA body composition (21-24). The next pressing issue is whether 3DO
72 could successfully capture *changes* in body composition as different modalities have not
73 validated well longitudinally to standard methods (25). However, the lack of available
74 longitudinal data has limited the assessment of 3DO in monitoring body composition changes.
75 The hypothesis of the study was that 3DO can monitor change with a similar sensitivity to DXA
76 given previous cross-sectional accuracy and precision. Therefore, the objective of this study was
77 to evaluate 3DO's accuracy for monitoring body composition changes across a variety of
78 intervention studies in comparison to DXA.

79 METHODS

80 Study Design

81 The current study was a retrospective analysis of six complimentary intervention studies
82 to the Shape Up! Adults study (NIH R01 DK109008, ClinicalTrials.gov ID NCT03637855), which
83 was originally a cross-sectional study with a planned longitudinal arm. In order to study 3DO's
84 ability to monitor body composition over time, collaborators adopted our DXA and 3DO
85 protocol. The studies included Time-restricted Eating on Weight Loss (TREAT) (26);
86 Macronutrients and Body Fat Accumulation: A Mechanistic Feeding Study (FB4: Framingham,
87 Boston, Bloomington, Birmingham, and Baylor) (27-29); Resistance Exercise and Low-Intensity
88 Physical Activity Breaks in Sedentary Time to Improve Muscle and Cardiometabolic Health Pilot
89 Study (REALPA, NIH R21AG058181, ClinicalTrials.gov ID NCT03771417) (30), Louisiana State
90 University (LSU) Athletes; Trial of Testosterone Undecanoate for Optimizing Performance
91 During Military Operations (OPS II) (31, 32); and patients with bariatric surgery. If available,
92 study-specific information (e.g., site, protocol, aims) can be found on ClinicalTrials.gov (**Table**
93 **1**). All study protocols were previously approved by their respective Institutional Review Board.

94 The TREAT participants were only allowed to consume food between 12 pm and 8 pm
95 with the goal of fat loss (26). The FB4 cohort was given a hypocaloric, low-carbohydrate diet
96 with the goal to lose $15\pm 3\%$ of baseline body weight (27-29). The REALPA study introduced
97 whole-body resistance exercise (2 days/week) alone (i), or with moderate intensity aerobic
98 exercise (50 min/day, 3 days/week at 4 METS) (ii), or low-intensity physical activity (LPA) breaks
99 in sedentary time (~10 min/break, 6 breaks/day, 5 days/week at 2METS) in adults 65-80 years

100 of age to observe changes to muscle and cardiometabolic health markers after the 16-week
101 long intervention. A subset of the REALPA participants who had both 3DO and DXA scans
102 available for analysis were included in this retrospective study. (30). The LSU athletes were
103 female basketball players that were measured at the beginning and through training camp
104 (ranging 1-5 months), evaluating body composition as a result of a preseason training program.
105 The OPS II study tested the effects of an intramuscular injection of testosterone undecanoate
106 compared to placebo on changes in muscle and fat mass in recreationally-active young males
107 undergoing simulated military operational stress consisting of sleep restriction, high exercise-
108 induced energy expenditure, and limited energy intake (31, 32). The bariatric patients who
109 were recruited from the University of California, San Francisco (UCSF) received bariatric surgery
110 (surgery type was not recorded) with the goal of weight-loss. Although weight loss or body
111 composition changes was not the goals of each study, the current aim of this analysis was to
112 evaluate 3DO's ability to monitor body composition changes in comparison to DXA.

113 **Participants**

114 All participants provided informed consent before participation. Participants were
115 deemed ineligible for this analysis if they were pregnant, breast feeding, had missing limbs,
116 non-removable metal (e.g., joint replacements), previous body-altering surgery (e.g., breast
117 augmentation), unable to stand still for one minute, or lie still for 3 minutes. Participants
118 received same-day whole-body 3DO and DXA scans at baseline and follow-up. If the study had
119 two follow-up appointments, the first of the two was used and will be considered as the
120 "follow-up" in the current analysis. Participants were excluded from the analysis if they were
121 missing either baseline or follow-up data from either DXA or 3DO.

122 **DXA**

123 Height and weight were measured prior to the DXA scan. Participants received a single
124 whole-body DXA scan with a Hologic Discovery/A system (Hologic Inc., Marlborough,
125 Massachusetts) according to International Society for Clinical Densitometry guidelines (33).
126 Participants laid supine on the scanning bed and were positioned by the DXA technician with
127 arms by the side and feet internally rotated. Scans took approximately three minutes for the
128 whole-body scan. All raw scans from each study were securely transferred to the University of
129 Hawaii Cancer Center and analyzed by a single certified technologist using Hologic Apex version
130 5.6 with the National Health and Nutrition Examination Survey Body Composition Analysis
131 calibration option disabled (34). After analysis, proprietary algorithms automatically generate
132 body composition values. DXA measurements used in this analysis included whole-body and
133 regional (i.e., arms, legs, trunk) fat and fat-free measures.

134 **Three-Dimensional Optical**

135 Participants changed into form-fitting tights, a swim cap, and a sports bra if female.
136 Female participants from the FB4 used form-fitting swimsuits. 3DO surface scans were taken on
137 the Fit3D ProScanner version 4.x (Fit3D Inc., San Mateo, California). Participants grasped
138 telescoping handles on the scanner platform and stood upright with shoulders relaxed and
139 arms positioned straight and abducted from their torso. The platform rotates once around and
140 takes approximately 45 seconds for the completion of the scan. Final point clouds were
141 converted to a mesh connected by triangles with approximately 300,000 vertices and 600,000
142 faces representing the body shape (22).

143 Fit3D meshes were sent to Meshcapade (Meshcapade GmbH, Tübingen, Germany) for
144 registration and to be digitally reposed. Their algorithm registers each mesh to a 110,000-vertex
145 template with complete anatomical correspondence. Each vertex corresponds to a specific
146 anatomical location across all registered meshes. All meshes were digitally reposed to a T-pose,
147 where the person was standing straight, arms were brought horizontal and in the plane with
148 the body, and arms and legs were straightened (35). The registered meshes were transformed
149 into principal component (PC) space from an established statistical shape model (24). Principal
150 component analysis orthogonalizes and reduces the dimensionality of the data so that fewer
151 variables are needed to describe the data's variance (21). Total FM and regional (i.e., arms, legs,
152 trunk, VAT) body composition estimates were derived previously using either exclusively PC
153 descriptors of shape or PC descriptors with demographic adjustments (24). In this study, PC
154 only body composition equations were used since complete demographics and
155 anthropometrics were not available on all cohorts. Total FFM and percent fat (%fat) were
156 derived dependently from total FM and total body mass. Appendicular lean mass (ALM) was
157 defined as the sum of lean soft tissue masses for legs and arms by convention (36). ALM index
158 (ALMI) was derived by dividing ALM by height-squared.

159 Average 3DO body shape representations were created to visualize the average shape at
160 baseline and follow-up for each intervention study. PCs were averaged by study intervention to
161 make an average vector of PCs, which was then inverted back into coordinate space (x, y, and z)
162 to acquire the image.

163 **Statistical Analysis**

164 Body composition change for all variables was defined as follow-up minus baseline. 3DO
165 body composition change was compared to DXA body composition change using linear
166 regression. Bland-Altman plots were made for the cross-sectional comparisons. Coefficient of
167 determination (R^2), Lin's concordance coefficient (CCC), and root mean square error (RMSE)
168 were used to report the relationship and accuracy of the comparison. Scatter plots were used
169 to visualize the comparison. Additional adjustments were made to explain any potential bias
170 using stepwise forward linear regression and five-fold cross-validation. Potential covariates
171 included ethnicity and baseline height, weight, BMI, and age as well as changes in weight,
172 height, and BMI. Covariates remained in the model if they had a p-value <0.05. The least
173 significant change (LSC) was used to determine if the change in the body composition measure
174 was significantly different (95% confidence) than zero (37). Student's t-test was used to
175 evaluate mean differences between 3DO and DXA outputs. P-value < 0.05 was considered
176 statistically significant.

177 The LSCs used in this analysis were derived from Wong et al. (24). LSC is defined as 2.77
178 x precision error (21, 37). Female LSC for FM, FFM, and %fat was 1.52 kg, 1.52 kg, and 2.27% for
179 3DO, while 0.64 kg, 0.75 kg, and 0.91% for DXA, respectively. Male LSC for FM, FFM, and %fat
180 was 1.22 kg, 1.22 kg, and 1.58% for 3DO, while 0.69 kg, 0.94 kg, and 0.78% for DXA,
181 respectively. Since ALM was not reported in the previous publication, in order to get the
182 precision error to calculate the LSC, the test-retest precision for ALM was derived using the
183 same sample from Wong et al. (24). Lastly, Cohen's kappa analysis was utilized to assess the
184 consistency between the DXA and 3DO report for each level, considering any agreement that
185 may have happened due to chance. The kappa scores can be interpreted as follows: 0-0.20 = no

186 agreement, 0.21-0.39 = fair agreement, 0.40-0.59 = moderate agreement, 0.60-0.79 =
187 substantial agreement, 0.80-0.99 = near perfect agreement, and 1 = perfect agreement (38).

188 To test Garrow's (12) and Prentice et al.'s (13) suggestion that 25% of weight loss is
189 FFM, change in total FFM was divided by the absolute change in weight multiplied by 100
190 ($[\Delta\text{FFM} / \text{absolute } \Delta\text{weight}] * 100$). This was applied to those who had a negative weight change
191 (weight loss). The absolute change in weight was used in order to evaluate gains or losses in
192 FFM.

193 Experimental models were created using the change in PCs (Δ PCs) to test if different
194 modeling methods would improve the body composition change predictions. Model 1 used the
195 Δ PCs, model 2 used the Δ PCs and baseline total fat mass, model 3 used the Δ PCs and baseline
196 PCs, model 4 used the baseline and follow-up PCs, and model 5 used the Δ PCs and change in
197 weight. These models were built with stepwise forward linear regression with five-fold cross-
198 validation. All statistical analyses were done in R version 4.2.1 (R Core Teams, Vienna, Austria).

199 **RESULTS**

200 One hundred and thirty-three participants (43 female and 85 male) were included in the
201 final analysis (**Table 2**). One-hundred sixty-four participants were excluded for dropout (n=4),
202 unavailable 3DO or DXA data at one or both timepoints (n=157), movement artifacts (n=2), or
203 mislabeled data (n=1) (**Supplemental Figure 1**). The time between baseline and follow-up DXA
204 scans across all studies ranged from three to twenty-three weeks (**Figure 1**). Females and males
205 lost on average 3.5 kg and 5.4 kg of total FM, 1.8 kg and 3.4 kg of total FFM, and 30 g and 100 g
206 of VAT as a result of interventions, respectively, according to DXA. The majority of the body
207 composition changes occurred in the trunk for both sexes. Average body shapes were
208 presented at baseline and follow-up for each study intervention (**Figure 2**).

209 In females (**Figure 3**), 3DO and DXA total FM and FFM at baseline (R^2 s; 0.91 and 0.84;
210 RMSE; 3.3 kg and 3.3 kg, respectively) and follow-up (R^2 s; 0.89 and 0.84; RMSE; 2.8 kg and 2.8
211 kg, respectively) were highly correlated, while %fat and ALM were moderately correlated (R^2 s:
212 0.61 – 0.79). Female Bland-Altman plots are shown in **Supplemental Figure 2**. Slight
213 proportional bias was observed in the baseline total FM from a single outlier with high leverage
214 but not seen in the follow-up. In the comparison of 3DO body composition changes to DXA
215 (**Supplemental Table 1**), 3DO achieved an R^2 of 0.86, 0.73, 0.23; and 0.70, CCC of 0.90, 0.82,
216 0.47, and 0.81; RMSE of 1.98 kg, 1.58 kg, 2.2%, 0.37 kg for total FM, total FFM, %fat, and ALM,
217 respectively. Mean differences were observed for VAT, ALM, ALMI, and leg FFM (30 g, 0.19 kg,
218 0.07 kg/m^2 , 0.16 kg, respectively, p-value < 0.05). After adjustments for possible covariates in
219 the stepwise linear regression models with forward selection (**Supplemental Table 2**), weight
220 change further explained variance in total FM, FFM, and ALM, while changes in BMI further

221 explained variance in %fat change, which modestly improved the R^2 s to 0.90, 0.80, 0.51, and
222 0.79, respectively. The adjustments with demographic covariates alleviated residual bias
223 between 3DO and DXA change. Mean differences (i.e., VAT, ALM, ALMI, and leg FFM) were no
224 longer observed after adjustments (p -value = 0.99). The green glyphs (**Figure 3 and 4**)
225 symbolizes the participants that exceeded both the DXA and 3DO LSCs, the purple glyphs were
226 those who did not exceed DXA nor 3DO's LSC, and the orange and blue glyphs exceeded one
227 LSC but not the other. The LSCs were depicted with the orange vertical lines (3DO) or the blue
228 horizontal lines (DXA).

229 In males (**Figure 4**), 3DO and DXA total FM and FFM at baseline (R^2 s; 0.88 and 0.84;
230 RMSE; 3.7 kg and 3.6 kg, respectively) and follow-up (R^2 s; 0.88 and 0.85; RMSE; 3.3 kg and 3.2
231 kg, respectively) were highly correlated, while %fat and ALM were moderately correlated (R^2 s:
232 0.60 – 0.75). Male Bland-Altman plots are shown in **Supplemental Figure 3**. Proportional bias
233 was observed in total FM and %fat due to a single outlier with high leverage. Not bias was
234 observed in total FFM and ALM. In the comparison of 3DO body composition changes to DXA
235 (**Supplemental Table 1**), 3DO achieved an R^2 of 0.75, 0.75, 0.25, and 0.52; CCC of 0.76, 0.78,
236 0.39, and 0.64; and RMSE of 2.3 kg, 1.8 kg, 2.4%, and 0.5 kg for total FM, total FFM, %fat, and
237 ALM, respectively. Of the observed variables in **Figure 4**, mean differences were observed for
238 total FM, total FFM, %fat, and ALM (p -value<0.001). After adjustments for possible covariates in
239 stepwise linear regression with forward selection (**Supplemental Table 2**), R^2 s modestly
240 improved for changes in total FM, FFM, %fat, and ALM to 0.80, 0.76, 0.42, and 0.56,
241 respectively. Mean differences in total FM, total FFM, %fat, and ALM were no longer observed
242 after adjustments (p -value = 0.99).

243 3DO predicted regional body composition changes (i.e., arms, legs, and trunk) with R^2 s
244 that ranged from 0.39 – 0.91 and RMSEs that ranged from 0.16 – 1.3 kg in females and males
245 (**Supplemental Table 1**). 3DO and DXA found significant changes in the majority of the sample
246 for total FM (69%), total FFM (78%), %fat (53%), and ALM (78%) (**Supplemental Table 3**).
247 Cohen's kappa showed that 3DO had a moderate agreement to DXA for most outputs.

248 From the total sample, 64% of females and 90% of males lost weight after intervention.
249 Of those that lost weight, % FFM loss relative to weight loss ranged from -1370% to 478% as
250 measured by DXA. Among those who lost FFM, 80% of females and 81% of males lost more
251 than 25% FFM relative to their weight loss (**Figure 5**). The % FFM loss relative to weight loss
252 according to study was on average 36% for Bariatric surgery, 37% for FB4, 27% for OPS II, 313%
253 for Athletes, 85% for TREAT, and 59% for REALPA. The very high percentages were for small
254 weight changes and/or large compartment changes with little overall weight change (e.g., Δ
255 FFM / absolute(Δ weight) * 100 for an Athlete = $(-1.96/0.36) * 100 = -544\%$).

256 The experimental models (**Supplemental Tables 4 and 5**) for females (models 1-5) and
257 males (models 1-3 and 5) predicted change better than the unadjusted results presented in
258 **Supplemental Table 1**. Female models 1, 2, and 3 were the same as male models 1, 2, and 3
259 after the stepwise regression. The prediction of total FM change improved in both females ($R^2 =$
260 0.94, RMSE = 1.29 kg) and males ($R^2 = 0.84$, RMSE = 1.79 kg).

261

262 DISCUSSION

263 These present findings support 3DO as a sensitive tool assessing changes in body
264 composition, compared to DXA. The current study evaluated 3DO's ability to monitor body
265 composition change with body shape via the 3DO mesh. Previously derived shape models and
266 equations were used to estimate body composition values (24), and 3DO change from baseline
267 to follow-up was compared to DXA change. Overall, 3DO change was highly correlated to DXA
268 change, and the two methods agreed on the statistical significance of the change in the
269 majority of the study population. Additional adjustments of demographics explained further
270 variance between 3DO and DXA change. The experimental models showed that specific
271 calibration with changes in shape might improve the body composition change prediction.
272 Additional longitudinal data may further validate the experimental models. Nevertheless, the
273 cross-sectional 3DO models produced comparable body composition estimates compared to
274 DXA, which supports our hypothesis that using 3DO imaging is a feasible method to monitor
275 body composition changes.

276 With respect to the "rule-of-thumb" by Garrow's (12) and Prentice et al. (13) that 25%
277 of weight loss comes from FFM, the proportion of FFM lost by the vast majority of our sample
278 was greater than 25% (**Figure 5**). Thus, this "rule-of-thumb" underestimated the loss of FFM for
279 most participants. The loss of FFM varied by study intervention due to differences in protocol,
280 type of activity, and objectives. This is a critical area of study given that loss of FFM is associated
281 with diminished strength and decrements in physical functioning, exercise endurance and
282 capacity, metabolic rate, and health (11). This further supports the idea that the relative
283 proportion of FM and FFM change is dependent on energy intake, diet composition, sex,

284 baseline adiposity, inactivity or type and level of physical activity level, and potentially
285 metabolic and hormonal responses (14). It should also be noted that the experimental group in
286 the OPS II study received testosterone, which has shown to improve muscle protein kinetics
287 during stress and was a contributing factor to the proportion of FFM loss (32). The LSU athletes
288 and OPS II participants were study examples that monitored body composition for
289 performance. 3DO can also provide an accessible and safe method to monitor body
290 composition changes since muscle build-up can play a pivotal role in strength, endurance, and
291 speed. Some participants had percent changes in FFM that were greater than 100%. This was
292 likely due to an undesirable recomposition where the person increased FM and lost FFM, which
293 resulted in a minimal weight change (small denominator). Hydration status may have impacted
294 individuals with minimal changes (15). Because DXA does not measure hydration status, DXA
295 has assumptions of hydration built-in, while 3DO estimates were regressed to DXA. The current
296 analysis further emphasized the importance of monitoring body composition, in addition to
297 weight, to better understand changes in health status and performance factors.

298 In the current study, the change in 3DO FM was underestimated while FFM was
299 overestimated in both females and males. These changes were not statistically significant in
300 females but were significant in males. A potential reason for the male discrepancies could be
301 that the current sample was beyond the adiposity range of the data used to train the original
302 3DO body composition models. The maximum percent fat of the training dataset was 38% (24),
303 while the current population's maximum was 47%. The training dataset lacked the shape
304 variance of males with extreme adiposity and underestimated FM and %fat in this study.
305 However, after adjusting for covariates, these discrepancies were no longer statistically

306 significant. Using either the 3DO changes models with demographics or retraining the model
307 with high adiposity participants could potentially address this issue.

308 Test-retest precision, also known as short-term precision, has an error that can be used
309 to derive the LSC (2.77 x precision error) (37). Metrics with changes that surpass their LSC
310 values were considered to be significantly greater than zero changes, with a 95% confidence.
311 3DO and DXA precision errors used for this study were taken from the Shape Up Adults group
312 (24). To reduce the 3DO precision error, to be closer to that of DXA, and track body composition
313 changes at the same significance, multiple 3DO scans could be taken and averaged at each time
314 point. For total FM, the 3DO precision errors were 0.44 kg and 0.55 kg as well as 0.25 kg and
315 0.23 kg for DXA in males and females, respectively (24). Assuming a normal distribution of the
316 precision errors, the precision of the averaged 3DO measure from multiple scans improves by
317 the square root of the number of scans (e.g., the average of four scans would lower the
318 precision error from 0.44 kg and 0.55 kg to 0.22 kg and 0.27 kg) (39). Unlike DXA, taking
319 multiple scans with 3DO is fast and without additional radiation concerns. Dizziness can be a
320 potential hazard, which is listed on the manufacturer warnings and our study consent form.
321 However, our participants did not report dizziness even with multiple scans.

322 3DO VAT change compared to DXA VAT change had a low correlation, but this is likely to
323 due to the compressed range of VAT change in our sample. In previous literature, DXA VAT has
324 been shown to be highly correlated to gold standard, magnetic resonance imaging (MRI), cross-
325 sectionally (R^2 range: 0.81 – 0.86) (40-42). However, DXA VAT often has proportional bias for
326 individuals with high VAT and has not validated well longitudinally compared to MRI (42, 43).
327 Some of the bias may be explained by technique differences as MRI and CT are analyzed by a

328 cross-section of the abdomen, while DXA is estimated from an X-ray's 2D area. Nevertheless,
329 given the cross-sectional correlation of DXA VAT to MRI and cardiometabolic markers, DXA VAT
330 can still be considered an accessible tool to characterize health risks that can prompt health
331 initiatives (44, 45). Since 3DO body shape has been shown to be correlated and predictive of
332 DXA VAT and cardiometabolic markers (21), 3DO may be considered an accessible tool that can
333 be used frequently due to its lack of ionizing radiation concerns. However, users should be wary
334 of DXA and 3DO's ability to accurately monitor VAT changes given the limited amount of
335 validation in the literature.

336 The 3DO vs DXA RMSEs for comparing baseline or follow-up scans were higher (worse)
337 than the RMSEs for the change comparisons (**Figures 3 and 4**). This may be due to the individual
338 systematic bias expressed in the comparison of one technique (3DO) to another (DXA) that is
339 subtracted away when looking at changes in the measures, and the fact that the precision error
340 of both techniques was much lower than this individual systemic bias found in inter-technique
341 comparisons of measures (46).

342 According to the authors' knowledge, one other study has reported 3DO's ability to
343 monitor body composition change. Tinsley et al. (25) compared the changes in proprietary body
344 composition estimates from 3DO scanners (Fit3D ProScanner, Size Stream SS20 (Size Stream
345 LLC, Cary, NC, USA), and Styku S100 (Styku Comp. Los Angeles, CA, USA)) with changes in 4-
346 compartment (4C) model measures in 21 volunteers. Changes in 3DO FM underperformed
347 (CCC, 0.22 to 0.40 for total FM) compared to the 4C model. CCC for total FM in this study was
348 much higher (CCC; 0.91 and 0.79) for females and males, respectively, showing better
349 agreement with DXA. However, there were major differences between the two studies. This

350 study had a larger sample size (n=128), greater ethnic-racial diversity, larger body composition
351 changes, publicly available equations, and a different body composition modality (DXA instead
352 of the 4C model, given that data for the 4C model was not available). Larger studies are
353 warranted to fully evaluate 3DO's ability to monitor change using the scanners' proprietary
354 body composition outputs.

355 This study had several strengths including sample size, variety and lengths of
356 interventions, and an ethnically diverse sample. A study with 128 participants is currently one
357 of the larger body composition change studies. Although the majority of the participants were
358 ethnically white, there was still a representation of other ethnic backgrounds. This study
359 included a variety of diet, physical activity, and surgical interventions that ranged from 3 to 23
360 weeks and included participants with large as well as little to no body composition changes.

361 This study is not without limitations. Although the sample size was rather large for a
362 longitudinal analysis, a representative test set could not be made to validate the adjusted and
363 experimental linear models. In addition, results from this study may not be generalizable to
364 populations with poor health status or children (< 18 years old). Future work would be required
365 to address gaps in knowledge in different populations such as infants and children, the accuracy
366 of change in specific subgroups (i.e., BMI, ethnicity, and age), and the assessment of more
367 accessible 3DO technology.

368 **CONCLUSION**

369 This study indicated that 3DO body composition changes were highly correlated to DXA
370 and show good feasibility to monitor changes over a variety of interventions. As the

371 accessibility and popularity of 3DO continues to grow, more people will be able to use this
372 technology to monitor their body composition in clinical and nonclinical settings. The findings of
373 this study extend the 3DO literature, which has been limited to cross-sectional performance.

374

375 **Acknowledgements**

376 We thank all the participants for graciously giving their time to be part of the studies, our
377 collaborators at each site for their part in the study, Tyler Carter and Greg Moore at Fit3D for
378 providing us the 3DO data, and Naureen Mahmood and Talha Zaman for providing us the
379 application program interface to register and repose our 3DO data. The data underlying this
380 study cannot be made publicly available because the data contains patient identifying
381 information. Data is available from the Shape Up! Studies for researchers who meet the criteria
382 for access to confidential data. For details and to request an application, please contact John
383 Shepherd johnshep@hawaii.edu or visit www.shapeup.shepherdresearchlab.org.

384 **Author Contributions**

385 MCW and JAS designed and conducted the research; MCW, LL, JB, IT, YEL, GM, and JAS were
386 part of the data analysis; JMWW, NNK, CBE, BI, MCS, JS, BD, NMJ, RM, CV, DSL, EW, SBH, and
387 JAS were in charge of their respective study recruitment and protocols; MCW and JAS drafted
388 the manuscript and were responsible over the final content. All authors reviewed and approved
389 the final manuscript.

390

REFERENCES

- 391 1. Wong MC, McCarthy C, Fearnbach N, Yang S, Shepherd J, Heymsfield SB. Emergence of the
392 obesity epidemic: six decade visualization with humanoid avatars. *The American journal of clinical*
393 *nutrition*. 2022.
- 394 2. Heymsfield SB, Wadden TA. Mechanisms, pathophysiology, and management of obesity. *New*
395 *England Journal of Medicine*. 2017;376(3):254-66.
- 396 3. Bennett J, Wong MC, McCarthy C, Fearnbach N, Queen K, Shepherd J, et al. Emergence of the
397 adolescent obesity epidemic in the United States: five-decade visualization with humanoid avatars.
398 *International Journal of Obesity*. 2022:1-4.
- 399 4. Hales CD CM, Fryar CD, and Ogden CL. Prevalence of Obesity and Severe Obesity Among Adults:
400 United States, 2017-2018 2020. Available from: [https://www.cdc.gov/nchs/data/databriefs/db360-](https://www.cdc.gov/nchs/data/databriefs/db360-h.pdf)
401 [h.pdf](https://www.cdc.gov/nchs/data/databriefs/db360-h.pdf).
- 402 5. Arem H, Irwin M. A review of web-based weight loss interventions in adults. *Obesity reviews: an*
403 *official journal of the International Association for the Study of Obesity*. 2011;12(5):e236-e43.
- 404 6. Wu T, Gao X, Chen M, Van Dam R. Long-term effectiveness of diet-plus-exercise interventions
405 vs. diet-only interventions for weight loss: a meta-analysis. *Obesity reviews*. 2009;10(3):313-23.
- 406 7. Flegal KM, Kit BK, Orpana H, Graubard BI. Association of all-cause mortality with overweight and
407 obesity using standard body mass index categories: a systematic review and meta-analysis. *Jama*.
408 2013;309(1):71-82.
- 409 8. Loveman E, Frampton G, Shepherd J, Picot J, Cooper K, Bryant J, et al. The clinical effectiveness
410 and cost-effectiveness of long-term weight management schemes for adults: a systematic review. NIHR
411 Health Technology Assessment programme: Executive Summaries. 2011.
- 412 9. Clifton PM, Bastiaans K, Keogh JB. High protein diets decrease total and abdominal fat and
413 improve CVD risk profile in overweight and obese men and women with elevated triacylglycerol.
414 *Nutrition, Metabolism and Cardiovascular Diseases*. 2009;19(8):548-54.
- 415 10. Dorner TE, Rieder A. Obesity paradox in elderly patients with cardiovascular diseases.
416 *International journal of cardiology*. 2012;155(1):56-65.
- 417 11. Kalyani RR, Corriere M, Ferrucci L. Age-related and disease-related muscle loss: the effect of
418 diabetes, obesity, and other diseases. *The lancet Diabetes & endocrinology*. 2014;2(10):819-29.
- 419 12. Garrow JS. *Obesity and related diseases*: Churchill Livingstone; 1988.
- 420 13. Prentice AM, Goldberg GR, Jebb SA, Black AE, Murgatroyd PR. Physiological responses to
421 slimming. *Proceedings of the Nutrition Society*. 1991;50(2):441-58.
- 422 14. Heymsfield S, Gonzalez M, Shen W, Redman L, Thomas D. Weight loss composition is one-fourth
423 fat-free mass: a critical review and critique of this widely cited rule. *Obesity reviews: an official journal*
424 *of the International Association for the Study of Obesity*. 2014;15(4):310-21.
- 425 15. Heymsfield SB, Ludwig DS, Wong JM, McCarthy C, Heo M, Shepherd J, et al. Are methods of
426 estimating fat-free mass loss with energy-restricted diets accurate? *European journal of clinical*
427 *nutrition*.
- 428 16. Shepherd JA, Ng BK, Sommer MJ, Heymsfield SB. Body composition by DXA. *Bone*.
429 2017;104:101-5.
- 430 17. Heymsfield SB, Bourgeois B, Ng BK, Sommer MJ, Li X, Shepherd JA. Digital anthropometry: a
431 critical review. *European journal of clinical nutrition*. 2018;72(5):680.
- 432 18. Bennett JP, Liu YE, Quon BK, Kelly NN, Wong MC, Kennedy SF, et al. Assessment of clinical
433 measures of total and regional body composition from a commercial 3-dimensional optical body
434 scanner. *Clinical Nutrition*. 2022;41(1):211-8.
- 435 19. Ng B, Hinton B, Fan B, Kanaya A, Shepherd J. Clinical anthropometrics and body composition
436 from 3D whole-body surface scans. *European journal of clinical nutrition*. 2016;70(11):1265.

- 437 20. Wong MC, Ng BK, Kennedy SF, Hwaung P, Liu EY, Kelly NN, et al. Children and Adolescents'
438 Anthropometrics Body Composition from 3-D Optical Surface Scans. *Obesity*. 2019;27(11):1738-49.
- 439 21. Ng BK, Sommer MJ, Wong MC, Pagano I, Nie Y, Fan B, et al. Detailed 3-dimensional body shape
440 features predict body composition, blood metabolites, and functional strength: the Shape Up! studies.
441 *The American journal of clinical nutrition*. 2019.
- 442 22. Tian IY, Ng BK, Wong MC, Kennedy S, Hwaung P, Kelly N, et al. Predicting 3D Body Shape and
443 Body Composition from Conventional 2D Photography. *Medical Physics*. 2020.
- 444 23. Tian IY WM, Kennedy S, Kelly NN, Yong EL, Garber AK, Heymsfield SB, Curless B, Shepherd JA. A
445 Device-Agnostic Shape Model for Automated Body Composition Estimates from 3D Optical Scans.
446 *Medical Physics*. 2022.
- 447 24. Wong MC, Ng BK, Tian I, Sobhiyeh S, Pagano I, Dechenaud M, et al. A pose-independent method
448 for accurate and precise body composition from 3D optical scans. *Obesity*. 2021;29(11):1835-47.
- 449 25. Tinsley GM, Harty PS, Stratton MT, Smith RW, Rodriguez C, Siedler MR. Tracking changes in body
450 composition: comparison of methods and influence of pre-assessment standardisation. *British Journal of*
451 *Nutrition*. 2021:1-19.
- 452 26. Lowe DA, Wu N, Rohdin-Bibby L, Moore AH, Kelly N, Liu YE, et al. Effects of time-restricted
453 eating on weight loss and other metabolic parameters in women and men with overweight and obesity:
454 the TREAT randomized clinical trial. *JAMA internal medicine*. 2020;180(11):1491-9.
- 455 27. Macronutrients and body fat accumulation: a mechanistic feeding study 2018. Available from:
456 <https://osf.io/m6v73/>.
- 457 28. Jansen LT, Yang N, Wong JM, Mehta T, Allison DB, Ludwig DS, et al. Prolonged Glycemic
458 Adaptation Following Transition From a Low-to High-Carbohydrate Diet: A Randomized Controlled
459 Feeding Trial. *Diabetes Care*. 2022;45(3):576-84.
- 460 29. Wong JM, Yu S, Ma C, Mehta T, Dickinson SL, Allison DB, et al. Stimulated insulin secretion
461 predicts changes in body composition following weight loss in adults with high BMI. *The Journal of*
462 *Nutrition*. 2022;152(3):655-62.
- 463 30. Scott MC. Exercise and Bone: Older Adults, Type II Diabetes, and Ketogenic Diets: Louisiana
464 State University and Agricultural & Mechanical College; 2021.
- 465 31. Varanoske AN, Harris MN, Hebert C, Howard EE, Johannsen NM, Heymsfield SB, et al. Effects of
466 testosterone undecanoate on performance during multi-stressor military operations: A trial protocol for
467 the Optimizing Performance for Soldiers II study. *Contemporary Clinical Trials Communications*.
468 2021;23:100819.
- 469 32. Varanoske AN, Harris MN, Hebert C, Howard EE, Johannsen NM, Heymsfield SB, et al.
470 Testosterone Undecanoate Administration Prevents Declines in Fat-Free Mass but not Physical
471 Performance During Simulated Multi-Stressor Military Operations. *Journal of Applied Physiology*. 2022.
- 472 33. Hangartner TN, Warner S, Braillon P, Jankowski L, Shepherd J. The Official Positions of the
473 International Society for Clinical Densitometry: acquisition of dual-energy X-ray absorptiometry body
474 composition and considerations regarding analysis and repeatability of measures. *Journal of Clinical*
475 *Densitometry*. 2013;16(4):520-36.
- 476 34. Ng BK, Liu YE, Wang W, Kelly TL, Wilson KE, Schoeller DA, et al. Validation of rapid 4-component
477 body composition assessment with the use of dual-energy X-ray absorptiometry and bioelectrical
478 impedance analysis. *The American journal of clinical nutrition*. 2018;108(4):708-15.
- 479 35. Loper M, Mahmood N, Romero J, Pons-Moll G, Black MJ. SMPL: A skinned multi-person linear
480 model. *ACM transactions on graphics (TOG)*. 2015;34(6):248.
- 481 36. Gallagher D, Visser M, De Meersman RE, Sepúlveda D, Baumgartner RN, Pierson RN, et al.
482 Appendicular skeletal muscle mass: effects of age, gender, and ethnicity. *Journal of applied physiology*.
483 1997;83(1):229-39.

- 484 37. Shepherd JA, Lu Y. A generalized least significant change for individuals measured on different
485 DXA systems. *Journal of Clinical Densitometry*. 2007;10(3):249-58.
- 486 38. Cohen J. A coefficient of agreement for nominal scales. *Educational and psychological*
487 *measurement*. 1960;20(1):37-46.
- 488 39. Taylor J. Introduction to error analysis, the study of uncertainties in physical
489 measurements 1997.
- 490 40. Maskarinec G, Shvetsov YB, Wong MC, Garber A, Monroe K, Ernst TM, et al. Subcutaneous and
491 visceral fat assessment by DXA and MRI in older adults and children. *Obesity*. 2022.
- 492 41. Neeland IJ, Grundy SM, Li X, Adams-Huet B, Vega GL. Comparison of visceral fat mass
493 measurement by dual-X-ray absorptiometry and magnetic resonance imaging in a multiethnic cohort:
494 the Dallas Heart Study. *Nutrition & Diabetes*. 2016;6(7):e221-e. doi: 10.1038/nutd.2016.28.
- 495 42. Taylor JL, Holland DJ, Coombes JS, Keating SE. Accuracy of dual-energy x-ray absorptiometry for
496 assessing longitudinal change in visceral adipose tissue in patients with coronary artery disease.
497 *International Journal of Obesity*. 2021;45(8):1740-50.
- 498 43. Dias KA, Ramos JS, Wallen MP, Davies PS, Cain PA, Leong GM, et al. Accuracy of longitudinal
499 assessment of visceral adipose tissue by dual-energy X-ray absorptiometry in children with obesity.
500 *Journal of Obesity*. 2019;2019.
- 501 44. Wu AJ, Rifas-Shiman SL, Taveras EM, Oken E, Hivert MF. Associations of DXA-measured
502 abdominal adiposity with cardio-metabolic risk and related markers in early adolescence in Project Viva.
503 *Pediatric obesity*. 2021;16(2):e12704.
- 504 45. Schousboe JT, Langsetmo L, Schwartz AV, Taylor BC, Vo TN, Kats AM, et al. Comparison of
505 Associations of DXA and CT Visceral Adipose Tissue Measures With Insulin Resistance, Lipid Levels, and
506 Inflammatory Markers. *Journal of clinical densitometry: the official journal of the International Society*
507 *for Clinical Densitometry*. 2017;20(2):256-64.
- 508 46. Cornell J, Berger R. Factors that influence the value of the coefficient of determination in simple
509 linear and nonlinear regression models. *Phytopathology*. 1987;77(1):63-70.

Figure Legends

Figure 1. Histograms of the time (weeks) between the baseline and follow-up DXA scans for females (top) and males (bottom). The numbers over the bars represent the count. Multiple numbers over a multi-color bar represent the count in the corresponding order.

Figure 2. Average baseline (left) and follow-up (right) body shapes for females (top) and males (bottom) by study.

Figure 3. (Female) Scatter plot comparisons between 3DO and DXA body composition at baseline, follow-up, and adjusted change. Blue, horizontal lines and orange, vertical lines are signifying the amount of change needed to pass the least significant change for DXA and 3DO, respectively. Purple (Zone 4) and green glyphs (Zone 1) represent agreement of significant or nonsignificant change by 3DO and DXA. The orange glyphs (Zone 3) represent significant change detected by 3DO but not DXA and vice versa for the blue glyphs (Zone 2). The zones and glyphs are consistent for the proceeding adjusted change plots.

Figure 4. (Male) Scatter plot comparisons between 3DO and DXA body composition at baseline, follow-up, and adjusted change. Blue, horizontal lines and orange, vertical lines are signifying the amount of change needed to pass the least significant change for DXA and 3DO, respectively. Purple (Zone 4) and green glyphs (Zone 1) represent agreement of significant or nonsignificant change by 3DO and DXA. The orange glyphs (Zone 3) represent significant change detected by 3DO but not DXA and vice versa for the blue glyphs (Zone 2). The zones and glyphs are consistent for the proceeding adjusted change plots.

Figure 5. Histograms to show the frequency of percent fat-free mass change relative to total weight change in the female (top) and male (bottom) samples.

Table 1. Descriptions of longitudinal studies.

Study Name (Acronym)	Sex (N)	Intervention	Clinical Trials Number
Athletes	Female 5	Basketball team during training	NA
Bariatric	Female 2	Bariatric surgery	NA
Macronutrients and Body Fat Accumulation: A Mechanistic Feeding Study (FB4: Framingham, Boston, Bloomington, Birmingham, and Baylor)	Female 15; Male 40	Hypocaloric diet with low carbohydrate intake	NCT03394664
Resistance Exercise and Low-Intensity Physical Activity Breaks in Sedentary Time to Improve Muscle and Cardiometabolic Health (REALPA)	Female 8; Male 5	16 weeks of resistance exercise with or without low intensity physical activity breaks in sedentary time or moderate intensity aerobic exercise in older adults	NCT03771417
Time Restricted Eating on Weight Loss (TREAT)	Female 15; Male 27	Time-restricted diet (16 hour fast, 8 hours to feed)	NCT03393195
Trial of Testosterone Undecanoate for Optimizing Performance During Military Operations (OPS II)	Male 16	Simulated military operational stress	NCT04120363

Complete study protocols can be found on <https://clinicaltrials.gov/>.

Table 2. Sample characteristics at baseline and follow-up visits

	Female (N=45)		Male (N=88)	
	Baseline	Follow-up	Baseline	Follow-up
Age (years)				
Mean (SD)	45.6 (15.7)	45.8 (15.7)	36.3 (13.3)	36.5 (13.3)
Median [Min, Max]	45.9 [18.5, 76.7]	46.2 [18.8, 77.1]	32.0 [19.0, 73.5]	32.0 [19.3, 73.9]
Ethnicity				
Asian	3 (6.7%)		11 (12.4%)	
Black	10 (22.2%)		7 (7.9%)	
Hispanic	5 (11.1%)		7 (7.9%)	
White	27 (60.0%)		64 (71.9%)	
Height (cm)				
Mean (SD)	165 (7.46)		177 (6.67)	
Median [Min, Max]	164 [149, 185]		176 [162, 194]	
Weight (kg)				
Mean (SD)	86.7 (17.5)	81.4 (13.2)	97.6 (17.1)	89.4 (17.3)
Median [Min, Max]	82.3 [63.5, 143]	80.6 [63.0, 116]	95.2 [61.5, 142]	87.3 [58.9, 175]
BMI (kg/m²)				
Mean (SD)	31.8 (6.61)	29.9 (5.00)	31.3 (5.42)	28.7 (5.54)
Median [Min, Max]	30.4 [21.8, 51.3]	29.5 [21.7, 43.7]	30.1 [19.8, 45.8]	27.9 [19.0, 57.3]
DXA Total FM (kg)				
Mean (SD)	34.8 (11.4)	31.3 (8.80)	28.6 (10.8)	23.2 (9.34)
Median [Min, Max]	34.6 [12.9, 68.4]	31.0 [14.0, 54.3]	28.6 [9.14, 67.3]	22.6 [6.00, 53.8]
DXA Total FFM (kg)				
Mean (SD)	52.3 (8.35)	50.5 (7.03)	69.6 (9.24)	66.2 (8.37)
Median [Min, Max]	52.4 [37.5, 77.0]	49.7 [35.9, 67.8]	68.0 [51.4, 91.6]	65.4 [49.6, 90.9]
DXA Percent Fat (%)				
Mean (SD)	39.3 (7.21)	37.8 (6.66)	28.3 (6.96)	25.2 (7.07)
Median [Min, Max]	41.8 [19.9, 51.5]	40.2 [20.6, 46.9]	29.1 [11.1, 47.4]	25.9 [7.81, 43.9]
DXA ALM (kg)				
Mean (SD)	11.3 (2.17)	11.0 (1.87)	15.8 (2.23)	15.1 (2.04)
Median [Min, Max]	11.6 [7.31, 15.6]	10.7 [7.52, 15.0]	15.8 [11.2, 20.9]	14.8 [10.8, 20.9]
DXA VAT (kg)				
Mean (SD)	0.56 (0.27)	0.53 (0.26)	0.56 (0.28)	0.46 (0.23)
Median [Min, Max]	0.51 [0.13, 1.10]	0.52 [0.12, 1.29]	0.50 [0.16, 1.40]	0.40 [0.13, 1.15]
DXA Arm FM (kg)				
Mean (SD)	2.14 (0.83)	1.92 (0.60)	1.78 (0.84)	1.46 (0.67)
Median [Min, Max]	2.04 [0.73, 5.32]	1.94 [0.79, 3.53]	1.61 [0.52, 4.94]	1.39 [0.36, 3.82]
DXA Arm FFM (kg)				
Mean (SD)	2.68 (0.51)	2.59 (0.44)	4.34 (0.73)	4.13 (0.66)
Median [Min, Max]	2.62 [1.79, 4.01]	2.49 [1.80, 3.74]	4.30 [2.93, 6.42]	4.10 [2.78, 6.19]
DXA Leg FM (kg)				
Mean (SD)	6.54 (2.18)	5.97 (1.75)	4.67 (1.70)	3.86 (1.42)

Median [Min, Max]	6.02 [2.90, 13.1]	5.89 [3.13, 10.8]	4.45 [1.65, 10.9]	3.88 [1.08, 9.16]
DXA Leg FFM (kg)				
Mean (SD)	8.64 (1.72)	8.37 (1.48)	11.4 (1.61)	11.0 (1.48)
Median [Min, Max]	8.75 [5.52, 12.0]	8.24 [5.72, 11.3]	11.3 [7.94, 15.1]	10.8 [7.62, 15.1]
DXA Trunk FM (kg)				
Mean (SD)	16.3 (6.16)	14.4 (4.79)	14.4 (6.10)	11.3 (5.48)
Median [Min, Max]	16.6 [4.61, 34.9]	14.8 [4.73, 24.5]	14.1 [3.52, 34.3]	10.8 [2.16, 26.6]
DXA Trunk FFM (kg)				
Mean (SD)	26.1 (4.31)	25.0 (3.81)	33.9 (4.88)	31.9 (4.44)
Median [Min, Max]	25.7 [19.2, 41.6]	23.9 [17.5, 37.4]	33.5 [25.1, 46.4]	31.7 [23.8, 44.3]

Abbreviations: FM (fat mass); FFM (fat-free mass); ALM (appendicular lean mass); DXA (dual-energy X-ray absorptiometry); VAT (visceral adipose tissue)









