

1     **A decision aid for policymakers to estimate the impact of e-cigarette flavour restrictions**  
2             **on population smoking and e-cigarette use prevalence among youth versus smoking**  
3                             **prevalence among adults**

4     Mark J Gibson<sup>1,2</sup>, Marcus R Munafò<sup>1,2,3</sup>, Angela S. Attwood<sup>1,2</sup>, Martin J. Dockrell<sup>4</sup>, Michelle  
5                             A. Havill<sup>4</sup>, Jasmine N Khouja<sup>1,2</sup>

6

7     \* Corresponding author: [mark.gibson@bristol.ac.uk](mailto:mark.gibson@bristol.ac.uk)

8

9     <sup>1</sup> MRC Integrative Epidemiology Unit (IEU) at the University of Bristol, Bristol, United  
10     Kingdom

11     <sup>2</sup> School of Psychological Science, University of Bristol, Bristol, United Kingdom

12     <sup>3</sup> Bristol Biomedical Research Centre (BRC), National Institute for Health and Care Research  
13     (NIHR), Bristol, United Kingdom

14     <sup>4</sup> Office of Health Improvement and Disparities, Department of Health and Social Care, UK  
15     SW1H 0EU

16     **Keywords:** E-cigarettes, Flavoured e-liquids, Nicotine, Tobacco, Youth Smoking, Policy,  
17     Public Health

18

19

20

21

1

2

## Abstract

3 **Background** Policy decisions should be evidence-based, but the magnitude of intended and  
4 unintended impacts cannot always be easily estimated from the available data. For example,  
5 banning flavours in electronic cigarettes (e-cigarettes) to reduce appeal to non-smoking  
6 young people could have the intended impact by reducing youth vaping but could have  
7 negative consequences for adult smokers and vapers.

8 **Methods** We developed a decision aid to help policymakers make informed decisions on the  
9 potential net impact of a ban on e-cigarette flavours. We estimated the number of non-  
10 smoking youth who would be deterred from ever vaping and subsequently ever smoking, and  
11 the number of smokers and ex-smokers who would be deterred from quitting or encouraged  
12 to relapse, to determine whether the benefits to youth outweigh the costs to existing smokers  
13 and vapers. This aid then outputs a report with the results graphically depicted to aid  
14 interpretability.

15 **Results** We demonstrated the value of this decision aid using data from various sources to  
16 estimate the impact of a flavour ban in three populations: the general UK population, low-  
17 socioeconomic position UK population, and the general US population. All three examples  
18 suggested a negative net population impact of a ban. These reports were then presented to the  
19 all-party parliamentary group for vaping.

20 **Conclusions** We demonstrate how decision aids can be used to help policymakers arrive at  
21 evidence-based decisions efficiently and can be used to quickly obtain up-to-date estimates as  
22 new data becomes available.

23

1

2 Policy decisions should be evidence-based and lead to positive, beneficial impacts in the  
3 affected population. However, sourcing relevant evidence that can be easily interpreted can  
4 be a difficult task for policymakers working under time constraints. Creating decision aids for  
5 policymakers that can quickly provide brief, digestible guidance can be particularly useful in  
6 areas where existing evidence suggests the proposed policy change may have positive and  
7 negative implications on the target population.

8 One example is electronic cigarette (e-cigarette) policy. Some jurisdictions have banned  
9 flavours in e-cigarettes to reduce appeal to non-smoking young people and the UK could do  
10 the same; this could have the intended impact by reducing youth vaping but could have  
11 negative consequences for adult smokers and e-cigarette users (vapers). Although e-cigarettes  
12 are considered to be less harmful than cigarettes [1], and can be used by smokers to help them  
13 quit [2], there have been concerns that the wide range of available flavours encourage non-  
14 smoking youth to vape and subsequently smoke. While there is some evidence to suggest that  
15 flavours encourage youth vaping in both the US and the UK, there is no clear evidence that  
16 they encourage subsequent smoking [3-6]. The emergence of disposable vapes, which are  
17 most popular and relatively accessible among young people in both the US and UK, has  
18 further fuelled concerns about flavours in e-cigarette products [7, 8]. These concerns have led  
19 to bans of e-cigarette flavours (i.e., all but unflavoured, tobacco and menthol) in several  
20 jurisdictions. Evidence of the actual and predicted effect of bans is conflicting with some  
21 studies suggesting a reduction of vaping rates [9, 10] and others suggesting no reduction [11]  
22 or an increase in smoking rates in both youth [12] and adults [10, 13].

23 Contrasting evidence on the effectiveness of a potential ban makes it difficult for  
24 policymakers to reach an informed decision. Therefore, to help policymakers make informed

1 decisions on a potential e-cigarette flavour ban, we aimed to develop a decision aid for  
2 policymakers to specifically estimate the impact of a ban in any given population, and to  
3 illustrate the potential value of such decision aids in general.

4

### 5 **Decision Aid for Policymakers**

6 We consulted with policymakers and researchers to create a decision aid to estimate the  
7 potential net impact of an e-cigarette flavour ban, for which population sizes and the  
8 proportions are used as inputs to estimate four numbers (See Figure 1 for calculations):

- 9 A. The number of non-smoking youth who could potentially be at risk of initiating  
10 vaping because of flavours.
- 11 B. The number of youth who could potentially be at higher risk of ever smoking because  
12 of the availability of flavoured e-liquids.
- 13 C. The number of potential lost additional quit attempts per year.
- 14 D. The number of ex-smokers who may relapse to smoking.

15 In this context, “youth” refers to those in the age range considered to be at risk of smoking  
16 (i.e.,11-17). From these four numbers the decision aid then calculates whether the number of  
17 non-smoking e-cigarette users introduced into the UK population because of flavoured e-  
18 liquid availability could outweigh the number of smokers and ex-smokers who might vape  
19 instead of smoke because of flavoured e-liquid availability ( $A - [C + D]$ ) and whether the  
20 number of ever smokers introduced into the UK population because of flavoured e-liquid  
21 availability could outweigh the number of smokers and ex-smokers who might vape instead  
22 of smoke because of flavoured e-liquid availability ( $B - [C + D]$ ). Positive values provide  
23 some support for a total ban on flavoured e-liquids, and negative values provide some

1 evidence that the negative consequences of a total ban would outweigh the potential benefits  
2 (i.e., a negative net impact of a ban).

### 3 **Applied Examples**

#### 4 **Methods**

5 We entered data from a variety of sources into our decision aid to estimate the effects of a  
6 smoking ban in the general UK population, the low-socioeconomic position UK population  
7 (specifically those from the C2, D and E categories of the National Readership Survey scale  
8 [14], which are those classified as skilled working class, working class or non-working) and  
9 the general US population. We used data from Action on Smoking and Health (ASH) [15],  
10 the Smoking Toolkit Study (STS) [16], the National Youth Tobacco Survey (NYST) [17] and  
11 publicly available figures from the Office of National Statistics (ONS) [18, 19], the  
12 Department for Work and Pensions [20], the Centres for Disease Control and Prevention  
13 (CDC) [21-23], census data [24-27], privately owned real-time statistics website  
14 Worldometer (estimates taken on 12/09/2022) [28, 29] and peer-reviewed academic research  
15 articles [30-32].

16 All population numbers or proportions and descriptions of data sources for these examples  
17 can be found in Supplementary Tables S1-3 and Supplementary Materials. Analysis code can  
18 be found at [https://github.com/MRCIEU/Estimation-of-the-impact-of-flavour-restrictions-on-](https://github.com/MRCIEU/Estimation-of-the-impact-of-flavour-restrictions-on-population-smoking-and-e-cigarette-use)  
19 [population-smoking-and-e-cigarette-use](https://github.com/MRCIEU/Estimation-of-the-impact-of-flavour-restrictions-on-population-smoking-and-e-cigarette-use).

20

#### 21 **Output**

22 For the UK general population, we calculated that 53,609 non-smoking youth were at risk of  
23 ever vaping due to flavours and 26,269 of those were at risk of ever smoking due to flavours.  
24 33,000 potential quit attempts would be lost a year and 295,403 ex-smokers would relapse to  
25 smoking.

1 For the UK low-socioeconomic position population, we calculated that 30,484 non-smoking  
2 youth were at risk of ever vaping due to flavours and 13,109 of those were at risk of ever  
3 smoking due to flavours. 19,096 potential quit attempts would be lost a year and 171,299 ex-  
4 smokers would relapse to smoking.

5 For the US general population, we calculated that 355,617 non-smoking youth were at risk of  
6 ever vaping due to flavours and 78,236 of those were at risk of ever smoking due to flavours.  
7 172,481 potential quit attempts would be lost a year and 1,369,341 ex-smokers would relapse  
8 to smoking.

9 For all three populations, the decision aid estimated that the number of adult smokers and ex-  
10 smokers who choose not to quit or relapse would outweigh the number of youth vapers who  
11 would be deterred from vaping or smoking if a ban was introduced.

12 The final report produced by the decision aid for the general UK population (including the  
13 resulting guidance for policymakers) is shown in Supplementary Material.

14  
15

## Discussion

16 We developed a decision aid to assist policymakers in decisions relating to a potential ban of  
17 e-cigarette flavours. We applied this decision aid to three different populations (with the  
18 output suggesting the ban would have a negative impact in the general UK, low-  
19 socioeconomic UK and general US populations). These reports were then presented to the  
20 UK All Party Parliamentary Group for Vaping  
21 (<https://publications.parliament.uk/pa/cm/cmallparty/210602/vaping.htm>), with the  
22 opportunity for clarification from, and feedback to, the researchers. This project demonstrates  
23 both the feasibility and utility of developing decision aids to increase the impact of science  
24 and facilitate the making of evidence-based decisions by policy makers.

1 While decision aids are useful, the output should always be used in combination with  
2 evidence from a range of studies and the interpretability of the output is dependent on the  
3 quality of the decision aid and the evidence supplied to it. There are some limitations of the  
4 decision aid demonstrated in this article. Firstly, much of the evidence is reliant on self-  
5 reported beliefs about how smokers and vapers think they will behave in the event of flavour  
6 restrictions being implemented, which may differ from their actual behaviour. Secondly,  
7 while there is a strong association between e-cigarette use and later smoking, there is no clear  
8 evidence that e-cigarette use *causes* smoking among youth. There is also evidence to suggest  
9 that at least part of this relationship could be explained by shared risk factors [33], so it is  
10 likely that this decision aid will overestimate the number of youth who are at risk of smoking  
11 due to e-liquid flavour availability. Furthermore, the decision aid does not account for  
12 displacement (i.e., the number of youths who do not smoke because e-cigarettes are  
13 available), meaning the benefit of a ban is likely overestimated [34, 35]. The decision aid also  
14 does not take into account negative consequences of a ban beyond smoking behaviour, such  
15 as the creation of a black market which could lead to the use of unregulated/homemade  
16 products [36] and death [37, 38]. It can also not predict the consequences of partial bans such  
17 as allowing flavoured e-liquid on prescription, banning only flavours which appeal to  
18 children or banning packaging which is appealing to children.

## 19 **Conclusions**

20 The development of decision aids for policymakers can increase the impact of research and  
21 facilitate the making of evidence-based decisions by policymakers and the decision aid  
22 outlined in this article can inform on the net impact of an e-cigarette flavours ban. Using the  
23 currently available data, the decision aid demonstrated here suggests that a flavour ban would  
24 have a negative impact on the UK general, UK low socioeconomic and the US general

1 populations. However, the output from the decision aid needs to be regularly updated to  
2 ensure it accurately represents the ever-changing sociocultural landscape. Future iterations  
3 should also investigate the effect of a potential ban on other at-risk populations (such as those  
4 who suffer from mental illness) and develop the decision aid to account for the effects of  
5 different e-cigarette flavours (i.e., sweet versus not), and the effect of disposables, as the  
6 relevant data becomes available.

## 7 **References**

- 8 1. McNeill A, Brose LS, Calder R, Hitchman SC, Hajek P, McRobbie H. E-cigarettes:  
9 an evidence update a report commissioned by Public Health England. London: UK, PHE;  
10 2015.
- 11 2. Hartmann-Boyce J, Begh R, Aveyard P. Electronic cigarettes for smoking cessation.  
12 *Bmj-Brit Med J*. 2018;360.
- 13 3. Pepper JK, Ribisl KM, Brewer NT. Adolescents' interest in trying flavoured e-  
14 cigarettes. *Tob Control*. 2016;25:ii62-ii6.
- 15 4. Notley C, Gentry S, Cox S, Dockrell M, Havill M, Attwood AS, et al. Youth use of e-  
16 liquid flavours-a systematic review exploring patterns of use of e-liquid flavours and  
17 associations with continued vaping, tobacco smoking uptake or cessation. *Addiction*.  
18 2022;117(5):1258-72.
- 19 5. Chan GCK, Stjepanovic D, Lim C, Sun TZ, Anandan AS, Connor JP, et al. Gateway  
20 or common liability? A systematic review and meta-analysis of studies of adolescent e-  
21 cigarette use and future smoking initiation. *Addiction*. 2021;116(4):743-56.
- 22 6. Khouja JN, Suddell SF, Peters SE, Taylor AE, Munafo MR. Is e-cigarette use in non-  
23 smoking young adults associated with later smoking? A systematic review and meta-analysis.  
24 *Tob Control*. 2021;30(1):8-15.



- 1 7. Park-Lee E, Ren CF, Sawdey MD, Gentzke AS, Cornelius M, Jamal A, et al. E-  
2 Cigarette Use Among Middle and High School Students - National Youth Tobacco Survey,  
3 United States, 2021. *Mmwr-Morbid Mortal W.* 2021;70(39):1387-9.
- 4 8. Tattan-Birch H, Jackson SE, Kock L, Dockrell M, Brown J. Rapid growth in  
5 disposable e-cigarette vaping among young adults in Great Britain from 2021 to 2022: a  
6 repeat cross-sectional survey. *Addiction.* 2022;Accepted Articles.
- 7 9. Rogers T, Brown EM, Siegel-Reamer L, Rahman B, Feld AL, Patel M, et al. A  
8 Comprehensive Qualitative Review of Studies Evaluating the Impact of Local US Laws  
9 Restricting the Sale of Flavored and Menthol Tobacco Products. *Nicotine & Tobacco*  
10 *Research.* 2022;24(4):433-43.
- 11 10. Yang Y, Lindblom EN, Salloum RG, Ward KD. The impact of a comprehensive  
12 tobacco product flavor ban in San Francisco among young adults. *Addictive Behaviors*  
13 *Reports.* 2020;11:100333.
- 14 11. Romm KF, Henriksen L, Huang JD, Le D, Clausen M, Duan ZS, et al. Impact of  
15 existing and potential e-cigarette flavor restrictions on e-cigarette use among young adult e-  
16 cigarette users in 6 US metropolitan areas. *Preventive Medicine Reports.* 2022;28.
- 17 12. Friedman AS. A Difference-in-Differences Analysis of Youth Smoking and a Ban on  
18 Sales of Flavored Tobacco Products in San Francisco, California. *Jama Pediatr.*  
19 2021;175(8):863-5.
- 20 13. Khouja J, Dyer ML, Havill MA, Dockrell MJ, Munafo M, Attwood AS. Exploring the  
21 opinions and potential impact of unflavoured e-liquid on smoking cessation among UK  
22 smokers and smoking relapse among UK e-cigarette users: Findings from a qualitative study.  
23 *Research Square.* 2022.
- 24 14. National Readership Survey scale. Social Grade [Available from:  
25 <https://www.nrs.co.uk/nrs-print/lifestyle-and-classification-data/social-grade/>.

- 1 15. Sandford A. ASH. Trends in Urology & Men's Health. 2012;3(2).
- 2 16. Fidler JA, Shahab L, West O, Jarvis MJ, McEwen A, Stapleton JA, et al. 'The  
3 smoking toolkit study': a national study of smoking and smoking cessation in England. BMC  
4 Public Health. 2011;11.
- 5 17. Centers for Disease Control and Prevention. National Youth Tobacco Survey (NYTS)  
6 [updated 14/03/2022. Available from:  
7 [https://www.cdc.gov/tobacco/data\\_statistics/surveys/nyts/index.htm](https://www.cdc.gov/tobacco/data_statistics/surveys/nyts/index.htm).
- 8 18. Office of National Statistics. Smoking prevalence in the UK and the impact of data  
9 collection changes: 2020 2021 [updated 07/12/2021. Available from:  
10 [https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/drugusealcohol  
11 andsmoking/bulletins/smokingprevalenceintheukandtheimpactofdatacollectionchanges/2020#](https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/drugusealcoholandsmoking/bulletins/smokingprevalenceintheukandtheimpactofdatacollectionchanges/2020#)  
12 .
- 13 19. Office of National Statistics. Smoking habits in the UK and its constituent countries  
14 2021 [updated 07/12/2021. Available from:  
15 [https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/healthandlifeex  
16 pectancies/datasets/smokinghabitsintheukanditsconstituentcountries](https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/healthandlifeexpectancies/datasets/smokinghabitsintheukanditsconstituentcountries).
- 17 20. Department for Work and Pensions. Households below average income: an analysis  
18 of the income distribution FYE 1995 to FYE 2020 [updated 25/03/2021. Available from:  
19 [https://www.gov.uk/government/statistics/households-below-average-income-for-financial-  
20 years-ending-1995-to-2020/households-below-average-income-an-analysis-of-the-income-  
21 distribution-fye-1995-to-fye-2020](https://www.gov.uk/government/statistics/households-below-average-income-for-financial-years-ending-1995-to-2020/households-below-average-income-an-analysis-of-the-income-distribution-fye-1995-to-fye-2020).
- 22 21. Centers for Disease Control and Prevention. Current Cigarette Smoking Among  
23 Adults in the United States [updated 17/03/2022. Available from:  
24 [https://www.cdc.gov/tobacco/data\\_statistics/fact\\_sheets/adult\\_data/cig\\_smoking/index.htm#](https://www.cdc.gov/tobacco/data_statistics/fact_sheets/adult_data/cig_smoking/index.htm#).

- 1 22. Kramarow EA. Health of Former Cigarette Smokers Aged 65 and Over: United  
2 States, 2018. Centers for Disease Control and Prevention; 2020.
- 3 23. Centers for Disease Control and Prevention. Electronic Cigarette Use Among U.S.  
4 Adults, 2018 [updated 30/04/2020. Available from:  
5 <https://www.cdc.gov/nchs/products/databriefs/db365.htm#>.
- 6 24. United States Census Bureau. Quick Facts [updated 01/07/2021. Available from:  
7 <https://www.census.gov/quickfacts/fact/table/US/PST045221>.
- 8 25. Childstats.gov. POP1 CHILD POPULATION: NUMBER OF CHILDREN (IN  
9 MILLIONS) AGES 0–17 IN THE UNITED STATES BY AGE, 1950–2020 AND  
10 PROJECTED 2021–2050 [Available from:  
11 <https://www.childstats.gov/americaschildren/tables/pop1.asp>.
- 12 26. Statista. Population of the United Kingdom from 1871 to 2020 [updated  
13 05/09/2022. Available from: <https://www.statista.com/statistics/281296/uk-population/>.
- 14 27. Statista. Population of the United Kingdom in 2020, by age [updated 05/09/2022.  
15 Available from: <https://www.statista.com/statistics/281174/uk-population-by-age/>.
- 16 28. Worldometer. United States Population [updated 12/09/2022. Available from:  
17 <https://www.worldometers.info/world-population/us-population/>.
- 18 29. Worldometer. U.K. Population [updated 12/09/2022. Available from:  
19 <https://www.worldometers.info/world-population/uk-population/>.
- 20 30. Berry KM, Fetterman JL, Benjamin EJ, Bhatnagar A, Barrington-Trimis JL,  
21 Leventhal AM, et al. Association of Electronic Cigarette Use With Subsequent Initiation of  
22 Tobacco Cigarettes in US Youths. *Jama Network Open*. 2019;2(2).
- 23 31. Chen RF, Pierce JP, Leas EC, Benmarhnia T, Strong DR, White MM, et al.  
24 Effectiveness of e-cigarettes as aids for smoking cessation: evidence from the PATH Study  
25 cohort, 2017-2019. *Tob Control*. 2022.

- 1 32. Gravelly S, Smith DM, Liber AC, Cummings KM, East KA, Hammond D, et al.  
2 Responses to potential nicotine vaping product flavor restrictions among regular vapers using  
3 non-tobacco flavors: Findings from the 2020 ITC Smoking and Vaping Survey in Canada,  
4 England and the United States. *Addict Behav.* 2022;125.
- 5 33. Khouja JN, Wootton RE, Taylor AE, Smith GD, Munafò MR. Association of genetic  
6 liability to smoking initiation with e-cigarette use in young adults: A cohort study. *Plos*  
7 *Medicine.* 2021;18(3).
- 8 34. Sokol NA, Feldman JM. High School Seniors Who Used E-Cigarettes May Have  
9 Otherwise Been Cigarette Smokers: Evidence From Monitoring the Future (United States,  
10 2009-2018). *Nicotine Tob Res.* 2021;23(11):1958-61.
- 11 35. Hallingberg B, Maynard OM, Bauld L, Brown R, Gray L, Lowthian E, et al. Have e-  
12 cigarettes renormalised or displaced youth smoking? Results of a segmented regression  
13 analysis of repeated cross sectional survey data in England, Scotland and Wales. *Tob Control.*  
14 2020;29(2):207-16.
- 15 36. Action on Smoking Health. Use of e-cigarettes (vapes) among adults in Great Britain.  
16 2020.
- 17 37. Werner AK, Koumans EH, Chatham-Stephens K, Salvatore PP, Armatas C, Byers P,  
18 et al. Hospitalizations and Deaths Associated with EVALI. *New Engl J Med.*  
19 2020;382(17):1589-98.
- 20 38. Ellington S, Salvatore PP, Ko J, Danielson M, Kim L, Cyrus A, et al. Update:  
21 Product, Substance-Use, and Demographic Characteristics of Hospitalized Patients in a  
22 Nationwide Outbreak of E-cigarette, or Vaping, Product Use-Associated Lung Injury -  
23 United States, August 2019-January 2020. *Mmwr-Morbid Mortal W.* 2020;69(2):44-9.

24

25

## Declarations

1 **Ethical approval and consent to participate:** This study only used secondary data. All  
2 methods were carried out by the original studies in accordance with relevant guidelines and  
3 regulations and informed consent was obtained from all subjects and/or their legal  
4 guardian(s). Ethical approval for the Smoking Toolkit Study was granted by the UCL ethics  
5 committee (ID 0498/001). All other data is publicly available or available upon request.

6 **Consent for publication:** NA

7 **Availability of data and materials:** The Smoking Toolkit Study data can be requested at  
8 <https://smokinginengland.info/resources/sts-documents>. Action on Smoking and Health can  
9 be contacted for data requests at <https://ash.org.uk/contact-us>. The National Youth Tobacco  
10 Survey data can be downloaded at  
11 [https://www.cdc.gov/tobacco/data\\_statistics/surveys/nyts/data/index.html](https://www.cdc.gov/tobacco/data_statistics/surveys/nyts/data/index.html)

12 **Competing interests:** All authors have completed the ICMJE uniform disclosure form at  
13 <http://www.icmje.org/disclosure-of-interest/>. The authors declare no support from any  
14 organisation for the submitted work; no financial relationships with any organisations that  
15 might have an interest in the submitted work in the previous three years; no other  
16 relationships or activities that could appear to have influenced the submitted work.

17 **Funding:** This work was originally supported by Public Health England (PHE) via an  
18 honorary contract awarded to ASA. There is no grant number for this research as it was  
19 commissioned by Public Health England via the honorary academic framework. Further  
20 support was received from the University of Bristol via an Economic and Social Research  
21 Council Impact Acceleration Award (A100111) awarded to JNK, ASA and MRM. MJG and  
22 MRM are supported by the Medical Research Council Integrative Epidemiology Unit  
23 (MC\_UU\_00011/7). JNK is supported by a Cancer Research UK programme grant (the  
24 Integrative Cancer Epidemiology Programme C18281/A29019). The funders had no role in

1 the study design, collection or analysis of data, or interpretation of results. The views  
2 expressed in this paper are those of the authors and not necessarily any funder or  
3 acknowledged person/institution.

4 **Author contributions:** The policy decision aid was designed by MRM, ASA, MJD, MAH  
5 and JNK who also provided feedback on this manuscript; The applied examples were  
6 conducted and the manuscript was written by MJG under the supervision of JNK.

7 **Acknowledgments:** We would like to thank the Smoking Toolkit Study and Smoking on  
8 Action Health for providing data.

9

# A

To what extent do flavours draw in non-smoking youth?

*a*

Number of youth at smoking age

*b*

Proportion of those who are non-smokers

*c*

Proportion of those non-smokers who have vaped

*d*

Proportion of those who state they vape because of flavours

Number of non-smoking youth who could potentially be at risk of initiating vaping because of flavours =  $a \times b \times c \times d$

**A =**

# B

How many non-smokers who vape subsequently smoke?

*a*

Number of non-smoking youth who may be at risk of vaping because of flavours

*b*

Proportion of non-smoking youth who try vaping and later try smoking

Number of youth who could potentially be at higher risk of ever smoking because of the availability of flavoured e-liquids =  $a \times b$

**B =**

# C

To what extent do flavours draw in adult smokers who would quit smoking using e-cigarettes as a resource?

*a*

Number of smokers

*b*

Proportion of smokers who quit smoking because of e-cigarettes

*c*

Proportion of smokers who state they would not quit/smoke more if flavours were not available

Number of potential lost additional quit attempts per year =  $a \times b \times c$

**C =**

# D

How many current adult vapers would relapse to smoking if flavours were not available?

*a*

Number of ex-smokers

*b*

Proportion of ex-smokers who vape

*c*

Proportion of ex-smokers who state they would relapse to smoking if flavours weren't available

Number of ex-smokers who may relapse to smoking =  $a \times b \times c$

**D =**