1 2 3	<i>Tcte1</i> knockout influence on energy chain transportation, apoptosis and spermatogenesis – implications for male infertility								
4	Running title: <i>Tcte1</i> gene in male infertility – mouse 'knockout' model								
6 7 8 9	Marta Olszewska ¹ , Agnieszka Malcher ¹ , Tomasz Stokowy ² , Nijole Pollock ³ , Andrea J. Berman ⁴ , Sylwia Budkiewicz ¹ , Marzena Kamieniczna ¹ , Hanna Jackowiak ⁵ , Joanna Suszynska-Zajczyk ⁶ , Piotr Jedrzejczak ⁷ , Alexander N. Yatsenko ³ , Maciej Kurpisz ^{1*.}								
10 11 12 13 14 15 16 17 18	 ¹ Institute of Human Genetics, Polish Academy of Sciences, Poznan, Poland ² Scientific Computing Group, IT Division, University of Bergen, Norway ³ Department of OB/GYN and Reproductive Sciences, School of Medicine, University of Pittsburgh, PA, USA ⁴ Department of Biological Sciences, University of Pittsburgh, PA, USA ⁵ Department of Histology and Embryology, Poznan University of Life Sciences, 60-625 Poznan, Poland ⁶ University of Life Sciences, Poznan, Poland ⁷ Division of Infertility and Reproductive Endocrinology, Department of Gynecology, Obstetrics and Gynecological Oncology, Poznan University of Medical Sciences, 60-535 Poznan, Poland 								
19	* Corresponding author:								
20 21 22 23 24 25 26 27	Maciej Kurpisz, PhD, MD Institute of Human Genetics, Polish Academy of Sciences Strzeszynska 32 60-479 Poznan, POLAND phone: +48616579202 e-mail: <u>maciej.kurpisz@igcz.poznan.pl</u>								
27 28 29	other authors' emails:								
30 31 32 33 34 35 36 37 38 39 40 41 42	Marta Olszewska, PhDmarta.olszewska@igcz.poznan.plAgnieszka Malcher, PhDagnieszka.malcher@igcz.poznan.plTomasz Stokowy, PhDtomasz.stokowy@uib.noNijole Pollock, BScNijole.pollock@gmail.comAndrea J. Berman, PhDAJB190@pitt.eduSylwia Budkiewicz, MScsylwia.budkiewicz@wp.plMarzena Kamieniczna, PhDmarzena.kamieniczna@igcz.poznan.plHanna Jackowiak, PhDhanna.jackowiak@up.poznan.plJoanna Suszynska-Zajczyk, PhDjoanna.suszynska-zajczyk@up.poznan.plPiotr Jedrzejczak, PhD, MDpiotrjedrzejczak@gmail.comAlexander N. Yatsenko, PhD, MDyatsenkoan@mwri.magee.edu								
43 44 45 46 47 48 49 50 51 52 53 54 55 56 57	ORCiD: Marta Olszewska, PhD 000-0002-2337-4545 Agnieszka Malcher, PhD 000-0002-3433-6990 Tomasz Stokowy, PhD 000-0003-0017-8338 Nijole Pollock, BSc - Andrea J. Berman, PhD 000-0003-1217-7412 Sylwia Budkiewicz, MSc - Marzena Kamieniczna, PhD 000-0002-9619-8220 Hanna Jackowiak, PhD 000-0002-7560-9350 Piotr Jedrzejczak, PhD, MD 0000-0002-7560-9350 Nijotr Jedrzejczak, PhD, MD 0000-0002-1690-0172 Alexander N. Yatsenko, PhD, MD 000-0002-1690-0172 Maciej Kurpisz, PhD, MD 000-0003-3275-3245								

2

58 Abstract

59 **STUDY QUESTION:** Is *Tcte1* mutation causative for male infertility?

60 SUMMARY ANSWER: Collected data underline the complex and devastating effect of the single-gene mutation

61 on testicular molecular network, leading to male reproductive failure.

62 WHAT IS KNOWN ALREADY: Latest data revealed mutations in genes related to axonemal dynein arms as 63 causative for morphology and motility abnormalities in spermatozoa of infertile males, including dysplasia of 64 fibrous sheath (DFS) and multiple morphological abnormalities in the sperm flagella (MMAF). The nexin-dynein 65 regulatory complex (N-DRC) coordinates the dynein arm activity, and is built from DRC1-DRC7 proteins. DRC5 66 (TCTE1) – one of N-DRC element, has been already reported as a candidate for abnormal sperm flagella 67 beating, however, only in restricted manner with no clear explanation of respective observations.

58 **STUDY DESIGN, SIZE, DURATION:** Using CRISPR/Cas9 genome editing technique, mouse knockout line of *Tcte1* 69 gene was created on the basis of C57BI/6J strain. Then, the mouse reproductive potential, semen 70 characteristics, testicular gene expression level, sperm ATP and testis apoptosis level measurements have been 71 performed, followed by visualization of N-DRC proteins in sperm, and protein modeling *in silico*. Also, a pilot 72 genomic sequencing study of samples from human infertile males (n=248) was applied for screening of *TCTE1* 73 variants.

74 PARTICIPANTS/MATERIALS, SETTING, METHODS: To check the reproductive potential of KO mice, adult 75 animals were crossed for delivery of three litters per caged pair, but no longer than for 6 months, in various 76 combinations of zygosity. All experiments were performed for wild type (WT – control group), heterozygous 77 Tcte1^{+/-}, and homozygous Tcte1^{-/-} male mice. Gross anatomy was performed on testis and epididymis, followed 78 by semen analysis. Sequencing of RNA (RNAseq; Illumina) has been done for mice testis tissues. STRING 79 interactions have been checked for protein-protein interactions, based on changed expression level of 80 corresponding genes identified in the mouse testis RNAseq experiments. Immunofluorescence in situ staining 81 was performed to detect the N-DRC complex proteins: Tcte1 (Drc5), Drc7, Fbxl13 (Drc6), and Eps8l1 (Drc3) in 82 mouse spermatozoa. To determine the ATP amount in spermatozoa, the luminescence level was measured. 83 Also, immunofluorescent in situ staining was performed to check the level of apoptosis via caspase 3 84 visualization on mouse testis samples. DNA from whole blood samples of infertile males (n=137 non-

3

85 obstructive with of azoospermia or cryptozoospermia, n=111 samples spectrum 86 oligoasthenoteratozoospermia, including n=47 with asthenozoospermia) has been extracted to perform 87 genomic sequencing (WGS, WES or Sanger). Protein prediction modeling of human identified variants and the 88 exon 3 structure deleted in mouse knockout has been also performed.

89 MAIN RESULTS AND THE ROLE OF CHANCE: No progeny at all was found for homozygous males with revealed 90 oligoasthenoteratozoospermia, while heterozygous animals (fertile) manifested oligozoospermia, suggesting 91 haploinsufficiency. RNA-sequencing of the testicular tissue showed the influence of *Tcte1* mutations on the 92 expression pattern of 21 genes responsible for mitochondrial ATP processing, linked with apoptosis, or 93 spermatogenesis. In *Tcte1^{-/-}* males the protein revealed only residual amounts in sperm head nucleus, and was 94 not transported to sperm flagella, as other N-DRC components. Decreased ATP level (2.4-fold lower) was found 95 in spermatozoa of homozygous mice, together with disturbed tail:midpiece ratio, leading to abnormal sperm 96 tail beating. Casp3-positive signals (indicating apoptosis) were observed in spermatogonia only, at similar level 97 in all three mouse genotypes. Mutation screening of human infertile males revealed 1 novel and 5 ultrarare 98 heterogeneous variants (predicted as disease causing) in 6.05% of patients studied. Protein prediction 99 modeling of identified variants revealed changes in the protein surface charge potential, leading to disruption 100 in helix flexibility or its dynamics, thus, suggesting the disrupted TCTE1 interaction with its binding partners 101 located within the axoneme.

- 102
- 103
- 104
- 105
- 106

107 **Keywords**

108 TCTE1, DRC5, N-DRC, mouse knockout model, male infertility, asthenozoospermia,
 109 oligoasthenoteratozoospermia, oligozoospermia, spermatogenesis, sperm motility, sperm mitochondria,
 110 mitochondrial respiratory complex, *mt-Co2*, *Fetub*, haploinsufficiency, MMAF, apoptosis

111

4

1	1	2
т	Ŧ	J

114

115 What does this mean for patients?

- 116 Abnormal semen parameters (sperm count, motility and morphology) are known as one of the first symptoms
- 117 that may be related to male fertility problems.
- 118 This study aimed to determine the role of *Tcte1* in male infertility using mouse knockout model. Tcte1 protein
- 119 is one of the structural elements building N-DRC, a complex within an axoneme that is responsible for
- 120 coordination of the sperm flagella elements activity, strictly related to sperm motility.
- 121 We have found that mutations in mouse *Tcte1* knockout model revealed two phenotypes, dependent on
- 122 zygosity: infertile oligoasthenoteratozoospermic homozygotes, and fertile oligozoospermic heterozygotic
- 123 males, suggesting haploinsufficiency mechanism. Pilot study on human samples with TCTE1 variants revealed a
- 124 wide spectra of semen quality (from non-obstructive azoospermia, via cryptozoospermia, to severe
- 125 oligoasthenozoospermia).
- 126 Thus, *TCTE1* gene is the next one that should be added to the 'male infertility list' because of its crucial role in
- 127 spermatogenesis influencing the variety of testicular molecular networks (incl. energy machinery processing)
- and proper sperm function.
- 129
- 130
- 131
- 132
- 133
- 134
- 135
- 136
- 137
- 138

-

- 139
- 140
- 141

5

142

143

144 INTRODUCTION

145 Infertility has been defined by World Health Organization as a social disease concerning approximately 146 10-18% of couples at reproductive age, which are unable to conceive within 1 year of regular unprotected 147 sexual intercourse [WHO, 2020]. Approximately 7% of males and ~12% of females worldwide reveal 148 reproductive problems, while the male factor is responsible for approximately 20-30% of all infertility cases 149 [Barratt et al., 2017; Vander Borght and Wyns, 2018; Agarwal et al., 2021]. Male infertility is a disease with 150 multifactorial etiology, including: genetic factors, chromosomal aberrations, epigenetic mutations, hormonal 151 abnormalities, infections or reproductive tract abnormalities [Agarwal et al., 2021; Matzuk and Lamb 2008; 152 Zorrilla and Yatsenko 2013; Krausz and Riera-Escamilla 2018]. Genetic causes of male infertility can be detected 153 in ~10-15% of cases, while in 25-50% of infertile males the etiology remains unexplained [Agarwal et al., 2021; 154 Matzuk and Lamb 2008; Zorrilla and Yatsenko 2013]. The male infertility phenotype mostly results in decreased 155 semen parameters, including: low sperm concentration, abnormal morphology or diminished motility. The 156 actual standards for genetic testing of male infertility rely on chromosomal screening for structural and 157 numerical abnormalities, followed by analyses of deletions in the AZF (azoospermia factor) regions of the Y 158 chromosome [Hwang et al., 2018]. However, it is still not sufficient due to the fact, that approximately 2,000 159 genes may be related to spermatogenesis, while only ~10% of them seems to have documented fertility role, 160 so far [Krausz and Riera-Escamilla 2018; Houston et al., 2021] Recent years NGS technology development, 161 including genomic sequencing of DNA (exome or genome; large cohorts of patients), and RNA (RNAseq, 162 scRNAseq on reproductive tissues), followed by animal models of infertility (mostly mouse models; more than 163 860 genes evaluated, so far [Mouse Genome Informatics, 24 June 2021]) have started the new era of 164 exploration on the massive scale [Houston et al., 2021; Oud et al., 2021; Wyrwoll et al., 2020; Hardy et al., 165 2021; Xavier et al., 2021].

Beside the genes related to spermatogenesis and low sperm count, also studies of candidates responsible for diminished sperm motility are pending. Thus, the main structural element responsible for motility – the sperm flagella, has to be of high-quality to guarantee sperm motility and then fertilization success. It is known that mutations in genes encoding factors involved in ciliogenesis process, lead to defects in structure and/or functioning of cilia [Girardet *et al.*, 2019; Sironen *et al.*, 2020; Smith *et al.*, 2021; Kumar and

6

171 Reiter 2021] There is a plenty of data concerning respiratory cilia, however, still little is known about the 172 axonemal dynein arms functioning in sperm tail. Latest data revealed mutations in genes related to axonemal 173 dynein arms as causative for morphology and motility abnormalities in spermatozoa of infertile males [Girardet 174 et al., 2019; Sironen et al., 2020; Aprea et al., 2021; Lorès et al., 2021; Touré et al., 2021; Martinez et al., 2022]. 175 Two definitions were established: (i) for sperm cells with disorganization in axonemal structures, so called 176 dysplasia of fibrous sheath (DFS) [Chemes et al., 1998] and (ii) for multiple morphological abnormalities in the 177 sperm flagella (MMAF), concerning all possible defects of sperm tail, mainly short and thick flagella, and a lack 178 of the central pair (CP) of microtubules or dynein arms [Khelifa *et al.*, 2014]. Spectrum of abnormal sperm tail 179 phenotypes observed allows qualification of a part of disorders into both listed groups (DFS and MMAF) 180 [Sironen et al., 2020; Touré et al., 2021]

181 The basic structural element of motile cilia and flagella is axoneme built from nine microtubules 182 arranged in doublets positioned radially and one central pair (CP) of singlet microtubules. This central pair is 183 connected to radial spokes (RS) - protein complexes of mechanic regulatory network responsible for regular 184 waveform beating of the cilia [Gui et al., 2019]. RS are linked to inner dynein arms (IDA) localized within each 185 outer dublet. Each outer doublet carries outer dynein arms (ODA) - multi-subunit motor protein complexes, 186 responsible for generation and regulation of sliding between dynein arms leading to ATP-dependent beating 187 [Satir et al., 2014; Wilson et al., 2015]. The 9+2 basic structure is evolutionary conservative [Girardet et al., 188 2019; Touré et al. 2021; Ishikawa 2017]. Doublets are linked with the nexin-dynein regulatory complex (N-DRC), 189 which coordinates the dynein arm activity and stabilizes the attachment of doublet microtubules. The N-DRC is 190 built from DRC1-DRC7 proteins, among which DRC1, DRC2 and DRC4 are responsible for physical linkage 191 between A and B microtubules of doublets [Gui et al., 2019; Morohoshi et al., 2020; Bowar et al., 2018; 192 Wirschell et al., 2013]. In terms of male infertility, the best studied gene among N-DRC is DRC7 (MIM:618769), which is amenable for proper assembly of the N-DRC structure (in $Drc7^{/2}$ male mice the length of the sperm tail 193 194 was reduced, followed by abnormalities of sperm head morphology) [Morohoshi et al., 2020]. Mouse model of 195 Drc6 (Fbx/13) gene (MIM:609080) revealed no essential role in sperm formation and motility [Morohoshi et al., 196 2020], however, its disruption together with CEP192 (MIM:616426) leads to defects in HEK293T cell motility 197 [Fung et al., 2018]. Mutations in Drc3 gene (MIM:618758) indirectly revealed male infertility as a lack of 198 pregnancy success with wild-type females [Ha et al., 2016].

7

199 One of the N-DRC structural genes is TCTE1 (T-Complex-associated-Testis-Expressed 1), known also as 200 DRC5, FAP155, D6S46 in humans (locus 6p21.1; MIM:186975) or Tcte1, D17S, Tcte-1, D17Sil1 in mouse (locus 201 17 B3; 17 22.54 cM; MGI:98640). TCTE1 is evolutionary conserved in most eukaryotic organisms, that contain 202 motility cilia or flagella in their life cycle (Figure 1), and is required for proper flagella functioning. Expression of 203 TCTE1 is being observed in testis (highest; with expression starting at the stage of spermatid), choroid plexus 204 and Fallopian tube. TCTE1 contains five leucine rich repeats (LRRs). To date, only two reports described a role 205 of *Tcte1* gene: as a mice model of infertility, where the *Tcte1-null* sperm cells exhibited asthenozoospermia or 206 in group of patients without clear explanation of the linkage between observed variant and sperm phenotype 207 [Zhou et al., 2022; Castaneda et al., 2017]. Thus, understanding of the nature of this specific motility type 208 defect in *Tcte1-null* observations has been limited, so far.

209 In our study we aimed to determine the role of *Tcte1* in male infertility using mouse knockout model. 210 We created homo- and heterozygous animals, which were next examined for their reproductive potential, 211 differences in anatomy and histology, and semen parameters. We also performed RNA sequencing (RNAseq) in 212 testis tissue to obtain an answer about the influence of *Tcte1* null mutation on the expression pattern of other 213 gonadal genes. Also, the ATP measurement of spermatozoa has been performed. Additionally, screening of 214 variants in human samples from males with severely decreased sperm count was made, followed by protein 215 prediction modeling of revealed variants. Thus, this study contains novel and the most complete data 216 concerning TCTE1 role in male infertility, so far.

217

218 MATERIALS AND METHODS

219 Animals

220 Using CRISPR/Cas9 genome editing technique, mouse heterozygous knockout line of Tcte1 gene 221 (Tcte1^{+/}) was created on the basis of C57BI/6J strain in the Cyagen Biosciences company (USA). The mouse 222 Tcte1 gene is located on chromosome 17 B3; 17 22.54 cM (GenBank: NM 013688.2; Ensembl: 223 ENSMUSG0000023949). Exon 3 of the Tcte1 gene was selected as target site (deletion of 1292 bp) (Figure 2A). 224 gRNAs targeting vectors (constructed, sequenced and generated by *in vitro* transcription) were injected into 225 fertilized oocytes (Supplementary file 1). The founders were genotyped and those with positive results were 226 bred to the next generation. Then, in the animal facility of the Institute of Human Genetics PAS (Poznan, Poland), adult Tcte1^{+/-} animals were mated to generate homozygous Tcte1^{-/-} mice. Wild type Tcte1^{+/+} (WT) 227

8

228 animals C57BI/6J were purchased from Charles River company (Germany). Relative expression of Tcte1 gene 229 was verified using Real-time PCR method (Figure 2B; Supplementary file 2). All mice were housed in pathogen-230 free animal facility with food and water *ad libitum*, stable temperature of 22°C, and a light:dark light cycle of 231 12:12. All procedures were in accordance with the guidelines and regulations for animal experimentations and 232 restricted usage of genetically modified organisms (GMO), and were approved by the Local Ethical Committee 233 for Animal Experiments at the Poznan University of Life Sciences (no. 1/2014, 18/2017), and Ministry of 234 Environment (no. 70/2017). Mouse models with infertility phenotypes were reviewed in the Mouse Genome 235 Informatics database (Jackson Laboratory; http://www.informatics.jax.org/) and through literature searches 236 (PubMed; https://pubmed.ncbi.nlm.nih.gov/).

237 Reproductive potential

To check the reproductive potential of KO mice, adult animals (7-8 weeks old) were crossed for delivery of three litters per caged pair, but no longer than for 6 months. Crossing combinations included: (i) 4 pairs: homozygous $Tcte1^{-/-}$ male × wild type (WT) female, (ii) 5 pairs: heterozygous $Tcte1^{+/-}$ male × heterozygous $Tcte1^{+/-}$ female, and (iii) 3 pairs: heterozygous $Tcte1^{+/-}$ male × wild type (WT) female. Vaginal plug presence was checked every morning to confirm mating incidence. Time period from mating start point to first pedigree appearance was checked. Number of pups was counted on the day after birth. For each crossed pair, the number of litters, pups, and sex ratio were estimated (also according to pups' genotypes) (Supplementary file 3).

245 3).

246 Genotyping

247 Genotyping of the pups delivered was performed via PCR assay, with two sets of primers: (i) to identify 248 wildtype and positive KO (het- and homozygous) animals: Tcte1-F: CGTTTTAAGAATGTATTGAGGGTTGGG, Tcte1-249 R: CTCCGTAGGCTCCTGCCAATATG; estimated product size: for WT 1915 bp, for KO ~630 bp (deleted ~1300 bp); 250 PCR results analysis: one band with 1915 bp or 0 bp determined WT, while one band with 630 bp evidenced 251 hetero- or homozygotes; (ii) to distinguish between hetero- and homozygotes: Tcte1-WT/He-F: 252 AGCTTGCCACACCCTCAAGGTACTA, Tcte1-R: CTCCGTAGGCTCCTGCCAATATG; estimated product size: 658 bp for 253 heterozygotes and 0 bp for homozygotes; PCR results analysis: one band with 658 bp determined 254 heterozygotes, while no one band (0 bp) indicated homozygotes (Figure 2C). The PCR mixture and reaction 255 conditions are described in Supplementary file 4.

256

Tissue gonadal preparation/collection

9

Gross anatomy was performed on testis and epididymis of WT (n=21), $Tcte1^{+/2}$ (n=22), and $Tcte1^{-/2}$ 257 258 (n=18) male animals (10 weeks old). Mice were weighted, then euthanized with carbon dioxide (CO_2) overdose 259 (euthanasia Minerve Easy-box; Minerve; France) with stable flow rate per minute. Testes and epididymides 260 were collected, weighted, and measured. For histological evaluation, two fixing approaches have been applied. 261 Both approaches started with anaesthetization under isoflurane (4%) and transcardial perfusion with 50 262 ml/animal of ice-cold PBS, followed by 50 ml/animal of 10% formalin solution, neutrally buffered (Sigma 263 Aldrich, USA, cat. no. HT501850) for 3 males per zygosity and technical variant. Next, the tissues have been 264 immersed in formalin solution for 24 hours, 4°C, in darkness. Then, in the first approach, Paraplast-embedded 265 sections were prepared (5 µm thickness), using Microtome Leica 2055. In the second technique, gonads were 266 sequentially immersed in 15%, 20% and 30% of sucrose (Sigma Aldrich, cat. no. S0389), each for 24 hours, at 267 room temperature. Thereafter, gonads were coated with Tissue-Tek O.C.T. Compound (Sakura, cat. no. 4583) 268 and frozen in cold (-80° C) 2-Methylbutane (Sigma Aldrich, USA, cat. no. M32631). Then, cryo-sections were 269 prepared (10 μ m thickness), using Leica Cryostat CM1950 and kept at -80°C until further stainings. To obtain 270 the histological picture, slides were stained with Masson-Goldner protocol [Romeis et al., 2010] for classic 271 paraffin blocks or standard progressive H&E staining method for frozen samples. Briefly, after fixation twice in 272 100% EtOH (10 min.), slides were first hydrated in the EtOH series (95%, 70% for 5 min.), washed in distilled 273 water and stained in Mayer's hematoxylin (Sigma) for 5 min., then washed in distilled water (3 times, 5 min.) 274 and dehydrated in 50%, 70 %, 80%, 96% EtOH (5 min. each). Counterstaining with eosin Y (Sigma) for 30 275 seconds was applied, followed by washing in 100% EtOH, clearing in xylene (5 min. each, two repeats), and 276 mounting in DPX (Sigma). Thereafter, slides were documented with Leica DM5500 light microscope (×40 dry or 277 ×63 oil immersive objectives, motorized stage), LASX software (with Navigator tool). All the evaluations have 278 been performed blindly by two experienced scientists per technique.

For each zygosity the geometrical features of the tubule sections have been evaluated, incl.: tubule diameter (D) and cell layer thickness (CL) (the mean of two measurements made at opposite sides of the tubule) and lumen diameter (I), using Fiji (Image J) measurement tool. To check density of the cells, ratios of: D:CL, D:I and CL:I were also estimated. In total, for each zygosity 133 -144 tubule sections were evaluated (3 random sections per zygosity, each min. 30 tubules).

284 Mouse semen samples

10

285 Mouse spermatozoa were collected from epididymis post mortem. Collected epididymis were placed 286 immediately in a 2.0 mL test tube with warm 500 µL of a non-capacitating Whitten's HEPES medium (pH 7.2-287 7.4, 37°C), then cutted into 5-7 fragments, and incubated in 37°C for 15 min. to allow the sperm cells to swim 288 out. Whittens-HEPES medium contained: 100.0 mM NaCl, 4.4 mM KCl, 1.2 mM KH2PO4, 1.2 mM MgSO4, 5.4 289 mM glucose, 0.8 mM pyruvic acid, 4.8 mM lactic acid (hemi-Ca), HEPES 20 mM [Wertheimer et al., 2013]. Then, 290 spermatozoa were collected and proceeded with seminological analysis (concentration, motility, morphology). 291 Next, semen samples were fixed in a fresh fixative solution (methanol:acetic acid, 3:1 v/v, $-20^{\circ}C$; $3\times$) and stored 292 until further use in -20°C.

293 Semen analysis

294 10 μ l of sperm suspension was applied on the Makler counting chamber (Sefi medical instrument Ltd., 295 Israel) to test the sperm concentration and total sperm count. Also, the ratio of progressive (fast and slow), 296 non-progressive (total and circular) and immotile spermatozoa were calculated [WHO, 2020]. For assessment 297 of sperm morphology, the Papanicolaou staining was performed. Briefly, the air-dried smears were fixed in 298 ethanol:diethyl ether (1:1 ratio, vol/vol) solution for 10 min. and left for air-drying. Then, slides were stained by 299 Papanicolau procedure according to WHO guidelines [WHO, 2020], and mounted in DPX Mountant (Sigma-300 Aldrich; USA; cat. no. 06522). A total of 200 spermatozoa were analyzed using Leica DM5500 light microscope 301 (×1000 magnification, oil immersion). Various morphological forms of spermatozoa were determined including 302 abnormalities of head, midpiece and tail. Also, the length of sperm tail, midpiece and the tail:midpiece ratio 303 were measured to check possible differences between males with various zygosity (6 males/per zygosity); for 304 each male at least 100 spermatozoa were measured with the 'length measurement' option by CellSense 305 Dimension software (Olympus).

306 Sequencing of RNA (RNAseq)

307 For RNA extraction testis tissue was collected into in RNAlater buffer (Invitrogen; Waltham, MA, USA) 308 and freezed immediately (-80°C). Samples were processed using the AllPrep DNA/RNA/Protein Mini Kit 309 (Qiagen; Hilden, Germany). Quality and quantity of extracted RNA samples were checked using NanoDrop 310 (Thermo Scientific) and Quantus (Promega), and 1 μg of RNA per sample has been used for sequencing 311 (Macrogen; South Korea). RNA sequencing was performed with Illumina NovaSeq 6000 platform using TruSeq 312 Stranded mRNA LT Sample Prep Kit, with specified conditions of paired-end reads at 100 bp, and throughput 313 100 M reads per sample. RNA-Seq raw reads were aligned to mouse reference genome mm10 using hisat 2.0.5

11

with Gencode vM13 transcriptome reference. Reads aligned in the coding regions of the genome were counted using Feature Counts. Then, read counts were normalized using DESeq2 and subjected to differential analysis. The differential analysis included computing of median based fold change, statistical significance using the Students' t-test and false discovery rates (correction for multiple testing) in the R/Bioconductor programming environment. Testis tissue expression was retrieved from GTEx (<u>https://gtexportal.org/home/</u>), BioGPS (<u>http://biogps.org/</u>), and AceView (<u>https://www.ncbi.nlm.nih.gov/IEB/Research/Acembly/</u>). All RNA-seq expression data have been deposited in Gene Expression Omnibus (GEO), accession number GSE207805.

321 STRING interactions

For analysis of protein-protein interactions, based on changed expression level of corresponding genes identified in the mouse testis RNAseq experiments, the STRING analysis was performed [version 11.0]. STRING database allows to delineate probability of the interactions between evaluated proteins using different evidence channels [Mering *et al.*, 2005; Szklarczyk *et al.*, 2019]. This tool enables the integration of all data available in public resources concerning protein-protein interactions.

327

Immunofluorescence on sperm – N-DRC visualization

328 Immunofluorescence in situ staining was performed to detect the N-DRC complex proteins: Tcte1 329 (Drc5), Drc7, Fbxl13 (Drc6), and Eps8l1 (Drc3) in mouse spermatozoa. Mouse spermatozoa were collected and 330 fixed as described above. Specific antibodies were applied: primary antibodies – rabbit anti-human-TCTE1 (cat. 331 no. orb357083), rabbit anti-mouse-Drc7 (orb586958), anti-mouse-Fbx|13 (orb314611), and anti-mouse-Eps8|1 332 (orb382538; all from Biorbyt; Cambridge, UK); secondary antibody – goat anti-rabbit-AF594 (Abcam; cat. no. 333 ab150160; Cambridge, UK). The UniProt IDs and homology of the immunization regions of primary antibodies 334 between human and mouse were described in Supplementary file 5. Antibodies were diluted in 0.5% PBST 335 (primary: 1:100, secondary: 1:500). First, slides with fixed sperm smears were washed in a series of washes in 336 1× PBST, and then incubated in 25 mM DTT/1 M Tris-HCl, pH 9.5, at room temperature for 20 min. Then, 337 washing in 1× PBST, and incubation in 6 N HCl for 30 min. was applied, followed by blocking step with 1% 338 BSA/1× PBST for 30 min. Next, overnight incubation with antibodies was performed at 4°C in a humidified 339 container. After washing, the samples with 1× PBST, secondary AF594-conjugated antibody was applied for 1 340 hr. Next, unconjugated antibodies were washed out (4× in 1× PBST, 5 min each, slight rotation). For the final 341 detection, 20 ul of Fluoroshield with DAPI (Sigma-Aldrich; cat. no. F6057) was added to the samples, and 342 microscopic analysis was performed. Images were acquired using a fluorescence microscope with a proper

12

filter set: Leica DM5500, filters: DAPI/TxR/Triple; objectives: 10× and 63× with oil immersion; DFC 7000T
camera; software: LASX.

345 Sperm ATP measurement

346 For measurement of ATP in spermatozoa the CellTiter-Glo 2.0 Assay (Promega; USA) was used. This 347 ready-to-use assay allows to determine the quantitation of the ATP amount, indicating the presence of 348 metabolically active cells. The assay is based on the thermostability of luciferase (Ultra-GloTM Recombinant 349 Luciferase), and was performed as described in manufacturers' protocol. Briefly, all laboratory work was done 350 in the stable temperature of 22°C. After collection of mouse spermatozoa (as described above), the amount of 351 approximately $1 \times 10^{\circ}$ of sperm cells was transferred to black opaque 1.5 mL test tube and proper amount of 352 CellTiter-Glo kit was added (1:1, v:v). Next, 2 min. of orbital mixing of the content was performed to activate 353 cell lysis. Then, after 10 min. of incubation (to stabilize the luminescence), the measurement of luminescent 354 signal was completed using Glomax 20/20 luminometer (Promega; USA). For each sample, three 355 measurements (repeats) were done, each in two time points of 0 and +60 min. Numbers of animals used were: WT (n=7), Tcte1^{+/-} (n=7), and Tcte1^{-/-} (n=6). Mean luminescence values obtained (represented as RLU – relative 356 357 luminescence unit) were compared between tested mice groups.

358 Immunofluorescence on testis

359 Immunofluorescence in situ staining on testis was performed to check: (i) mean cell count of 360 spermatogonia and spermatocytes (preleptotene and pachytene) in tubules' sections (cryo-sections), and (ii) to 361 estimate the level of apoptosis via caspase 3 visualization on mouse testicular samples (paraffine embedded 362 sections) from all the three zygosities. Caspases play important role in the induction, transduction, and 363 amplification of intracellular apoptotic signals, and thus, caspase-3 can be used as one of the apoptotic 364 indicators [Lej et al, 2022]. Mouse testes were collected and fixed as described above. Specific antibodies were 365 applied: for (i) primary antibodies – mouse anti-MLH1 (Abcam; Cambridge, UK, cat. no. ab14206), 1:30 in 0.5% 366 PBST, rabbit anti-RAD51 (Abcam; Cambridge, UK, cat. no. ab63801), 1:200 in 0.5% PBST; (ii) primary antibody – 367 rabbit anti-Casp3, (Biorbyt; Cambridge, UK, cat. no. orb10237), 1:100 in 0.5% PBST; secondary antibodies - goat 368 anti-mouse-FITC (Sigma-Aldrich, cat. no. F2012), 1:400 and goat anti-rabbit-AF594 (Abcam; Cambridge, UK, cat. 369 no. 150080), 1:500 in 0.5% PBST. First, slides with paraffine fixed testis cross sections were deparaffinized in 3 370 changes of xylene (10 mins. each), while for cryo-sections this step has been omitted. Then slides were 371 incubated in a series of ethanol: 100%, 96%, and 70%, twice for 5 mins. in each solution. Next, after rinsing in

13

372 distilled water, tissue sections were incubated sequentially in: a freshly prepared 2% NaBH₄/1× PBS, and in 373 0.1M glycine (30 min., room temp. each). Heat-induced epitope retrieval was performed by immersing the 374 slides in a 0.01M sodium citrate/0.05% Tween 20, pH 6.0 for 30 min., followed by cooling down in a room 375 temp. for next 20 min. Blocking step was done in an aliquot of freshly prepared 1% BSA/1× PBST, at 37 °C for 1 376 h. Then, tissue sections were incubated with primary antibody in a humidified container overnight at 4°C. 377 Secondary antibody was applied for 1 hr, followed by the washing out of the unconjugated antibody (4× in 1× 378 PBST, 5 min each, slight rotation). For the final detection, 20 µl of Fluoroshield with DAPI (Sigma-Aldrich; cat. 379 no. F6057) was added. Images were captured using a fluorescence Leica DM5500 microscope with a proper 380 filter set: DAPI/TxR/SpG/Triple; objectives: 10× and 63× with oil immersion; DFC 7000T camera, LASX software. 381 The analysis was performed for 3 sections per zygosity, each min. 30 tubules (for cell count) or 2 cross-sections 382 per zygosity, 15-20 tubules per tissue section (for Casp-3 evaluation).

383 TUNEL assay

384 TUNEL assay was performed to check the level of apoptosis and necrosis on the testicular tissue 385 according to the manufacturer's protocol (Abcam, ab66110 TUNEL Assay Kit – BrdU-Red). Breifly, paraffine 386 embedded sections (or 2 cross-sections per zygosity, with 15-20 random tubules analyzed/slide) have been 387 deparaffinized in xylene (twice, each 5 min.), followed by EtOH series: 100% (twice, 5 min.), 95%, 85%, 70% and 388 50% (3 min.). Next, slides were incubated in 0.85% NaCl (5 min.) and washed in PBS. After incubation with 389 Proteinase K solution (20 ug/mL in Tris-HCl-EDTA; 5 min.) and washing with PBS, slides were immersed in 4% 390 formaldehyde/PBS. Then, Wash Buffer was applied twice for 10 min. with plastic coverslip on the top. Next, 391 DNA Labeling Solution was added and incubation in a dark humidified incubator for 1,5 hr in 37°C was 392 processed. After next two washes with PBS, Antibody Solution was applied and slides were incubated for 30 393 min. at room temp. Last steps included: ddH_2O washing (5 min., twice) and application of 20 μ l of Fluoroshield 394 with DAPI (Sigma-Aldrich; cat. no. F6057) for the final detection. Slides were analyzed within 1 hr using the 395 fluorescent Leica DM5500 microscope with a proper filter set: DAPI/TxR/Triple; objectives: 10× and 63× with oil 396 immersion; DFC 7000T camera, LASX software.

Sequencing of the human samples

397

A pilot genomic sequencing study of samples from human infertile males was also applied. Bioethical
 committee approvals were received for the study (Local Bioethics Committee of Poznan University of Medical
 Sciences, approval no. 1003/18; National Bioethical Committee, Ministry of Health, Warsaw, Poland, approval

14

401 no. OKB-5-2/15). All participants were notified about the aim of the study, and provided written informed 402 consent. All experiments were performed in accordance with relevant guidelines and regulations. DNA from 403 whole blood samples (collected into sterile single test tubes with EDTA) was extracted using standard protocols 404 of MagCore HF16 Automated Nucleic Acid Extractor (RBC Biosciences, Taiwan), or Gentra Puregene kit (Qiagen, 405 USA). Total number of sequenced patients was n=248, including: 137 with non-obstructive azoospermia (lack of 406 spermatozoa in ejaculate) or cryptozoospermia (spermatozoa detectable in ejaculate after centrifugation), 407 followed by 111 samples with spectrum of seminal abnormalities in concentration, and/or motility, and/or 408 morphology. Among those samples, 47 revealed decreased motility (asthenozoospermia). Semen evaluation 409 was done according to WHO guidelines [WHO, 2020], followed by AUA and ASRM guidelines for infertile male 410 diagnosis [Schleger et al., 2021]. All men revealed normal hormonal levels, had no chromosomal abnormalities 411 and a lack of AZF microdeletions. For whole exome samples (n=76 from Pittsburgh), sequencing libraries were 412 constructed using the Agilent technologies (Santa Clara, USA), including: SureSelect XT All Exon V4 + UTRs 413 capture kit and SureSelect XT Target Enrichment System. Libraries were sequenced with average read coverage 414 of 62-100x (~6Gb per sample), using 100 bp paired-end sequencing with TruSeq PE Cluster Kit v3 cBot HS and 4 415 × 50 TruSeq SBS V3 HS sequencing kit (Illumina, San Diego, USA) on the HiSeq2000 (Illumina, San Diego, CA). 416 Whole genome sequencing (n=42, Poznan) was performed using Illumina HiSeqX with the coverage of at least 417 30x (100-120 Gb per sample). Sanger sequencing for all exomes of *TCTE1* gene (n=130, Poznan) was performed 418 using LightRun96 system of Eurofins Genomics (Eurofins Genomics AT GmbH, Vienna, Austria). Samples were 419 prepared with QIAquick Gel Extraction (cat. no. 28704) or PCR Purification (cat. no. 28104) kits (Qiagen, USA), 420 and analyzed with CLC Main Workbench 8 software (Qiagen, USA).

421 Filtering of whole genome sequencing results

422 The quality of raw NGS data was evaluated using the FastQC and MultiQC packages. Reads were aligned to the 423 reference human genome GRCh37 using bwa-mem aligner included in the Speedseq package [Chiang et al., 424 2015]. For calling of single nucleotide variants (SNVs) and small indels the freeBayes v0.9.21 (speedseq-var) 425 was implemented, following previously described pipeline [Supernat et al., 2018]. Variant disease association 426 and missense mutation prediction effects were determined using: dbSNP (https://www.ncbi.nlm.nih.gov/snp/), 427 OMIM (https://www.ncbi.nlm.nih.gov/omim/), ClinVar (https://www.ncbi.nlm.nih.gov/clinvar/), 428 MutationTaster (http://www.mutationtaster.org/), SIFT (https://sift.bii.a-star.edu.sg/), Polyphen 429 (http://genetics.bwh.harvard.edu/pph2/), CADD (https://cadd.gs.washington.edu/snv), MetaDome

15

430 (https://stuart.radboudumc.nl/metadome/), REVEL (rare exome variant ensemble learner) [loannidis et al., 431 2016], Ensembl Variant Effect Predictor [Mclaren et al., 2016], and Human Gene Mutation Database (HGMD; 432 Qiagen, Redwood City, CA). SNV pathogenicity was assessed on the basis of American College of Medical 433 Genetics and Genomics (ACMG) guidelines [Richards et al., 2015]. PhyloP (Cold Spring Harbor Laboratory) and 434 Clustal Omega were used for determination of conservation (https://www.ebi.ac.uk/Tools/msa/clustalo/). All 435 NGS variants were confirmed via the Integrative Genomics Viewer (Broad Institute). Variant frequencies have 436 been obtained from the gnomAD database (frequencies from more than 125k exomes and 15k genomes, 437 http://gnomad.broadinstitute.org; v2.1.1), and were filtered in search of rare variants (with minor allele 438 frequency (MAF) <1%).

439 Protein prediction

440 The amino acid sequence of TCTE1 was input into Phyre2 [Kelley et al., 2015] for both intensive and 441 fast mode homology modeling, I-TASSER [Roy et al., 2010; Yang et al., 2015, Yang and Zhang 2015], and 442 AlphaFold [Jumper et al., 2021; Varadi et al., 2022] which utilize sequence alignment-based algorithms for 443 generating predicted secondary and tertiary structures; AlphaFold additionally uses machine learning to 444 integrate biological principles of protein folding with the alignment-based algorithms. Confidence scores were 445 obtained from these programs; AlphaFold additionally generates predicted aligned error plots for assessing the 446 quality of the model's long-range interactions. Models were inspected manually and aligned using the 447 secondary structure matching (SSM) [Krissinel and Henrick, 2004] tool in COOT [Emsley et al., 2010] to 448 minimize user input bias. Docking of the structures into electron tomography data was performed manually in 449 COOT [Emsley et al., 2010]. No refinement was performed to minimize bias. All structure model figures and 450 mutations without post refinement, were generated using Pymol (Schrödinger). I-TASSER was used also for 451 prediction of the exon 3 structure deleted in our mouse KO model (Supplementary file 6).

452 Statistical analysis

453 Statistical analyses included the following: normality test (D'Agostino and Pearson or Kolmogorov-454 Smirnoff), unpaired two-tailed t-test with Welch's correction or Mann-Whitney test, one-way ANOVA test and 455 Fisher's exact, and two-tailed Pearson correlation. All tests were performed with a significance level of α =0.05 456 using GraphPad Prism (v.7.0e) software.

457 **RESULTS**

458 Reproductive potential of KO mice

16

459 Reproductive potential results were shown in Figure 3 and Supplementary file 3. Only pairs with homozygous Tcte1^{-/-} males revealed no progeny (Tcte1^{-/-} male x WT female). Other combinations (Tcte1^{+/-} male 460 \times *Tcte1*^{+/-} female; *Tcte1*^{+/-} male \times WT female) showed similar results, when considering; number of litters, pups, 461 and sex ratio. Mean time of caging to delivery of the 1st pedigree was longer in $Tcte1^{+/}$ male $\times Tcte1^{+/}$ female 462 463 combination by 25.45% (statistically not significant). When considering genotypes of the pups, statistically 464 significant differences (p<0.05) were observed in the number of $Tcte1^{-7}$ male and female pups delivered – no homozygous animals were obtained from combination of $Tcte1^{+/-}$ male × WT female, while in mating of $Tcte1^{+/-}$ 465 male \times *Tcte1*^{+/-} female, homozygous males (33.25% of all pups) and females (24.67%) were received. 466

467

Histopathology and seminal parameters

468 Histopathological evaluation of testicular and epididymal sections revealed no visual (structural) 469 differences between WT, $Tcte1^{+/}$, and $Tcte1^{-/}$ males confirming preserved spermatogenesis (Figure 4A, 470 Supplementary file 7). No odds were found for body mass of the animals and the length of the epididymis (Figure 4B). In the Tcte1^{-/-} males testis weight was decreased by approximately 41% (p<0.0001) followed by the 471 ~22% decline of its size (p<0.0001), while in *Tcte1^{+/.}* males only testis weight was slightly decreased (approx. 472 473 7%; p<0.05) when compared to control animals. Differences in dimensions of the tubule sections were noted 474 (Figure 4C), with increasing tubule and lumen diameters while decreasing cell layer thickness in Tcte1⁻². In 475 Tcte^{+/-} the cell layer thickness and tubule diameter were also increased. When comparing germ cell count 476 (Figure 5) reduced numbers of spermatogonia (~22%) and pachytene spermatocytes (13-17%) were 477 documented for $Tcte1^{-4}$ and $Tcte^{+4}$ (p<0.001), while preleptotene spermatocytes level was elevated (p<0.001). 478 Mean sum of evaluated the cell types was similar between WT and homozygotes (65.95 and 64.39 of 479 cells/tubule section, respectively). Such data suggest some kind of time shift at the first stages of prophase -480 germ cells from KO mice seem to be delayed at preleptotene stage, while WT males enters pachytene stadium 481 at higher ratio.

When considering seminal parameters, both: in $Tcte1^{-t}$ as well as in $Tcte1^{+t}$ males the sperm concentration was importantly decreased (3.27, and 2.47-fold, respectively), when collating to control results (p<0.0001; Figure 6A). Additionally, in $Tcte1^{-t}$ males the frequency of progressively motile spermatozoa was residual, with progressive type of motility in only 0.22% of spermatozoa (decreased ~150-fold vs. control 33.52%, (p<0.0001)), followed by 4.89% of slow progressive spermatozoa (decreased ~3-fold vs. control 15.50%) (p<0.0001). Interestingly, ~2.4-fold increase in circular type of motility (around own axis and non-

17

progressive) was observed for $Tcte1^{-/}$ males when compared to WT and $Tcte1^{+/}$ animals (p<0.01) 488 489 (Supplementary file 8; Supplementary file 9; Supplementary file 10). The frequency of immotile spermatozoa also increased in Tcte1^{-/-} males (2-fold to control; p<0.0001). The frequency of morphologically normal 490 491 spermatozoa in $Tcte1^{-7}$ animals was significantly decreased (2-fold to control; p<0.0001; Figure 6A, B, C); 492 Supplementary file 11). The majority of morphological defects was observed for sperm head, mostly of decapitated spermatozoa in Tcte1^{-/-} animals - 52.55% vs. 19.16% for Tcte1^{+/-} and 23.33% for WT, and 493 amorphous head shape with decreased frequency in Tcte1 $^{\prime \prime}$ (26.11%) and Tcte1 $^{*\prime \prime}$ (33.72%) males vs. WT 494 animals (48.88%) (Supplementary file 11). Midpiece defects were found in Tcte1^{+/-} (9.71%) and WT (8.20%) 495 496 animals, while Tcte1^{-/-} mice revealed only low frequency (0.38%) of such defects. On the other hand, an increased frequency (~5-fold) of coiled tail was observed for $Tcte1^{+/-}$ animals when compared to other mice 497 498 zygosity (p<0.01) (Supplementary file 11). Summarizing, seminological changes in mice with Tcte1 mutation resulted in oligoasthenoteratozoospermia in homozygous $Tcte1^{-/-}$ or oligozoospermia in heterozygous $Tcte1^{+/-}$ 499 500 animals.

501 Increased mean length of tail and tail:midpiece ratio in $Tcte1^{-/2}$ animals (139.30 ± 7.71 um; 5.16, 502 respectively) was measured when comparing to WT (122.00 ± 6.375 um; 4.51; p<0.002) and $Tcte1^{+/2}$ (123.50 ± 503 1.89; 4.83; p<0.008) (Figure 6D). Also, in the $Tcte1^{+/2}$ the midpiece length differed from $Tcte1^{-/2}$ animals 504 (p=0.0334), followed by tail:midpiece ratio differences when compared to WT (p=0.0435).

505

Sequencing of RNA (RNAseq) and STRING pathway analysis

506 RNA sequencing analysis of gene expression in testes of KO mice revealed statistically significant changes (p<0.05) in 21 genes, including 5 genes both for $Tcte1^{-1}$ vs. $Tcte1^{+1}$ and WT, 8 genes only for $Tcte1^{-1}$ vs. 507 WT, 7 genes only for Tcte1^{-/-} vs. Tcte1^{+/-}, and 1 gene both for Tcte1^{+/-} vs. Tcte1^{-/-} and WT (Figure 7). In Tcte1^{-/-} 508 509 animals vs. both: Tcte1 $^{+/-}$ and WT, the expression level of 4 genes from kallikrein subfamily was decreased 2-8-510 fold, including genes involved in semen liquefaction and degradation of extracellular matrix proteins in the 511 interstitial area surrounding Leydig cells of the adult mouse testes (Klk1b27, Klk1b21, Klk1b24), or neuronal 512 activity (Klk1b22) (Figure 7; Supplementary file 12) [Michaelis et al., 2017; Marques et al., 2016; Du Plessis et 513 al., 2013; Fink et al., 2007]. 2-fold decrease of expression of genes related to mitochondrial cytochrome 514 oxidase subunits and asthenozoospermia was documented (mt-Co2, mt-Co3) (Figure 7) [Heidari et al., 2016; 515 Mostafa et al., 2016; Baklouti-Gargouri et al., 2013]. In a scheme: Tcte1^{-/-} vs. WT, 8 genes revealed changes in 516 expression level, including 4.7-fold rise for Fetub gene, which is known as required for egg fertilization (via

18

517 specific inhibitor activity of the *zona pellucida* (ZP) ovastacin, the cysteine protease responsible for ZP2 518 cleveage) [Binsila *et al.*, 2021; Karmilin *et al.*, 2019], and plays also a role of a secreted factor during 519 reprogramming [Bansho *et al.*, 2017].

520 Two-fold rise for Igf2 gene (strongly linked to spermatogenesis, sex determination, testis 521 development, infertility and prostate cancer; [Matsuzaki et al., 2020; Cannarella et al., 2020; Neirijnck et al., 522 2019; Damaschke et al., 2017]) has been also noticed. Igf2 and Klk proteins are enclosed in a common network 523 consisted of proteins directly linked to fertility (Figure 8; Supplementary file 12; Supplementary file 13). One of 524 them is Ambp, a fusion protein, which proteolysis generates bikunin – a protein which deficiency results in 525 infertility, due to the diminished formation of the stable cumulus-oocyte complex required for maturation and 526 ovulation [Sato and Kan,2001; Naz and Rajesh, 2005]. Bikunin is one of the elements of ITIH domains (Inter- α -527 Trypsin Inhibitor Heavy-chain), responsible for hyaluronian metabolism [Hull et al., 2012; Salustri et al., 2019]. 528 ITIH is also linked to Plg, which has the protease activity and activates metalloproteins, such as: Mmp8 and 529 Mmp9, also included in the network. Mmp8 is engaged in degradation/stabilization of occluding and blood-530 testis barrier [Chen et al., 2018], while Mmp9 (with Igf1 and Igf2), is important for normal placentation [Ghosh 531 et al., 2011]. Igf1 and Igf2, are related to Igfbp3, which interacts with humanin expressed in Leydig cells. 532 Humanin together with IGFBP3 and Bax prevents the activation of germ cell apoptosis [Jia et al., 2015; Lue et 533 al., 2021]. What is important, expression of IGFBP3 is regulated by HOXD10 gene [Xue et al., 2013; Pan et al., 534 2021], of which expression has been found with 2.9-fold rise in our study. *Hoxd10* is involved in prostate cancer 535 processing, cell differentiantion and morphogenesis [Jonkers et al., 2020; Mo et al., 2017; Zhang et al., 2019].

Another gene with 2.2-fold expression rise is *Slc45a3*, a well-known marker of prostate cancer progression, when a loss of the protein determined poorer prognosis for cancer patients [Hernández-Llodrà *et al.* 2017; Pin *et al.*, 2017; Perner *et al.*, 2013]. An overexpression of another gene identified in our study (12fold rise for *Usp39* gene) is linked with promotion of prostate or ovarian tumorigenesis. *Usp39* is also involved in pre-mRNA splicing, cytokinesis and spindle checkpoint working, and is crucial for apoptosis [Huang *et al.*, 2016; Yuan *et al.*, 2021; Zhao *et al.*, 2016].

Also, 2.2-2.5-fold rise for ncRNA genes (*Gm26848*, *Gm45648*; no data concerning their role) and 2-fold reduction for one pseudogene (*Gm28661*; also no data) were found (Figure 7). Following the latest data (November 2021) from Gene NCBI database, the Gm45648 has been annotated as one with the testisrestricted expression in mouse but no function has been suggested. The second ncRNA *Gm26848* found is

19

suggested as protein coding organic cation transporter-like protein across membranes in *Drosophila sechellia*. There were also two other genes with expression change (3.6-fold decrease – pseudogene *Gm44126*, and 2.5fold increase – *4933430L12Rik* probably involved in antisense transcription). 'Gene' database data showed that the *Gm44126* in mice is rather a protein coding gene with the function of UDP-glucuronic acid decarboxylase, and contains two domains involved in NAD(P)H processing. Thus, it can be suggested that this gene is as a novel candidate involved in mitochondrial respiratory chain. Our data with 3.6-fold expression decrease of this gene seem to fit into observed diminished functioning of mitochondrial machinery in *Tcte1*^{-/-} mice.

553 Another three genes manifested 2-fold decrease of their testis expression when comparing $Tcte1^{-r}$ vs. 554 Tcte1^{+/-} (but not to WT): Agp2 (aquaporin-2 involved in osmotic transportation of the water), with well-known 555 role in spermatogenesis, female endometrium cycle and urinary incontinence; [Klein et al., 2013; Ramli et al., 556 2019], Mt3 (metallothionein-3, committed in homeostasis and detoxification of toxic metals, also in testis) 557 important in apoptosis regulation of Bax and Caspase pathways [Gao et al., 2021; Honda et al., 2010]), and 558 Rps27rt (ribosomal protein) (Figure 5; Supplementary file 12). It was also found that Rps27rt may interact 559 indirectly with Usp39 and mitochondrial respiratory complex linking the mitochondrial energy processing with 560 apoptosis (Supplementary file 12). Last gene – Ube2l6, was found to have 2-fold increased expression level in testis in Tcte1^{+/-} animals vs. Tcte1^{-/-} and WT. Its role has been established as ubiquitin enzyme engaged in: 561 562 stress damage response, cell cycle progression, embryo development, immune response, and factors linked to 563 ATP machinery (Figure 7) [Huang et al., 2016; Yuan et al., 2021; Zhao et al., 2016; Sandy et al., 2020; Cai et al., 564 2021]. Increased expression of Ube216 was linked to degradation/suppression of cancer cells and affected 565 downstream apoptotic factors (i.e., Caspase 3, Caspase 9, Bcl-2, Bax) [Li et al., 2018].

566 Immunofluorescence on sperm

567 An example of immunofluorescence in situ staining on mouse spermatozoa used to detect the N-DRC 568 proteins has been presented in Figure 9. WT mouse spermatozoa showed Tcte1 protein presence in the sperm head nucleus, midpiece, and a part of the tail (Figure 9A). Immunofluorescent signal of Tcte1 in Tcte1^{+/-} mice 569 was similar to WT animals, while in $Tcte1^{-r}$ was localized only in the nucleus (Figure 9A). It seems to confirm 570 571 that in Tcte1^{-/-} males the protein has not been produced properly, revealing only residual amounts in sperm 572 head nucleus, and not transported to sperm flagella. The other N-DRC components (Drc7, Fbxl13 and Eps8l1) 573 were localized in the sperm nucleus and among the whole sperm tail (Fbx|13, Eps8|1) or only within its 574 fragment (Drc7), with exception of the midpiece (Figure 9B). Results obtained seem to suggest co-localization

20

of evaluated N-DRC proteins in various combinations, not common among the midpiece and the terminal partof the sperm tail.

577 Sperm ATP level

578 Measurements performed in two time points (0 and +60 min.) revealed that in $Tcte1^{+/-}$ samples the 579 luminescence level was approximately 2.4-fold lower when compared to WT and $Tcte1^{+/-}$ mice (p<0.0001) 580 (Figure 10; Supplementary file 14). There were no differences between WT and $Tcte1^{+/-}$ males. Additionally, the 581 decrease of the luminescent signal after 60 min. was similar in all three groups studied – approximately by 35% 582 (WT: 35.27%, $Tcte1^{+/-}$: 33.88%, $Tcte1^{-/-}$ 37.76%).

583 Apoptosis in testicular tissue

Evaluation of the immunofluorescent staining results revealed Casp3-positive signals (indicating apoptosis) in spermatogonia only, in all three mouse genotypes (Figure 11A). In each zygosity the numbers of tubules and positively stained spermatogonia (intensity of the signal) were similar suggesting equal levels of apoptosis. Also lack of differences between zygosities were found in case of TUNEL assay, for all testicular cell types (Figure 11B). Apoptosis-positive signals were observed in all tubules for both techniques.

589 Sequencing – human male samples

590 Potentially associated/important 6 heterozygous SNVs for TCTE1 gene has been found in 15 out of 248 591 (6.05%) infertile males with drastically decreased semen parameters, ranging from azoospermia to severe 592 oligoasthenozoospermia (Table 1). 3 variants were identified with ultrarare allele frequency (MAF \leq 0.0005 in 593 the general population; gnomAD) in three males with NOA: c.862C>T (p.Arg288Ter) predicted as stop-gained 594 mutation leading to non-mediated decay, and thus to protein loss of function with high probability; c.185G>A 595 (p.Arg62His), or c.1048G>A, (p.Gly350Ser), defined as missense ones and probably damaging. Next two SNVs 596 were found as the rare missense ones with MAF <0.005: c.397C>T (p.Arg133Cys) (one NOA case) and c.470A>C 597 (p.Glu157Ala) (four NOA cases), and were defined as probably damaging. The last 1 SNVs (1 NOA, 1 crypto-, 598 and 5 with severe oligoasthenozoospermia) was not found in databases, and thus specified as the novel one 599 (c.374T>G; p.Ile125Arg) with missense prediction (Table 1; Supplementary file 15).

600 Protein prediction

To understand the molecular implications of identified mutations on the structure and function of TCTE1, we performed homology modeling experiments using the Phyre2, I-TASSER, and AlphaFold tools [Kelley *et al.*, 2015; Roy *et al.*, 2010; Yang *et al.*, 2015; Yang and Zhang 2015; Tunyasuvunakool *et al.*, 2021]. TCTE1 is

21

604 predicted to contain at least five tandem leucine-rich repeats (LRR) in its C-terminus [UniProt 2019], while no 605 known three-dimensional protein folds have been assigned to its N-terminus. All homology searches yielded a 606 polypeptide with tandem leucine-rich repeats with nearly 100% confidence (Figure 12A; Supplementary file 607 16). Modeling the N-terminal region of TCTE1 was more difficult; although both Phyre2 and I-TASSER 608 confidently predicted a largely α -helical secondary structure for the N-terminal half of TCTE1 (Figure 12A; 609 Supplementary file 16), neither tertiary structure was predicted with high confidence; indeed, comparison of 610 the two demonstrates vastly different folds for the N-terminus from the programs. However, the confidence of 611 the AlphaFold-generated model was high for the region just N-terminal to the LRR. Further, based on the 612 predicted aligned error plot for calculated for its models, the confidence of the predicted structure of residues 613 100-200 was high relative to the position of residues 220-480, providing support for the modeled through-614 space three-dimensional interactions. That said, we interpreted these models very conservatively.

615 We attempted to hypothesize how mutations of interest could affect the tandem LRRs and putative 616 secondary structure of the N-terminus of the protein. Residue R64 resides in a predicted unstructured region 617 and as such most likely is solvent exposed in the tertiary structure (Figure 12A, B). Mutation of this residue 618 from arginine to histidine could therefore affect the surface potential of this disordered region, making the it 619 less positively charged. Indeed, this mutation would be expected to impact the interactions that the N-620 terminus of TCTE1 can have with other molecules, including other members of the N-DRC. The interaction of 621 microtubule-associated proteins, including the N-DRC, with binding partners is known to be largely charge-622 based [Kubo et al., 2017; Drechsler et al., 2019].

623 Three mutations of interest occur in the globular region just N-terminal to the LRR. The AlphaFold 624 model clearly shows how mutation 125R would affect the structure of this region Figure 12A, C). Residue 625 isoleucine 125 is predicted to be nestled within the hydrophobic core including L97 L109, L112, F91, W131, 626 L197, Y130, and the methyl group of the T121 sidechain. A substitution of positively charged arginine for 627 isoleucine in this position would not be accommodated in this hydrophobic region, likely resulting in local 628 unfolding that could potentially propagate beyond the pocket. Likewise, residue R133 supports the structure of 629 this putative globular domain by forming hydrogen bonds with the backbone of residues 109, 110, and/or 112, 630 thereby possibly contributing to the overall organization of this domain and its positioning relative to the LRR 631 (Figure 12D). Mutation of this residue to a cysteine would remove the hydrogen bonding groups supporting 632 these interactions, likely leading to a local unfolding that could possibly propagate farther. Interestingly,

22

residue E157 is predicted to interact with residues in the LRR, likely stabilizing positively charged residues around it, including several from the linker leading up to the first β-strand of the LRR (Figure 12B, E). Mutation of this residue to an alanine would lead to a tertiary destabilization, most likely affecting the overall structure of the N-terminal region of the LRR and the globular region just N-terminal to it.

637 Two mutations of interest affect the LRR helices. Mutation R288* results in premature truncation, 638 eliminating most of the LRR region from the final protein product, although it is predicted that this premature 639 termination codon would lead to nonsense mediated decay. Mutation G350S would be expected to result in 640 the destabilization of inter-repeat interactions (Figure 12F), possibly even unfolding the protein from that 641 position on; the lack of a side chain in glycine allows for the close approach of these tandem structural 642 elements, whereas the addition of the serine side chain would abrogate the tight packing of residues, 643 regardless of which rotamer it adopts. This could result in a restructuring its C-terminus and/or deforming the 644 surface of the module, therefore affecting its interactions with its binding partners.

645 Since the N-DRC is conserved from single-celled eukaryotes to mammals, we took advantage of 646 published cryo-electron tomography studies examining the N-DRC from model organism Chlamydomonas [Gui 647 et al., 2019]. This study was of great interest since it explicitly located the position of the C-terminus of DRC5 648 with a nanogold particle; importantly, it was consistent with earlier cryo-ET experiments positioning the N-DRC 649 [Heuser et al., 2009]. While the resolution of this structure is only 40Å, it was still useful for placing the 650 modeled TCTE1 structure in the context of the entire N-DRC (Supplementary file 16B). Manual docking of the 651 AlphaFold predicted TCTE1 structure is consistent with the cryo-EM tomography data [Gui et al., 2019; Heuser 652 et al., 2009].

653

654 **DISCUSSION**

In this study we have investigated the role of *Tcte1* in male infertility using mouse knockout model. Using homo- and heterozygous animals, their reproductive potential has been examined, noting no progeny by homozygous males. The size of the gonads varied between the zygosity of animals, followed by two semen phenotypes revealed within the sperm quality evaluation. RNAseq analysis of the mutant testicular tissue demonstrated the influence of *Tcte1* mutations on the expression pattern of genes related to mitochondrial cytochrome oxidase activity, apoptosis and spermatogenesis. The ATP level estimate of spermatozoa showed its decrease in homozygous males. Additionally, protein prediction modeling of identified variants in human

23

samples revealed changes in the protein surface charge potential, leading to disruption in helix flexibility or dynamics, and thus, suggesting the disrupted TCTE1 interaction with its binding partners within the axoneme. The major findings concerning observed abnormal sperm tail beating in KO *Tcte1* mice were summarized in Figure 13. We have also identified and discussed novel functions of genes with testicular expression significantly disturbed in the *Tcte1* knockout. Therefore, this study contains novel and the most complete data concerning *TCTE1* role in male infertility, so far.

668

Phenotypes of homozygotes and heterozygotes

669 There was only one published study so far, concerning knockout mouse model of *Tcte1* [Castaneda et 670 al., 2017]. It was shown that the expression of *Tcte1* starts at the stage of early haploid round spermatids and 671 continues into the elongated spermatids. Authors have shown the infertility of homozygous males due to 672 asthenozoospermia derived from limited and circular flagellar beating, followed by fertile heterozygotes. The 673 study did not find any differences between homozygous and WT males in the testicular histology, followed by 674 similar testis weight, sperm concentration and sperm morphology [Castaneda et al., 2017]. We also have not 675 found any structural differences in the testicular and epididymal tissues histology. However, the weight of the 676 testes decreased significantly (~41% in $Tcte1^{-4}$, 7% in $Tcte1^{+4}$), followed also by testis size reduction (by 22%) 677 and lowered germ cell layer thickness in $Tcte1^{-L}$ animals (Figure 4B, C). When comparing semen parameters 678 (Figure 6A-C), we have found enormous decline in sperm concentration in both: $Tcte1^{-4}$ (3.3-fold) or $Tcte1^{+4}$ 679 (2.5-fold) males, and double increase in abnormal sperm morphology in Tcte1^{-/-} animals. Similarly, to Castaneda 680 et al. [2017], we have documented the impaired sperm motility representing circular mode of the sperm tail 681 beating. Also, the increased ratio of immotile spermatozoa in homozygous males was documented in our 682 study. We can suggest that these discrepancies may result from various number of animals used for evaluation: 683 3 males in study of Castaneda et al., [2017] vs. at least 18 per each zygosity in our study, when comparing 684 semen parameters and tissue characteristics (except of sperm morphology evaluation with at least 5 animals 685 used). Probably, also the differences in knockout construction used could be causative for observed variances: 686 reporter-tagged insertion with conditional potential vs. the exon 3 deletion. Thus, we have observed two 687 various phenotypes dependent on the zygosity of the animals: homozygous males manifested 688 oligoasthenoteratozoospermia (OAT) and were infertile, while heterozygous males demonstrated 689 oligozoospermia (O) and remained fertile. Data collected in International Mouse Phenotypic Consortium (IMPC; 690 a repository of all available mouse phenotypes created, followed with the possible linkage to human disease

24

691 defects; <u>https://www.mousephenotype.org</u>), concern only one *Tcte1* mouse study [Castaneda *et al.*, 2017],

692 while we reveal an unmatched phenotypes of decreased sperm count.

693 We suggest that similar effect – decreased sperm count linked to TCTE1 mutations, could be observed 694 in humans. Our pilot mutation screening of 248 human infertile males with severely decreased sperm count 695 (from oligo-, via crypto-, to azoospermia) revealed 6 heterogeneous SNVs (1 novel, 3 ultrarare, and 2 rare; all 696 predicted as disease causing) with the total frequency of 6.05% (Table 1). We propose that molecular 697 epidemiology studies of infertile patients should reveal the existence of homozygous or a compound 698 heterozygous TCTE1 variants with clear direct downgrading effect on spermatogenesis. However, we claim that 699 also heterozygous variants can be causative, similarly to SPINK2 which deficiency has been documented as 700 responsible for oligoasthenozoospermia in heterozygous mice vs. azoospermia in homozygous animals, 701 followed by the azoospermia in human homozygous brothers vs. their heterozygous father with 702 oligozoospermia [Kherraf et al., 2017]. It can be explained by the probable haploinsufficiency occurrence, when 703 the single copy of the WT allele in heterozygous combination of which a variant allele is insufficient to produce 704 the WT phenotype. Our observations, followed by the SPINK2 data and data related to infertility (i.e.: P1/P2 705 protamines, Pde8b, Trim28, Tcfl5), and to the other diseases (i.e.: PIKFYVE in cataract, NFIA, SCN1A in 706 neurological or epilepsy disturbances, KLF1 in hemoglobin persistenece, NR5A2 in pancreatic cancer), seem to 707 confirm that the severity of the phenotype depends on the gene expression level [Jodar and Oliva, 2014; Mei et 708 al., 2022; Ogura et al., 2022; Heshusius et al., 2022; Valassina et al., 2022; Sandhu et al., 2021; Leal et al., 2021; 709 Tan et al., 2020; Xu et al., 2022]. We have documented variable phenotypes for a novel TCTE1 variant c.374T>G 710 (p.lle125Arg) that has been documented in seven males, including: azoospermia, crypto-, and severe 711 oligoasthenozoospermics. Thus, it seems to suggest the reasonable candidacy of haploinsufficiency as the 712 mechanism responsible for observed phenotypic spectra also in case of the TCTE1 gene. There is an only one 713 study available documenting the TCTE1^{-/-} in human case, so far [Zhou et al., 2022]. Among 130 714 asthenozoospermic patients, authors have found frameshift mutation c.396-397insTC (p.Arg133Serfs*33) in 715 one of them that resulted in premature translational arrest of the forming peptide. Successful birth delivery 716 was available only after in vitro fertilization (IVF). What is important, authors also checked IVF rate in 717 homozygous mice, but without pregnancy success. The unresolved question is whether homozygous TCTE1 718 mutation in human causes only asthenozoospermia or rather the semen quality phenotype is mixed, including 719 also abnormal morphology, but this was not evaluated by the authors.

25

720 Kallikreins, Aquaporin 2

721 Our pathway analysis revealed a network including two genes with changed expression level in testis: 722 Igf2 (2-fold increased) and Klk1b22 (8-fold decreased), supported by an interesting set of proteins linked to 723 apoptosis and cellular processing (Figure 7; Figure 8; Supplementary file 12; Supplementary file 13). We can 724 suggest that observed Igf2 overexpression in our study can be one of the associated factors leading to 725 decreased sperm count and testis parameters in evaluated KO mice model because of its known relation to 726 spermatogenesis and testis development [Matsuzaki et al., 2020; Cannarella et al., 2020; Neirijnck et al., 2019; 727 Damaschke et al., 2017]. Documented reductions can be supported by the observed constrained expression of 728 Klk1b22 – kallikrein mostly known from its neuronal activity; however, it cannot be excluded that we have 729 revealed additional role of this protein, similar to the other members of the kallikrein family - role in semen 730 liquefaction and degradation of extracellular matrix proteins in the interstitial area surrounding Leydig cells of 731 the adult mouse testes (Klk1b27, Klk1b21, Klk1b24) [Michaelis et al. 2017; Marques et al., 2016, Du Plessis et 732 al., 2013; Fink et al., 2007]. Notably, Klk1b22 expression is significantly enriched in testis. While expression of 733 all four kallikreins was decreased 2.5-8-fold in our *Tcte1* KO model, indicating their participation in the 734 reduction of sperm parameters measured (Fig.5).

735 We have also documented 2-fold decrease of expression of Aqp2 (Aquaporin 2), involved in fluid 736 resorption and water movement within the duct system, with known role in spermatogenesis [Klein et al., 737 2013; Ramli et al., 2019]. It was documented that in stallions Aqp2 is being expressed in Leydig cells, round and 738 elongated spermatids [Klein et al., 2013]. Also, this protein works within a network of molecules that are linked 739 to cell membrane trafficking (i.e., Rab11a, Myo5b), followed by activity of protein kinases (Prkacb, Prkaca) that 740 are known as regulators of cell proliferation, chromatin condensation/decondensation, and nuclear envelope 741 disassembly/reassembly, and all is supported by the ATP metabolism (Hspa8) [Supplementary file 12] [Markou 742 et al., 2022; Wagner et al., 2022]. Considering literature data, we suggest Aqp2 play an important role in 743 reduction of the spermatids volume during spermatogenesis and in proper sperm morphology assessment and 744 observed morphological differences between Tcte1 mice zygosities in our study. However, such measurements 745 require high-resolution microscopy, thus, we can only speculate Aqp2 role in our model.

746 Disruptions in sperm cell structure

747 TCTE1 protein contains five leucine rich repeats (LRRs) that are known to be directly involved in 748 receptor-mediated signaling on the structural basis for interaction between proteins [Ng *et al.*, 2011; Kobe and

26

749 Kajava 2001]. It was documented that variants in genes encoding LRRs are commonly implicated in the 750 pathogenesis of more than 60 human diseases [Ng et al., 2011; Matsushima et al., 2019]. LRR mutations mostly 751 occur within regions that are protective for the hydrophobic core of the LRR domains. Thus, mutations in LRRs 752 may affect the charge of the surface potential, leading to conformational changes of the protein or induction of 753 protein cumulation. Our prediction model of potential changes in TCTE1 protein resulting from mutations 754 found in human samples seems to point out the prospective impact into the interactions of the TCTE1 with 755 other molecules (binding partners) known to be largely charge-base, including N-DRC elements or microtubule-756 associated proteins [Kubo et al., 2017; Drechsler et al., 2019]. In the case of TCTE1 protein observed amino acid 757 substitutions promote the abnormal aggregation of TCTE1 protein, and decrease its structural stability, then 758 probably leading to the observed disturbances. Beating motion of flagella is maintained by electrostatic cross-759 bridge between negatively charged tubulins and positively charged N-DRC [Kubo et al., 2017]. Thus, changes in 760 the charge potential of TCTE1 surface, which is one of the N-DRC components, seem to provide modified 761 (reduced) flagellar beating, similarly as in case of mutations in other N-DRC element – DRC4 protein, of which 762 changes in charge were essential for proper flagellar motility [Kubo et al., 2017; Lewis et al., 2016]. Also, the 763 inflexibility of the midpiece observed in spermatozoa of Tcte1 knockout mice seems to be a consequence of 764 changed structure of LRRs in Tcte1 protein.

765 It is also known that the lengths of the sperm midpiece and tail may be crucial in determination of 766 sperm swimming velocity, followed then by decreased fertilization rate [Firman and Simmons 2010; Gu et al., 767 2019]. Available data showed that sperm with shorter midpiece may swim slower and were observed in 768 asthenozoospermic cases [Firman and Simmons 2010; Gu et al., 2019; Anderson et al., 2002]. The meta-769 analysis of sperm shape features according to sperm competition theory has also shown, that in majority of 770 vertebrates the longer sperm cell (including midpiece or flagellum) has been attributed to faster swimming, 771 followed by the statement that the sperm size may be increased when sperm number is lowered [Lüpold et al., 772 2020]. In our study we have documented longer sperm tails and midpleces in Tcte1^{-/-} animals, that in fact were 773 infertile. Thus, we can suggest that: (i) the observed longer sperm tails and midpieces may be co-responsible 774 for aberrant motility observed, and (ii) the documented longer sperm tails and midpleces may be an indirect 775 result of constrained sperm count observed, because of some sort of compensation for aberrant motility. 776 However, this issue remains only a suggestion because of a restricted literature data relative to similar 777 observations. Bennison et al. [2016] revealed than in a passerine bird the sperm tail velocity was correlated to

27

total sperm length, but only to some point of the length value at which the velocity started to be decreased. On
the other hand, Gu et al. [2019] have shown that in mammalian species there were some changes in the
axonemal slopes when the flagellar length increases. The results obtained within our study, and supported by
other research concerning sperm tail beating pathways, should be a good start point for creation of further
mathematical computations for better understanding of the asthenozoospermia.

783 Constrained energy metabolism

784 The flagellar motility is determined by coordinated activity of asymmetrically distributed dyneins in 785 the axoneme. The driving factor for this activity is the hydrolysis of ATP released from OXPHOS system 786 (Mitochondrial Oxidative Phosphorylation System) [Tourmente et al., 2015; Ford 2006]. ATP is being produced 787 in mitochondria placed within sperm midpiece. In our study ATP level measurements revealed its decreased 788 level in sperm cells of Tcte1^{-/-} mice, what can result in faulty dynein functioning, leading to asthenozoospermia 789 via abnormal sperm tail beating. We have documented that Tcte1 KO resulted in significant changes in 790 expression level of testicular genes involved in mitochondrial energy transport. That leads to the conclusion 791 that homozygous *Tcte1* mutation results not only in inflexible midplece leading to asymmetric circular motility, 792 as presented in this study and by others [Fung et al., 2018], and then to asthenozoospermia, but also to 793 aberrant functioning of mitochondrial OXPHOS machinery leading to disruption of ATP production. Thus, the 794 sperm tail has also insufficient power for beating because of the decreased ATP content. Castaneda et al. 795 suggested that the reasons of the observed ATP decrease may result from disturbances in a glycolytic pathway 796 [Castaneda et al., 2017]. It is already known, that glycolytic energy is enough for spermatozoa to survive 797 [Nascimento et al., 2008; Zhang et al., 2022]. However, for sperm motility and then fertilization, the OXPHOS 798 energy production is required [Tourmente et al., 2015; Ford 2006; Zhang et al., 2022]. In our study, RNAseq 799 results clearly revealed the influence of *Tcte1^{-/-}* mutation on genes involved directly in electron transport chain, 800 such as: mt-Co2 and mt-Co3, crucial components of the respiratory chain complex IV, one of the OXPHOS 801 elements. Observed reduced expression seems to clearly evidence the decreased ATP production via disturbed electron chain transportation, and then revealing the sperm motility reduction observed in Tcte1^{-/-} males, 802 803 leading to observed asthenozoospermia. Detailed STRING analysis revealed definite interactions between both 804 listed proteins identified in our study, and a list of other mitochondrial ones (mt-Co1, mt-Cytb, Cox5a, Cox6a, 805 Cox7c, mt-Atp6; Supplementary file 12; Supplementary file 13) clearly related to proper processing of 806 cytochrome energetic activity [Brischigliaro and Zeviani 2021; Ganetzky et al., 2019]. Overexpression or

28

807 mutations in COX1, COX2, and COX3 were previously observed in infertile men with varicocele and OAT, or 808 diabetes [Heidari et al., 2016; Mostafa et al., 2016; Perrotta et al., 2012]. Additionally, it was shown that 809 system of COX2 and prostaglandins plays key role in development of male gonad, spermatogenesis and 810 steroidogenesis, especially in functioning of Sertoli and Leydig cells, and also was found to be significant in the 811 background basis of idiopathic infertility status [Frungieri et al., 2015]. It seems that in our study the reason of 812 observed disturbed motility, represented as circular movement of spermatozoa, supported by increased ratio 813 of immotile sperm cells in Tcte^{-/-} mice, leading to asthenozoospermia, resulted from two factors: (i) inflexibility 814 of sperm midpiece resulted from changed protein charge level in the Tcte1 protein, followed by elevated 815 lengths of sperm tail and midpiece, and (ii) decreased amount of ATP produced, that may be an outcome of 816 inability to get sufficient energy for proper sperm motility.

817

818

819 Protection and/or compensation?

820 We have also documented an important role of the *Fetub* gene for the first time in the male testis. 821 Besides of its known role in female fertility [Karmilin et al., 2019; Bansho et al., 2017], the latest study pay 822 attention also in the context of prostate cancer is a detailed way [Zhan et al., 2020]. It was found that 823 overexpression of FETUB increases the apoptosis of prostate cancer cells, followed by the inhibition of their 824 proliferation, migration and invasion, and also slower growth of tumor when compared to controls [Zhan et al., 825 2020]. Thus, in the overexpression mode, the Fetub plays protective role for the prostate. It seems to explain results obtained in our study, where the Fetub expression was increased almost 5-fold in Tcte1^{-/-} mice, but no 826 827 tumor signs have been observed in the mouse testis at the tissue level. Zhan et al. [2020] have found the 828 connection between Fetub overexpression, and induction of Bax (Bcl-2-associated X protein) and capases, 829 resulting in promoting the apoptosis. It is known that the Bax/Bcl-2 induction/inhibition is directly related to 830 the release of cytochrome c (Cyt c) in the caspase-dependent signaling [Dadsena et al., 2021; Samaiya et al., 831 2021; Li et al., 2021]. And cytochrome c is one of the two electron carriers in electron transport chain 832 responsible for transport directly to the respiratory chain complex IV [Yin and O'Neil 2021; Alvarez-Paggi et al., 833 2017]. Due to Cyt c release, complex IV is essential for apoptosis and cell fate [Samaiya et al., 2021; Li et al., 834 2021]. Following of all of those facts, our results seem to reveal the connection between Fetub and diminished 835 mitochondrial chain IV activity in mice testis, what have not been reported so far. It can be also supported by

29

the fact of the known relation between FETUB and the inactivation of the PI3K/AKT signaling pathway (key
player in regulation of cell proliferation and apoptosis in cancerogenesis), in which the Bax/Bcl-2 are involved
[Zhan *et al.*, 2020].

839 The novel testis-protective role in our study can be also credited to three other genes with increased 840 expression level in testicular tissue: Hoxd10 (3-fold increase), Slc45a3 (2.2-fold), and Ube21 (2-fold). 841 Overexpression/activation of HOXD10 has been already documented as inhibitor of tumorigenesis [Jonkers et 842 al., 2020; Mo et al., 2017; Zhang et al., 2019]. Expression of Slc45a3, known also as 'prostein', has been 843 documented as weaker in cancer tissues [Hernández-Llodrà et al. 2017; Pin et al., 2017; Perner et al., 2013]. 844 Elevated expression level of Ube21 is known to be connected with the degradation/suppression of tumor cells 845 [Li et al., 2018]. On the other hand, the 12-fold increased gene expression level for Usp39 (ubiquitin specific 846 peptidase 39) - an oncogenic splicing factor which expression is positively linked with cancerogenesis' 847 promotion in prostate or ovaries, while its silencing induces apoptosis and cell cycle arrest, has been noticed 848 [Huang et al., 2016; Yuan et al., 2021; Zhao et al., 2016].

849 Taken together data obtained for genes related to cancerogenesis and/or apoptosis described above, 850 we can hypothesize that we observe an interesting novel network of interactions between two groups of genes 851 that are reciprocally neutralizing potential influence onto the testis tissue. Mainly, in the first group we have 852 genes with suggested novel anti-cancer role in the testis (Fetub, Hoxd10, Slc45a3, Ube2l6), which increased 853 expression level inhibits the cancer cell development, supported by protective role of Mt3; while in the second 854 corner of the ring the Usp39 gene with elevated expression level (the most from all the observed changes) 855 should lead to reveal some tumor-like changes within the testis tissue. Following the fact, that both: 856 histological evaluation, as well as apoptosis stainings (Fig. 11), have shown no differences between hetero- and 857 homozygous mutant mice, our hypothesis seems to be justifiable. On the other hand, there is a prediction that 858 many of the genes (especially so-called cancer-testis associated antigens, CTA) related to spermatogenesis may 859 function in tumorigenesis [Aurriere et al., 2021; Yang et al., 2020; Babatunde et al., 2017; Wang et al., 2016; 860 Bode et al., 2014; Cheng et al., 2011; Nagirnaja et al., 2018]. Due to known similarities between oncogenic and 861 spermatogenic processes, especially in the energy metabolism, we can carefully suggest/hypothesize that in 862 our KO model the genes with changed expression profile found, known as highly involved in tumorigenesis, 863 may also be important in spermatogenesis. Obviously, there is a need for further studies in this direction to 864 confirm these suggestions or to reveal the other (still unknown) mechanisms.

30

865 Conclusions

866 We have documented that TCTE1 gene is the next one that should be added to the 'male infertility list' 867 of molecular background because of its crucial role in spermatogenesis and proper sperm functioning. The 868 mutations in mouse Tcte1 knockout model revealed two phenotypes, depending from zygosity with infertile 869 oligoasthenoteratozoospermic homozygotes, and fertile oligozoospermic heterozygotic males, suggesting 870 haploinsufficiency mechanism involvement. All collected data pointing that the Tcte1 protein is functioning in 871 spermatogenesis and show how its mutations add the level of complexity to the clinical manifestation of 872 decreased semen parameters. The disrupted energy machinery of the spermatozoa, followed by newly 873 suggested roles of genes related to cancerogenesis and/or apoptosis (here in the context of their 874 neutralizing/compensation effect on the testicular tissue), seem to underline the complex and devastating 875 effect of the single-gene mutation, on the variety of testicular molecular networks, leading to reproductive 876 failure. Further advanced studies are required to reveal the details of mechanisms responsible for the 877 molecular pathways of the interactions between Tcte1 (the structural protein) and other non-structural 878 networks of which expression have been changed remarkably in this study.

879

880 ACKNOWLEDGEMENTS

We are grateful to Oliwia Kordyl, MSc, and Katarzyna Kurek, MSc (IHG PAS, Poznan) for their technical help in preparation of the samples for sequencing, genotyping, and immunofluorescence staining.

883

884

885 Authors' contributions:

886 MO design of the study, manuscript drafting and editing, coordination of the tasks, data interpretation, gene 887 selection, preparation of human blood and semen samples, mice: mating, genotyping, gross sections and 888 preparation of biological samples, RNAseq analysis, immunofluorescence, Sanger sequencing, ATP 889 measurements; AM gene selection, data interpretation, genotyping, RNAseq analysis, collection and 890 preparation of human testis samples, WGS and Sanger sequencing, manuscript editing; TS bioinformatic 891 analysis of RNAseq and WGS results; NP WES of azoospermia samples (incl. bioinformatics); AB protein 892 prediction modeling and its description; SB preparation of: DNA for sequencing, RNA for RNAseq, screening of 893 databases for RNAseq results; MKa collection of human blood and semen samples, semen analyses of human

31

- and mouse samples, detailed morphology of mouse spermatozoa; JS-Z and HJ histopathology of mouse tissue
- sections; PJ recruitment of patients and their medical history; ANY gene selection, interpretation of
- azoospermia WES results, data interpretation; MKu data interpretation, gene selection, recruitment of patients
- and their medical history, correction of the manuscript, funds collection and supervision.
- All authors critically reviewed and approved the final version of the manuscript.
- 899
- 900 LARGE SCALE DATA: All data generated or analyzed during this study are included in this published article and
- 901 its supplemental information files.
- RNAseq data are available in GEO database (<u>https://www.ncbi.nlm.nih.gov/geo/</u>) under the accession number:
 GSE207805.
- 904 The results described in the publication are based on whole-genome or exome sequencing data which includes
- 905 sensitive information in the form of patient specific germline variants. Information regarding such variants
- 906 must not be shared publicly following the European Union legislation described in the following document:

907 https://publications.jrc.ec.europa.eu/repository/bitstream/JRC113479/policy_report_-

908 review of eu national legislation on genomics - with identifiers 1.pdf, therefore the access to raw data

- 909 that support findings of this study are available from the corresponding author upon reasonable request.
- 910 LIMITATIONS, REASONS FOR CAUTION: In the study, the *in vitro* fertilization ability of homozygous male mice
- 911 was not checked.
- 912 WIDER IMPLICATIONS OF THE FINDINGS: This study contains novel and the most complete data concerning
- 913 TCTE1 role in male infertility. TCTE1 gene is the next one that should be added to the 'male infertility list' of
- 914 molecular background because of its crucial role in spermatogenesis and proper sperm functioning.
- 915 STUDY FUNDING/COMPETING INTEREST(S): This work was supported by National Science Centre in Poland,
- 916 grants no.: 2015/17/B/NZ2/01157 and 2020/37/B/NZ5/00549 (to MK), 2017/26/D/NZ5/00789 (to AM), and
- 917 HD096723, GM127569-03, NIH SAP #4100085736 PA DoH (to AY). The authors declare that there is no conflict
- 918 of interest that could be perceived as prejudicing the impartiality of the research reported.

920 **REFERENCES**

- 922 Agarwal A, Baskaran S, Parekh N, Cho CL, Henkel R, Vij S, Arafa M, Panner Selvam MK, Shah R. Male infertility.
- 923 Lancet 2021;397:319-333 https://doi.org/10.1016/S0140-6736(20)32667-2
- 924 Alvarez-Paggi D, Hannibal L, Castro MA, Oviedo-Rouco S, Demicheli V, Tórtora V, Tomasina F, Radi R, Murgida
- 925 DH. Multifunctional Cytochrome c: Learning New Tricks from an Old Dog. *Chem Rev* 2017;**117**:13382-13460
- 926 <u>https://doi.org/10.1021/acs.chemrev.7b00257</u>
- 927 Anderson MJ, Dixson AF. Sperm competition: motility and the midpiece in primates. *Nature* 2002;**416**:496
- 928 <u>https://doi.org/10.1038/416</u>496a
- 929 Aprea I, Raidt J, Höben IM, Loges NT, Nöthe-Menchen T, Pennekamp P, Olbrich H, Kaiser T, Biebach L,
- 930 Tüttelmann F. et al. Defects in the cytoplasmic assembly of axonemal dynein arms cause morphological
- abnormalities and dysmotility in sperm cells leading to male infertility. *PLoS Genet* 2021;17:e1009306
- 932 https://doi.org/10.1371/journal.pgen.1009306
- Aurrière J, Goudenège D, Baris OR, Boguenet M, May-Panloup P, Lenaers G, Khiati S. Cancer/Testis Antigens
- 934 into mitochondria: a hub between spermatogenesis, tumorigenesis and mitochondrial physiology adaptation.
- 935 Mitochondrion 2021;56:73-81 https://doi.org/10.1016/j.mito.2020.11.002
- Babatunde KA, Najafi A, Salehipour P, Modarressi MH, Mobasheri MB. Cancer/Testis genes in relation to sperm
- 937 biology and function. Iran J Basic Med Sci 2017; 20:967-974 https://doi.org/10.22038/IJBMS.2017.9259
- 938 Baklouti-Gargouri S, Ghorbel M, Mahmoud AB, Mkaouar-Rebai E, Cherif M, Chakroun N, Sellami A, Fakhfakh F,
- 939 Ammar-Keskes L. Mitochondrial DNA mutations and polymorphisms in asthenospermic infertile men. *Mol Biol*
- 940 Rep 2013;40:4705-4712 https://doi.org/10.1007/s11033-013-2566-7
- Bansho Y, Lee J, Nishida E, Nakajima-Koyama M. Identification and characterization of secreted factors that are
- 942 upregulated during somatic cell reprogramming. *FEBS Letters* 2017;**591**:1584-1600
 943 https://doi.org/10.1002/1873-3468.12665
- 944 Barratt CLR, Björndahl L, De Jonge CJ, Lamb DJ, Martini FO, McLachlan R, Oates RD, van der Poel S, St John B,
- 945 Sigman M, et al. The diagnosis of male infertility: an analysis of the evidence to support the development of
- 946 global WHO guidance-challenges and future research opportunities. Hum Reprod Update 2017;23:660-680
- 947 <u>https://doi.org/10.1093/humupd/dmx021</u>

- 948 Bennison C, Hemmings N, Brookes L, Slate J, Birkhead T. Sperm morphology, adenosine triphosphate (ATP)
- 949 concentration and swimming velocity: unexpected relationships in a passerine bird. Proc R Soc B
- 950 2016;**283**:20161558 <u>https://doi.org/10.1098/rspb.2016.1558</u>
- 951 Binsila BK, Archana SS, Ramya L, Swathi D, Selvaraju S, Shivakumar Gowda NKS, Pal DT, Rafay A, Bhattaet R.
- 952 Elucidating the processes and pathways enriched in buffalo sperm proteome in regulating semen quality. Cell
- 953 Tissue Res 2021;383:881-903 https://doi.org/10.1007/s00441-020-03303-9
- 954 Bode PK, Thielken A, Brandt S, Barghorn A, Lohe B, Knuth A, Moch H. Cancer testis antigen expression in
- 955 testicular germ cell tumorigenesis. *Modern Pathol* 2014;**27**: 899-905
- 956 https://doi.org/10.1038/modpathol.2013.183
- 957 Bower R, Tritschler D, VanderWaal Mills K, Heuser T, Nicastro D, Porter ME. DRC2/CCDC65 is a central hub for
- assembly of the nexin-dynein regulatory complex and other regulators of ciliary and flagellar motility. *Mol Biol*
- 959 *Cell* 2018;**29**:137-153 https://doi.org/10.1091/mbc.E17-08-0510
- Brischigliaro M, Zeviani M. Cytochrome c oxidase deficiency. *BBA-Bioenergetics* 2021;**1862**:148335
 https://doi.org/10.1016/j.bbabio.2020.148335
- 962 Cai J, Huang H, Hu X, Lang W, Fu W, Xu L, Qiu Z, Zhong H, Chen F. Homoharringtonine Synergized with
- 963 Gilteritinib Results in the Downregulation of Myeloid Cell Leukemia-1 by Upregulating UBE2L6 in FLT3-ITD-
- 964 Mutant Acute Myeloid (Leukemia) Cell Lines. J Oncol 2021;2021:3766428
- 965 https://doi.org/10.1155/2021/3766428
- 966 Cannarella R, Condorelli RA, La Vignera S, Bellucci C, Luca G, Calafiore R, Calogero AE. IGF2 and IGF1R mRNAs
- 967 Are Detectable in Human Spermatozoa. World J Mens Health 2020;38:545-551
 968 https://doi.org/10.5534/wjmh.190070
- 969 Castaneda JM, Hua R, Miyata H, Oji A, Guo Y, Cheng Y, Zhou T, Guo X, Cui Y, Shen B, et al. TCTE1 is a conserved
- 970 component of the dynein regulatory complex and is required for motility and metabolism in mouse
- 971 spermatozoa. PNAS 2017;114:E5370-E5378 https://doi.org/10.1073/pnas.1621279114
- 972 Chemes HE, Olmedo SB, Carrere C, Oses R, Carizza C, Leisner M, Blaquier J. Ultrastructural pathology of the
- 973 sperm flagellum: association between flagellar pathology and fertility prognosis in severely asthenozoospermic
- 974 men. Hum Reprod 1998;13:2521-2526 https://doi.org/10.1093/humrep/13.9.2521

34

- 975 Chen Y, Wang J, Pan C, Li D, Han X. Microcystin-leucine-arginine causes blood-testis barrier disruption and
- 976 degradation of occludin mediated by matrix metalloproteinase-8. *Cell Mol Life Sci* 2018;75:1117-1132
- 977 https://doi.org/10.1007/s00018-017-2687-6
- 978 Cheng YH, Wong EWP, Cheng CY. Cancer/testis (CT) antigens, carcinogenesis and spermatogenesis.
- 979 Spermatogenesis 2011;1:209-220 <u>https://doi.org/10.4161/spmg.1.3.17990</u>
- 980 Chiang C, Layer RM, Faust GG, Lindberg MR, Rose DB, Garrison EP, Marth GT, Quinlan AR, Hall IM. SpeedSeq:
- 981 Ultra-fast personal genome analysis and interpretation. *Nat Methods* 2015;**12**:966-968
- 982 https://doi.org/10.1038/nmeth.3505.
- 983 Dadsena S, King LE, Garcia-Saez AJ. Apoptosis regulation at the mitochondria membrane level. BBA-
- 984 Biomembranes 2021;1863:183718 https://doi.org/10.1016/j.bbamem.2021.183716
- Damaschke NA, Yang B, Bhusari S, Avilla M, Zhong W, Blute ML, Huang W, Jarrard DF. Loss of Igf2 Gene
- 986 Imprinting in Murine Prostate Promotes Widespread Neoplastic Growth. Cancer Res 2017;77:5236-5247
- 987 https://doi.org/10.1158/0008-5472.CAN-16-3089
- 988 Drechsler H, Xu Y, Geyer VF, Zhang Y, Diez S. Multivalent electrostatic microtubule interactions of synthetic
- 989 peptides are sufficient to mimic advanced MAP-like behavior. *Mol Biol Cell* 2019;**30**:2953-2968
- 990 <u>https://doi.org/10.1091/mbc.E19-05-0247</u>
- 991 Du Plessis SS, Gokul S, Agarwal A. Semen hyperviscosity: causes, consequences, and cures. Front Biosci
- 992 2013;5:224-231 https://doi.org/10.2741/E610
- 993 Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of Coot. Acta Crystallogr D 2010;66:486-
- 994 501 <u>https://doi.org/10.1107/S0907444910007493</u>

995 Fink E, Bhoola KD, Snyman C, Neth P, Figueroa CD. Cellular expression of plasma prekallikrein in human tissues.

- 996 Biol Chem 2007;**388**:957-963 https://doi.org/10.1515/BC.2007.104
- 997 Firman RC, Simmons L. W. Sperm midpiece length predicts sperm swimming velocity in house mice. Biol Letters
- 998 2010;**6**:513-516 <u>https://doi.org/10.1098/rsbl.2009</u>.1027
- 999 Ford WC. Glycolysis and sperm motility: does a spoonful of sugar help the flagellum go round? Hum Reprod
- 1000 Update 2006;12:269-274 <u>https://doi.org/10.1093/humupd/dmi053</u>
- 1001 Frungieri MB, Calandra RS, Mayerhofer A, Matzkin ME. Correlation between testicular mast cell count and
- 1002 spermatogenic epithelium in non-obstructive azoospermia. *Reproduction* 2015;**149**:R169-R180
- 1003 <u>https://doi.org/10.1111/iep.12261</u>

- 1004 Fung E, Richter C, Yang HB, Schäffer I, Fischer R, Kessler BM, Bassermann F, D'Angiolella V. FBXL13 directs the
- 1005 proteolysis of CEP192 to regulate centrosome homeostasis and cell migration. EMBO Rep 2018;19:e44799
- 1006 https://doi.org/10.15252/embr.201744799
- 1007 Ganetzky RD, Stendel C, McCormick EM, Zolkipli-Cunningham Z, Goldstein AC, Klopstock T, Falk MJ. MT-ATP
- 1008 mitochondrial disease variants: phenotypic and biochemical features analysis in 218 published cases and
- 1009 cohort of 14 new cases. *Hum Mutation* 2019;**40**:499-515 <u>https://doi.org/10.1002/humu.23723</u>
- 1010 Gao Y, Hong J, Guo Y, Chen M, Chang AK, Xie L, Ying X. Assessment spermatogenic cell apoptosis and the
- 1011 transcript levels of metallothionein and p53 in Meretrix meretrix induced by cadmium. *Ecotoxicol Environ Saf*
- 1012 2021;217:112230 https://doi.org/10.1016/j.ecoenv.2021.112230
- 1013 Ghosh D, Najwa AR, Khan MA, Sengupta J. IGF2, IGF binding protein 1, and matrix metalloproteinases 2 and 9 in
- 1014 implantation-stage endometrium following immunoneutralization of vascular endothelial growth factor in the
- 1015 rhesus monkey. *Reproduction* 2011;**141**:501-509 https://doi.org/10.1530
- 1016 Girardet L, Augière C, Asselin MP, Belleannée C. Primary cilia: biosensors of the male reproductive tract.
- 1017 Andrology 2019;7:588-602 https://doi.org/10.1111/andr.12650
- 1018 Gu NH, Zhao WL, Wang GS, Sun F. Comaprative analysis of mammalian sperm ultrastructure reveals
- 1019 relationships between sperm morphology, mitochondrial functions and motility. Reprod Biol Endocrinol
- 1020 2019; 17:66 https://doi.org/10.1186/s12958-019-0510-y
- 1021 Gui L, Song K, Tritschler D, Bower R, Yan S, Dai A, Augspurger K, Sakizadeh J, Grzemska M, Ni T, et al. Scaffold
- 1022 subunits support associated subunit assembly in the Chlamydomonas ciliary nexin-dynein regulatory complex.
- 1023 Proc Natl Acad Sci USA 2019;116:23152-23162 https://doi.org/10.1073/pnas.1910960116
- 1024 Ha S, Lindsay AM, Timms AE, Beier DR. Mutations in Dnaaf1 and Lrrc48 Cause Hydrocephalus, Laterality
- 1025 Defects, and Sinusitis in Mice. G3 2016;6:2479-2487 https://doi.org/10.1534/g3.116.030791
- 1026 Hardy JJ, Wyrwoll MJ, Mcfadden W, Malcher A, Rotte N, Pollock NC, Munyoki S, Veroli MV, Houston BJ,
- 1027 Miguel J et al. XavierVariants in GCNA, X-linked germ-cell genome integrity gene, identified in men with
- 1028 primary spermatogenic failure. Hum Genet 2021;140:1169-1182 https://doi.org/10.1007/s00439-021-02287-y
- 1029 Heidari MM, Khatami M, Danafar A, Dianat T, Farahmand G, Talebi AR. 4Mitochondrial Genetic Variation in
- 1030 Iranian Infertile Men with Varicocele. Int J Fertil Steril 2016;10:303-309 doi: 10.22074/ijfs.2016.5047.
- 1031 Hernández-Llodrà S, Juanpere N, de Muga S, Lorenzo M, Gil J, Font-Tello A, Agell L, Albero-González R, Segalés
- 1032 L, Merino J, et al. ERG overexpression plus SLC45A3 (prostein) and PTEN expression loss: Strong association of

- 1033 the triple hit phenotype with an aggressive pathway of prostate cancer progression. *Oncotarget* 2017;8:74106-
- 1034 74118 https://doi.org/10.18632/oncotarget.18266
- 1035 Heshusius, S, Grech L, Gillemans N, Brouwer RWW, den Dekker XT, van IJcken WFJ, Nota B, Felice AE, van Dijk
- 1036 TB, von Lindern M, et al. Epigenomic analysis of KLF1 haploinsufficiency in primary human erythroblasts. Sci
- 1037 Rep 2022;12:336 https://doi.org/10.1038/s41598-021-04126-6
- 1038 Heuser T, Raytchev M, Krell J, Porter ME, Nicastro D. The dynein regulatory complex is the nexin link and a
- 1039 major regulatory node in cilia and flagella. J Cell Biol 2009;187:921-933 https://doi.org/10.1083/jcb.200908067
- 1040 Honda A, Komuro H, Shimada A, Hasegawa T, Seko Y, Nagase H, Hozumi I, Inuzuka T, Hara H, Fujiwara Y, et al.
- 1041 Attenuation of cadmium-induced testicular injury in metallothionein-III null mice. Life Sci 2010;87:545-550
- 1042 https://doi.org/10.1016/j.lfs.2010.09.009
- 1043 Houston BJ, Riera-Escamilla A, Wyrwoll MJ, Salas-Huetos A, Xavier MJ, Nagirnaja L, Friedrich C, Conrad DF, Aston
- 1044 AI, Krausz C et al. A systematic review of the validated monogenic causes of human male infertility: 2020
- 1045 update and a discussion of emerging gene-disease relationships. Hum Reprod Update 2021;28:15-29
- 1046 https://doi.org/10.1093/humupd/dmab030
- 1047 Huang Y, Huang Y, Pan XW, Li L, Lu Chen, Liu X, Lu JL, Zhu XM, Huang H, Yang QW, Ye JQ et al. Overexpression
- 1048 of USP39 predicts poor prognosis and promotes tumorigenesis of prostate cancer via promoting EGFR mRNA
- 1049 maturation and transcription elongation. *Oncotarget* 2016;7:22016-22030
- 1050 <u>https://doi.org/10.18632/oncotarget.7882</u>
- 1051 Hull RL, Johnson PY, Braun KR, Day AJ, Wight TN. Hyaluronan and hyaluronan binding proteins are normal
- 1052 components of mouse pancreatic islets and are differentially expressed by islet endocrine cell types. J
- 1053 Histochem Cytochem 2021;60:749-760 https://doi.org/10.1369/002215541245704
- 1054 Hwang K, Smith JF, Coward RM, Penzias A, Bendikson K, Butts S, Coutifaris C, Falcone T, Fossum G, Gitlin S et al
- 1055 (Practice Committee of the American Society for Reproductive Medicine in collaboration with the Society for
- 1056 Male Reproduction and Urology) Evaluation of the azoospermic male: a committee opinion. Fertil Steril
- 1057 2018;109:777-782 https://doi.org/10.1016/j.fertnstert.2018.01.043
- 1058 Ioannidis NM, Rothstein JH, Pejaver V, Middha S, McDonnell SK, Baheti S, Musolf A, Li Q, Holzinger E, Karyadi D,
- 1059 et al. REVEL: An Ensemble Method for Predicting the Pathogenicity of Rare Missense Variants. Am J Hum Genet
- 1060 2016;99:877-885 https://doi.org/10.1016/j.ajhg.2016.08.016

- 1061 Ishikawa T. Axoneme structure from motile cilia. *Cold Spring Harbor Perspectives in Biology* 2017;**9**,a028076
- 1062 https://doi.org/10.1101/cshperspect.a028076
- 1063 Jia Y, Ohanyan A, Lue YH, Swerdloff RS, Liu PY, Cohen P, Wang C. The effects of humanin and its analogues on
- 1064 male germ cell apoptosis induced by chemotherapeutic drugs. *Apoptosis* 2015;**20**:551-561
- 1065 https://doi.org/10.1007/s10495-015-1105-5
- 1066 Jodar M, Oliva R. Protamine alterations in human spermatozoa. Adv Exp Med Biol 2014;**791**:83-102
- 1067 https://doi.org/10.1007/978-1-4614-7783-9_6
- 1068 Jonkers J, Pai P, Sukumar S. Multiple roles of HOX proteins in Metastasis: Let me count the ways. *Cancer*
- 1069 Metastasis Rev 2020;39:661-679 https://doi.org/10.1007/s10555-020-09908-4
- 1070 Jumper J Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A,
- 1071 Potapenko A et al. Highly accurate protein structure prediction with AlphaFold. *Nature* 2021;**596**:583-589
- 1072 https://doi.org/10.1038/s41586-021-03819-2
- 1073 Karmilin K, Schmitz C, Kuske M, Körschgen H, Olf M, Meyer K, Hildebrand A, Felten M, Fridrich S, Yiallouros I, et
- 1074 al. Mammalian plasma fetuin-B is a selective inhibitor of ovastacin and meprin metalloproteinases. Sci Rep
- 1075 2019;9:546 https://doi.org/10.1038/s41598-018-37024-5
- 1076 Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. The Phyre2 web portal for protein modeling,
- 1077 prediction and analysis. Nat Protoc 2015;10:845-858 https://doi.org/10.1038/nprot.2015.053.
- 1078 Khelifa MB, Coutton C, Zouari R, Karaouzène T, J, Bidart' M, Yassine SV, Delaroche J, Hennebicq S, et al.
- 1079 Mutations in DNAH1, which encodes an inner arm heavy chain dynein, lead to male infertility from multiple
- 1080 morphological abnormalities of the sperm flagella. Am J Hum Genet 2014;**94**:95-104
- 1081 <u>http://dx.doi.org/10.1016/j.ajhg.2013.11.017</u>
- 1082 Kherraf ZE, Christou-Kent M, Karaouzene T, Amiri-Yekta A, Martinez G, S Vargas AS, Lambert E, Borel C, Dorphin
- 1083 B, Aknin-Seiferet I, et al. SPINK2 deficiency causes infertility by inducing sperm defects in heterozygotes and
- 1084 azoospermia in homozygotes. EMBO Mol Med 2017;9:1132-1149 https://doi.org/10.15252/emmm.201607461
- 1085 Klein C, Troedsson MHT, Rutllant J. Region-Specific Expression of Aquaporin Subtypes in Equine Testis,
- 1086 Epididymis, and Ductus Deferens. Anatomical Records (Hoboken) 2013;296:1115-1126
- 1087 https://doi.org/10.1002/ar.22709
- 1088 Kobe B, Kajava AV. The leucine-rich repeat as a protein recognition motif. Curr Opin Struct Biol 2001;11:725-
- 1089 732 https://doi.org/10.1016/S0959-440X(01)00266-4

- 1090 Krausz C, Riera-Escamilla A. Genetics of male infertility. *Nat Rev Urol* 2018;15: 369-384 1091 https://doi.org/10.1038/s41585-018-0003-3
- 1092 Krissinel E, Henrick K. Secondary-structure matching (SSM), a new tool for fast protein structure alignment in
- 1093 three dimensions. Acta Crystallogr D Biol Crystallogr 2004;60:2256-2268
- 1094 https://doi.org/10.1107/S0907444904026460
- 1095 Kubo T, Oda T. Electrostatic interaction between polyglutamylated tubulin and the nexin-dynein regulatory
- 1096 complex regulates flagellar motility. Mol Biol Cell 2017;28:2260-2266 https://doi.org/10.1091/mbc.e17-05-
- 1097 <u>0285</u>
- 1098 Kumar D. Reiter J. How the centriole builds its cilium: of mothers, daughters, and the acquisition of
- 1099 appendages. *Curr Opin Strucl Biol* 2021;66:41-48 <u>https://doi.org/10.1016/j.sbi.2020.09.006</u>
- 1100 Leal LF, Szarek E, Berthon A, Nesterova M, Faucz FR, London E, Mercier C, Abu-Asab M, Starost MF, Dye L, et al.
- 1101 Pde8b haploinsufficiency in mice is associated with modest adrenal defects, impaired steroidogenesis, and
- 1102 male infertility, unaltered by concurrent PKA or Wnt activation. *Mol Cellular Endocrinol* 2021;**522**:111117
- 1103 https://doi.org/10.1016/j.mce.2020.111117
- 1104 Lei Q, Huanga X, Zhenga L, Zhenga F, Donga J, Chena F, Zeng W. Biosensors for Caspase-3: From chemical
- 1105 methodologies to biomedical applications. *Talanta* 2022;240:123198
- 1106 https://doi.org/10.1016/j.talanta.2021.123198
- 1107 Lewis WR, Malarkey EB, Tritschler D, Bower R, Pasek RC, Porath JD, Birket SE, Saunier S, Antignac C, Knowles
- 1108 MR, et al. Mutation of Growth Arrest Specific 8 Reveals a Role in Motile Cilia Function and Human Disease.
- 1109 PLoS Genet 2016; 12:e1006220 https://doi.org/10.1371/journal.pgen.1006220
- 1110 Li H, Xing L, Zhao N, Wang J, Zheng N. Furosine Induced Apoptosis by the Regulation of STAT1/STAT2 and
- 1111 UBA7/UBE2L6 Genes in HepG2 Cells. Int J Mol Sci 2018; 19:1629 https://doi.org/10.3390/ijms19061629
- 1112 Li K, van Delft MF, Dewson G. Too much death can kill you: inhibiting intrinsic apoptosis to treat disease. EMBO
- 1113 J 2021;40:e107341 https://doi.org/10.15252/embj.2020107341
- 1114 Lorès P, Kherraf ZE, Amiri-Yekta A, Whitfeld M, Daneshipour A, Stouvenel L, Cazin C, Cavarocchi E, Coutton C,
- 1115 Llabador MA et al. A missense mutation in IFT74, encoding for an essential component for intraflagellar
- 1116 transport of Tubulin, causes asthenozoospermia and male infertility without clinical signs of Bardet-Bied
- 1117 syndrome. *Hum Genet* 2021;**140**:1031-1043 <u>https://doi.org/10.1007/s00439-021-02270-7</u>

- 1118 Lue Y, Swerdloff R, Jia Y, Wang C. The emerging role of mitochondrial derived peptide humanin in the testis.
- 1119 Biochim Biophys Acta Gen Subj 2021;1865:130009 https://doi.org/10.1016/j.bbagen.2021.130009
- 1120 Lüpold S, de Boer RA, Evans JP, Tomkins JL, Fitzpatrick JL. How sperm competition shapes the evolution of
- 1121 testes and sperm: a meta-analysis. Philos Trans R Soc Lond B Biol Sci 2020;375:20200064
- 1122 https://doi.org/10.1098/rstb.2020.0064
- 1123 Markou, A, Unger L, Abir-Awan M, Saadallah A, Halsey A, Baklava Z, Conner M, Törnroth-Horsefield S, Greenhill
- 1124 SD, Alex Conneret A, et al. Molecular mechanisms governing aquaporin relocalisation. Biochim Biophys Acta –
- 1125 Biomembr 2022; 1864: 183853 https://doi.org/10.1016/j.bbamem.2021.183853
- 1126 Marques PI, Fonseca F, Carvalho AS, Puente DA, Damião I, Almeida V, Barros N, Barros A, Carvalho F,
- 1127 Azkargortaet M et al. Sequence variation at KLK and WFDC clusters and its association to semen hyperviscosity
- 1128 and other male infertility phenotypes. *Hum Reprod* 2016;**31**:2881-2891
- 1129 https://doi.org/10.1093/humrep/dew267
- 1130 Martinez G, Coutton C, Loeuillet C, Cazin C, Muroňová J, Boguenet M, Lambert E, Dhellemmes M, Chevalier G,
- 1131 Hograindleur JP et al. Oligogenic heterozygous inheritance of sperm abnormalities in mouse. Elife
- 1132 2022;11:e75373 https://doi.org/10.7554/eLife.75373
- 1133 Matsushima N, Takatsuka S, Miyashita H, Kretsinger RH. Leucine rich repeat proteins: sequences, mutations,
- 1134 structures and diseases. Protein Pept Lett 2019;**26**:108-131
- 1135 https://doi.org/10.2174/0929866526666181208170027
- 1136 Matsuzaki, H., Hitomi Matsuzaki, Kuramochi D, Okamura E, Hirakawa K, Ushiki A, Tanimoto K. Recapitulation of
- 1137 gametic DNA methylation and its post-fertilization maintenance with reassembled DNA elements at the mouse
- 1138 lgf2/H19 locus. Epigenetics Chromatin 2020:13:2 https://doi.org/10.1186/s13072-019-0326-1
- 1139 Matzuk MM, Lamb DJ. The biology of infertility: research advances and clinical challenges. Nat Med
- 1140 2008;14:1197-1213 <u>https://doi.org/10.1038/nm.f.1895</u>
- 1141 McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, Flicek P, Cunningham F. The Ensembl Variant
- 1142 Effect Predictor. Genome Biol 2016; 17:122 https://doi.org/10.1186/s13059-016-0974-4
- 1143 Mei S, Wu Y, Wang Y, Cui Y, Zhang M, Zhang T, Huang X, Yu S, Yu T. Disruption of PIKFYVE causes congenital
- 1144 cataract in human and zebrafish. *Elife* 2022;11:e71256 <u>https://doi.org/10.7554/eLife.71256</u>

- 1145 Michaelis M, Sobczak A, Koczan D, Langhammer M, Reinsch N, Schoen J, WeitzeL JM. Selection for female traits
- 1146 of high fertility affects male reproductive performance and alters the testicular transcriptional profile. BMC
- 1147 Genomics 2017;18:889 https://doi.org/10.1186/s12864-017-4288-z
- 1148 Mo RJ, Lu JM, Wan YP, Hua W, Liang YX, Zhuo YJ, Kuang QW, Liu YL, He HC, Zhong WD. Decreased HoxD10
- 1149 Expression Promotes a Proliferative and Aggressive Phenotype in Prostate Cancer. Curr Mol Med 2017;17:70-78
- 1150 https://doi.org/10.2174/1566524017666170220104920
- 1151 Morohoshii A, Miyataid H, Shimadaid K, Nozawa K, Matsumura T, Yanaseid R, Shibaid K, Inabaid K, Ikawa M.
- 1152 Nexin-dynein regulatory complex component DRC7 but not FBXL13 is required for sperm flagellum formation
- 1153 and male fertility in mice. *PloS Genet* 2020;16:e1008585 https://doi.org/10.1371/journal.pgen.1008585
- 1154 Mostafa T, Rashed L, Taymour M. Seminal cyclooxygenase relationship with oxidative stress in infertile
- 1155 oligoasthenoteratozoospermic men with varicocele. Andrologia 2016;48:137142
- 1156 <u>https://doi.org/10.1111/and.12423</u>
- 1157 Nagirnaja L, Aston KI, Conrad DF. Genetic intersection of male infertility and cancer. *Fertil Steril* 2018; 109:20-26
- 1158 https://doi.org/10.1016/j.fertnstert.2017.10.028
- 1159 Nascimento JM, Shi LZ, Tam J, Chandsawangbhuwana C, Durrant B, Botvinick EL, Berns MW. Comparison of
- 1160 glycolysis and oxidative phosphorylation as energy sources for mammalian sperm motility, using the
- 1161 combination of fluorescence imaging, laser tweezers, and real-time automated tracking and trapping. J Cell
- 1162 Physiol 2008; 217: 745-51 https://doi.org/10.1002/jcp.21549
- 1163 Naz RK, Rajesh C. Gene knockouts that cause female infertility: search for novel contraceptive targets. Front
- 1164 Biosci 2005;10:2447-2459 https://doi.org/10.2741/1709
- 1165 Neirijnck Y, Kühne F, Mayère C, Pavlova E, Sararols P, Foti M, Atanassova N, Nef S. Tumor Suppressor PTEN
- 1166 Regulates Negatively Sertoli Cell Proliferation, Testis Size, and Sperm Production In Vivo. Endocrinology
- 1167 2019;160:387-398 https://doi.org/10.1210/en.2018-00892
- 1168 Ng ACY, Eisenberg JM, Heath RJ, Huett A, Robinson CM, Nau GJ, Xavier RJ. Human leucine-rich repeat proteins:
- 1169 a genome-wide bioinformatic categorization and functional analysis in innate immunity. Prot Nat Acad Sci
- 1170 2011; 108: 4631-4638 https://doi.org/10.1073/pnas.100009310
- 1171 Ogura, Y., Uehara T, Ujibe K, Yoshihashi H, Yamada M, Suzuki H, Takenouchi T, Kosaki K, Hirata H. The
- 1172 p.Thr395Met missense variant of NFIA found in a patient with intellectual disability is a defective variant.
- 1173 American Journal of Medical Genetics A 2022;188:1184-1192 https://doi.org/10.1002/ajmg.a.62638

- 1174 Oud MS, Houston BJ, Volozonoka L, Mastrorosa FK, Holt GS, Alobaidi BKS, deVries PF, Astuti G, Ramos L,
- 1175 Mclachlan RI et al. Exome sequencing reveals variants in known and novel candidate genes for severe sperm
- 1176 motility disorders. Hum Reprod 2021;36:2597-2611 https://doi.org/10.1093/humrep/deab099
- 1177 Pan W, Wang K, Li J, Li H, Cai Y, Zhang M, Wang A, Wu Y, Gao W, Weng W. Restoring HOXD10 Exhibits
- 1178 Therapeutic Potential for Ameliorating Malignant Progression and 5-Fluorouracil Resistance in Colorectal
- 1179 Cancer. Front Oncol 2021;11:771528 https://doi.org/10.3389/fonc.2021.771528
- 1180 Perner S, Rupp NJ, Braun M, Rubin MA, Moch H, Dietel M, Wernert N, Jung K, Stephan C, Kristiansen G. Loss of
- 1181 SLC45A3 protein (prostein) expression in prostate cancer is associated with SLC45A3-ERG gene rearrangement
- 1182 and an unfavorable clinical course. Int J Cancer 2013;132;807-812 https://doi.org/10.1002/ijc.27733
- 1183 Perrotta I, Santoro M, Guido C, Avena P, Tripepi S, De Amicis F, Gervasi MC, Aquila S. Expression of
- 1184 cyclooxygenase-1 (COX-1) and COX-2 in human male gametes from normal patients, and those with varicocele
- and diabetes: a potential molecular marker for diagnosing male infertility disorders. *J Anat* 2012;**221**:209-220
- 1186 https://doi.org/10.1111/j.1469-7580.2012.01534.x
- 1187 Pin E, Frauke Henjes F, Hong MG, Wiklund F, Patrik Magnusson P, Anders Bjartell A, Uhlen M, Nilsson P,
- 1188 Schwenk JM. Identification of a Novel Autoimmune Peptide Epitope of Prostein in Prostate Cancer. J Proteome
- 1189 *Res* 2017;**16**:04-216 https://doi.org/10.1021/acs.jproteome.6b00620
- 1190 Ramli NSK, Giribabu N, Karim K, Salleh N. Hormonal control of vas deferens fluid volume and aquaporin
- 1191 expression in rats. J Mol Histol 2019;50:21-34 https://doi.org/10.1007/s10735-018-9804-1
- 1192 Richards S, Aziz N, Sherri Bale S, Bick D, Das S, Gastier-Foster J, Grody WW, Hegde M P, Lyon E, Spector E, et al.
- 1193 Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the
- 1194 American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*
- 1195 2015; 17:405-424 https://doi.org/doi:10.1038/gim.2015.30
- 1196 Romeis B, Mulisch M, Welsch U. Romeis Mikroskopische Technik. Dordrecht: Springer. 2010; xii, 550 s. p.
- 1197 Roy A, Kucukural A, Zhang Y. I-TASSER: a unified platform for automated protein structure and function
- 1198 prediction. *Nat Protoc* 2010;5:725-738 https://doi.org/10.1038/nprot.2010.5
- 1199 Salustri A, Campagnolo L, Klinger FG, Camaioni A. Molecular organization and mechanical properties of the
- 1200 hyaluronan matrix surrounding the mammalian oocyte. *Matrix Biol* 2019;78-79:11-23
- 1201 https://doi.org/10.1016/j.matbio.2018.02.002

- 1202 Samaiya PK, Krishnamurthy S, Kumar A. Mitochondrial dysfunction in perinatal asphyxia: role in pathogenesis
- 1203 and potential therapeutic interventions. *Mol Cell Biochem* 2021;**476**:4421-4434
- 1204 <u>https://doi.org/10.1007/s11010-021-04253-8</u>
- 1205 Sandhu N, Rana S, Meena K. Nuclear receptor subfamily 5 group A member 2 (NR5A2): role in health and
- 1206 diseases. *Mol Biol Rep* 2021;48:8155-8170 <u>https://doi.org/10.1007/s11033-021-06784-1</u>
- 1207 Sandy Z, da Costa IC, Schmidt CK. More than Meets the ISG15: Emerging Roles in the DNA Damage Response
- 1208 and Beyond. *Biomolecules* 2020;**10**:1557 https://doi.org/10.3390/biom10111557
- 1209 Satir P, Heuser T, Sale WS. A structural basis for how motile cilia beat. *Bioscience* 2014;64:1073-1083
- 1210 https://doi.org/10.1093/biosci/biu180
- 1211 Sato JD, Kan M. Media for culture of mammalian cells. *Curr Protoc Cell Biol* 2001; Chapter 1:Unit 1.2.
- 1212 Schlegel PN, Sigman M, Collura B, De Jonge CJ, Eisenberg ML, Lamb DL, Sandlow JI, Sokol RZ, Spandorfer SD,
- 1213 Tanrikut C, et al. Diagnosis and treatment of infertility in men: AUA/ASRM guideline part I. Fertil Steril
- 1214 2021;115, 54–61 <u>https://doi.org/10.1016/j.fertnstert.2020.11.016</u>
- 1215 Sironen A, Shoemark A, Patel M, Loebinger MR, Mitchison HM. Sperm defects in primary ciliary dyskinesia and
- related causes of male infertility. Cell Mol Life Sci 2020;77:2029-2048 https://doi.org/10.1007/s00018-019-
- 1217 <u>03389-7</u>
- 1218 Smith CEL, Lake ARV, Johnson CA. Primary Cilia, Ciliogenesis and the Actin Cytoskeleton: A Little Less
- 1219 Resorption, A Little More Actin Please. *Front Cell Dev Biol* 2020;8:622822 1220 https://doi.org/10.3389/fcell.2020.622822
- Supernat A, Vidarsson OV, Steen VM, Stokowy T. Comparison of three variant callers for human whole genome
 sequencing. *Sci Rep* 2018;8:17851 https://doi.org/10.1038/s41598-018-36177-7
- Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Jaie Huerta-Cepas J, Simonovic M, Doncheva NT, Morris JH,
 Bork P et al. STRING v11: protein-protein association networks with increased coverage, supporting functional
 discovery in genome-wide experimental datasets. *Nucleic Acid Res* 2019;47:D607-D613
 <u>https://doi.org/10.1093/nar/gky1131</u>
- 1227 Tan JHL, Wollmann H, van Pelt AMM, Kaldis P, Messerschmidt DM. Infertility-Causing Haploinsufficiency 1228 Reveals TRIM28/KAP1 Requirement in Spermatogonia. *Stem Cell Rep* 2020;**14**:818-827
- 1229 <u>https://doi.org/10.1016/j.stemcr.2020.03.013</u>

- 1230 Touré A, Martinez G, Kherraf ZE, Cazin C, Beurois J, Arnoult C, Ray PF, Coutton C. The genetic architecture of
- 1231 morphological abnormalities of the sperm tail. Hum Genet 2021;140:21-42 https://doi.org/10.1007/s00439-
- 1232 020-02113-x
- 1233 Tourmente M, Villar-Moya P, Rial E, Roldan ER. 2015 Differences in ATP generation via glycolysis and oxidative
- 1234 phosphorylation and relationships with sperm motility in mouse species. J Biol Chem 2015;290:20613–20626
- 1235 https://doi.org/10.1074/jbc.M115.664813
- 1236 Tunyasuvunakool K, Adler J, Wu Z, Green T, Zielinski M, Žídek A, Bridgland A, Cowie A, Meyer C, Laydon A, et al.
- 1237 Highly accurate protein structure prediction for the human proteome. *Nature* 2021;**596**:590-596
- 1238 https://doi.org/10.1038/s41586-021-03828-1
- 1239 UniProt, C. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res* 2019;**47**:D506-D15
- 1240 <u>https://doi.org/10.1093/nar/gky1049</u>
- 1241 Valassina N, Brusco S, Salamone A, Serra L, Luoni M, Giannelli S, Bido S, Massimino L, Ungaro F, Mazzara PG, et
- 1242 al. Scn1a gene reactivation after symptom onset rescues pathological phenotypes in a mouse model of Dravet
- 1243 syndrome. Nat Commun 2022;13:161 https://doi.org/10.1038/s41467-021-27837-w
- 1244 Vander Borght M, Wyns C. Fertility and infertility: Definition and epidemiology. *Clin Biochem* 2018; 62: 2-10
- 1245 Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe O, Wood G, Laydoet A et al.
- 1246 AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space
- 1247 with high-accuracy models. Nucleic Acids Res 2022;50:D439-D444 https://doi.org/10.1093/nar/gkab1061
- 1248 von Mering C, Jensen LJ, Snel B, Hooper SD, Krupp M, Foglierini M, Jouffre N, Huynen MA, Bork P. STRING:
- 1249 known and predicted protein-protein associations, integrated and transferred across organisms. Nucleic Acid
- 1250 Res 2005;33:D433-443 https://doi.org/10.1093/nar/gki005
- 1251 Wagner K, Unger L, Salman MM, Kitchen P, Bill RM, Yool AJ. Signaling Mechanisms and Pharmacological
- 1252 Modulators Governing Diverse Aquaporin Functions in Human Health and Disease. Int J Mol Sci 2022;23:1388
- 1253 https://doi.org/10.3390/ijms23031388
- 1254 Wang C, Gu Y, Zhang K, Xie K, Zhu M, Dai N, Jiang Y, Guo X, Liu M, Dai J, et al. Systematic identification of genes
- 1255 with a cancer-testis expression pattern in 19 cancer types. Nat Commun 2016;7:10499
- 1256 <u>https://doi.org/10.1038/ncomms10499</u>

- 1257 Wertheimer E, Krapf D, de la Vega-Beltran, Sánchez-Cárdenas C, Navarrete F, Haddad D, Escoffier J, Salicioni
- 1258 AM, Levin LR, Buck J, Mager J, et al. Compartmentalization of distinct cAMP signaling pathways in mammalian
- 1259 sperm. J Biol Chem 2013; 288:35307-35320 https://doi.org/10.1074/jbc.M113.489476
- 1260 WHO 2020: https://www.who.int/news-room/fact-sheets/detail/infertility
- 1261 Wilson KS, Gonzalez O, Dutcher SK, Bayly PV. Dynein-deficient flagella respond to increased viscosity with
- 1262 contrasting changes in power and recovery strokes. *Cytoskeleton (Hoboken)* 2015;**72**:477-490
- 1263 https://doi.org/10.1002/cm.21252
- 1264 Wirschell M, Olbrich H, Werner C, Tritschler D, Bower R, Sale WS, Loges NT, Pennekamp P, Lindberg S, Stenram
- 1265 U, et al. The nexin-dynein regulatory complex subunit DRC1 is essential for motile cilia function in algae and
- 1266 humans. Nat Genet 2013;45:262-268 https://doi.org/10.1038/ng.2533
- 1267 Wyrwoll MJ, Temel SG, Nagirnaja L, Oud MS, Lopes AM, van der Heijden GW, Heald JS, Rotte N, Wistuba
- 1268 J, Marius Wo"ste M et al. Bi-allelic Mutations in M1AP Are a Frequent Cause of Meiotic Arrest and Severely
- 1269 Impaired Spermatogenesis Leading to Male Infertility. Am J Hum Genet 2020;107:342-351
- 1270 https://doi.org/10.1016/j.ajhg.2020.06.010
- 1271 Xavier MJ, Salas-Huetos A, Oud MS, Aston KI, Veltman JA. Disease gene discovery in male infertility: past,
- 1272 present and future. Hum Genet 2021;140:7-19 https://doi.org/10.1007/s00439-020-02202-x
- 1273 Xu W, Zhang Y, Qin D, Gui Y, Wang S, Du G, Yang F, Li L, Yuan S, Wang M, et al. Transcription factor-like 5 is a
- 1274 potential DNA- and RNA-binding protein essential for maintaining male fertility in mice. J Cell Sci
- 1275 2022;135:jcs259036 https://doi.org/10.1242/jcs.259036
- 1276 Xue M, Xue M, Fang Y, Sun G, Zhuo W, Zhong J, Qian C, Wang L, Wang L, Si J, Shujie Chen. IGFBP3, a
- 1277 transcriptional target of homeobox D10, is correlated with the prognosis of gastric cancer. PLoS One
- 1278 2013;8:e81423 https://doi.org/10.1371/journal.pone.0081423
- 1279 Yang J, Yan R, Roy A, Xu D, Poisson J, Zhang Y. The I-TASSER Suite: protein structure and function prediction.
- 1280 Nat Methods 2015;12;7-8 https://doi.org/10.1038/nmeth.3213
- 1281 Yang J, Zhang Y. I-TASSER server: new development for protein structure and function predictions. Nucleic
- 1282 Acids Res 2015;43:W174-181 https://doi.org/10.1093/nar/gkv342
- 1283 Yang LF, Yang F, Zhang FL, Xie YF, Hu ZX, Huang SL, Shao ZM, Li DQ. Discrete functional and mechanistic roles of
- 1284 chromodomain Y-like 2 (CDYL2) transcript variants in breast cancer growth and metastasis. Theranostics
- 1285 2020; 10:5242-5258 https://doi.org/10.7150/thno.43744

- 1286 Yin, M., O'Neil LAJ. The role of the electron transport chain in immunity. FASEB J 2021;35:e21974
- 1287 <u>https://doi.org/10.1096/fj.202101161R</u>
- 1288 Yuan J, Li X, Zhang G, Cheng W, Wang W, Lei Y, Ma Q, Song G. USP39 mediates p21-dependent proliferation
- 1289 and neoplasia of colon cancer cells by regulating the p53/p21/CDC2/cyclin B1 axis. *Mol Carcinog*enesis
- 1290 2021;60:265-278 https://doi.org/10.1002/mc.23290
- 1291 Zhan K, Liu R, Tong H, Gao S, Yang G, Hossain A, Li T, He W. Fetuin B overexpression suppresses proliferation,
- 1292 migration, and invasion in prostate cancer by inhibiting the PI3K/AKT signaling pathway. *Biomed Pharmacother*
- 1293 2020;131:110689 https://doi.org/10.1016/j.biopha.2020.110689
- 1294 Zhang J, Liu S, Zhang D, Ma Z, Sun L. Homeobox D10, a tumor suppressor, inhibits the proliferation and
- 1295 migration of esophageal squamous cell carcinoma. J Cell Biochem 2019;120:13717-13725
- 1296 https://doi.org/10.1002/jcb.28644
- 1297 Zhang Z, Miao J, Wang Y. Mitochondrial regulation on spermatogenesis. Reproduction 2022;163:R55-R69
- 1298 <u>https://doi.org/10.1530/REP-21-0431</u>
- 1299 Zhao Y, Zhang B, Lei Y, Sun J, Zhang Y, Sen Yang S, Zhang X. Knockdown of USP39 induces cell cycle arrest and
- 1300 apoptosis in melanoma. *Tumour Biol* 2016;**37**:13167-13176 <u>https://doi.org/10.1007/s13277-016-5212-x</u>
- 1301 Zhou S, Wu H, Zhang J, He X, Liu S, Zhou P, Hua R, Cao X, Liu M. Bi-allelic variants in human TCTE1/DRC5 cause
- asthenozoospermia and male infertility. Eur J Hum Genet 2022;30:721-729 https://doi.org/10.1038/s41431-
- 1303 <u>022-01095-w</u>
- 2011 Zorrilla M, Yatsenko AN. The Genetics of Infertility: Current Status of the Field. Curr Genet Med Rep
- 1305 2013;1:247–260 <u>https://doi.org/10.1007/s40142-013-0027-1</u>
- 1306
- 1307
- 1308
- 1309
- 1310
- 1311
- 1312
- 1313
- 1314

46

- 1315
- 1316
- 1317 Figures
- 1318

Figure 1: Characteristics of *Tcte1* gene. A Putative homologs of *Tcte1* gene conserved in Bilateria, according to
 HomoloGene (8434), including collation of proteins and their conserved domains; B Sequence alignement
 between human, mouse and rat; C NCBI Gene schematic representation of *Tcte1* gene in *Mus musculus*.

1322

Figure 2: Characteristics of *Tcte1* mice knockout. A Schematic representation of gRNA localization in CRISPR/Cas9 KO creation; B Relative expression of *Tcte1* gene in wild type (WT), hetero-^(+/-) and homozygous mice ^(-/-) (Real-time PCR); **** p<0.0001, ** p<0.01; C Example of genotyping result. Lane 1 – 1kb marker (GeneRuler), lanes 2, 3 – wild type (WT), lanes 4, 5 – hetero-^(+/-), and lanes 6, 7 – homozygous mice ^(-/-); lanes: 2, 4, 6 identified the lack (0 bp) or presence of mutation (630 bp), lanes: 3, 5, 7 differentiated hetero- (658 bp) and homozygotes (0 bp).

1329

Figure 3: Reproductive potential of KO *Tcte1* **mice.** Mating combinations and their results including: time of breeding, number of litters and pups, number of male and female pups, and genotypes observed in pups of each combination. **A** Mean time: from mating to 1st pedigree, and between particular litters; **B** Ratio of males and females per litter in each mating combination; **C** Frequency of males and females and their genotype observed each mating combination; left panel: according to sex of pups; right panel: according to mating combinations; **D** Ratio of males and females per litter and their genotype observed each mating combination.

1336 *** p < 0.0001, ** 0.0001 < p < 0.01, * 0.01 < p < 0.05

1337

Figure 4: Characteristics of *Tcte1* wild type (WT), hetero- ^(+/-) and homozygous mice ^(-/-). A Histopathology of testes and epididymis (caput); staining: classic hematoxyline-eosine staining (testis) or Masson-Goldner protocol (epididymis); microscope: Leica DM5500, objective 40×, LASX software with Navigator function; **B** Comparison of body mass, testis weight and size, and epididymis length, followed by gross morphology of gonads (n – no. of mice analyzed); **C** Comparison of the dimensions of testicular tubules' sections (n – no. of sections analyzed); *** p < 0.0001, ** 0.0001 < p < 0.01, * 0.01 < p < 0.05

1344	
1345	
1346	Figure 5: Evaluation of germ cell types in testes of <i>Tcte1</i> wild type (WT), hetero- $(+/-)$ and homozygous mice $(-/-)$
1347	⁾ . Three germ cell types were evaluated: spermatogonia (stained with intensive green color), preleptotene
1348	spermatocytes (intensive red color), and pachytene spermatocytes (medium green and red).
1349	Immunofluorescent staining with antibodies – primary: mouse anti-MLH1 (Abcam; Cambridge, UK, cat. no.
1350	ab14206), 1:30, rabbit anti-RAD51 (Abcam; Cambridge, UK, cat. no. ab63801), 1:200; secondary: goat anti-
1351	mouse-FITC (Sigma-Aldrich, cat. no. F2012), 1:400 and goat anti-rabbit-AF594 (Abcam; Cambridge, UK, cat. no.
1352	150080), 1:500.
1353	
1354	Figure 6: Characteristics of spermatozoa of <i>Tcte1</i> wild type (WT), hetero- ^(+/-) and homozygous mice ^(-/-) .
1355	A Sperm parameters: concentration, motility and morphology comparison; statistical marks indicated
1356	differences according to WT animals; n – number of mice; B Frequency of morphological defects according to
1357	three part of the spermatozoa; C Examples of main sperm morphological abnormalities found (all abnormalities
1358	observed available in Supplementary file 7); Leica DM5500 microscope, objective 63x, CytoVision software; D
1359	Comparison of the mean length of: sperm tail, midpiece and ratio tail:midpiece; n – no. of animals, each with at
1360	least 100 spermatozoa analyzed; *** p < 0.0001, ** 0.0001 < p < 0.01, * 0.01 < p < 0.05
1361	
1362	Figure 7: RNAseq results of changes in gene expression level found in testes of KO <i>Tcte1^{+/-}</i> and <i>Tcte1^{-/-}</i> mice
1363	vs. control (WT). Volcano plots: black dots indicated all analyzed genes within the genome, red dots
1364	represented genes with significantly changed expression level (followed by the heatmap and the table).
1365	Filtering criteria: p val <0.05, log ratio >1 or <-1 (adequate to fold change value of 2 (expression level increased)
1366	or 0.5 (expression level decreased)), expression level >8. nd – no data concerning function or disease.
1367	
1368	Figure 8: STRING analyzes of potential interactions between proteins with observed differences in gene
1369	expression level in RNAseq for mouse testis of the KO model for Tcte1 gene. Genes revealed in RNAseq were
1370	bracketed. (STRING database, 24th Feb 2021).
1371	

48

1372	Figure 9: Immunolocalization of the nexin-dynein regulatory complex (N-DRC) proteins: Tcte1 (Drc5), Drc7,
1373	Fbxl13 (Drc6), and Eps8l1 (Drc3) within mouse sperm cells. A. Localization of Tcte1 in spermatozoa from wild
1374	type (WT), hetero- ($^{+/-}$) and homozygous ($^{-/-}$) animals; B. Localization of other proteins (Drc7, Fbxl13, Eps8l1)
1375	building the nexin-dynein regulatory complex (N-DRC), responsible for coordination of the dynein arm activity
1376	and stabilization of the doublet microtubules attachment; wild type (WT) spermatozoa. Antibodies: primary (all
1377	anti-rabbit 1:100; Biorbyt, UK): Tcte1 (orb357083); Drc7 (orb58695); Fbxl13 (orb678278); Eps8l1 (orb382538);
1378	secondary: goat anti-rabbit-AF594 conjugated, 1:500, ab150160, Abcam, UK. Fluorescent microscope: Leica
1379	DM5500, filters – DAPI, TxR, FITC, BGR, magnification 630× (with immersion); software: LASX or CytoVision. Bar
1380	represents 10 μm.

1381

Figure 10: ATP level in wild type (WT), heterozygous (^{+/-}) and homozygous (^{-/-}) *Tcte1* mice. Measurement done in two time points (0 and +60 min.) showed decrease of approximately 35% of luminescence in all three groups. Mean measured values in homozygous group were approximately 2.4-fold lower. Dots – particular animals; long dash – mean value for the group; short dashes – standard deviation; RLU – relative luminescence unit. *** p<0.0001, ** 0.0001 < p < 0.01.

1387

1388 Figure 11: Immunolocalization of caspase 3 within mouse testicular tissue of KO Tcte1 model. Caspase 3 is 1389 one of the apoptotic indicators becuase of its engagement in the induction, transduction, and amplification of 1390 intracellular apoptotic signals. Caspase 3 signals were found only on spermatogonia in all three genotypes evaluated (wild type (WT), hetero- (+) and homozygous (-)). No differences between the level of fluorescent 1391 1392 signals were found between evaluated mouse genotypes, indicating similar level of apoptosis. Antibodies: 1393 primary (1:100) rabbit anti-Casp3 (Biorbyt, UK); secondary (1:500): goat anti-rabbit-AF594 conjugated 1394 (ab150160, Abcam, UK). Fluorescent microscope: Leica DM5500, filters – DAPI, TxR, magnification 630× (with 1395 immersion); software: LASX. Bar represents 5 μ m. Analysed in 3 independent males per zygosity, on 15-20 1396 tubules each.

1397

Figure 12: Homology modeling of TCTE1. A Secondary structure diagrams and associated confidence schema
 generated by I-TASSER [46,47,48], Phyre2[45], and AlphaFold [49,50] intensive mode. Positions of amino acids
 of interest are indicated. Green, predicted unstructured regions; yellow rectangles, predicted α-helices; orange

49

1401 arrows and arrowheads, predicted β -strands. Confidence scale ranges from white to cyan, with the probability 1402 of the secondary structure increasing with increasing intensity of cyan; B Cartoon diagram of homology model 1403 of human TCTE1 generated by AlphaFold with residues of interest shown as spheres and labeled. N, amino 1404 terminus; C, carboxy terminus; C-F Zoomed views of indicated residues and mutations: C lle-125 resides deep 1405 within a hydrophobic pocket, which would be disturbed upon substitution with arginine; D Mutation R133C 1406 would impact the hydrogen bond interactions that organize the architecture of the globular domain N-terminal 1407 to the tandem leucine-rich repeats; E Mutation of Glu-157 to alanine would destabilize the charge of the 1408 surface of this globular region, likely leading to unfolding; F Gly-350 allows for the compact interactions of 1409 tandem α -helices of the LRR. Depending on the rotamer it adopted, a serine side chain in this position would 1410 sterically clash with residues D321, A324, or V346 in the neighboring helix. 1411 1412 Figure 13: Summary of findings revealed for abnormal sperm tail beating in mouse knockout model of Tcte1 1413 gene, a component of the N-DRC complex in sperm tail. Three major groups of factors required for proper

sperm tail beating were marked with bold colour arrows (green, yellow and blue). Documented connections (solid grey arrows) revealed on the basis of: RNAseq data for testis tissues, protein *in silico* predictioning for

- 1416 mutated Tcte1 protein, and measurements of ATP level and sperm cell components in mouse spermatozoa.
- 1417 Unresolved possible linkages (cause/effect) marked with question (?) mark.

1419	Supplementary files legend
1420	
1421	Supplementary File 1: Vectors used for knockout creation of Tcte1 gene.
1422	
1423	Supplementary File 2: Real time PCR conditions used for determination of relative expression level of Tcte1
1424	gene in mouse knockout animals.
1425	
1426	Supplementary File 3: Reproductive potential of KO Tcte1 mice. Mating combinations and their results
1427	including: time of breeding, number of litters and pups, number of male and female pups, and genotypes
1428	observed in pups of each combination.
1429	
1430	Supplementary File 4: Genotyping protocol for <i>Tcte1</i> knockout model.
1431	
1432	Supplementary File 5: Characteristics of antibodies used for N-DRC complex staining in mouse sperm. Human
1433	antigen sequence identity between human and mouse orthologs was also shown (blast comparison;
1434	information from Biorbyt producer).
1435	
1436	Supplementary File 6: Schematic models of exon 3 deleted in KO mice. A Tcte1 protein exon 3 (deleted in KO)
1437	prediction of 5 top models (software: I-TASSER). C-score is the confidence score of the prediction. C-score
1438	ranges [0-1], where a higher score indicates a more reliable prediction; TM-score is a measure of global
1439	structural similarity between query and template protein., RMSD - root-mean-square deviation of the average
1440	distance between the atoms. B The deletion sequence used for modeling, including exon 3 (yellow).
1441	
1442	Supplementary File 7: Examples of HE staining on testicular tissue.
1443	
1444	Supplementary File 8: An example of circular type movement pathway of spermatozoa from Tcte1 ^{-/-} male
1445	mice.
1446	Supplementary File 9: An example of sperm cells motility in <i>Tcte1^{+/-}</i> male mice.
1447	Supplementary File 10: An example of sperm cells motility in <i>WT</i> male mice.

1448	Supplementary File 11: Detailed parameters of mouse sperm morphology. Statistically significant values: p <								
1449	0.05; ns – statistically not significant (p \ge 0.05). For samples with normal distribution (positive Shapiro-Wilk								
1450	normality test), t-test with Welch's correction was applied; for samples without normal distribution (midpiece								
1451	and tail defects), Kolmogorov-Smirnoff test was applied (GraphPad Prism v. 7.0e). n=200 spermatozoa were								
1452	analyzed per each mouse; for WT and <i>Tcte1^{-/-}</i> n=5 animals were evaluated, while for <i>Tcte1</i> ^{-/+} the number of								
1453	animals was n=7. Spermatozoa stained with the classic Papanicolau staining procedure. Bright field microscope:								
1454	Leica DM5500, magnification 630x (with immersion), software: CytoVision.								
1455									
1456	Supplementary File 12: STRING analyzis data for each gene with changed expression pattern in <i>Tcte1</i> KO								
1457	mouse model.								
1458									
1459	Supplementary File 13: STRING analyzis of potential interactions between proteins with observed major								
1460	differences in gene expression level in mouse testis of the homozygous <i>Tcte1</i> mice (Usp39, Fetub, Klk1b22).								
1461	According to genes revealed in RNAseq analysis (marked with arrows), two groups of interactions were formed								
1462	for genes with highest expression changes (red arrows); (STRING database, 11th March 2021).								
1463									
1464	Supplementary File 14: ATP measurement data obtained for KO Tcte1 male mice. wt = wild type, het =								
1465	heterozygous, ho - homozygous. RLU = relative luminescence unit.								
1466									
1467	Supplementary File 15: Examples of potentially disease-causing rare heterozygous TCTE1 variants found in								
1468	patients with disturbed spermatogenesis. Results obtained from screening of n=248 participants. A novel								
1469	variant (Sanger sequencing); B ultra rare variants (WES).								
1470									
1471	Supplementary File 16: The C-terminal region of TCTE1 was consistently modeled and fits well in published								
1472	maps. A Superposition of the cartoon representation of three models of human TCTE1 generated by Phyre2								
1473	[45] intensive (green) or fast mode (grey), I-TASSER [46,47,48] (yellow), and AlphaFold [49,50] (cyan). N- and C-								
1474	termini are labeled. B. Manual docking of the model of full-length TCTE1 generated by AlphaFold in maps								
1475	calculated from a cryo-electron tomography experiment performed on <i>C. reinhardtii</i> using nanogold to locate								
1476	the C-terminus of DRC5 (EMD-20821) [23].								

amino Deper	acid; S ident De	SNV – single nucleo epletion tool	otide va	ariant; PLof	= – Loss of Fu	unction pr	obability; rs – Ref	erence SNP	accession	number;	CADD – C	ombined Ann	otation												
Sample ID	[if ava concentr mot	Phetotype ilable – sperm count: ration/ volume/ total // ility (%): a+b/ total	origin	NT change	cDNA position	prediction	Mutation T@ster	GnomAD	rs	length of protein	CADD	position(s) of altered AA [HGVSp]	Polyphen/ Sift												
10vel varia	nts																								
·52	SOAs	4.3 x10 ⁶ /ml/ 3.0 ml/ 12.9 x10 ⁶ // 15.0/20.0	Poland	c.374T>G	cDNA.497T>G	disease causing	amino acid sequence changed; protein	not found	-	normal	missense	p. e125Arg	unknown												
P88	SOAs	3.4 x10 ⁶ /ml/ 1.5 ml/ 5.1 x10 ⁶ // 2.0/5.0					features (might be) affected; splice site																		
P105	SOAs	0.2 x10 ⁶ /ml/ 1.5 ml/ 0.3 x10 ⁶ // 17.0/23.0										cnanges													
111	SOAs	3.7 x10 ⁶ /ml/ 3.0 ml/ 11.1 x10 ⁶ // 15.0/20.0																							
2161	SOAs	2.3 x10 ⁶ /ml/ 3.0 ml/ 6.90 x10 ⁶ // 10.0/13.0																							
94	Crypto	0.05 x10 ⁶ /ml/ 2.0 ml/ 0.1 x10 ⁶ // 0.0/0.0	-																						
P149	NOA	0.0 x10 ⁶ /ml/ 2.0 ml/ 0.0 x10 ⁶ // 0/0																							
ultrarare v	ariants (M	AF <0.005)	P	- F						1	T														
J13 10 biopsy	NOA	low volume	USA	c.185G>A	cDNA.308G>A	d isease cau sing	amino acid sequence changed	0.00007778	rs201144307	normal	missense	p.A rg62 His	probably damaging/ deletoriou												
J17	NOA	late meiotic arrest	USA	c.397C>T	cDNA.520C>T	disease causing	amino acid sequence changed; heterozygous in TGP or ExAC; protein features (might be) affected; splice site changes	0.001722	rs145304614	normal	missense	p.Arg133Cys	probably damaging/ deletorious												
J50, J57, J117, J235	NOA	low volume	USA	c.470A>C	cDNA.593A>C	disease causing	amino acid sequence changed; heterozygous in TGP or ExAC; protein	0.002681	rs45591736	normal	missense	p.Glu157Ala	probably damaging/ deletoriou:												

													54
							features (might be) affected; splice site changes						
U11	NOA earl	ly meiotic arrest	USA	c.862C>T	cDNA.985C>T	d isease cau sing	NMD amino acid sequence changed; protein features (might be) affected; splice site changes	0.0002840	rs138414421	NMD	stop_gained	p.A rg288Ter	stopgain SNV; PLoF: high confidence
U76	NOA		USA	c.1048G>A	cDNA.1171G>A	disease causing	amino acid sequence changed; protein features (might be) affected; splice site changes	0.00006382	rs570334918	normal	missense	p.Gly350Ser	probably damaging/ deletorious ਲਾ

				Indentity (%)		conserved domains
species	gene symbol	Gene ID/ NCBI	UniProt	protein	DNA	Leucine-rich repeats (LRRs), cd00116 Leucine-rich repeats (LRRs), cl15309 CemA family (cl03855)
H. sapiens	TCTE1	202500	Q5JU00			– NP 872345.2.501 aa
vs.						
P. troglodytes	TCTE1	462732	H2QT36	99.2	99.2	– XP_518509.2, 501 aa
M. mulatta	TCTE1	702738	F7CLN3	97.0	96.9	— XP_001099848.1, 501 aa
C. lupus	TCTE1	609664	-	85.4	87.0	– XP_852058.1, 501 aa
B. taurus	TCTE1	523600	-	77.3	80.3	XP_002697348.1, 246 aa
M. musculus	Tcte1	21645	A6H639	77.3	77.5	- NP_038716.2, 498 aa
R. norvegicus	Tcte1	316242	D3ZQW5	78.0	78.0	— NP_001101676.1, 498 aa
G. gallus	TCTE1	421437	-	63.8	66.8	– XP_004935446.1, 457 aa
X. tropicalis	tcte1	100487694	-	57.6	63.5	— XP_002942200.1, 460 aa
D. rerio	tcte1	100331698	131538	55.5	61.0	— XP_002667399.3, 448 aa
D. melanogaster	CG14325	42123	Q9VEH4	31.6	43.6	— NP_650641.2, 460 aa
A. gambiae	AgaP_AP002751	1273219	Q7QCP9	28.3	45.5	——————————————————————————————————————





Genomic Sequence: NC_000083.6 Chromosome 17 Reference GRCm38.p6 C57BL/6J

В





D

males

testis

epididymis

50 µm

	WT	Tcte1+/-	Tcte1 ^{-/-}
body mass (g)	27.43 ±3.32	28.37 ±4.66	26.01 ±3.39
testis weight (mg)	134 ±0.01	125* ±0.01	78*** ±0.01
testis size (mm)	7.56 ±0.31/	7.56 ±0.39/	5.92*** ±0.23/
long (a)/short (b) axis	6.28 ±0.15	6.30 ±0.19	4.96*** ±0.17
epididymis length (mm)	12.76 ±0.70	12.84 ±0.68	12.52 ±1.05
n	21	22	18

В

Α

С

	WT	Tcte1+/-	Tcte1 ^{-/-}
tubule diameter [D]	191.60 ±29.58	201.50** ±30.03	211.00 ^{***/**} ±24.25
cell layer thickness [CL]	68.28 ±12.79	75.01***/** ±10.67	70.75*** ^{/***} ±10.01
lumen diameter [l]	125.20 ±24.59	126.50 ±22.25	140.2 ^{***} ±19.92
D:CL	2.84 ±0.36	2.70** ±0.24	3.01***/*** ±0.36
D:l	1.56 ±0.10	1.60** ±0.09	1.51***/*** ±0.09
CL:I	0.56 ±0.10	0.60 ^{**} ±0.09	0.51***/*** ±0.09
n	133	144	143

WT Tcte1^{+/-} Tcte1^{-/-} WT Tcte1^{+/-} Tcte1^{-/-} WT Tcte1^{+/-} Tcte1^{-/-}

	CELL TYPE:	WT	<i>Tcte1+/-</i>	<i>Tcte1^{-/-}</i>
	spermatogonia	17.23 ±5.37	13.44 ^{***} ±4.25	13.57 ^{***} ±5.26
medRxiv preprint doi: https://doi.org/10.1101/2022.11.1 (which was not certified by peer review) is the a It is made av	Azzazzas; this version posted Februard 6, 2024. The copyright helder for this preprint thor/funder, whe has used of mellow a cense to display the preprint in personal in personal to the preprint ine	12.96 ±5.19	25.39 ^{***/**} ±7.06	21.24 ^{***} ±9.46
	pachytene spermatocytes	35.76 ±14.30	31.23 ^{**} ±9.81	29.58 ^{***} ±13.00
	n	90	90	180

Tcte1+/-

Tcte1-/-

sperm parameters		WT	<i>Tcte1^{+/-}</i>	<i>Tcte1^{-/-}</i>	
concentration (×10 ⁶ /mL)		14.22 ±4.23	5.76 ^{***} ±1.52	4.35 ^{***} ±3.41	
motility (%)	a – progressive	33.52 ±4.07	33.14 ±3.75	0.22 ^{***} ±0.43	
	b – slow progressive	15.50 ±3.80	14.95 ±2.34	4.89 ±5.41 ^{***}	
	c – non-progressive (total)	23.43 ±3.87	24.27 ±5.44	37.89 [*] ±21.23	
	c – non-progressive (circular)	12.43 ±4.20	13.00 ±6.52	29.94 ^{**} ±16.94	
	d – immotile	27.55 ±4.80	27.73 ±6.18	57.00 ^{***} ±24.25	
n		21	22	18	
normal morphology (%)		70.81 ±11.78	71.94 ±8.97	35.41 ±9.10 ^{***}	
n		5	7	5	

B

medRxiv preprint doi: https://doi.org/10.1101/2022.11.17.22282339; this version poster Feynary 1 201, espoyute the deriver this reprint up provide the preprint in perpetuit (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to tala the preprint in perpetuit (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to tala the preprint in perpetuit (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to tala the preprint in perpetuit (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to tala the preprint in perpetuit (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to tala the preprint in perpetuit (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to tala the preprint in perpetuit (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to tala the preprint in perpetuit (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to tala the preprint in perpetuit (which was not certified by peer review) is the author/funder a CC-Bit 4.0 International license to tala the preprint in perpetuit (which was not certified by peer review) is the author/funder a CC-Bit 4.0 International license to tala the preprint in perpetuit (which was not certified by peer review) is the author/funder a CC-Bit 4.0 International license to tala the preprint in perpetuit (which was not certified by peer review) is the author/funder a CC-Bit 4.0 International license to tala the preprint in perpetuit (which was not certified by peer review) is the author/funder a CC-Bit 4.0 International license to tala the preprint in perpetuit (which was not certified by peer review) is the author/funder a CC-Bit 4.0 International license to tala the preprint in perpetuit (which was not certified by peer review		WT	<i>Tcte1+/-</i>	<i>Tcte1-/-</i>	
	tail	122.00 ±6.38	123.50 ±1.89	139.30** ±7.71	
	midpiece	27 ±2.21	25.58 ±0.53	26.97** ±1.17	
	tail:midpiece ratio	4.51 ±0.30	4.83* ±0.09	5.17 ^{**} ±0.20	
	n	6	6	6	

25₇

MIDPIECE TAIL ** ** ** Tcte1^{+/-} Tcte1^{-/-} WT Tcte1^{+/-} Tcte1^{-/-} WT Tcte1^{+/-} Tcte1^{-/-}

Tcte1^{-/-} vs. WT

	gene ID /NCBI	gene type	fold change			
gene symbol			Tcte1 ^{-/-}	Tcte1-/-	Tcte1+/-	known function/characteristics/disease
			vs.	vs.	vs.	
			WT	Teto 1+/-	WT	

			VV I	nier	VV I	
Klk1b27	16619	protein coding	0.2305	0.2937	-	٦
Klk1b21	16616	protein coding	0.2333	0.2626	-	
Klk1b24	16617	protein coding	0.3101	0.3102	-	
Klk1b16	16615	protein coding	-	0.3277	-	-
Klk1b22	13646	protein coding	0.1266	0.1494	-	
mt-Co2	17709	protein coding	0.4966	0.4734	-	٦
mt-Co3	17710	protein coding	-	0.4859	-	J
Fetub	59083	protein coding	4.6789	-	-	
Hoxd10	15430	protein coding	2.8503	-	-	
lgf2	16002	protein coding	2.0400	-	-	
medRxiv preprint doi: https://doi.org/10.1101/2022.11.17 (which was not certified by peer review) is the au It is made avail	22282339; this version posted February 16, 202 thor/funder, who has granted medRxiv a license t able under a CC-BY 4.0 International license .	4. The copyright holder for this preprint to display the preprint in perpetuity.				
Slc45a3	212980	protein coding	2.2164	-	-	
Usp39	28035	protein coding	12.0339	-	-	
Gm28661	-	unprocessed pseudogene	0.4576	-	-	
Gm26848	-	protein coding	2.2287	-	-	
Gm45648	102637934	ncRNA	2.5642	-	-	
Aqp2	11827	protein coding	-	0.4813	-	
Mt3	17751	protein coding	-	0.4868	-	
	400040040			~		
Rps2/rt	100043813	protein coding	-	0.4842		
Gm44126	-	protein coding (?)	-	0.2745		
4933430L12Rik	71251	protein coding	-	2.5102	-	
Ube2l6	56791	protein coding	-	0.4969	2.1110	
					11	-

degradation of extracellular matrix proteins in the interstitial area surrounding the Leydig cells of the adult mouse testis, seminal liquefaction seminal liquefaction neurons' differentiation mitochondrial cytochrome oxidase subunit, asthenozoospermia, cancer diabetes, deafness, apoptosis fertilization marker, sperm binding to *zona pellucida*, diabetes, coronary disease, inhibition and apoptosis of prostate cancer cells prostate cancer, cell differentiation and morphogenesis spermatogenesis, sex determination, testis development, infertility,

sperm DNA damage, prostate cancer, apoptosis

prostate cancer, myelin maintenance

spliceosomal complex, prostate cancer, apoptosis, cytokinesis, mitotic spindle checkpoint, vascular remodeling

nd

nd (membrane trasporter-like protein in D. sechellia)

nd (testis-restricted expression in mouse)

spermatogenesis, female endometrium cycle, stress urinary incontinence testis protection from cadmium, apoptosis, VEGF induction, retinal damage, Alzheimer disease ribosomal protein nd (suggested: UDP-glucuronic acid decarboxylase in mouse) nd cancer, apoptosis, autophagy, obesity

empty nodes – unknown 3D protein structure filled nodes – known/predicted 3D protein structure

experimentally determined

Tcte1 mouse

Tcte1+/-

head, midpiece + part of tail

Tcte1^{-/-}

Β

Drc7 mouse (WT)

Fbxl13 mouse (WT)

Eps8l1 mouse (WT)

head + whole tail

+60 min.

A

WT

medRxiv preprint doi: https://doi.org/10.1101/2022.11.17.22282339; this version posted February 16, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Tcte1+/-

Tcte1-/-

WT

DAPI

Tcte1+/-

Tcte1-/-

PROTEIN SURFACE POTENTIAL

DISTURBED protein conformation

ttps://doi.org/10.1101/2022.11.17.22282339; this version posted February 16, 2024. The copyright holder for this p rtified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetu

DISTURBED mt oxidase cytochrome complex