

Monoclonal antibody levels and protection from COVID-19

5 Eva Stadler¹, Martin T Burgess², Timothy E Schlub³, Khai Li Chai⁴, Zoe K McQuilten^{4,5}, Erica M
Wood^{4,5}, Mark N Polizzotto^{6,7}, Stephen J Kent^{8,9}, Deborah Cromer¹, Miles P Davenport^{1*},
David S Khoury^{1*}.

Affiliations:

1. Kirby Institute, University of New South Wales, Sydney, Australia
2. School of Mathematics and Statistics, University of New South Wales, Sydney, Australia
- 10 3. Sydney School of Public Health, Faculty of Medicine and Health, University of Sydney, Sydney, New South
Wales, Australia
4. Transfusion Research Unit, School of Public Health and Preventive Medicine, Monash University,
Melbourne, VIC, Australia
5. Department of Clinical Haematology, Monash Health, Clayton, VIC, Australia
- 15 6. Clinical Hub for Interventional Research, College of Health and Medicine, The Australian National University,
Canberra, ACT, Australia
7. Department of Clinical Haematology, Canberra Region Cancer Centre, The Canberra Hospital, ACT, Australia
8. Department of Microbiology and Immunology, University of Melbourne at the Peter Doherty Institute for
Infection and Immunity, Melbourne, VIC, Australia
- 20 9. Melbourne Sexual Health Centre and Department of Infectious Diseases, Alfred Hospital and Central Clinical
School, Monash University, Melbourne, VIC, Australia

*corresponding authors:

25 David Khoury
L6 Wallace Wurth Building,
Kirby Institute, UNSW Sydney
Kensington, NSW 2052
dkhoury@kirby.unsw.edu.au

30 Miles Davenport
L6 Wallace Wurth Building,
Kirby Institute, UNSW Sydney
Kensington, NSW 2052
35 m.davenport@unsw.edu.au

Abstract

Multiple monoclonal antibodies have been shown to be effective for both prophylaxis and therapy for SARS-CoV-2 infection. Here we aggregate data from randomized controlled trials
40 assessing the use of monoclonal antibodies in preventing symptomatic SARS-CoV-2
infection. We use data on changes in the *in vivo* concentration of monoclonal antibodies,
and the associated protection from COVID-19, over time to model the dose-response
relationship of monoclonal antibodies for prophylaxis. We estimate that 50% protection
from COVID-19 is achieved with a monoclonal antibody concentration of 54-fold of the *in*
45 *vitro* IC50 (95% CI: 16 – 183). This relationship provides a quantitative tool allowing
prediction of the prophylactic efficacy and duration of protection for new monoclonal
antibodies administered at different doses and against different SARS-CoV-2 variants.
Finally, we compare the relationship between neutralization titer and protection from
COVID-19 after either monoclonal antibody treatment or vaccination. We find no evidence
50 for a difference between the 50% protective titer for monoclonal antibodies and
vaccination.

Introduction

Vaccination has been shown to be highly effective at preventing both symptomatic and severe COVID-19 (reviewed by Cromer et al.¹). However, vaccination is less effective in many immune compromised and elderly individuals where immunogenicity and clinical data show considerably impaired responses to vaccination^{2,3}. Multiple monoclonal antibody products have been shown to be effective as pre- and post-exposure prophylaxis against pre-Omicron variants⁴⁻⁶, as well as when administered therapeutically early in infection⁷⁻¹¹. We recently analysed the available data on antibody treatment of symptomatic SARS-CoV-2 infection to determine the dose-response relationship between the antibody dose administered (after conversion to a neutralizing dose equivalence) and the protection from progression to hospitalisation¹². However, the dose-response curve for monoclonal antibody administration as prophylaxis of COVID-19 has not yet been determined, since there have to-date been fewer randomized control trials assessing the prophylactic effect. Here we adopt an alternative approach, comparing the loss of antibody *in vivo* with the loss of efficacy of monoclonal antibodies over time following administration. In addition, we use data on the loss of neutralisation and protection observed to new variants to inform this relationship¹³. Using this data on temporal changes in monoclonal antibody concentration and changes in potency to new variants, we estimate the relationship between *in vivo* antibody concentration and protection, which may provide a valuable clinical tool for predicting the efficacy of new monoclonal products and existing products against new variants¹². Finally, we assess whether neutralizing antibodies mediate protection or merely correlate with protection by comparing the relationship between neutralization titer and protection after vaccination¹⁴ and in naïve individuals receiving monoclonal antibodies. Together this work provides a quantitative framework for dissecting the mechanisms of protection in vaccination and informing the use of critical immunotherapies.

Results

Aggregating studies of monoclonal antibodies as prophylaxis

We searched MEDLINE, PubMed, Embase and the Cochrane COVID-19 Study Register for randomized placebo-control trials of SARS-CoV-2-specific monoclonal antibodies (mAbs) used as pre-exposure and peri-exposure prophylaxis for COVID-19. We included only studies where both protection from symptomatic infection and pharmacokinetic information of the

monoclonal antibody were provided within the same study. We identified six eligible studies assessing monoclonal antibodies as pre-exposure and peri-exposure prophylaxis for COVID-
90 19^{4-6,15,16} (see Table S1). The antibodies used in these studies were casirivimab/imdevimab (3 studies), bamlanivimab, tixagevimab/cilgavimab, and adintrevimab. One of these studies did not provide data on the pharmacokinetics of the antibody (bamlanivimab)¹⁶ and was excluded. Of the remaining four studies, three reported a break-down of cases in treatment and control arms by time since administration and two studies had data on the timing of
95 cases that could be extracted from the publication^{4-6,15}. Four of these five studies assessed protection at a time before the omicron variants were the dominant circulating variants and against variants where the antibody had been shown to be able to neutralize the virus *in vitro*¹⁷. One study assessed protection in two time periods; firstly in a pre-Omicron period when the Delta variant was the dominant circulating variant, and separately later when
100 Omicron variants BA.1 and BA.1.1 were the dominant variants¹³. The overall efficacies against pre-Omicron variants in the included studies ranged from between 66.7 – 92.4%. We identified a trend for lower efficacies with increasing time since administration and against the escaped variant (Figure 1).

105 **A significant dose-response relationship between protection and mAb concentration**

To investigate whether declining efficacy with time and new variants was indicative of a dose-response relationship between mAb concentration and efficacy, we compared the antibody concentrations reported within different time intervals in each study with the reported efficacy at the corresponding time interval (Figure 2). To compare between
110 antibodies of different potencies and against different variants, we normalized antibody concentration using the *in vitro* IC50 for each antibody/variant combination from a meta-analysis of *in vitro* studies (Table S2 and Figure S1, as previously reported¹² using data from Tao et al.¹⁸). We found a significant relationship between efficacy and mAb concentration (as a fold of the *in vitro* IC50) ($p < 0.0001$, generalized linear mixed model (GLMM) and chi-squared test, see Methods). In a leave-one-out analysis, we found this relationship
115 remained significant when any one study is omitted (Table S3). Fitting a logistic dose-response relationship to this data, we estimated a peak efficacy of 94% (95% CI: 84 – 106%), and concentration for 50% of efficacy of 54-fold *in vitro* IC50 (95% CI: 16 – 183) (Table S4). Against the ancestral virus, this equates to a plasma antibody concentration of 0.22 mg/L

120 concentration for tixagevimab/cilgavimab, 0.18 mg/L for casirivimab/imdevimab and 0.36 mg/L for adintrevimab being required for 50% protection against COVID-19.

Predicting monoclonal antibody efficacy against new variants

A major challenge in the COVID-19 pandemic has been in decision-making around whether
125 pharmaceuticals shown to be effective against ancestral SARS-CoV-2 should continue to be used when new variants emerge. This is especially true for monoclonal antibody therapies, where recommendations on the use of mAb therapeutics have changed numerous times with the emergence of each Omicron subvariant^{19,20}. This has been particularly difficult when a mAb loses partial, but not complete, recognition of a new variant. If we assume that
130 the relationship defined here between antibody concentration (normalised to *in vitro* neutralising IC50) and efficacy will continue to hold for different variants of concern, as it has for vaccine-induced neutralizing antibodies^{1,21}, we can use the dose-response relationship in Figure 2 to estimate the loss of efficacy and duration of protection of monoclonal antibodies to new variants. For example, tixagevimab/cilgavimab administered
135 intramuscular at a dose of 300 mg is predicted to maintain >50% protection for 670 days (95% CI: 503 – 836 days) against the ancestral variant where the *in vitro* IC50 is 4.00 ng/mL and the half-life of antibody 95 days (Figure 3, Tables S2, S5 and S6). However, if the *in vitro* IC50 against a new variant were increased (but still detectable), as is the case for some mAbs against the Omicron subvariants (Table S2), the model allows us to predict the
140 reduction in the time above 50% protection. In the meta-analysis used here on mAb IC50's against different variants, tixagevimab/cilgavimab was found to maintain detectable *in vitro* neutralization titers against the BA.1, BA.2 and BA.4/5 Omicron subvariants (see Table S2, meta-analysis as described in¹², using data from Tao et al.¹⁸). The increase in IC50 for this antibody combination against is predicted to reduce efficacy at the peak antibody
145 concentration from 88.8% (95% CI: 77.3 – 93.0) against ancestral virus, to 56.7% (95% CI: 38.9 – 71.4), 77.0% (95% CI: 62.8 – 85.4) and 43.4% (95% CI: 21.1 – 58.5) efficacy against the BA.1, BA.2 and BA.4/5 subvariants, respectively (Table S7). Using the dose-response relationship here (Figure 2), this would predict that the duration of >50% protection of tixagevimab/cilgavimab against BA.1 and BA.2 will be 97 (95% CI: 0 – 263) and 360 (95% CI:
150 194 – 526) days, respectively (Table S5). On the other hand, casirivimab/imdevimab and

adintrevimab have larger increases in the IC50 against these subvariants (Table S2) and are not predicted to maintain >50% efficacy to these variants (Tables S5 and S7).

Another formulation of this question is to ask “What is the maximum increase of IC50 (drop in neutralization titer) that can be tolerated while still maintaining a minimum duration of protection?”. For example, if we wish to provide a period of at least 30 days with >50% protection, then tixagevimab/cilgavimab, casirivimab/imdevimab and adintrevimab, at the current doses, can tolerate at most 106.8-fold (95% CI: 31.7 – 360.0), 319.5-fold (95% CI: 94.8 – 1076.5) and 89.6-fold (95% CI: 26.6 – 302.0) increases in *in vitro* IC50, respectively (Table S8). Figure 3d shows the predicted duration of >50% protection for casirivimab/imdevimab, tixagevimab/cilgavimab and adintrevimab for any given change in *in vitro* IC50. This analysis provides a tool to determine how regularly monoclonal antibodies may need to be re-administered to provide high confidence that subjects will maintain a serum antibody concentration that will provide more than 50% efficacy against the circulating variant.

165

Comparing monoclonal antibody prophylaxis with vaccine-induced protection

Multiple lines of evidence have established that neutralizing antibody titers correlate with protection from COVID-19 in vaccinated individuals^{14,22-24}. An important question is whether neutralizing antibodies are mechanistic in mediating this protection, or merely correlate with protection²⁵. Similarly, if antibodies are able to directly mediate protection, identifying the magnitude of their contribution to overall protection (compared to other mechanisms) is an important question. One way to address this is to compare the level of protection achieved after administration of antibodies alone with that achieved after vaccination. Antibody administration alone should reflect the antibody-related contribution to protection, while vaccination should incorporate both antibody- and cell-mediated protection. Recently, Schmidt et al., reported the loss of *in vitro* neutralisation of adintrevimab to Omicron BA.1 and BA.1.1 resulted in a corresponding loss of efficacy that was not inconsistent with the relationship between reports of vaccine effectiveness¹³. Here we address this question by integrating the available data across currently available studies on monoclonal antibody prophylaxis. We then used an established correlate of vaccine protection for COVID-19 to analyse whether prophylaxis against COVID-19 after passive

180

antibody administration is achieved at similar levels of neutralisation to protection observed after vaccination (Figure S3).

185 Figure 4a compares the efficacy of high-potency mRNA vaccines^{26,27} and monoclonal
antibody prophylaxis in prevention of symptomatic SARS-CoV-2 infection with the ancestral
or delta variant. We find that the observed mean protection achieved with mAbs of 84.8%
(95% CI: 76.0 – 90.8) was significantly different to the efficacy achieved with vaccination of
94.5% (95% CI: 91.6 – 96.7) ($p=0.002$, GLMM, see Methods). However, a similar efficacy may
190 be possible by administering monoclonal antibodies at much higher neutralizing antibody
titers compared with the neutralising antibody titers achieved in vaccinated individuals.
Thus, we next compared the level of protection achieved for a given neutralizing antibody
titer after either vaccination (from Khoury et al.¹⁴) or after treatment with monoclonal
antibodies (Figure 4b). To compare titers between vaccination and monoclonal antibody
195 administration, we normalized the titers of each to a scale relative to the geometric mean
titer of neutralizing antibodies seen in convalescent individuals against ancestral virus, after
the first wave of COVID-19 (a ‘fold-convalescent’ scale, as previously described for the
studies of vaccination¹⁴, and as described in the Methods for mAbs). In order to test
whether the neutralizing antibody titer associated with a given level of protection for
200 vaccines and mAbs were consistent or different, we fitted these data together using the
same parameters for the monoclonal antibody and vaccine dose-response relationships. We
then tested whether the model fitted better with separate parameters for mAb treatment
and vaccination (likelihood ratio test, see Methods and Figure S3). There was no evidence of
a difference in the neutralisation titre required for 50% protection ($p = 0.48$, Figure S4).
205 However, this analysis showed that the best-fit model was one where the same dose-
response relationship existed for both vaccination and mAbs but with the estimated slope
being higher for vaccination. Alternatively, a model with the same parameters except a
different maximum efficacy had a nearly equivalently good fit (AIC difference 1.9, Table S9).
In these two nearly equivalently good models, at neutralisation titres above the 50%
210 protective level, the efficacy of mAb tended lower than that achieved by vaccination (Figure
4b, Figure S3, Table S9). Together, these results indicate that similar levels of neutralising
antibodies from either vaccination or the administration of monoclonal antibody are
associated with 50% protection from COVID-19, but with a trend towards a lower maximum

215 protection from COVID-19 that can be achieved with monoclonal antibodies compared to
vaccination.

Discussion

Here we demonstrate a relationship between the monoclonal antibody concentration and efficacy to prevent COVID-19. Further, we estimate the concentration of antibody required to have a high confidence of maintaining at least 50% protection. Our model fitting enabled us to quantify the uncertainty in this relationship and estimate that if a treated population can maintain a mean *in vivo* monoclonal antibody concentration of > 54-fold (95% CI: 16 – 183) of the *in vitro* IC50 of the antibody to the circulating variant, they should maintain > 50% efficacy against COVID-19. Analysis of the dose-response curve for monoclonal antibodies allows prediction of the level and duration of protection against different SARS-CoV-2 variants (Figure 3). Our results suggest that casirivimab/imdevimab, tixagevimab/cilgavimab and adintrevimab would provide >50% protection against the ancestral SARS-CoV-2 strain for 269, 670 and 903 days respectively (Table S5). At the current doses, casirivimab/imdevimab and adintrevimab are not predicted to achieve 50% protection against the Omicron BA.2 variant (Table S7). For tixagevimab/cilgavimab, the protective interval against the BA.2 variant is dramatically reduced from 670 days to 360 days, and none of these antibodies were predicted to achieve 50% protection against the Omicron BA.4/5 variant (Tables S5 and S7). Counterintuitively, although antibodies with a longer half-life are expected in general to provide protection for longer, these are also expected to lose more ‘days of protection’ for a given fold-increase in IC50 (to a new variant), compared to mAbs with shorter half-lives (Figure 3). The higher susceptibility of therapeutics with longer half-lives to fold-shifts in the IC50 has been discussed previously for antimalarial products²⁸, and can be explained by considering that when antibodies lose 2-fold neutralization against a new variant it is equivalent to the mAb losing one half-life of time above a threshold. Therefore, a mAb with a 100-day half-life will lose 100 days above a specified threshold, whereas a mAb with only a 30-day half-life will lose 30 days above the same threshold.

The estimated *in vivo* concentration of antibody required for 50% protection from COVID-19 is much higher than the level of antibody required to neutralize virus *in vitro* (approx. 50-fold), suggesting that *in vivo* neutralization may be much less efficient than the observed neutralization *in vitro*. This difference between *in vitro* IC50 and the *in vivo* 50% protective titer is not unexpected, given the major differences between infection in these

environments. For example, *in vitro* neutralization assays usually involve pre-incubation of
250 antibody and virus for an hour before exposure to cells. Similarly, the *in vitro* IC50 in plaque
reduction assays estimates the antibody concentration required to neutralize 50% of virions.
However, the dose required to completely neutralize large inocula may be considerably
higher²⁹. In addition, *in vivo* antibody titers are assessed in the serum. However, antibody
concentration at the mucosa is lower than the plasma level³⁰, and thus higher (serum) titers
255 may be required to achieve neutralization on mucosal surfaces.

We and others have previously shown that neutralizing antibodies are a correlate of
protection from COVID-19^{14,22,31-33}. A major question in understanding vaccine-mediated
immunity is whether neutralizing antibodies are simply a surrogate marker of protection or
260 are mechanistic in protecting individuals from symptomatic infection¹⁴. To-date it has only
been possible to consider this question indirectly. For example, we have noted that the drop
in neutralizing antibodies against new variants and over time both provide good predictions
of the change in efficacy of vaccines over time and against new variants^{1,34}. Schmidt et al.
recently showed that the monoclonal antibody adintrevimab lost neutralisation to Omicron
265 BA.1 and BA.1.1, and a corresponding loss of prophylactic efficacy compared with that
predicted by the relationship between neutralisation titre and vaccine effectiveness
studies¹³. Here, we extracted data on antibody PK and temporal changes in efficacy from
five monoclonal antibody studies (including from Schmidt et al.¹³), and compared this with
an established immune correlate of protection after vaccination that has been validated
270 across a number of settings^{1,14,21,35,36}. This allows a direct comparison of the protection
provided by passively administered antibodies versus vaccine-induced. We have been able
to show that the neutralization titer required for 50% protection by vaccination or
monoclonal antibodies is comparable (Figure 4), although a predicted trend towards higher
vaccine efficacy at high neutralising antibody titers was observed (Figure S4). This suggests
275 that neutralizing antibodies are sufficient to explain much of the protection from
symptomatic infection induced by vaccination.

The difference in protection at a given neutralization titer between vaccination and
monoclonal antibody therapy may be due to the additional benefit in vaccinees of a
280 polyclonal antibody response, other non-neutralizing functions of antibodies, recall of

immune memory and/or other cellular immune responses. These functions may contribute to the estimated trend towards higher maximal protection for vaccines. Also, the efficacy studies of tixagevimab/cilgavimab and adintrevimab had infection events with other pre-Omicron variants, in particular the delta variant, and we have assumed that the *in vitro* IC50's of these antibodies (especially) against the delta variant were equivalent to the IC50s against ancestral virus, for simplicity. However, it is likely that there was some increase in IC50 against the delta variant which we have not accounted for and this may explain some of the apparent lower efficacy in these studies (Figure 4a). While our analysis has shown that neutralizing antibodies are sufficient for protection from COVID-19, it is not possible to conclude from this analysis that neutralizing antibodies are necessary for protection. We note that evidence in animal models supports the findings that neutralising antibodies mediate protective immunity³⁷, with some showing an additional benefit of Fc-receptor function³⁸.

The analysis presented here has a number of limitations. Firstly, our dose-response analysis requires comparison of the *in vivo* measured antibody concentrations and the estimated *in vitro* IC50s. We also assume the population average half-life of antibodies in the treated population, rather than assessing individual titers over time. Further, we are reliant upon summary data on monoclonal antibody protection broken down by time as reported in, or extracted from, the published studies, and thus could not account fully for subjects lost to follow-up (although fortunately these numbers are relatively small). Additionally, the analysis is strongly influenced by the results from the Herman et al.¹⁵ and Schmidt et al.¹³ studies given their contribution of data at lower antibody concentrations (Figure 2). To gain more robust estimates of the dose-response curve for the use of monoclonal antibodies for prophylaxis, studies with longer follow-up times would be greatly beneficial where this is ethical.

Another factor affecting our analysis is the different mode of delivery of the monoclonal antibodies. Casirivimab/imdevimab was administered subcutaneously in the Isa⁵, O'Brien⁶ and Herman¹⁵ studies, whereas tixagevimab/cilgavimab and adintrevimab were administered intramuscularly^{4,13}. Plasma antibody concentrations had much slower increases following intramuscular administration compared with subcutaneous

administration of casirivimab/imdevimab, and thus it is possible that there is a delay until protective antibody concentrations are achieved (Figure 1). To avoid this difference and also
315 to account for the risk of infection at the time of antibody administration, we omitted the earliest time interval from our analysis (which encompassed the first 7, 10, 28 or 90 days across different studies).

Studies of *in vitro* neutralization of different SARS-CoV-2 variants provide a range of
320 different IC50's for each antibody. We therefore used a meta-analysis of *in vitro* IC50's to inform the mean IC50 for each antibody and to normalize antibody levels to the average convalescent titer for comparison with vaccine titers¹⁸. Only a subset of these studies reported the geometric mean neutralization titer of a cohort of convalescent plasma from early in the pandemic (n=13). This meta-analysis aggregated IC50's estimated from a wide
325 range of neutralization assays, and the definition of convalescent sera was specified differently in each study, introducing some potential confounders to these aggregated estimates. However, our multiple regression model included random effects for study, which can account for systematic differences in neutralization in the studies – for example due to assay differences.

330

Conclusion

Vaccination has provided a high level of population immunity to COVID-19. However, there remain a number of subgroups in which vaccination is either not possible or is ineffective (largely due to immunodeficiency). The use of monoclonal antibodies for prophylaxis in
335 these cohorts has the potential to provide long-term protection from both symptomatic and severe COVID-19 for these vulnerable groups^{4-11,15,16}. However, the frequent observation of novel SARS-CoV-2 variants that escape antibody recognition has raised significant challenges in predicting monoclonal antibody protection against new variants. Further work is required to obtain more data on protection at low antibody levels, as well as to validate predictions
340 of prophylactic efficacy against SARS-CoV-2 variants. Within this context, the work presented here provides a quantitative and evidence-based framework for predicting monoclonal antibody efficacy that can be used in the assessment of novel therapeutics or in designing optimal regimes for new SARS-CoV-2 variants.

345 **Methods**

Search strategy for studies of COVID-19 prophylaxis with monoclonal antibodies

We searched MEDLINE, PubMed, Embase and the Cochrane COVID-19 Study Register from inception to 30 November 2022, for randomized control trials of monoclonal antibodies for
350 the prevention of COVID-19. We identified six studies. We included only the five studies where monoclonal antibody concentration was also reported for the cohort (Table S1). These studies were in a mixture of true pre-exposure prophylaxis and peri-exposure prophylaxis settings (Table S1).

Two studies, O'Brien et al.⁶ and Herman et al.¹⁵, reported results from the same clinical trial
355 over different follow-up intervals (4 weeks and 8 months respectively). Thus, to avoid duplication of the same trial results, we integrated the results from these studies. In particular, the O'Brien trial reported outcomes on a weekly basis for 4 weeks whereas the Herman trial reported outcomes on a monthly basis for 8 months. Therefore, for these trials the weekly outcomes reported in O'Brien were used for the weeks 2 to 4 after
360 administration (the initial week was omitted due to rising antibody levels in this period) and the results from Herman et al., were used for the months 2 to 8 only. In addition, antibody concentration data for the cohort was extracted from Figure S4 of O'Brien et al. from 0-168 days, whereas the Herman reported only pharmacokinetic model predictions of the concentration over the interval of 30-240 days. Therefore, the raw O'Brien et al. antibody
365 concentration data was used from 0-168 days and the predicted concentrations from Herman et al. were used for the remaining interval, 168-240 days after treatment. This is indicated in Figure 1b by a different line type (solid for *in vivo* concentration data from O'Brien et al. and dashed for modelled concentration data from Herman et al.). In Figure 1d, the dashed line for the concentration data also denotes modelled concentration data rather
370 than *in vivo* measurements of concentration.

Meta-analysis of monoclonal antibody IC50 and neutralization titer for early pandemic convalescent sera

A previous systematic review and meta-analysis of *in vitro* studies of monoclonal antibodies
375 IC50 against ancestral SARS-CoV-2 and BA.1, BA.2 and BA.1.1 Omicron subvariants has been reported¹⁸, and provided the raw data from this review. In addition, we have previously

extended this systematic review to include studies of *in vitro* IC50 against the BA.4/5 subvariants¹². Thus, using this previously developed dataset of *in vitro* IC50's we modified a previously performed meta-analysis of the *in vitro* IC50 of antibodies (bamlanivimab, casirivimab, cilgavimab, imdevimab, sotrovimab, tixagevimab, etesevimab, regdanvimab and bebtelovimab, as these were the antibodies of interest in our previous analysis¹²) to include adintrevimab, as well as the tixagevimab/cilgavimab and casirivimab/imdevimab combinations (Table S2). This meta-analysis involved performing a mixed-effects linear regression (with censoring if IC50 above 10,000) using the *lme4* package^{39,40}, as described previously¹². From this meta-regression we derive a central estimate of the IC50 for each antibody-variant combination (casirivimab/imdevimab, tixagevimab/cilgavimab, and adintrevimab, see Figure S1) considered here, which we use to calculate the *in vivo* antibody concentration as a fold of the *in vitro* IC50 as described below. In addition, we searched all studies included in the systematic review described above for additional data on whether the study reported neutralizing antibody titers for a panel of convalescent subjects from early in the pandemic. We identified 11 studies in which a panel of convalescent sera (obtained from individuals with infections that occurred early in the pandemic, when ancestral viral lineages dominated transmission) were assessed for their neutralizing titers⁴¹⁻⁵¹. Additionally, we identified a further 2 studies where a panel of convalescent sera from individuals infected pre-delta variant of concern were assessed^{52,53}. We included the (inverse-transformed) geometric mean neutralization titers reported for these panels of convalescent sera in our meta-regression in order to estimate a mean neutralization titer of early pandemic convalescent sera across the same set of studies as was used to estimate a mean IC50 (Figure S1). This mean convalescent titer across the studies was used to estimate the neutralization titer (on the fold-convalescent scale) for each monoclonal antibody based on the concentration data reported in each prophylaxis study as is described below.

Estimation of antibody concentration on fold-IC50 scale and estimate of neutralization titer on fold-convalescent scale

Antibody concentrations in each study were extracted and normalized as a ratio to the IC50 from the meta-regression described above (Table S2, reported as antibody concentration, fold-IC50). Note that since some of these studies used antibody combination therapies, the total antibody concentration (sum of both antibody components) was used for the antibody

concentration. When comparing the efficacy of these monoclonal antibody trials against
410 that seen after vaccination, we used the estimated neutralization titer in individuals treated
with mAbs, converted to a fold-of-convalescent scale since all vaccine trials has been aligned
based on this scale previously¹⁴. This was estimated by taking the antibody concentration as
a fold of the IC50 described here, and dividing by the mean convalescent neutralization titer
(as described previously¹²).

415

Test of concentration effect on the efficacy of prophylactic mAb treatment

Since the concentration data is very sparse below a concentration of 1,000-fold *in vitro* IC50,
we tested if there is a significant effect of the antibody concentration on the efficacy in the
data. For this analysis, we used a log-binomial regression model, a generalised linear mixed
420 model (GLMM) with a binomial error family and logarithmic link function (using the *glmer*
function of the *lme4* package⁵⁴ in R version 4.2.1⁵⁵). The model includes random intercepts
for different trials, the covariate “treatment” and the interaction of treatment with
concentration. The significance of the interaction of treatment and concentration was
tested with a chi-squared test (with the function *drop1*).

425

Dose-response fitting with maximum likelihood approach

To estimate the dose-response curve, we fitted a logistic function to the efficacy by
concentration data (Figure 2, Figure 4). As we aimed to estimate the concentration that
gives 50% protection, we scaled the logistic function such that this concentration is directly
430 estimated as a parameter of the function. The efficacy for a treatment with mAb
concentration c is then given by

$$E(c | m, k, c_{50}) = \frac{m}{1 + (2m - 1) \times \exp\left(-k \times (\log_{10}(c) - \log_{10}(c_{50}))\right)},$$

435 where m denotes the maximal efficacy, k is a slope parameter and c_{50} is the concentration
that gives 50% efficacy. Note that this rescaled logistic function will be ill-defined when m is
below 50%.

We fitted this dose-response curve to the data with a maximum likelihood approach (as
previously described¹²). Briefly, the likelihood function used in this optimisation was,

440

$$L(p) = \prod_{\tau} \text{Binom}(e_c^{\tau}, n_c^{\tau}, b^{\tau}) \times \text{Binom}(e_t^{\tau}, n_t^{\tau}, b^{\tau} \times (1 - E(c(\tau) | m, k, c_{50}))),$$

where p denotes the parameters of the likelihood function, i.e. the three parameters of the efficacy function (m, k, c_{50}) and the baseline risk b^{τ} for each trial/time interval combination (τ), Binom is the probability mass function of the binomial distribution, and for each trial/time interval combination, e_c^{τ} and n_c^{τ} are the numbers of events (symptomatic infections) and total number of individuals in the control group, e_t^{τ} and n_t^{τ} are the numbers of events and total number of individuals in the treatment group in the corresponding trial time interval, and b^{τ} is the baseline risk which is reduced by the efficacy of treatment for the treatment group in the same interval. The initial guess of the trial/time interval combinations baseline risk used in the optimization were $b^{\tau} = e_c^{\tau}/n_c^{\tau}$. The parameter $c(\tau)$ is the (\log_{10}) concentration of monoclonal antibodies (in the fold IC50 scale) in the trial time interval τ .

The negative log-transform of this likelihood function was minimised using the *nlm* optimiser in the R statistical software package to estimate the log-transform of the model parameter k, c_{50} and m . The optimiser was run 25 times using randomly generated initial parameters for the transformed parameters m, k, c_{50} drawn uniformly from the ranges $\log(m) = [\log(0.6), \log(0.99)]$, $\log(k) = [\log(0.1), \log(100)]$ and $\log(c_{50}) = [\log(10), \log(5000)]$.

460

Model fitting was used for parameter estimation and hypothesis testing, the latter using a likelihood ratio test of nested models, and Akaike Information Criterion (AIC) for non-nested models. In all fitting, we excluded the earliest time interval of each study to account for the rapid change of the antibody concentration over this time interval and ensured exclusion of treatments which occurred after an unidentified infection (see Table S1 for the time intervals included in the analysis).

The confidence intervals of the fitted model and the fitted model parameters were estimated by parametric bootstrapping. The covariance matrix of the fitted parameters was calculated from the inverse of the Hessian from the model fit²¹. The log-transform of the

470

three model parameters m , k , and c_{50} were drawn randomly from a multinomial Gaussian distribution 10,000 times, using the covariance matrix (and the function *rmvnorm* from the package *mvtnorm* in the R statistic package⁵⁶). The 2.5% and 97.5% percentiles of the estimated parameters provided an estimate of the 95% confidence intervals. Similarly,
475 evaluating the confidence intervals of the fitted model in Figure 2 was estimated by evaluating the model at each neutralisation titre using each of the 10,000 bootstrapped parameter estimates and taking the 2.5% and 97.5% percentiles of the evaluated models at each antibody concentration.

480 **Timing of protection**

Using the dose-response curve, we can predict the protection over time and how long the protection remains above 50% protection (Figure 3). With the meta-analysis of IC₅₀'s against different variants, the timing of protection can be predicted not only for the ancestral strain but also for variants (Figure 3).
485 We assumed that the concentration of mAbs declines exponentially from the time of the peak concentration for each antibody (extracted from the data) with a half-life fitted to the data (linear fit to log-transformed concentration, see supplementary methods). With the concentration over time and the efficacy by concentration (Figure 2), we computed the protection over time. For variants, we scaled the concentration data by dividing the fold-
490 IC₅₀ concentration from the meta-analysis by the fold-increase in the IC₅₀ and computed the protective efficacy in the same way as against ancestral. The time from treatment to 50% protection is then the time until the concentration falls to the concentration that gives 50% protection which we computed from the concentration over time (using a linear fit to the log-transformed concentration data, Figure S2). The uncertainty in the time to 50%
495 protection is due to the uncertainty in the concentration that gives 50% protection. The upper and lower bounds for the time to 50% protection are the time to reach the upper or lower bound of the 95% CI of the concentration that gives 50% protection, respectively.


500 **Comparison of the relationship between neutralizing antibodies and protection for vaccination and monoclonal antibody prophylaxis**

To compare the mAb data with the data from vaccine studies (Figure 4a), we aimed to match the mAb data as closely as possible with the vaccine studies. Thus, we restricted the

mAb data to 2-3 months after treatment, patients who are PCR-negative at baseline, and cases later than 1-2 weeks after treatment (the start of follow-up in vaccine studies) (Figure 4a). We then used a generalised linear mixed model (GLMM) with a binomial error family and logarithmic link function (see above). The model included random intercepts for different trials and a treatment variable with the factors “control”, “mAb” and “vaccine”. The treatment effect of mAbs and vaccination was compared by testing if there is a significant difference between the coefficients for mAb treatment and vaccination (using the *glht* function from the *multcomp* package⁵⁷).

To further compare the efficacy of vaccination and prophylactic mAb treatment, we normalized the concentration to a common ‘fold-convalescent’-scale (as previously described^{12,14}). Using a maximum likelihood approach (see above), we fitted logistic dose-response curves to the monoclonal antibody and vaccine data (Figure 4). We tested whether there is a significant difference between the monoclonal antibody prophylaxis and vaccination by fitting all data with the same parameters for the two types of treatment (Figure 4b, black curve) and compared this fit to fits which have different parameters for antibody treatment and vaccination (e.g., Figure 4b red and blue curves). Different models were compared with a likelihood ratio test. The p-values for different model comparisons are reported in Figure S3 and parameter values for different fits in Table S6. Confidence intervals were calculated using bootstrapping, as described above.

Figures and Legends



525

Figure 1: Reported protection and antibody concentration from three RCTs of monoclonal

antibodies in preventing COVID-19. The efficacy at each time interval is shown in blue (horizontal error bars indicate time interval and vertical error bars represent 95% CI). The antibody concentration is shown in black. (a) Antibody concentration data for

530

tixagevimab/cilgavimab was extracted from Levin et al. (b) Single administration of casirivimab/imdevimab data is a combination of data from O'Brien et al. and Herman et al. who report on the same clinical trial over different follow-up intervals. Efficacy data was

reported frequency over the first four weeks in O'Brien et al. (diamonds), and monthly for eight months in Herman et al. (circles). Antibody concentration data was reported up to day

535

168 in O'Brien et al. (solid line, panel b), and modelling of the pharmacokinetic profile of the antibody concentration between 168 and 240 days (dashed line, panel b). (c) Isa et al. reported

efficacy and *in vivo* concentration after repeated administration of 1.2g of casirivimab/imdevimab every four weeks. Hence, the antibody concentration did not

540

decline as in the other studies. (d) The modelled concentration of adintrevimab after a single administration was extracted from the study by Schmidt et al. The efficacy of

adintrevimab was reported both when the delta variant was dominant (circles) and when omicron variants BA.1 and BA.1.1 were dominant (triangles).

545




Figure 2: Dose-response relationship between antibody concentration and protection. The estimated geometric mean antibody concentration and protective efficacy in the matching study and time interval (expressed as a fold of the *in vitro* IC50 of each antibody) is shown.

550

Horizontal error bars indicate the range of the (means of the) antibody concentrations reported during each time interval, and vertical error bars indicate the 95% confidence interval of the efficacy. We estimate a dose-response relationship (black line) by fitting a logistic model to the data and estimate the 95% confidence intervals using parametric bootstrapping (grey shading). Efficacy data reported early (in the first 7, 10, 28 or 90 days,

555

depending on the first time point reported in the study) after treatment were excluded from the model fitting (low opacity data points), as antibody concentration changed rapidly over this time interval and ensured exclusion of incidental treatments which occurred after an unidentified infection.

560





Figure 3: Duration of protection against SARS-CoV-2 variants. Using the dose-response relationship of antibody concentration and protective efficacy in Figure 2 and the estimated half-life of antibodies after treatment (28.8 days (95% CI: 26.6 – 31.3) for casirivimab/imdevimab, 94.9 days (95% CI: 84.2 – 108.9) for tixagevimab/cilgavimab and 134.7 days (95% CI: 132.7 – 136.7) for adintrevimab, see Figure S2 and Table S5), we predict the efficacy over time for these antibody combinations (black line, a, b and c). We also estimate the protection over time curves of these antibody combinations given different fold-increases in the IC50 that may be experienced to new variants (coloured lines). (d) For each hypothetical loss of potency of these antibodies (i.e. fold increase in IC50, x-axis), we predict the number of days each antibody will remain above 50% protection (i.e. the number of days the mean antibody concentration will remain above the level associated with 50% protection of 54-fold-IC50, Figure 2). The shaded regions indicate the 95% confidence interval of the duration of protection (using the 95% CI for 50% protection: 16 – 183). We note that the casirivimab/imdevimab and the tixagevimab/cilgavimab combinations and adintrevimab are expected to maintain more than 50% protection for approximate 269, 670 and 903 days against a variant with the same IC50 as the ancestral

virus, respectively, and can tolerate up to a 319.6-, 106.8- and 89.6-fold drop in potency to a new variant, respectively, and still be expected to maintain 30 days of more than 50%
 580 protective efficacy. The increase in IC50 against BA.4/5 is sufficient to reduce efficacy below 50% for all three products.



585 **Figure 4: Comparison of dose response curves after vaccination and monoclonal antibody administration.** (a) We compared the overall efficacies of prophylactic mAb treatment and high-potency RNA vaccines and found a significantly difference in efficacy ($p = 0.002$, GLMM, see Methods). The mean efficacy in the mAb studies was 84.8% (95% CI: 76.0 – 90.8%) and for the high-potency mRNA vaccines the mean efficacy was 94.5% (95% CI: 91.6 – 96.7%). (b) We normalized the neutralization titers to a common scale of ‘fold-of-convalescent’ titer to allow direct comparison taking differences in the neutralization into account. We then fitted the combined data with a logistic model in which the same three parameters were used for maximum efficacy, neutralization for 50% efficacy, and slope (b, black line). Next, we allowed parameters to vary between the monoclonal and vaccination data fitting. Here we show the best fit to the data was with a model where the slope is allowed to vary between mAb studies (blue) and vaccine studies (red) but the maximal efficacy and the neutralization giving 50% protection were the same for both types for treatment (based on model comparisons with the likelihood-ratio test, see Figure S3).
 595

Acknowledgements:

Ethics statement:

605 This work was approved under the UNSW Sydney Human Research Ethics Committee (approval HC200242).

Funding statement:

610 This work is supported by an Australian government Medical Research Future Fund awards GNT2002073 (to MPD, SJK), MRF2005544 (to SJK and MPD), MRF2005760 (to MPD), an NHMRC program grant GNT1149990 (SJK and MPD), and the Victorian Government (SJK). DSK and SJK are supported by NHMRC fellowships. DC, ZKM and EMW and MPD are supported by NHMRC Investigator grants. KLC is supported by PhD scholarship from the Leukaemia Foundation, the Haematology Society of Australia and New Zealand (HSANZ) and
615 Monash University. The Melbourne WHO Collaborating Centre for Reference and Research on Influenza is supported by the Australian Government Department of Health.

Competing Interests statement:

620 The authors declare no competing interests.

Data Availability Statement:

All data and code will be made available on GitHub upon publication.

References

- 625 1. Cromer, D., *et al.* Neutralising antibodies predict protection from severe COVID-19. *medRxiv*, 2022.2006.2009.22275942 (2022).
2. Teh, J.S.K., *et al.* Immunogenicity of COVID-19 vaccines in patients with hematologic malignancies: a systematic review and meta-analysis. *Blood Adv* **6**, 2014-2034 (2022).
- 630 3. Collier, D.A., *et al.* Age-related immune response heterogeneity to SARS-CoV-2 vaccine BNT162b2. *Nature* **596**, 417-422 (2021).
4. Levin, M.J., *et al.* Intramuscular AZD7442 (Tixagevimab-Cilgavimab) for Prevention of Covid-19. *N Engl J Med* **386**, 2188-2200 (2022).
5. Isa, F., *et al.* Repeat subcutaneous administration of casirivimab and imdevimab in adults is well-tolerated and prevents the occurrence of COVID-19. *Int J Infect Dis* **122**, 585-592 (2022).
- 635 6. O'Brien, M.P., *et al.* Subcutaneous REGEN-COV Antibody Combination to Prevent Covid-19. *N Engl J Med* **385**, 1184-1195 (2021).
7. Streinu-Cercel, A., *et al.* Efficacy and Safety of Regdanvimab (CT-P59): A Phase 2/3 Randomized, Double-Blind, Placebo-Controlled Trial in Outpatients With Mild-to-Moderate Coronavirus Disease 2019. *Open Forum Infect Dis* **9**, ofac053 (2022).
- 640 8. Weinreich, D.M., *et al.* REGEN-COV Antibody Combination and Outcomes in Outpatients with Covid-19. *N Engl J Med* **385**, e81 (2021).
9. Chen, P., *et al.* SARS-CoV-2 Neutralizing Antibody LY-CoV555 in Outpatients with Covid-19. *N Engl J Med* **384**, 229-237 (2021).
- 645 10. Dougan, M., *et al.* Bamlanivimab plus Etesevimab in Mild or Moderate Covid-19. *N Engl J Med* **385**, 1382-1392 (2021).
11. Bushell, E.S., *et al.* Paternal effect of the nuclear formin-like protein MISFIT on Plasmodium development in the mosquito vector. *PLoS Pathog* **5**, e1000539 (2009).
- 650 12. Stadler, E., *et al.* Determinants of passive antibody effectiveness in SARS-CoV-2 infection. *medRxiv*, 2022.2003.2021.22272672 (2022).
13. Schmidt, P., *et al.* Antibody-mediated protection against symptomatic COVID-19 can be achieved at low serum neutralizing titers. *medRxiv*, 2022.2010.2018.22281172 (2022).
- 655 14. Khoury, D.S., *et al.* Neutralizing antibody levels are highly predictive of immune protection from symptomatic SARS-CoV-2 infection. *Nat Med* **27**, 1205-1211 (2021).
15. Herman, G.A., *et al.* Efficacy and safety of a single dose of casirivimab and imdevimab for the prevention of COVID-19 over an 8-month period: a randomised, double-blind, placebo-controlled trial. *Lancet Infect Dis* **22**, 1444-1454 (2022).
- 660 16. Cohen, M.S., *et al.* Effect of Bamlanivimab vs Placebo on Incidence of COVID-19 Among Residents and Staff of Skilled Nursing and Assisted Living Facilities: A Randomized Clinical Trial. *JAMA* **326**, 46-55 (2021).
17. Cox, M., *et al.* SARS-CoV-2 variant evasion of monoclonal antibodies based on in vitro studies. *Nat Rev Microbiol* (2022).
- 665 18. Tao, K., Tzou, P.L., Kosakovsky Pond, S.L., Ioannidis, J.P.A. & Shafer, R.W. Susceptibility of SARS-CoV-2 Omicron Variants to Therapeutic Monoclonal Antibodies: Systematic Review and Meta-analysis. *Microbiol Spectr* **10**, e0092622 (2022).
19. FDA. Fact Sheet for Healthcare Providers Emergency Use Authorization (EUA) of Sotrovimab. (2022).
- 670

20. FDA. FDA updates Sotrovimab emergency use authorization Update [4/5/2022]. (2022).
21. Cromer, D., *et al.* Neutralising antibody titres as predictors of protection against SARS-CoV-2 variants and the impact of boosting: a meta-analysis. *The Lancet Microbe* **3**, e52-61 (2021).
- 675 22. Feng, S., *et al.* Correlates of protection against symptomatic and asymptomatic SARS-CoV-2 infection. *Nat Med* **27**, 2032-2040 (2021).
23. Corbett, K.S., *et al.* Immune correlates of protection by mRNA-1273 vaccine against SARS-CoV-2 in nonhuman primates. *Science* **373**, eabj0299 (2021).
- 680 24. Gilbert, P.B., *et al.* Immune correlates analysis of the mRNA-1273 COVID-19 vaccine efficacy clinical trial. *Science* **375**, 43-50 (2022).
25. Plotkin, S.A. & Gilbert, P.B. Nomenclature for immune correlates of protection after vaccination. *Clin Infect Dis* **54**, 1615-1617 (2012).
26. Polack, F.P., *et al.* Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med* **383**, 2603-2615 (2020).
- 685 27. Baden, L.R., *et al.* Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N Engl J Med* **384**, 403-416 (2021).
28. Khoury, D.S., Cao, P., Zaloumis, S.G., Davenport, M.P. & Interdisciplinary Approaches to Malaria, C. Artemisinin Resistance and the Unique Selection Pressure of a Short-acting Antimalarial. *Trends Parasitol* **36**, 884-887 (2020).
- 690 29. Khoury, D.S., *et al.* Measuring immunity to SARS-CoV-2 infection: comparing assays and animal models. *Nat Rev Immunol* **20**, 727-738 (2020).
30. Selva, K.J., *et al.* Tear antibodies to SARS-CoV-2: implications for transmission. *Clin Transl Immunology* **10**, e1354 (2021).
- 695 31. Bergwerk, M., *et al.* Covid-19 Breakthrough Infections in Vaccinated Health Care Workers. *N Engl J Med* **385**, 1474-1484 (2021).
32. Earle, K.A., *et al.* Evidence for antibody as a protective correlate for COVID-19 vaccines. *Vaccine* **39**, 4423-4428 (2021).
- 700 33. Gilbert, P.B., *et al.* Immune correlates analysis of the mRNA-1273 COVID-19 vaccine efficacy clinical trial. *Science*, eab3435 (2021).
34. Cromer, D., *et al.* SARS-CoV-2 variants: levels of neutralisation required for protective immunity. *medRxiv*, 2021.2008.2011.21261876 (2021).
35. Cromer, D., *et al.* Relating in vitro neutralisation level and protection in the CVnCoV (CUREVAC) trial. *Clin Infect Dis In Press*, ciac075 (2022).
- 705 36. Khoury, D.S., *et al.* Correlates of protection, thresholds of protection, and immunobridging in SARS-CoV-2 infection. *medRxiv*, 2022.2006.2005.22275943 (2022).
37. McMahan, K., *et al.* Correlates of protection against SARS-CoV-2 in rhesus macaques. *Nature* **590**, 630-634 (2021).
- 710 38. Schafer, A., *et al.* Antibody potency, effector function, and combinations in protection and therapy for SARS-CoV-2 infection in vivo. *J Exp Med* **218**(2021).
39. Vaida, F., Fitzgerald, A.P. & Degruittola, V. Efficient Hybrid EM for Linear and Nonlinear Mixed Effects Models with Censored Response. *Comput Stat Data Anal* **51**, 5718-5730 (2007).
- 715 40. Vaida, F. & Liu, L. Fast Implementation for Normal Mixed Effects Models With Censored Response. *J Comput Graph Stat* **18**, 797-817 (2009).

41. Cameroni, E., *et al.* Broadly neutralizing antibodies overcome SARS-CoV-2 Omicron antigenic shift. *Nature* **602**, 664-670 (2022).
- 720 42. Dejnirattisai, W., *et al.* SARS-CoV-2 Omicron-B.1.1.529 leads to widespread escape from neutralizing antibody responses. *Cell* **185**, 467-484 e415 (2022).
43. Hoffmann, M., *et al.* The Omicron variant is highly resistant against antibody-mediated neutralization: Implications for control of the COVID-19 pandemic. *Cell* **185**, 447-456 e411 (2022).
- 725 44. Iketani, S., *et al.* Antibody evasion properties of SARS-CoV-2 Omicron sublineages. *Nature* **604**, 553-556 (2022).
45. Liu, L., *et al.* Striking antibody evasion manifested by the Omicron variant of SARS-CoV-2. *Nature* **602**, 676-681 (2022).
46. Lusvarghi, S., *et al.* SARS-CoV-2 BA.1 variant is neutralized by vaccine booster-elicited serum but evades most convalescent serum and therapeutic antibodies. *Sci Transl Med* **14**, eabn8543 (2022).
- 730 47. Gruell, H., *et al.* mRNA booster immunization elicits potent neutralizing serum activity against the SARS-CoV-2 Omicron variant. *Nat Med* **28**, 477-480 (2022).
48. Gruell, H., *et al.* Delineating antibody escape from Omicron sublineages. *bioRxiv*, 2022.2004.2006.487257 (2022).
- 735 49. Ju, B., *et al.* Immune escape by SARS-CoV-2 Omicron variant and structural basis of its effective neutralization by a broad neutralizing human antibody VacW-209. *Cell Res* **32**, 491-494 (2022).
50. Sheward, D.J., *et al.* Neutralisation sensitivity of the SARS-CoV-2 omicron (B.1.1.529) variant: a cross-sectional study. *Lancet Infect Dis* **22**, 813-820 (2022).
- 740 51. Yamasoba, D., *et al.* Virological characteristics of the SARS-CoV-2 Omicron BA.2 spike. *Cell* **185**, 2103-2115 e2119 (2022).
52. Ikemura, N., *et al.* SARS-CoV-2 Omicron variant escapes neutralization by vaccinated and convalescent sera and therapeutic monoclonal antibodies. *medRxiv*, 2021.2012.2013.21267761 (2021).
- 745 53. Tada, T., *et al.* Increased resistance of SARS-CoV-2 Omicron variant to neutralization by vaccine-elicited and therapeutic antibodies. *EBioMedicine* **78**, 103944 (2022).
54. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* **67**, 1 - 48 (2015).
- 750 55. R Core Team. R: A Language and Environment for Statistical Computing. *R Foundation for Statistical Computing Vienna, Austria*, <https://www.R-project.org/> (2019).
56. Genz, A., *et al.* mvtnorm: Multivariate Normal and t Distributions., <https://CRAN.R-project.org/package=mvtnorm> (2020).
- 755 57. Hothorn, T., Bretz, F. & Westfall, P. Simultaneous inference in general parametric models. *Biom J* **50**, 346-363 (2008).