perpetuit	y.
It is made available under a CC-B	Y 4.0 International license .

1	The relationship between blastocyst morphology and the sex ratio of singleton
2	births in single blastocyst transfer cycles: A retrospective cohort study
3	Running title: Relationship between blastocyst morphology and the sex ratio of
4	singleton births
5	
6	Jijun Hu, Yichun Guan [*] , Rujing Yang, Lijun Sun, Hua Lou, Xingling Wang
7	The Third Affiliated Hospital of Zhengzhou University, Reproductive Medicine
8	Center, Zhengzhou, China
9	
10	*Corresponding Author:
11	Yichun Guan
12	The Third Affiliated Hospital of Zhengzhou University, Reproductive Medicine
13	Center, Zhengzhou, China
14	Tel: +86-13673459605
15	E-mail: lisamayguan@163.com
16	
17	Abstract
18	Background: To study the relationship between blastocyst morphology and the sex
19	ratio of singleton births in single blastocyst transfer cylces.
20	Methods: This retrospective cohort study included women who underwent in vitro
21	fertilization/intracytoplasmic sperm injection-embryo transfer (IVF/ICSI-ET) from
22	January 2017 to December 2019 and obtained a single live baby following a single
23	blastocyst transfer. The sex ratio which influenced by many possible factors were
24	studied. Age-stratified analysis was performed for the maternal and paternal groups
25 NO	younger and older than 35 years. The 3108 cycles included here were grouped by sex TE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

at birth (male: 1777 cycles and female: 1331 cycles). Blastocysts were conceived via
2834 IVF and 274 ICSI treatments, and were used for 1054 fresh ET cycles and 2054
frozen-thawed embryo transfer (FET) cycles.

Results: Sex ratio among singleton of poor blastocyst decreased significantly 29 compared to those of good blastocyst (51.5% vs 60%, P<0.001). The results of 30 multivariate logistic analysis showed that high-quality blastocysts had a 151% higher 31 32 probability of being male singleton than poor-quality blastocysts (P < 0.001). Further stratified analysis shows that in both group younger than 35 the sex ratio increased 33 34 significantly in grades A and/or B trophectoderm (P < 0.001) and good-quality blastocysts (P<0.001). While grade A inner cell mass (ICM) degree was significantly 35 associated with a lower sex ratio than grade B ($P \le 0.05$). 36

37 **Conclusions:** A single high-quality blastocyst transfer was significantly associated 38 with a higher neonatal sex ratio. Grade A trophectoderm and grade B ICM contributed 39 to improving the implantation potential of male blastocysts, especially in infertile men 40 or women \leq 35 years old.

41

42 Key Words: sex ratio; single blastocyst transfer; single live birth; blastocyst quality;
43 inner cell mass; trophectoderm

44

45 Background

IVF was first developed over 40 years ago. Millions of babies have since been born as a result of infertility treatment worldwide. Routine blastocyst culture and transfer using sequential culture media in IVF/ICSI is associated with a significant increase in implantation rates and clinical pregnancy rate.However,there are some adverse effects in blastocyst transfer such as a male-biased imbalance in the sex ratio

of newborns and an increased incidence of monozygotic twinning (MZT). [1-3]

A three-part scoring system, which based on blastocyst expansion, inner cell mass, and trophectoderm development formulated by Gardner and Schoolcraft [4], aiming to grade human blastocysts before transfer. The scoring system brought great convenience to embryologists and researchers' studies on the development and viability of blastocyst. In many cases, blastocysts achieving the best morphologic scores often achieve implantation and produce a live birth, and those with high scores on trophectoderm have relatively easy access to a male child (2021).

59 The human sex ratio is often expressed as the number of males per 100 female live births [5]. In Western gender-neutral countries, the sex ratio at birth is estimated 60 to be approximately 1.06 [6]. In Japan, the overall neonatal sex ratio has steadily 61 62 declined from 1.079 in 1970 to as low as 1.056 in 2016 (Ministry of Health, Labour and Welfare Vital Statistics of Japan 2016). In the 2000 census in China, the reported 63 sex ratio at birth was 117 per 100 female births, which appears to be due to a 64 declining fertility rate and a traditional preference for males, a phenomenon that is 65 common to many other Asian countries [7]. A recent study by Fan et al. reported that 66 the sex ratio at birth gradually declined to normal levels (1.047 with the universal 67 child policy) following the gradual rollout of the two-child policy in China [8]. 68

The sex ratio at birth from assisted artificial technology cycles ranges from 1.03–1.25. However, there are discrepancies in the sex ratio between embryos generated by standard IVF/ICSI, and between cleavage and single blastocyst transfer (SBT) embryos. As SBT is the most effective approach for avoiding multiple pregnancies, SBT has gradually become the preferred embryo transfer strategy throughout the world [9-14]. However, a male-biased imbalance in the sex ratio is one of the important potential limitations in blastocyst transfer [1, 3]. Several studies have

reported a significantly higher sex ratio at birth among infants born after SBT (54.1%)

than among those born after cleavage-stage (day 3) transfer cycles (49.9%) [15, 16].

In practice, embryologists tend to select preferentially better morphology and more developmentally advanced blastocysts for transfer, because good-quality blastocyst has higher implantation rate and clinical pregnancy rate. The contribution of these parameters to the implantation potential or sex ratio is not fully known. Some studies have reported that trophectoderm morphology may be predictive of pregnancy outcomes and sex ratio related to SBT [17, 18].

We intended to study the relationship between blastocyst morphology and the sex ratio of singleton births in single blastocyst transfer. To that end, here we analyzed clinical data for the SR of IVF/ICSI children conceived after SBT between 2017 and 2019. The variables considered in this research included: assisted reproductive technologies (ART) such as IVF, ICSI, FET, or ET, maternal or paternal age, and year of infertility.

90

91 Methods

92 Study subjects

This retrospective cohort study included the singleton offspring born to mothers 93 94 after successful treatment cycles following the autologous SBT of 95 non-preimplantation genetic testing (PGT) blastocysts at the Reproductive Medicine Center of the Third Affiliated Hospital of Zhengzhou University between January 96 2017 and December 2019. The study was carried out after the protocol was approved 97 98 by the Institutional Review Board of The Third Affiliated Hospital of Zhengzhou University. Informed consent was waived due to retrospective design of the study 99 100 based on our local ethics committee protocols.

101 **Ovarian stimulation protocol**

Each female patient underwent a conventional ovarian hyperstimulation procedure 102 [19, 20] involving a gonadotrophin releasing hormone agonist or antagonist or mild 103 104 stimulation protocol. The physician adjusted the starting dose according to the 105 patient's age, body mass index (BMI) and ovarian reserve. Ovarian follicle development was monitored based on serum estradiol and transvaginal 106 107 ultrasonographic measurements. Oocytes were retrieved transvaginally 36-38h after human chorionic gonadotrophin (Serono, Aubonne, Switzerland) administration when 108 109 at least 40% of the follicles had reached or exceeded an average diameter of 18mm as determined by ultrasound. The follicles were aspirated using a single-lumen needle 110 111 attached to a syringe under transvaginal ultrasound guidance. The oocytes were then 112 inseminated via conventional IVF/ICSI.

113 Laboratory protocol

Embryos were placed in an incubator (K-MINC-1000, Cook, USA) and cultured 114 at 37°C under 6% CO₂ and 5% O₂. Embryos were cultured in G-1TM plus (Vitrolife, 115 Sweden) from the pronucleate stage to day 3. On day 3 of culture, the embryo cell 116 number and morphology were scored. Day 3 embryos are scored from I to IV. Classes 117 I-III represent embryos that are available for transfer. After scoring, the embryos 118 were subsequently transferred to G-2TM plus (Vitrolife, Sweden) medium for an 119 120 additional 2-3 days of culture (days 5-6 of development) [21]. On day 5, the 121 percentage of blastocyst formation was determined, and one blastocyst was selected for fresh cycle transfer, and the untransferred blastocysts were cryopreserved. On day 122 123 5 or 6 after insemination, blastocyst morphology was evaluated according to that described by Gardner and Schoolcraft [22]. Blastocysts expansion degree > 3, inner 124 cell mass (ICM) grade A/B, and trophectoderm grade A/B (AA, AB, BA, BB) were 125

considered high-quality blastocysts. Blastocysts expansion degree > 3, ICM grade A/B and trophectoderm grade C (AC, BC) were considered low-quality blastocysts [23]. All gamete and embryo manipulations occurred in a clean bench designed to control humidity, temperature, and pH fluctuation.

A laser hole caused the blastocysts to collapse, and the shrunken blastocyst was 130 vitrified using a Cryotop device (Kitazato BioPharmao. Shizuoka, Japan). 131 132 Vitrification and warming kits were used as media. For vitrification, the blastocysts were first equilibrated in solution I (7.5% v/v ethylene glycol [EG] and 7.5% v/v 133 134 dimethyl sulfoxide [DMSO]) at room temperature for 10 min and then placed into vitrification solution II (15% v/v EG, 15% v/v DMSO, and 0.5 M sucrose) for 1 min. 135 Subsequently, blastocysts were individually loaded onto Cryotop in a volume of 136 137 <0.1µL and quickly plunged into liquid nitrogen. The blastocyst was warmed by placing the Cryotop in thawing solution (1.0 M sucrose) for 1 min at 37°C. The 138 blastocyst was then transferred to dilution solution (0.5 M sucrose) for 3 min at room 139 140 temperature, then washed twice in washing solution at room temperature for 3 min 141 each. The blastocyte was transferred into a 20-µL droplet of culture medium (G-2; Vitrolife) under mineral oil. Blastocysts were transferred 1–2 h post-warming. 142

143

144 Embryo transfer

Fresh single blastocyst transfer was performed 5 days after fertilization in our Reproductive Center. ET was routinely performed under abdominal ultrasound guidance. Fresh ET was cancelled in cases with severe ovarian hyperstimulation syndrome (OHSS), an endometrial thickness of ≤ 7 mm, progesterone levels $\geq 2\mu g/L$ on the human chorionic gonadotropin (HCG) trigger day, and some other special conditions; all the embryos were cryopreserved and FETs were subsequently

151 performed to decrease the rate of OHSS and to improve the clinical pregnancy rate.

FET and clinical pregnancy confirmed were performed referring to the descriptions of the previous study[17]. The singleton SR was primary outcome measure of this study. The sex ratio at birth was calculated as the number of males per 100 female live births.

156 Statistical analysis

157 All data analyses were performed using Statistical Package for Social Sciences (SPSS) software package version 22.0 (IBM Corp, Armonk, NY, USA). For 158 159 continuous variables, normally distributed variables are expressed as the mean and standard deviation, while non-normally distributed variables are shown as the median 160 and interquartile range. Categorical variables are summarized as frequencies with 161 162 proportions. The unpaired Student's t test or Mann–Whitney U test was used for 163 comparing the difference between the two groups. Univariate regression analysis was performed for factors, including maternal age, paternal age, mean duration of 164 infertility, mean maternal BMI, type of infertility, type of infertility (%), day of 165 blastocyst transfer, type of fertilization, quality of transferred blastocyst, ICM, and 166 trophectoderm grade. Both crude and adjusted multivariable logistic regression 167 models for sex ratio of live singleton were established (enter). For the adjusted 168 169 multivariate logistic regression model, factors including maternal age, paternal age, 170 BMI, duration of infertility, type of infertility, and day of blastocyst transfer were further adjusted. P value <0.05 was set as the threshold of statistical significance. 171

172

173 **Results**

174 Study participants and baseline characteristics

175 A

A total of 5632 SBT cycles of women who underwent IVF/ICSI-ET were

176 screened. Of these, 177 cycles that underwent PGT were excluded. Of the 5285 autologous cycles remaining, there were 328 miscarriages, 29 ectopic pregnancies, 177 and 56 cycles that resulted in monozygotic twins. After excluding these cycles, there 178 179 were 3108 cycles that resulted in singleton live births, which were included in the analysis. There were 1054 (33.9%) fresh blastocyst transfer cycles and 2054 (66.1%) 180 frozen blastocyst transfer cycles, 2070 (66.6%) high-quality and 1038 (33.4%) 181 poor-quality single blastocyst FETs. The blastocysts were conceived via 2834 182 (91.2%) IVF and 274 (8.8%) ICSI treatments. The analyzed cycles yielded 1777 male 183 184 infants and 1331 female infants, and the sex ratio was 133% (133/100). All the SBT cycles of each patient was analyzed to avoid repeated measures bias. Patients who 185 underwent PGT or used donor oocytes or donor sperm were excluded from this study. 186 187 Figure 1 presents the identification and collation process of the study sample. Table 1 188 describes the baseline characteristics of the included treatment cycles.

The results of the univariable analysis for the potential factors influencing sex 189 190 ratio among singleton live births revealed that transfer of a single high-quality blastocyst resulted in a significantly higher sex ratio than SBT of a poor-quality 191 blastocyst (60% vs. 51.5%, P < 0.001, Fig. 2). Transfer of blastocysts with a higher 192 TE grade and lower ICM grade resulted in a significantly higher sex ratio. Maternal 193 194 age, paternal age, duration of infertility, maternal BMI, main infertility cause, type of 195 fertilization, or day of embryo transfer had no significant influence on the sex ratio 196 (Table 2).

197 Multivariate analysis

Multivariate analysis was performed after adjusting for potential confounding factors. Based on the results of the analysis, we identified significant associations of sex ratio among singleton births with the morphological characteristics of blastocysts.

201 The results of the multivariate analysis revealed that high-quality blastocysts have a 190% higher probability of being male than poor-quality blastocysts (aOR, 1.964; 202 95% CI, 1.559–2.474; P < 0.001). However, blastocysts with grade A ICM had a 37% 203 lower probability of being male than in blastocysts with grade B trophectoderm (aOR 204 0.632; 95% CI, 0.529–0.757; P < 0.001). Moreover, when the trophectoderm was of 205 grade A, there was a 190% higher probability of being male than when the 206 trophectoderm was of grade C (aOR, 1.964; 95% CI, 1.559–2.474; P < 0.001; Table 207 3). 208

209 Stratified analysis by maternal and paternal age

We also determined relationship between quality classification of ICM grade, TE 210 grade of transferred blastocysts and sex ratio according to parents aged 35 and older. 211 212 When the maternal age was \leq 35, blastocysts with grade A ICM had a significantly 213 lower likelihood of being male than with grade B ICM (OR, 0.803; 95% CI, 0.677-0.951; P = 0.011). Similarly, when the paternal age was <35 years, blastocysts with 214 215 grade A ICM had a significantly lower likelihood of being male than blastocysts with grade B ICM (OR, 0.791; 95% CI, 0.663–0.942; P = 0.009). Blastocysts with grades 216 A and B TE in women \leq 35 years and grade B TE in men \leq 35 years had a significantly 217 higher sex ratio than those with grade C TE (women \leq 35 years grade B TE: OR, 218 219 1.401; 95% CI, 1.172–1.676 and grade A TE: OR, 1.589; 95% CI, 1.282–1.970; P 220 <0.001 for both) (men \leq 35 years grade B TE: OR, 1.555; 95% CI, 1.245–1.942; P <0.001). Furthermore, good-quality blastocysts were significantly more likely to be 221 male in both women and men aged \leq 35 years (OR: 1.461, 95% CI, 1.238–1.725; P 222 <0.001 for women and OR: 1.442; 95% CI, 1.214–1.712; and P <0.001 for men) 223 (Table 4) 224

225

226 Discussion

In the present study, SBT was used to avoid the complications associated with 227 multiple pregnancies and maintain higher rates of implantation, pregnancy, and live 228 229 birth in couples undergoing ART. Our results following the analysis of 3108 cycles revealed that high-quality blastocysts had a significantly higher probability of being 230 male than poor-quality blastocysts. However, blastocyst culture conditions 231 232 contributed to preferential selection of male blastocysts for transfer because of the advanced developmental rates in vitro. It is worthwhile to note that some studies 233 234 explored the relationship between the stage at which ET occurs and the SSR, and found that ET at the blastocyst stage tends to increase the SSR compared with ET at 235 the cleavage stage, and this effect is independent of other factors [24]. 236

237 Among the 3108 IVF/ICSI cycles over a three-year period, there were 1777 males and 1331 females, making the SR 133 males per 100 females, which was 238 consistent with the ratios reported in previous studies. A large nationwide cohort 239 study in Japan showed a higher sex ratio after blastocyst transfer relative to 240 spontaneous conception (OR: 1.095; 95% CI: 1.001-1.198) [3]. The study also found 241 that implantation with high-quality blastocysts was associated with an increased sex 242 ratio during SBT cycles. The similar result that blastocyst transfer was associated with 243 increased sex ratio was also found in a retrospective cohort study [25]. 244

We also explored the effect of FET on SSR. Very few studies have reported differences in the SSR following fresh and frozen ET, and the results are controversial [26, 27]. However, the results of our preliminary univariable analysis after controlling for related factors, including maternal age, paternal age, mean duration of infertility, mean maternal BMI, type of infertility, type of infertility (%), day of blastocyst transfer, type of fertilization, quality of transferred blastocyst, ICM, and

trophectoderm grade, showed that the SSR was not related with FET (OR = 0.871, 95% CI: 0.750-1.013, P = 0.074). In most reproductive medical centers in China, blastocyst transfers are usually performed in FET cycles, and our center was no exception. It may be that blastocyst transfer and not FET is responsible for the skewing of the SSR toward males.

Previous studies have explored the effect of the IVF/ICSI insemination method 256 257 on sex ratio. [28] ICSI, which is primarily used to treat male factor infertility, is now widely used all over the world. Data from other countries have shown that ICSI use 258 259 significantly reduces the proportion of male babies compared with IVF (48.8% versus 51.4% in America: 50.0% vs 53.0% in Australia and New Zealand) [15, 29]. 260 Although the sex ratio with ICSI insemination was found to be lower than that with 261 262 IVF insemination in our study, this difference was not statistically significant. One study showed that ICSI reduces SSR only in cleavage-stage embryo transfer cycles, 263 but not in blastocyst transfer cycles [24]. Currently, it is unclear why ICSI 264 insemination reduces SSR. Tarín et al. compared sex ratios at implantation and birth 265 in cleavage- and blastocyst-stage transfer cycles to propose molecular mechanisms 266 accounting for differences in post-implantation mortality in male and female offspring 267 and consequent variations in the sex ratios at birth in ART cycles. They reported that 268 in vitro culture-induced precocious X-chromosome inactivation together with 269 270 ICSI-induced decrease in the number of trophectoderm cells in female blastocysts 271 may account for the preferential female mortality at early post-implantation stages, resulting in the observed variations in sex ratios at birth in ART cycles [30]. 272

The similar effect was observed in previous study that diverse blastocyst parameter grades were related to the sex ratio of singleton live births in SBT cycles [17]. The proportion of male childbirth from high-quality blastocysts transfer was

significantly increased than those from poor-quality blastocysts transfer, which had the trend of elevation with the advance of blastocysts quality. In the present study, we also found that blastocysts with better morphological scores were much more likely to produce male infants, and the trophectoderm grade and ICM grade were both identified as important factors affecting the newborn sex ratio after a single-blastocyst transfer cycle.

282 The trophectoderm grade A blastocysts had statistically significant association with the sex ratio among newborns conceived via blastocyst transfer, which was 283 284 supported by M.B. Borgstrøm's investigation [31]. That is to say, blastocyst morphology was closely related to the sex ratio of livebirth when blastocysts with 285 higher grade trophectoderm were transferred. The contribution of trophectoderm 286 287 during the implantation stage could be attributed to its impact as a promoter of incubation and endometrial invasion [32]. The study of Ebner found that the 288 morphology of trophectoderm was associated with the sex ratio of offspring, with the 289 rates of implantation, pregnancy, and miscarriage. What matters most was that male 290 blastocysts were 2.53 times more likely than female ones with grade A TE [33]. 291 Previous study have shown that human female embryos consume much higher levels 292 of glucose than male embryos [34], which might reflect the need for more energy in 293 294 humans to inactivate the second X-chromosome [35] thereby reducing the mitotic 295 progression. Interestingly, we demonstrated that after controlling for the trophoblast grade, ICM grade B was positively correlated with the higher sex ratio among 296 singleton births but not ICM grade A. Thus, disparities in the morphological scores 297 298 between male and female embryos were likely to reflect some difference in development. Previous studies had different conclusions about the relationship 299 between ICM grade and sex ratio. Lou et al. found that the ICM quality had no 300

301 obvious effect on the newborn sex ratio after single-blastocyst FET cycles [17]. Kate et al. demonstrated that newly expanded normally fertilized blastocysts had an 302 average of 58.3 ± 8.1 cells on day 5, which increased to 84.4 ± 5.7 and 125.5 ± 19 on 303 days 6 and 7, respectively. The number of TE cells was basically the same on days 5 304 and 6 (37.9 ± 6.0 and 40.3 ± 5.0 , respectively), and doubled on day 7 (80.6 ± 15.2). In 305 contrast, the number of ICM cells doubled between days 5 and 6 (20.4 ± 4.0 and 41.9306 307 \pm 5.0, respectively) and remained almost unchanged on day 7 (45.6 \pm 10.2) [36]. This was in line with our results, i.e., male embryos grew more rapidly than female ones, 308 309 contributing to the higher proportion of male blastocysts seen on day 5. However, there were fewer ICM cells than TE cells on day 5, leading to a lower ICM grade. 310 Inversely, the proportion of male embryos declined on day 6 compared to that on day 311 312 5; however, the ICM cells increased, which may account for a higher ICM grade. In 313 China, there were studies concerning the ICM grade in relation to the newborn sex ratio, clinical pregnancy rate, live birth rate, and pregnancy loss rate. However, not all 314 315 results were consistent. In our reproductive center, embryo transfer strategy required that the two best-quality cleavage-stage embryos be chosen for fresh cycle transfer or 316 for cryopreservation on day 3, with the remaining embryos being cultured to the 317 blastocyst stage for possible cryopreservation. 318

Therefore, we had very few good embryos (grade AA) and a large percentage of "fair" BB and "poor" BC embryos, and the strategy required an ICM grade of grade B or higher for optimal cryosurvival. As we all know, ICM developed into a fetus and TE develops into the placenta eventually,We does not transfer the blastocyst with grade C ICM but TE with grade C can be transferred routinely when the ICM is grade A or B in our center. When an ICM grade was C, the embryos were abandoned. Therefore, we only evaluated the outcomes associated with embryos having an ICM

grade of A or B. In this situation, data concerning the ICM had the risk of information bias, which is one of the limitations of the study. However, blastocyst transfer is more advantageous than cleavage-stage embryo transfer in terms of successful implantation rate and pregnancy rates. We prefer high-quality blastocyst transfer to obtain a better clinical pregnancy rate. Although we determined that the transfer of high-quality blastocysts may skew the sex ratio in favor of male embryos, it is difficult to change perceptions regarding popular blastocyst transfer strategies.

In clinical practice, age was a very important factor in considering infertility, 333 334 which was strongly associated with gametes, embryo, and clinical outcomes. This study was stratified by age of infertile patients, and the results revealed that the sex 335 ratio was significantly affected by blastocyst quality in young patients (≤35 years), 336 337 which is in agreement with the results of previous studies [1, 3]. Dean et al. referred 338 to female age as being one of the significant factors affecting the probability of having a male birth. They found that a lower age was associated with a higher likelihood of 339 male birth [15]. Bahadur et al. studied a population in the United Kingdom and drew a 340 similar conclusion [37]. This may because of the higher oxidative stress level in 341 advanced maternal age, but male infants had poor survivability when subjected to 342 high oxidative stress injury compared to female infants. Therefore, male infants may 343 be miscarried in early pregnancy among women of older age. Furthermore, our results 344 345 found that a higher sex ratio was associated with a lower age in men. However, only few studies have reported the effect of paternal age on sex ratio. Jeremy M found that 346 the sex ratio significantly decreased with increasing paternal age and remained so 347 348 even after adjusting for maternal age in offspring with autism spectrum disorders, suggesting that the reduction in the sex ratio with increasing paternal age consistent 349 with de novo genetic or genomic anomalies, which arise more frequently as older men 350

351 conceive [37].

This study has some limitations. As the study was only conducted in a single center, the results may not be generalizable. In addition, it is uncertain that the data accurately reflect the real sex ratio at birth finally under the influence of Chinese culture, which is one limitations of this study. The retrospective nature of this study is another limitation. In future, prospective multicenter studies should perform stratified analysis according to maternal or paternal age ≤ 35 years in order to validate the findings of our study.

359

360 Conclusions

In summary, we demonstrated that the morphological criteria associated with 361 higher-quality blastocysts resulted in a higher sex ratio after SBT. In our study, not 362 only the trophectoderm grade but also the ICM grade was significantly associated 363 with the sex ratio among offspring. The sex ratio increased as the trophectoderm 364 grade increased, whereas a higher ICM grade (grade A) resulted in a lower sex ratio 365 than ICM of grade B, especially in infertile couples in which either partner was below 366 35 years of age. Our findings may shed light on potential factors underlying the 367 skewed sex ratio among offspring from blastocyst transfer, which provided reference 368 for embryo transfer strategy. In future studies, we would further explore the factors 369 370 affecting the sex ratio with the aim of maintaining a balanced sex ratio among 371 offspring in IVF/ICSI, without affecting the pregnancy rate.

372

373 List of abbreviations

374 in vitro fertilization (IVF)

375 intracytoplasmic sperm injection-embryo transfer (ICSI-ET)

- 376 frozen-thawed embryo transfer (FET)
- 377 primary sex ratio (PSR)
- 378 secondary sex ratio (SSR)
- 379 single blastocyst transfer (SBT)
- 380 inner cell mass (ICM)
- 381 trophectoderm (TE)
- 382 assisted reproductive technologies (ART)
- 383 preimplantation genetic testing (PGT)
- 384 body mass index (BMI)
- 385 dimethyl sulfoxide (DMSO)
- 386 ovarian hyperstimulation syndrome (OHSS)
- 387 human chorionic gonadotropin (HCG)
- 388 hormone replacement therapy (HRT)
- 389 Statistical Package for Social Sciences (SPSS)
- 390
- 391 Declarations

392 Ethics approval and consent to participate

The study was carried out after the protocol was approved by the Institutional Review Board of The Third Affiliated Hospital of Zhengzhou University. I confirm that all methods were performed in accordance with the relevant guidelines. The requirement for informed consent was waived by the Institutional Review Board of The Third Affiliated Hospital of Zhengzhou University because of the retrospective nature of the study.

399 Consent for publication

400 Not applicable.

401 Availability of data and materials

402 All data generated or analysed during this study are included in this published article

403 **Competing interests**

- 404 The authors declare that they have no competing interests
- 405 Funding
- 406 Not applicable.

407 Authors' contributions

- 408 Study design: JJH, YCG, and RJY. Data acquisition and analysis: JJH and RJY.
- 409 Drafting of the manuscript and interpretation: YCG, HL. Revision of the manuscript:
- 410 LJS ,XLW and RJY. The authors read and approved the final manuscript.

411 Acknowledgments

- 412 The authors would like to thank all of the participants and the staff of the
- 413 Reproductive Center, the Third Affiliated Hospital of Zhengzhou University, for their
- 414 selfless assistance.
- 415

416

417 **References**

Supramaniam PR, Mittal M, Ohuma EO, Lim LN, McVeigh E, Granne I, et al. Secondary sex
 ratio in assisted reproduction: an analysis of 1 376 454 treatment cycles performed in the UK. Human
 reproduction open. 2019;2019(4):hoz020. Epub 2019/10/11. doi: 10.1093/hropen/hoz020. PubMed
 PMID: 31598568.

Lin PY, Huang FJ, Kung FT, Wang LJ, Chang SY, Lan KC. Comparison of the offspring sex ratio
between cleavage stage embryo transfer and blastocyst transfer. Taiwanese journal of obstetrics &
gynecology. 2010;49(1):35-9. Epub 2010/05/15. doi: 10.1016/s1028-4559(10)60006-x. PubMed PMID:
20466290.

426 3. Hattori H, Kitamura A, Takahashi F, Kobayashi N, Sato A, Miyauchi N, et al. The risk of

secondary sex ratio imbalance and increased monozygotic twinning after blastocyst transfer: data from
the Japan Environment and Children's Study. Reproductive biology and endocrinology : RB&E.

- 429 2019;17(1):27. Epub 2019/02/24. doi: 10.1186/s12958-019-0471-1. PubMed PMID: 30795788.
- 430 4. Gardner DK, WB S. In vitro culture of human blastocysts. In: Jansen, R. and Mortimer, D., Eds.,
- 431 Towards Reproductive Certainty: Infertility and Genetics Beyond 1999: The Plenary Proceedings of

432 the 11th World Congress on In Vitro Fertilization and Human Reproductive Genetics, Parthenon Press, Pearl River. 1999:378-88. 433 434 5. Jacobsen R, Møller H, Mouritsen A. Natural variation in the human sex ratio. Human 435 reproduction (Oxford, England). 1999;14(12):3120-5. Epub 1999/12/22. doi: 436 10.1093/humrep/14.12.3120. PubMed PMID: 10601107. 437 Hesketh T, Xing ZW. Abnormal sex ratios in human populations: causes and consequences. 6. 438 Proceedings of the National Academy of Sciences of the United States of America. 439 2006;103(36):13271-5. Epub 2006/08/30. doi: 10.1073/pnas.0602203103. PubMed PMID: 16938885. 440 Wu Z, Viisainen K, Hemminki E. Determinants of high sex ratio among newborns: a cohort study 7. 441 from rural Anhui province, China. Reproductive health matters. 2006;14(27):172-80. Epub 2006/05/23. 442 doi: 10.1016/s0968-8080(06)27222-7. PubMed PMID: 16713892. 443 Fan SL, Xiao CN, Zhang YK, Li YL, Wang XL, Wang L. How does the two-child policy affect 8. 444 the sex ratio at birth in China? A cross-sectional study. BMC Public Health. 2020;20(1):789. doi: 445 10.1186/s12889-020-08799-v. PubMed PMID: 32460822. 446 9. De Croo I, Colman R, De Sutter P, Tilleman K. Blastocyst transfer for all? Higher cumulative live 447 birth chance in a blastocyst-stage transfer policy compared to a cleavage-stage transfer policy. Facts, 448 views & vision in ObGyn. 2019;11(2):169-76. Epub 2019/12/12. PubMed PMID: 31824637. 449 10. Chen H, Lv JQ, Wu XM, Xiao Y, Xi HT, Zhu CF, et al. Blastocyst-stage versus cleavage-stage 450 embryo transfer in the first frozen cycles of OHSS-risk patients who deferred from fresh embryo 451 transfer. Gynecological endocrinology : the official journal of the International Society of 452 Gynecological Endocrinology. 2015;31(9):698-701. Epub 2015/07/21. doi: 453 10.3109/09513590.2015.1062858. PubMed PMID: 26190533. 454 11. Zhu L, Xi Q, Zhang H, Li Y, Ai J, Jin L. Blastocyst culture and cryopreservation to optimize 455 clinical outcomes of warming cycles. Reproductive biomedicine online. 2013;27(2):154-60. Epub 456 2013/06/19. doi: 10.1016/j.rbmo.2013.04.006. PubMed PMID: 23769665. 457 12. Hur YS, Ryu EK, Song SH, Yoon SH, Lim KS, Lee WD, et al. A retrospective study of single 458 frozen-thawed blastocyst transfer. Clinical and experimental reproductive medicine. 2016;43(2):106-11. 459 Epub 2016/07/01. doi: 10.5653/cerm.2016.43.2.106. PubMed PMID: 27358829. 460 13. Maheshwari A, Hamilton M, Bhattacharya S. Should we be promoting embryo transfer at 461 blastocyst stage? Reproductive biomedicine online. 2016;32(2):142-6. Epub 2015/12/18. doi: 462 10.1016/j.rbmo.2015.09.016. PubMed PMID: 26673100. 463 14. Monteleone PA, Petersen PG, Peregrino PF, Miorin J, Gomes AP, Fujii MG, et al. Should single 464 embryo transfer be used in patients with any kind of infertility factor? Preliminary outcomes. JBRA 465 assisted reproduction. 2019;23(3):200-4. Epub 2019/03/27. doi: 10.5935/1518-0557.20190006. 466 PubMed PMID: 30912516. 467 15. Dean JH, Chapman MG, Sullivan EA. The effect on human sex ratio at birth by assisted reproductive technology (ART) procedures -- an assessment of babies born following single embryo 468 469 transfers, Australia and New Zealand, 2002-2006. BJOG : an international journal of obstetrics and 470 gynaecology. 2010;117(13):1628-34. Epub 2010/09/30. doi: 10.1111/j.1471-0528.2010.02731.x. 471 PubMed PMID: 20875033. 472 16. Sharara FI, Abdo G. Incidence of monozygotic twins in blastocyst and cleavage stage assisted 473 reproductive technology cycles. Fertility and sterility. 2010;93(2):642-5. Epub 2009/02/17. doi: 474 10.1016/j.fertnstert.2008.12.130. PubMed PMID: 19217097.

medRxiv preprint doi: https://doi.org/10.1101/2022.11.28.22282845; this version posted November 30, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in

perpetuity. It is made available under a CC-BY 4.0 International license .

475 17. Lou H, Li N, Zhang X, Sun L, Wang X, Hao D, et al. Does the sex ratio of singleton births after 476 frozen single blastocyst transfer differ in relation to blastocyst development? Reproductive biology and 477 endocrinology : RB&E. 2020;18(1):72. doi: 10.1186/s12958-020-00623-x. PubMed PMID: 32669110. 478 18. Chen X, Zhang J, Wu X, Cao S, Zhou L, Wang Y, et al. Trophectoderm morphology predicts 479 outcomes of pregnancy in vitrified-warmed single-blastocyst transfer cycle in a Chinese population. 480 Journal of assisted reproduction and genetics. 2014;31(11):1475-81. Epub 2014/08/16. doi: 481 10.1007/s10815-014-0317-x. PubMed PMID: 25123128. 482 19. Sun LJ, Zhao B, Hu JJ. Application of three kinds of ovulation induction in vitro 483 fertilization/intracytoplasmic sperm injection patients with poor ovarian response. Chin J Reprod 484 Contracep. 2018;38(03):186-91. 485 20. Wang Ll, Sun LJ, Feng YY. A comparative study of ovulation promotion effect between three 486 kinds of down regulation long protocol in infertile IVF/ICSI patients with normal ovarian function. 487 Prog Obstet Gynecol. 2019;28(08):597-601. 488 21. Papanikolaou EG, D'Haeseleer E, Verheyen G, Van de Velde H, Camus M, Van Steirteghem A, et 489 al. Live birth rate is significantly higher after blastocyst transfer than after cleavage-stage embryo 490 transfer when at least four embryos are available on day 3 of embryo culture. A randomized 491 prospective study. Human reproduction (Oxford, England). 2005;20(11):3198-203. Epub 2005/08/02. 492 doi: 10.1093/humrep/dei217. PubMed PMID: 16055454. 493 22. Schoolcraft WB, Gardner DK, Lane M, Schlenker T, Hamilton F, Meldrum DR. Blastocyst 494 culture and transfer: analysis of results and parameters affecting outcome in two in vitro fertilization 495 programs. Fertility and sterility. 1999;72(4):604-9. Epub 1999/10/16. doi: 496 10.1016/s0015-0282(99)00311-8. PubMed PMID: 10521095. 497 23. Herbemont C, Chekroune S, Bonan S, Cedrin-Durnerin I, Vivot A, Sonigo C, et al. Impact of 498 post-warming culture duration on clinical outcomes of vitrified good-quality blastocyst transfers: a 499 prospective randomized study. Fertility and sterility. 2018;110(7):1290-7. Epub 2018/12/07. doi: 500 10.1016/j.fertnstert.2018.07.1153. PubMed PMID: 30503128. 501 24. Bu Z, Chen ZJ, Huang G, Zhang H, Wu Q, Ma Y, et al. Live birth sex ratio after in vitro 502 fertilization and embryo transfer in China--an analysis of 121,247 babies from 18 centers. PloS one. 503 2014;9(11):e113522. Epub 2014/11/21. doi: 10.1371/journal.pone.0113522. PubMed PMID: 504 25412419. 505 25. Fernando D, Halliday JL, Breheny S, Healy DL. Outcomes of singleton births after blastocyst 506 versus nonblastocyst transfer in assisted reproductive technology. Fertility and sterility. 507 2012;97(3):579-84. Epub 2012/01/28. doi: 10.1016/j.fertnstert.2011.12.032. PubMed PMID: 508 22281036. 509 26. Lin PY, Huang FJ, Kung FT, Wang LJ, Chang SY, Lan KC. Comparison of the offspring sex ratio 510 between fresh and vitrification-thawed blastocyst transfer. Fertility and sterility. 2009;92(5):1764-6. 511 Epub 2009/06/16. doi: 10.1016/j.fertnstert.2009.05.011. PubMed PMID: 19523616. 512 27. Wikland M, Hardarson T, Hillensjö T, Westin C, Westlander G, Wood M, et al. Obstetric 513 outcomes after transfer of vitrified blastocysts. Human reproduction (Oxford, England). 2010;25(7):1699-707. Epub 2010/05/18. doi: 10.1093/humrep/deg117. PubMed PMID: 20472913. 514 515 28. Wang M, Liu X, Zhang H, Li L, Liu R, Zhang H, et al. Associated factors of secondary sex ratio 516 of offspring in assisted reproductive technology: a cross-sectional study in Jilin Province, China. BMC 517 pregnancy and childbirth. 2020;20(1):666. Epub 2020/11/06. doi: 10.1186/s12884-020-03373-1. 518 PubMed PMID: 33148206.

medRxiv preprint doi: https://doi.org/10.1101/2022.11.28.22282845; this version posted November 30, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in

perpetuity. It is made available under a CC-BY 4.0 International license .

519 29. Luke B, Brown MB, Grainger DA, Baker VL, Ginsburg E, Stern JE. The sex ratio of singleton

520 offspring in assisted-conception pregnancies. Fertility and sterility. 2009;92(5):1579-85. Epub

521 2008/10/28. doi: 10.1016/j.fertnstert.2008.08.107. PubMed PMID: 18950756.

- 522 30. Tarín JJ, García-Pérez MA, Hermenegildo C, Cano A. Changes in sex ratio from fertilization to
- 523 birth in assisted-reproductive-treatment cycles. Reproductive biology and endocrinology : RB&E.
- 524 2014;12:56. Epub 2014/06/25. doi: 10.1186/1477-7827-12-56. PubMed PMID: 24957129.
- 525 31. Borgstrøm MB, Kesmodel US, Klausen TW, Danielsen AK, Thomsen T, Gabrielsen A, et al.
- 526 Developmental stage and morphology of the competent blastocyst are associated with sex of the child
- 527 but not with other obstetric outcomes: a multicenter cohort study. Human reproduction (Oxford,
- 528 England). 2021;37(1):119-28. Epub 2022/01/06. doi: 10.1093/humrep/deab242. PubMed PMID:
- 529 34986219.
- 530 32. Ahlström A, Westin C, Reismer E, Wikland M, Hardarson T. Trophectoderm morphology: an
- 531 important parameter for predicting live birth after single blastocyst transfer. Human reproduction
- 532 (Oxford, England). 2011;26(12):3289-96. Epub 2011/10/06. doi: 10.1093/humrep/der325. PubMed 533 PMID: 21972253.
- 534 33. Ebner T, Tritscher K, Mayer RB, Oppelt P, Duba HC, Maurer M, et al. Quantitative and
- 535 qualitative trophectoderm grading allows for prediction of live birth and gender. Journal of assisted
- 536 reproduction and genetics. 2016;33(1):49-57. Epub 2015/11/18. doi: 10.1007/s10815-015-0609-9. 537 PubMed PMID: 26572782.
- 538 34. Gardner DK, Wale PL. Analysis of metabolism to select viable human embryos for transfer.
- 539 Fertility and sterility. 2013;99(4):1062-72. Epub 2013/01/15. doi: 10.1016/j.fertnstert.2012.12.004. PubMed PMID: 23312219. 540
- 541 35. Epstein CJ, Smith S, Travis B, Tucker G. Both X chromosomes function before visible
- 542 X-chromosome inactivation in female mouse embryos. Nature. 1978;274(5670):500-3. Epub
- 543 1978/08/03. doi: 10.1038/274500a0. PubMed PMID: 672979.
- 544 36. Hardy K, Handyside AH, Winston RM. The human blastocyst: cell number, death and allocation
- 545 during late preimplantation development in vitro. Development (Cambridge, England).
- 546 1989;107(3):597-604. Epub 1989/11/01. PubMed PMID: 2612378.
- 547 37. Anello A, Reichenberg A, Luo X, Schmeidler J, Hollander E, Smith CJ, et al. Brief report:
- parental age and the sex ratio in autism. Journal of autism and developmental disorders. 548
- 549 2009;39(10):1487-92. Epub 2009/05/20. doi: 10.1007/s10803-009-0755-y. PubMed PMID: 19452267.
- 550

Figure legends 551

- Figure. 1 Data selection process for the analysis of perinatal outcomes in singleton 552
- live births resulting from single blastocyst transfer (SBT) cycles 553
- Note: Flowchart showing exclusions made to singleton cycles for analysis data to 554
- identify component subsets that were merged to construct the final study dataset. 555
- PGT=preimplantation genetic testing 556

557 **Figure. 2** Sex ratio among singleton births following single blastocyst transfer (SBT)

558 cycles with different blastocyst qualities.

- 559 Note: The sex ratio at birth is calculated as the proportion of liveborn males per 100
- females. CI = confidence interval; P value* was the P value after adjustment (P<0.001).
- 562

563 **Table 1.** Characteristics of 3108 cycles, including sex ratio, among singleton births

from January 2017 to December 2019.

Live birth sex	Female	Male	P value
N	1331	1777	
Maternal age, years (mean ± SD)	30.29 ± 4.32	30.12 ± 4.25	0.288
Paternal age, years (mean \pm SD)	31.30 ± 5.05	31.19 ± 5.02	0.578
Mean duration of infertility (years SD)	± 3.11± 2.48	3.17 ± 2.50	0.511
Mean maternal BMI (kg/m ²)	23.80 ± 3.27	23.58 ± 3.21	0.061
Type of infertility (%)			0.605
Primary	604 (45.38%)	823 (46.31%)	
Secondary	727 (54.62%)	954 (53.69%)	
Factor of infertility (%)			0.881
Female factor	687 (51.62%)	929 (52.28%)	
Male factor	324 (24.34%)	436 (24.54%)	
Mixed factor	240 (18.03%)	301 (16.94%)	
Unexplained infertility	80 (6.01%)	111 (6.25%)	
Type of blastocyst transfer (%)			0.073
ET	428 (32.16%)	626 (35.23%)	

	FET	903 (67.84%)	1151 (64.77%)	
	Day of blastocyst transfer (%)			0.655
	D5	1039 (78.06%)	1399 (78.73%)	
	D6	292 (21.94%)	378 (21.27%)	
,	Type of fertilization (%)			0.832
	ICSI	1212 (91.06%)	1622 (91.28%)	
	IVF	119 (8.94%)	155 (8.72%)	
	Blastocyst quality of transfer (%)			<0.001
	Poor	503 (37.79%)	535 (30.11%)	
	Good	828 (62.21%)	1242 (69.89%)	
	ICM			0.007
	А	414 (31.10%)	474 (26.67%)	
	В	917 (68.90%)	1303 (73.33%)	
,	TE			<0.001
	А	261 (19.61%)	418 (23.52%)	
	В	567 (42.60%)	824 (46.37%)	
	С	503 (37.79%)	535 (30.11%)	

565 Note: The sex ratio at birth is calculated as the proportion of males among all566 livebirths.

567 CI, confidence interval; BMI, body mass index; D, day; IVF, in vitro fertilization;

ICSI, intracytoplasmic sperm injection; ET, fresh embryo transfer; FET, frozenembryo transfer

570

Clinical features	Classification	Univariable a	analysis			
		Wald	OR	95%CI	Р	
Maternal age		1.129	0.991	0.975-1.008	0.288	
Paternal age		0.335	0.996	0.982-1.010	0.563	
BMI		3.497	0.979	0.958-1.001	0.061	
Duration of infertility		0.432	1.010	0.981-1.039	0.511	
Type of infertility						
	Primary		Reference			
	Secondary	0.268	0.963	0.835-1.111	0.605	
Factor of infertility						
	Female factor					
	Male factor	0.003	0.995	0.836-1.185	0.956	
	Mixed factor	0.566	0.927	0.762-1.129	0.452	
	Unexplained infertility	0.028	1.026	0.757-1.390	0.868	

Table 2. Univariable analysis for the potential factors influencing sex ratio among singleton live births

Type of blastocyst transfer

	ET		Reference		
	FET	3.202	0.871	0.750-1.013	0.074
Type of fertilization					
	ICSI		Reference		
	IVF	0.045	0.973	0.758-1.250	0.832
Day of blastocyst transfer					
	D5		Reference		
	D6	0.200	0.961	0.809–1.142	0.655
ICM					
	В		Reference		
	А	7.308	0.806	0.689–0.942	0.007
TE					
	С		Reference		
	В	14.255	1.366	1.162-1.607	0.001

		Α	16.617	1.506	1.237–1.833	0.001
]	Blastocyst quality of transfer					
		Poor		Reference		
		Good	20.134	1.410	1.214–1.639	0.001

572 Note: The sex ratio at birth is calculated as the proportion of males among all live births.

573 CI = confidence interval; BMI = body mass index; D = day; IVF = in vitro fertilization; ICSI = intracytoplasmic sperm injection; ET = fresh

574 embryo transfer; FET = frozen embryo transfer; ICM = Inner cell mass; TE = Trophectoderm

Table 3. Multivariate logistic regression model of sex ratio (male) among singleton births following single blastocyst transfer cycles in relation

Blastocyst	Mutivariable Logistic Regression analysis (not adjusted) Mutivariable L		Mutivariable Log	istic Regression analysis	(adjusted)	
	OR	95% CI	Р	aOR	95% CI	Р
ICM						
В	Reference			Reference		
А	0.638	0.534-0.761	<0.001	0.632	0.529–0.757	<0.001
TE						
С	Reference					
В	1.492	1.263-1.62	0.011	1.510	1.269–1.796	<0.001
Α	1.937	1.551-2.420	<0.001	1.964	1.559–2.474	<0.001
Blastocyst quality	7					
Poor	Reference			Reference		
Good	1.506	1.237–1.833	<0.001	1.552	1.259–1.915	<0.001

577 to blastocyst quality and morphology.

578 Note: Sex ratio among singleton births following single blastocyst transfer cycles by different grades of trophectoderm (TE) and inner cell mass

579 (ICM). P< 0.05 was considered statistically significant. CI= confidence interval;

	Stratified	Blastocyst	Quality classification	OR	95% CI	Р
		ICM				
Maternal age	≤35		В	Reference		
			А	0.803	0.677-0.951	0.011
	>35		В	Reference		
			А	0.802	0.533-1.206	0.289
Paternal age	≤35		В	Reference		
			А	0.791	0.663-0.942	0.009
	>35		В	Reference		
			А	0.862	0.609-1.219	0.401
		TE				
Maternal age	≤35		С	Reference		
			В	1.401	1.172–1.676	<0.001
			А	1.589	1.282-1.970	<0.001

Table 4. Effect of stratified maternal and paternal age and quality classification of ICM grade, TE grade, transferred blastocysts on sex ratio.

	>35		С	Reference		
			В	1.208	0.820-1.778	0.339
			А	1.082	0.653-1.794	0.759
Paternal age	≤35		С	Reference		
			В	1.555	1.245-1.942	<0.001
			А	1.333	0.867-2.049	0.19
	>35		С	Reference		
			В	1.389	1.154-1.671	0.001
			А	1.298	0.927-1.818	0.129
		Transferred blastocyst				
Maternal age	≤35		Poor	Reference		
			Good	1.461	1.238-1.725	<0.001
	>35		Poor	Reference		
			Good	1.17	0.818-1.672	0.39
Paternal age	≤35		Poor	Reference		

	Good	1.442	1.214-1.712	<0.001
>35	Poor	Reference		
	Good	1.309	0.959–1.786	0.09

582 Note: CI = confidence interval; OR = odd ratios; ICM = Inner cell mass; TE = Trophectoderm



Figure



Figure