

1 **Depression and its associated factors among COVID-19 survivors in a middle income country**

2 *Foong Ming Moy¹, Eugene Ri Jian Lim², Noran Naqiah Hairi¹ and Awang Bulgiba¹

3 ¹Centre for Epidemiology and Evidence-based Practice, Department of Social & Preventive Medicine,
4 Faculty of Medicine, University of Malaya, 50603 Kuala Lumpur, Malaysia

5 ²School of Medicine, International Medical University, 57000 Kuala Lumpur, Malaysia

6

7 *Corresponding author (Email: moyfm@ummc.edu.my)

8

9 **Abstract**

10 **Introduction:** COVID-19 survivors who have mental health issues are more likely to have a lower
11 quality of life, reduced work productivity, social troubles, and other health issues. However,
12 information on the mental health of COVID-19 survivors is scarce. Therefore, we aimed to determine
13 the COVID-19 survivors' mental health status in the form of depression and its associated factors.

14 **Methods:** This was a cross-sectional study conducted in Malaysia, from July to September 2021,
15 during a nationwide lockdown. Data was collected using an online questionnaire shared on social and
16 news media. Socio-demographic variables, comorbidities, self-perception of health, information on
17 the person's acute condition during COVID-19 infection, symptoms and duration of symptoms post-
18 COVID, and state of depression were gathered. The Patient Health Questionnaire 9 was used to assess
19 depression. Factors associated with mild to severe depression were analysed using both univariable
20 and multivariable logistic regression analyses.

21 **Results:** A total of 732 COVID-19 survivors responded to the survey. The respondents were mainly
22 females and of younger age (in their 20s and 30s). Two-thirds perceived themselves to be in good
23 health. One in five reported to have experienced Long COVID. Slightly less than half (47.3%) of the
24 respondents had mild to severe depression (total PHQ-9 score of 5 -27). In the multivariable analysis,
25 being female (aOR: 1.68; 95% CI: 1.08,2.62), of younger age (20s – aOR: 3.26; 95% CI: 1.47, 7.25; 30s
26 – aOR: 2.08; 95% CI: 1.05, 4.15; and 40s – aOR: 2.43; 95% CI: 1.20, 4.90; compared to those in the 50s
27 and above), being overweight/obese (aOR: 1.83; 95% CI: 1.18, 2.83), having Long COVID (aOR: 2.45;
28 95% CI: 1.45, 4.16) and perceiving to have poorer health (aOR: 4.54; 95% CI: 2.89, 7.13) were
29 associated with mild to severe depression.

30 **Conclusion:** Females, younger age groups, being overweight/obese, having Long COVID and perceiving
31 to be in poor health were factors associated with higher odds for mild to severe depression.

32 (308 words)

33

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

34 **Introduction:**

35 SARS-CoV-2 has quickly spread all over the world. Globally, the COVID-19 pandemic has had numerous
36 catastrophic consequences in terms of economic loss, morbidity, and mortality. As of 8 December
37 2022, the COVID-19 pandemic has caused 6.6 million deaths and 652 million confirmed cases
38 worldwide. SARS-CoV-2 affects the respiratory, nervous, renal, and cardiovascular systems[1] . With
39 the increase in COVID-19 cases, mental health issues are also now being recognized as a consequence
40 of COVID-19 infection [2, 3].

41
42 Both acute and long-term mental health problems have been linked to COVID-19 infection [4]. The
43 aetiology of mental health consequences during acute COVID-19 is complex. Direct impacts of viral
44 infection, cerebrovascular illness, physiological impairment, immunological response, medical
45 treatments, social isolation, the psychological impact of severe and possibly deadly COVID-19, worry
46 about infecting others, and stigma may all play a role [5].

47
48 COVID-19 survivors are also more likely to have mental health problems in the post-COVID phase [6,
49 7] compared to the general population. These COVID-19 survivors may experience worry, fear, guilt
50 or helplessness as they are more likely to be affected by the stigma of being labelled as an ex-COVID-
51 19 patient. They face uncertainty about their prognosis and future, which predispose them to higher
52 risk of mental health problems. According to a review by Thye et al [4], acute COVID-19 patients had
53 a significant prevalence of anxiety (38.5%) and depression (35.9%), with post-infection anxiety ranging
54 from 6.5 to 63%. COVID-19 survivors who have mental health issues are more likely to have a worse
55 quality of life, reduced work productivity, social problems, abuse vulnerability, and other health issues
56 [2, 8].

57
58 Depression is a common mental health problem globally. The frequency of depressive symptoms
59 reported more than 12 weeks following SARS-CoV-2 infection range from 11 to 28% [9]. Being female,
60 having lower education levels or with chronic diseases are risk factors for depression among COVID-
61 19 survivors [10]. Life events such as loss of a loved one (i.e., parents, spouse or siblings) may also
62 contribute to depression [11]. Lifestyle factors such as poor sleep, poor nutrition, stress and substance
63 abuse are other risk factors for depression [12].

64
65 In Malaysia, there are more than 5 million COVID-19 survivors in the community [13], but there is little
66 data on the mental health status of these patients recovering from COVID-19. In contrast, there have
67 been a number of reports from other countries [14-16]. Considering the negative effects of depression

68 on health-related quality of life and daily functioning, it is crucial to examine if COVID-19 survivors
69 experience depression so that timely intervention can be provided. Therefore, we aimed to
70 investigate the mental health status in the form of depression among COVID-19 survivors in the
71 community and its associated factors.

72

73 **Methods:**

74 From July to September 2021, a cross-sectional study was undertaken in Malaysia, an upper middle-
75 income country, during the execution of a nationwide lockdown. Data was gathered utilizing an online
76 questionnaire, the REDCap electronic data capture tools [17, 18] hosted at our university (University
77 of Malaya).

78

79 To accommodate the multiethnic community, the questionnaire was written in English and translated
80 into Malay and Chinese. All COVID-19 survivors were invited to participate via the social media, the
81 COVID-19 support group's website, and in the news media in the aforementioned languages. The
82 survey was anonymous, and no personally identifying information such as name, phone number, or
83 email address were collected. However, respondents were given the option of being contactable (by
84 leaving their email addresses) in case the researchers determined that their condition warranted extra
85 attention or follow-up with additional activities or support.

86

87 Ethical clearance was obtained from the University of Malaya Research Ethics Committee (Reference
88 number: UM.TNC2/UMREC_1439). Written informed consent was obtained from all participants. The
89 inclusion criteria were all Malaysian adults (aged 18 years and above) who had previously been
90 infected by COVID-19. In 2020-2021, all COVID-19 cases were confirmed via reverse transcription
91 polymerase chain reaction (RT-PCR) tests.

92

93 Referring to the study by Mei et al [10] on depression among COVID-19 survivors, using risk factors
94 for depression such as females (OR=3.4), those with a low income level (OR=2.4) and those with a
95 comorbid chronic disease (OR=2.8), the required calculated sample size with a power of 80% and 95%
96 confidence interval, ranged from 276 to 606.

97

98 The questionnaire asked about socio-demographic variables, comorbidities, self-perception of health,
99 information on the person's acute condition during COVID-19 infection, symptoms and duration of
100 symptoms post-COVID, and mental health state (depression). The severity of condition during the
101 acute infection was self-perceived, with categories of no symptoms, mild, moderate and severe

102 (requiring oxygen supplementation). Long COVID was defined as a state of poor health with post-
103 COVID symptoms that lasted more than 12 weeks after an acute COVID-19 infection which could not
104 be explained by any other illness[19].

105

106 Depression was measured using the Patient Health Questionnaire 9 (PHQ-9)[20]. The PHQ-9 is made
107 up of nine items that asks about sleep, tiredness, changes in appetite, concentration problems, and
108 suicidal thoughts in the last two weeks. The total score was determined by summing the raw scores
109 of each item for a total ranging from 0 to 27. The total scores were divided into five categories namely
110 normal/minimal with score of 0 – 4, mild 5-9, moderate 10-14, moderately severe 15-19 and severe
111 20 -27. In the current analysis, the total PHQ-9 scores were divided into two categories which were
112 normal (0–4) and abnormal (5–27) scores indicating mild to severe depression, similarly used by other
113 researchers [21-23] .

114

115 Data from RedCap were exported into the SPSS version 23 software for data analysis. Normally
116 distributed data was presented using mean with standard deviation (SD), while skewed data were
117 presented using median with inter-quartile range (IQR). Categorical variables were presented using
118 frequency with percentage. Independent variables were cross-tabulated with the categories of PHQ-
119 9 (normal vs abnormal). Chi-square tests were conducted for each variable, and the p-values obtained
120 were presented. For each independent variable, univariable logistic regression models were used to
121 obtain crude Odds Ratios with 95% confidence intervals. We then performed multivariable logistic
122 regression to determine the adjusted Odds Ratios with 95% confidence intervals. To obtain the final
123 model, we included all independent variables which had p-values of 0.25 or less during univariable
124 analysis and any variables which were deemed to be clinically significant. This is in keeping with the
125 “purposeful selection algorithm” proposed by Hosmer and Lemeshow [24]. Significance level was
126 preset at $p < 0.05$.

127

128 **Results:**

129 A total of 732 COVID-19 survivors responded to the survey. The mean (\pm standard deviation) duration
130 between COVID-19 infection and data collection was 27.3 (\pm 12.5) weeks. The respondents were
131 mainly females, of younger age (in their 20s and 30s) and had at least a bachelor’s degree (Table 1).
132 About 50% of them were overweight or obese and the majority were free of comorbidities (74%).
133 About half of them had no symptoms or only had mild symptoms during their acute COVID-19
134 infection, with 25.9% being hospitalized. One-third of respondents perceived themselves to be in poor

135 health at the time of survey. One in five were found to be experiencing Long COVID, based on their
136 self-reported duration of symptoms experienced post-COVID.
137
138

139 Table 1: Socio-demographic characteristics, Body Mass Index (BMI) categories, comorbidities and
140 medical history of respondents

		n (%)
Sex (n=732)	Males	302 (41.3)
	Females	430 (58.7)
Age groups (n=611)	Twenties	156 (25.5)
	Thirties	224 (36.7)
	Forties	33 (5.4)
	Fifties	84 (13.7)
Mean age = 40.2±10.9	Sixties	114 (18.7)
Education levels (n=732)	Secondary and lower	103 (14.1)
	Diploma	142 (19.4)
	Bachelor's degree	341 (46.6)
	Master's degree or higher	146 (19.9)
BMI* categories (n=727)	Underweight	40 (5.5)
	Normal weight	281 (38.7)
	Overweight	219 (30.1)
	Obese	187 (25.7)
Comorbidities** (n=728)	Free of comorbidities	541 (73.9)
	Diabetes mellitus	50 (6.9)
	Hypertension	109 (15.0)
	Heart disease	16 (2.2)
	Hypercholesterolemia	123 (16.9)
	Cancers	10 (1.4)
Categories of acute COVID-19 (n= 634)	No symptom	71(11.2)
	Mild	287 (45.3)
	Moderate	202 (31.9)
	Severe (with oxygen)	74 (11.7)
Long COVID (n=598)	Yes	126 (21.1)
	No	472 (78.9)
Perceived in current good health (n=732)	Yes	494 (67.5)
	No	238 (32.5)

141 Footnote:

142 *underweight (<18.5kg/m²), normal weight (18.5-24.9kg/m²), overweight (25.0–29.9kg/m²), obese
143 (>30.0kg/m²)

144 **the numbers did not add up to 100% as some respondents experienced more than one
145 comorbidity

146 A total of 567 respondents (77.5%) answered the section on PHQ-9. There were no significant
147 differences ($p>0.05$) in terms of sex, age groups, BMI groups, education levels, comorbidities, and
148 experience of Long COVID between those who answered the PHQ-9 compared to those who did not.
149 However, those who had mild symptoms during their acute COVID-19 infection were less likely to
150 answer PHQ-9. The median PHQ-9 total score was 4 with an Inter Quartile Range (IQR) of 9. A total
151 of 52.7% of the respondents had normal PHQ-9 scores (0 -4). There were 21.9% of the respondents
152 with mild depression (5 – 9), followed by 13.8% with moderate (10 – 14), 7.2% moderately severe (15
153 – 19) and 4.4% with severe depression (20 -27), which made up a total of 47.3% with abnormal scores
154 for PHQ-9. The subsequent analyses used a binary outcome with normal (0 – 4) and abnormal (5 –
155 27) categories, which the abnormal category indicated mild to severe depression.

156

157 In the bivariable analysis (Table 2), sex, age groups, BMI categories, severity of acute COVID-19, Long
158 COVID, perception of poor health at the time of survey were significantly associated with mild to
159 severe depression ($p<0.05$). On the other hand, comorbidities and education levels were not
160 associated with mild to severe depression. In the multivariable analysis (Table 2), being female, of
161 younger age (20s, 30s and 40s compared to those in the 50s and above), being overweight/obese,
162 having Long COVID and having perception of poorer health at the time of survey were associated with
163 higher odds for mild to severe depression.

164

165

166

167

168

169

170

171

172

173

174

175

176 Table 2: Factors associated with PHQ-9 categories

n=567	PHQ-9				
	Normal	Abnormal	p-value	Crude OR (95% CI)	*Adjusted OR (95% CI)
Sex					
Male	136 (60.4)	89 (39.6)	0.003	Reference	Reference
Female	163 (47.7)	179 (52.3)		1.68 (1.19, 2.36)	1.68 (1.08,2.62)
Age group					
Twenties	44 (46.3)	51 (53.7)	0.037	2.19 (1.19, 4.04)	3.26 (1.47, 7.25)
Thirties	95 (52.8)	85 (47.2)		1.69 (0.98, 2.92)	2.08 (1.05, 4.15)
Forties	56 (46.7)	64 (53.3)		2.16 (1.21, 3.87)	2.43 (1.20, 4.90)
Fifties and above	53 (65.4)	28 (34.6)		Reference	Reference
Education levels					
Secondary and below	38 (45.2)	46 (54.8)	0.329	1.42 (0.89, 2.29)	1.73 (0.95, 3.29)
Diploma	62 (53.9)	53 (46.1)		1.01 (0.66,1.53)	1.33 (0.78, 2.25)
Degree and above	199 (54.1)	169 (45.9)		Reference	Reference
Perceived in current good health					
Yes	251 (66.4)	127 (33.6)	<0.001	Reference	Reference
No	48 (25.4)	141 (74.6)		5.81 (3.93, 8.58)	4.54 (2.89, 7.13)
BMI categories					
Under-Normal weight	142 (57.7)	104 (42.3)	0.026	Reference	Reference
Overweight-Obese	152 (48.3)	163 (51.7)		1.46 (1.05, 2.05)	1.83 (1.18, 2.83)
Any comorbidities					
Yes	76(49.4)	78(50.6)	0.324	1.21 (0.83, 1.75)	1.29 (0.75, 2.20)
No	223 (54.0)	190 (46.0)		Reference	Reference
Severity of acute COVID					
No symptom	40 (72.7)	15 (27.3)	0.001	Reference	Reference
Mild	146 (55.9)	115 (44.1)		2.10 (1.11,3.99)	1.49 (0.73, 3.08)
Moderate	82 (45.6)	98 (54.4)		3.18 (1.64, 6.18)	2.19 (1.02, 4.69)
Severe (with oxygen)	29 (42.6)	39 (57.4)		3.59 (1.67, 7.69)	2.41 (0.93, 6.22)
Long COVID					
No	262 (58.5)	286 (41.5)	<0.001	Reference	Reference
Yes	37 (31.3)	82 (68.9)		3.12 (2.03, 4.81)	2.45 (1.45, 4.16)

177

178 **Discussion:**

179 Less than half of our respondents reported to be experiencing mild to severe depression. Females,
180 younger respondents, being overweight/obese, having Long COVID and perception of poorer health
181 were factors associated with mild to severe depression among these COVID-19 survivors.

182

183 As we used an online questionnaire, respondents were more likely to be of younger age with higher
184 levels of education. People who were older and had less education were therefore under-represented.
185 There were also more females who answered the survey. The majority of respondents were free of
186 comorbidities such as hypertension, diabetes or heart disease, since they were younger. On the other
187 hand, half of them were overweight or obese, which is similar to what has been found in a Malaysian
188 national survey [25].

189

190 In comparison to before they were infected with COVID-19, one-third of the respondents perceived
191 themselves to be in poor health at the time of survey. Furthermore, one in five respondents
192 experienced Long COVID based on their reported post-COVID symptoms and duration. Fatigue, brain
193 fog, depression, anxiety, insomnia, arthralgia, and myalgia were the most often reported symptoms
194 of Long COVID among our respondents, as reported in another publication [26] . A living systematic
195 review on Long COVID [27] reported that weakness, general malaise, fatigue, concentration
196 impairment and breathlessness were the most common symptoms. Feeling anxious and depressed
197 were also among the common symptoms experienced by COVID-19 survivors [27, 28]. The underlying
198 mechanism is likely to be multifactorial and might include the direct effects of viral infection,
199 immunological response, corticosteroid therapy, ICU stay, social isolation, and stigma [5].

200

201 Similar to other studies among COVID-19 patients, females were consistently found to be more prone
202 to depression [4, 29]. Females appear to be protected against severe symptoms and deaths, and they
203 recover more quickly from COVID-19, presumably due to oestrogen protection [30]. On the other hand,
204 they may suffer severe economic and psychological consequences as a result of COVID-19 infection in
205 the short and long term. Females may have more responsibilities towards their families, children, and
206 the elderly in addition to being in employment. Thus, long-term stressors that happened during the
207 COVID-19 pandemic may have a greater impact on them [31, 32].

208

209 Compared to the oldest age group, those in their 20s, 30s and 40s had higher odds for mild to severe
210 depression. The younger age groups may have more concerns about their future, their employment
211 during lockdown (data collection was conducted during the nationwide lockdown). They may also

212 have poorer mental resilience compared to the older age groups [29, 33]. Older respondents maybe
213 more likely to have faced a number of major life events than the younger respondents. Therefore,
214 the older respondents may have more experience to face adversity better [33].

215

216 Respondents who were overweight or obese also had higher odds for mild to severe depression.
217 There is a reciprocal link between depression and obesity. Obesity is found to increase the risk of
218 depression, in addition, depression was also found to be predictive of obesity [34]. Obesity is an
219 inflammatory condition, and it has been proven that gaining weight activates inflammatory pathways.
220 Inflammation, in turn, has been linked to depression.

221

222 The severity of acute COVID-19, Long COVID experience and the perception of poorer health were
223 significantly associated with mild to severe depression. Higher odds for Long COVID have been seen
224 in respondents who had more severe acute COVID-19 [35]. Respondents having Long COVID may
225 perceived their current health to be poor as they were still experiencing post-COVID symptoms. A
226 collinearity diagnostic was carried out to test if these variables were collinear with each other.
227 However, we did not find any collinearity among these variables. The severity of acute COVID-19 and
228 the experience of Long COVID may be in the past, while the perception of poor health was current at
229 the time of survey. These three factors may have some association with abnormal PHQ-9 scores
230 regardless of their timing. Those who had severe acute COVID-19 may still experience some post-
231 COVID symptoms or Long COVID which may cause them to perceive their health to be poorer at the
232 time of survey. They may also be worried and fearful of their slow recovery which may affect their
233 work productivity and health-related quality of life [5].

234

235 There might be a bidirectional association between subjectively poorer health perception and
236 depression. On one hand, a negative or middling view of one's health may be linked to more severe
237 depression. Poor health perception, on the other hand, can lead to health-related worry, which can
238 lead to depression and other mental health difficulties [36, 37].

239

240 Our findings may need to be interpreted cautiously considering the following issues. The study
241 respondents were not screened for pre-existing mental health disorders. Patients with a history of
242 mental disorder would be more likely to have depression or poor mental health post-COVID [38]. The
243 lockdown implemented during the pandemic may also affected the mental health status of the general
244 population. According to an online survey from Yee et al [39], mild-to-severe depression was found
245 in 28% of the Malaysian community.

246

247 We did not investigate the death or serious illness due to COVID-19 of a loved one or loss of income,
248 which may also have a great influence on mental health [11, 40]. Since our study sample consisted of
249 respondents recruited through online platforms, there may be selection bias where the older and less
250 educated individuals were not adequately represented. Future research should be designed as a
251 prospective cohort study where risk factors of depression among these COVID-19 survivors could be
252 established before depression.

253

254 On the other hand, our study may be one of the few that investigated the state of mild to severe
255 depression of COVID-19 survivors from the Asian region as well as being the first in Malaysia. Our
256 findings support earlier studies that mental health state, especially in the form of depression among
257 COVID-19 survivors should not be overlooked. Even a small percentage of them having mental health
258 disorders might result in a big number of individuals in our nation and throughout the world. Mental
259 disorders are frequently overlooked throughout both the acute and chronic phases of COVID-19,
260 despite the fact that depressive disorders are linked to a much higher risk of death from other causes
261 [41].

262

263 Our findings should be utilised to raise awareness of COVID-19 survivors' increased risk of mental
264 health issues and to advocate for mental healthcare integration as a major component of post-COVID-
265 19 treatment plans. The programmes should be tailored based on the above-mentioned risk factors.
266 Clinicians treating COVID-19 survivors should also be educated on the vulnerability of these patients
267 for depression. As the pandemic progresses, new virus variants emerge, acute COVID-19 treatment
268 options improve, and vaccination uptake rises, the epidemiology of mental health outcomes in the
269 post-acute phase of COVID-19 is anticipated to change over time [42].

270

271 **Conclusions:**

272 A substantial proportion of COVID-19 survivors were found to have mild to severe depression.
273 Females, younger age groups, overweight/obesity, Long COVID and perceived poor health were
274 factors associated with the odds of mild to severe depression. Strengthening access to mental health
275 services such as early assessment and prompt treatment should be incorporated as a core component
276 in the post-COVID-19 care strategies, targeting at those risk factors.

277

278 **References:**

- 279 1. Yuki K, Fujiogi M, Koutsogiannaki S. COVID-19 pathophysiology: A review. *Clin Immunol.*
280 2020;215:108427. Epub 2020/04/24. doi: 10.1016/j.clim.2020.108427.
- 281 2. Amsalem D, Dixon LB, Neria Y. The Coronavirus Disease 2019 (COVID-19) Outbreak and
282 Mental Health: Current Risks and Recommended Actions. *JAMA Psychiatry.* 2021;78(1):9-10. Epub
283 2020/06/25. doi: 10.1001/jamapsychiatry.2020.1730. PubMed PMID: 32579160.
- 284 3. Xiang Y, Yang Y, Li W, Zhang L, Zhang Q, Cheung T, et al. Timely mental health care for the
285 2019 novel coronavirus outbreak is urgently needed. *Lancet Psychiatry.* 2020;7:228–9.
- 286 4. Thyre AY, Law JW, Tan LT, Pusparajah P, Ser HL, Thurairajasingam S, et al. Psychological
287 Symptoms in COVID-19 Patients: Insights into Pathophysiology and Risk Factors of Long COVID-19.
288 *Biology.* 2022;11(1). Epub 2022/01/22. doi: 10.3390/biology11010061. PubMed PMID: 35053059;
289 PubMed Central PMCID: PMCPCMC8773222.
- 290 5. Rogers JP, Chesney E, Oliver D, Pollak TA, McGuire P, Fusar-Poli P, et al. Psychiatric and
291 neuropsychiatric presentations associated with severe coronavirus infections: a systematic review
292 and meta-analysis with comparison to the COVID-19 pandemic. *Lancet Psychiatry.* 2020;7(7):611-27.
293 Epub 2020/05/22. doi: 10.1016/S2215-0366(20)30203-0. PubMed PMID: 32437679; PubMed Central
294 PMCID: PMCPCMC7234781.
- 295 6. Al-Aly Z, Xie Y, Bowe B. High-dimensional characterization of post-acute sequelae of COVID-
296 19. *Nature.* 2021;594(7862):259-64. Epub 2021/04/23. doi: 10.1038/s41586-021-03553-9. PubMed
297 PMID: 33887749.
- 298 7. Taquet M, Dercon Q, Luciano S, Geddes JR, Husain M, Harrison PJ. Incidence, co-occurrence,
299 and evolution of long-COVID features: A 6-month retrospective cohort study of 273,618 survivors of
300 COVID-19. *PLoS Med.* 2021;18(9):e1003773-e. doi: 10.1371/journal.pmed.1003773. PubMed PMID:
301 34582441.
- 302 8. Xie Y, Xu E, Al-Aly Z. Risks of mental health outcomes in people with covid-19: cohort study.
303 *BMJ (Clinical research ed).* 2022;376:e068993. Epub 2022/02/18. doi: 10.1136/bmj-2021-068993.
304 PubMed PMID: 35172971.
- 305 9. Renaud-Charest O, Lui LMW, Eskander S, Ceban F, Ho R, Di Vincenzo JD, et al. Onset and
306 frequency of depression in post-COVID-19 syndrome: A systematic review. *Journal of Psychiatric*
307 *Research.* 2021;144:129-37. doi: <https://doi.org/10.1016/j.jpsychires.2021.09.054>.
- 308 10. Mei Q, Wang F, Bryant A, Wei L, Yuan X, Li J. Mental health problems among COVID-19
309 survivors in Wuhan, China. *World psychiatry : official journal of the World Psychiatric Association*
310 *(WPA).* 2021;20(1):139-40. Epub 2021/01/13. doi: 10.1002/wps.20829. PubMed PMID: 33432745;
311 PubMed Central PMCID: PMCPCMC7801855.
- 312 11. Mousavi SA, Hooshyari Z, Ahmadi A. The Most Stressful Events during the COVID-19
313 Epidemic. *Iran J Psychiatry.* 2020;15(3):220-7. Epub 2020/11/17. doi: 10.18502/ijps.v15i3.3814.
314 PubMed PMID: 33193770; PubMed Central PMCID: PMCPCMC7603585.
- 315 12. Hoveling LA, Liefbroer AC, Schwaren LJS, Bültmann U, Smidt N. Socioeconomic differences in
316 major depressive disorder onset among adults are partially explained by lifestyle factors: A
317 longitudinal analysis of the Lifelines Cohort Study. *J Affect Disord.* 2022. Epub 2022/07/20. doi:
318 10.1016/j.jad.2022.06.018. PubMed PMID: 35850289.

- 319 13. KKMNOW, COVID-19 -The latest data on the pandemic in Malaysia (as of 08.12.2022)
320 [Internet]. 2022 [cited 09.12.2022]. Available from: <https://data.moh.gov.my/covid>.
- 321 14. Houben-Wilke S, Goërtz YM, Delbressine JM, Vaes AW, Meys R, Machado FV, et al. The
322 Impact of Long COVID-19 on Mental Health: Observational 6-Month Follow-Up Study. *JMIR mental*
323 *health*. 2022;9(2):e33704. Epub 2022/02/25. doi: 10.2196/33704. PubMed PMID: 35200155;
324 PubMed Central PMCID: PMCPCMC8914795.
- 325 15. Villalpando JMG, Forcelledo HA, Castillo JLB, Sastre AJ, Rojop IEJ, Hernandez VO, et al.
326 COVID-19, Long COVID Syndrome, and Mental Health Sequelae in a Mexican Population. *Int J*
327 *Environ Res Public Health*. 2022;19(12). Epub 2022/06/25. doi: 10.3390/ijerph19126970. PubMed
328 PMID: 35742219; PubMed Central PMCID: PMCPCMC9222368.
- 329 16. Titze-de-Almeida R, da Cunha TR, Dos Santos Silva LD, Ferreira CS, Silva CP, Ribeiro AP, et al.
330 Persistent, new-onset symptoms and mental health complaints in Long COVID in a Brazilian cohort of
331 non-hospitalized patients. *BMC infectious diseases*. 2022;22(1):133. Epub 2022/02/10. doi:
332 10.1186/s12879-022-07065-3. PubMed PMID: 35135496; PubMed Central PMCID:
333 PMCPCMC8821794.
- 334 17. Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, et al. The REDCap consortium:
335 Building an international community of software platform partners. *J Biomed Inform*.
336 2019;95:103208. Epub 2019/05/13. doi: 10.1016/j.jbi.2019.103208. PubMed PMID: 31078660;
337 PubMed Central PMCID: PMCPCMC7254481.
- 338 18. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data
339 capture (REDCap)--a metadata-driven methodology and workflow process for providing translational
340 research informatics support. *J Biomed Inform*. 2009;42(2):377-81. Epub 2008/10/22. doi:
341 10.1016/j.jbi.2008.08.010. PubMed PMID: 18929686; PubMed Central PMCID: PMCPCMC2700030.
- 342 19. National Institute for Health and Care Excellence. COVID-19 rapid guideline: managing the
343 long-term effects of COVID-19 NICE guideline; <https://www.nice.org.uk/guidance/ng188>. 2020.
- 344 20. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity
345 measure. *Journal of general internal medicine*. 2001;16(9):606-13. Epub 2001/09/15. doi:
346 10.1046/j.1525-1497.2001.016009606.x. PubMed PMID: 11556941; PubMed Central PMCID:
347 PMCPCMC1495268.
- 348 21. Wu C, Hu X, Song J, Yang D, Xu J, Cheng K, et al. Mental health status and related influencing
349 factors of COVID-19 survivors in Wuhan, China. *Clinical and translational medicine*. 2020;10(2):e52.
350 Epub 2020/06/09. doi: 10.1002/ctm2.52. PubMed PMID: 32508037; PubMed Central PMCID:
351 PMCPCMC7300592.
- 352 22. Xiao X, Yang X, Zheng W, Wang B, Fu L, Luo D, et al. Depression, anxiety and post-traumatic
353 growth among COVID-19 survivors six-month after discharge. *European journal of*
354 *psychotraumatology*. 2022;13(1):2055294. Epub 2022/04/12. doi: 10.1080/20008198.2022.2055294.
355 PubMed PMID: 35401948; PubMed Central PMCID: PMCPCMC8986234.
- 356 23. Levis B, Benedetti A, Thombs BD. Accuracy of patient health questionnaire-9 (PHQ-9) for
357 screening to detect major depression: Individual participant data meta-analysis. *BMJ (Clinical*
358 *research ed)*. 2019;365:l1476. doi: 10.1136/bmj.l1476.

- 359 24. Hosmer D, Lemeshow S. Model-building strategies and methods for logistic regression. . In:
360 Shewhart W, Wilks S, editors. Applied Logistic Regression. Hoboken, New Jersey: John Wiley & Sons;
361 2005.
- 362 25. Institute for Public Health. National Health and Morbidity Survey (NHMS) 2019: Non-
363 communicable diseases, healthcare demand, and health literacy—Key Findings. Putrajaya: 2020.
- 364 26. Moy FM, Hairi NN, Lim ERJ, Bulgiba A. Long COVID and its associated factors among COVID
365 survivors in the community from a middle-income country-An online cross-sectional study. PloS one.
366 2022;17(8):e0273364. Epub 2022/08/31. doi: 10.1371/journal.pone.0273364. PubMed PMID:
367 36040960; PubMed Central PMCID: PMCPCMC9426885.
- 368 27. Michelen M, Manoharan L, Elkheir N, Cheng V, Dagens A, Hastie C, et al. Characterising long
369 COVID: a living systematic review. BMJ Glob Health. 2021;6(9):e005427. doi: 10.1136/bmjgh-2021-
370 005427.
- 371 28. van Kessel SAM, Olde Hartman TC, Lucassen P, van Jaarsveld CHM. Post-acute and long-
372 COVID-19 symptoms in patients with mild diseases: a systematic review. Fam Pract. 2022;39(1):159-
373 67. Epub 2021/07/17. doi: 10.1093/fampra/cmab076. PubMed PMID: 34268556; PubMed Central
374 PMCID: PMCPCMC8414057.
- 375 29. Staneva A, Carmignani F, Rohde N. Personality, gender, and age resilience to the mental
376 health effects of COVID-19. Soc Sci Med. 2022;301:114884. Epub 2022/03/29. doi:
377 10.1016/j.socscimed.2022.114884. PubMed PMID: 35344776; PubMed Central PMCID:
378 PMCPCMC8915456.
- 379 30. Dupuis ML, Maselli A, Pagano MT, Pierdominici M, Ortona E. Immune response and
380 autoimmune diseases: a matter of sex. Italian Journal of Gender-Specific Medicine. 2019;5(1):11-20.
- 381 31. Bucciarelli V, Nasi M, Bianco F, Seferovic J, Ivkovic V, Gallina S, et al. Depression pandemic
382 and cardiovascular risk in the COVID-19 era and long COVID syndrome: Gender makes a difference.
383 Trends in cardiovascular medicine. 2022;32(1):12-7. Epub 2021/10/08. doi:
384 10.1016/j.tcm.2021.09.009. PubMed PMID: 34619336; PubMed Central PMCID: PMCPCMC8490128.
- 385 32. Cohen S, Murphy M, Prather A. Ten Surprising Facts About Stressful Life Events and Disease
386 Risk Annu Rev Psychol. 2019;70:577- 97.
- 387 33. Rossi R, Jannini TB, Socci V, Pacitti F, Lorenzo GD. Stressful Life Events and Resilience During
388 the COVID-19 Lockdown Measures in Italy: Association With Mental Health Outcomes and Age.
389 Front Psychiatry. 2021;12:635832. Epub 2021/03/26. doi: 10.3389/fpsy.2021.635832. PubMed
390 PMID: 33762980; PubMed Central PMCID: PMCPCMC7982412.
- 391 34. Luppino FS, de Wit LM, Bouvy PF, Stijnen T, Cuijpers P, Penninx BWJH, et al. Overweight,
392 Obesity, and Depression: A Systematic Review and Meta-analysis of Longitudinal Studies. Archives of
393 General Psychiatry. 2010;67(3):220-9. doi: 10.1001/archgenpsychiatry.2010.2 %J Archives of General
394 Psychiatry.
- 395 35. Han Q, Zheng B, Daines L, Sheikh A. Long-Term Sequelae of COVID-19: A Systematic Review
396 and Meta-Analysis of One-Year Follow-Up Studies on Post-COVID Symptoms. Pathogens. 2022;11(2).
397 Epub 2022/02/27. doi: 10.3390/pathogens11020269. PubMed PMID: 35215212; PubMed Central
398 PMCID: PMCPCMC8875269.

- 399 36. Cappeliez P, Sèvre-Rousseau S, Landreville P, Prévaille M, Scientific Committee of ESAS.
400 Physical health, subjective health, and psychological distress in older adults: Reciprocal relationships
401 concurrently and over time. *Ageing International*. 2004;29(3):247-66. doi: 10.1007/s12126-996-
402 1001-y.
- 403 37. Lee K, Feltner F, Bailey A, Lennie T, Chung M, Smalls B, et al. The relationship between
404 psychological states and health perception in individuals at risk for cardiovascular disease *Psychol*
405 *Res Behav Manag*. 2019;12:317-24. doi: <https://doi.org/10.2147/PRBM.S198280>.
- 406 38. Henssler J, Stock F, van Bohemen J, Walter H, Heinz A, Brandt L. Mental health effects of
407 infection containment strategies: quarantine and isolation—a systematic review and meta-analysis
408 *Eur Arch Psychiatry Clin Neurosci*. 2021:[https:// doi. org/ 10. 1007/ s00406- 020- 1196-x](https://doi.org/10.1007/s00406-020-1196-x).
- 409 39. Yee A, Hodori NM, Tung YZ, Ooi PL, Latif S, Isa HM, et al. Depression level and coping
410 responses toward the movement control order and its impact on quality of life in the Malaysian
411 community during the COVID-19 pandemic: a web-based cross-sectional study. *Ann Gen Psychiatry*.
412 2021;20(1):31. Epub 2021/05/26. doi: 10.1186/s12991-021-00352-4. PubMed PMID: 34030704;
413 PubMed Central PMCID: PMCPCMC8142880.
- 414 40. Ferrando SJ, Klepacz L, Lynch S, Shahar S, Dornbush R, Smiley A, et al. Psychiatric
415 emergencies during the height of the COVID-19 pandemic in the suburban New York City area. *J*
416 *Psychiatr Res*. 2021;136:552-9. Epub 2020/11/08. doi: 10.1016/j.jpsychires.2020.10.029. PubMed
417 PMID: 33158555; PubMed Central PMCID: PMCPCMC7992036.
- 418 41. Cuijpers P, Vogelzangs N, Twisk J, Kleiboer A, Li J, Penninx BW. Comprehensive Meta-Analysis
419 of Excess Mortality in Depression in the General Community Versus Patients With Specific Illnesses.
420 *Am J Psychiatry*. 2014;171(4):453-62. doi: 10.1176/appi.ajp.2013.13030325. PubMed PMID:
421 24434956.
- 422 42. Cai M, Bowe B, Xie Y, Al-Aly Z. Temporal trends of COVID-19 mortality and hospitalisation
423 rates: an observational cohort study from the US Department of Veterans Affairs. *BMJ open*.
424 2021;11(8):e047369. Epub 2021/08/18. doi: 10.1136/bmjopen-2020-047369. PubMed PMID:
425 34400452; PubMed Central PMCID: PMCPCMC8370839.

426

427 **Declaration:**

428 **Ethics approval and consent to participate**

429 Ethical clearance was obtained from the University of Malaya Research Ethics Committee
430 (Reference number: UM.TNC2/UMREC_1439). Written informed consent was obtained from all
431 participants. The study was conducted in compliance with recognized international standards,
432 including the principles of the Declaration of Helsinki. All procedures were performed in
433 accordance with relevant guidelines.

434 **Consent for publication**

435 Not applicable

436 **Availability of data and materials**

437 All data generated or analysed during this study are included in this published article [and its
438 supplementary information files].

439 **Competing interests**

440 The authors declare that they have no competing interests.

441 **Funding**

442 This study is part of the COVID-19 Epidemiological Analysis and Strategies (CEASe) Project with
443 funding from the Ministry of Science, Technology and Innovation (UM.0000245/HGA.GV).

444 **Authors' contributions**

445 FMM, NNH and AB contributed to the conception of the work; FMM worked on the acquisition,
446 analysis, and interpretation of data; FMM, NNH, ERJL and AB contributed to the drafting,
447 finalizing and approval of the submitted version. All authors agreed to be personally
448 accountable for the author's own contributions and to ensure that questions related to the
449 accuracy or integrity of any part of the work.

450 **Acknowledgements**

451 The respondents and the support from the University Malaya are acknowledged.