It is made available under a CC-BY 4.0 International license .

1	- Review -
2	The epidemiology of human <i>Taenia solium</i> infections: a
3	systematic review of the distribution in Eastern and
4	Southern Africa.
5	Short title: Epidemiology of human <i>T. solium</i> infections in Eastern and
6	Southern
7	Gideon Zulu ^{1, 2*} , Dominik Stelzle ³ , Kabemba E. Mwape ² , Tamara M. Welte ^{4,5} , Hilde
8	Strømme ⁶ , Chishimba Mubanga ² , Wilbroad Mutale ⁷ , Annette Abraham ^{3,8} , Alex
9	Hachangu ² , Veronika Schmidt ^{3,8} , Chummy. S. Sikasunge ⁹ , Isaac. K. Phiri ^{2&} and
10	Andrea S. Winkler ^{3,8&}
11 12	¹ Department of Clinical Studies, School of Veterinary Medicine, University of Zambia, Lusaka, Zambia
13	² Ministry of Health, Government of the Republic of Zambia
14	³ Center for Global Health, Department of Neurology, Klinikum rechts der Isar, Technical
15	University Munich, Munich, Germany
16	⁴ Department of Neurology, Center for Global Health, Faculty of Medicine, Technical University
17	of Munich, Munich, Germany.

It is made available under a CC-BY 4.0 International license .

- ⁵ Epilepsy Center, Department of Neurology, University Hospital Erlangen, Germany.
- ⁶ University of Oslo, University Library, Medical Library, Norway
- 20 ⁷ School of Public Health, University of Zambia, Lusaka, Zambia
- ⁸ Centre for Global Health, Institute of Health and Society, University of Oslo, Oslo, Norway
- ⁹ Department of Para-clinical studies, School of Veterinary Medicine, University of Zambia,
- 23 Lusaka, Zambia
- 24 * Corresponding author
- 25 Email: gideonzulu@yahoo.com (GZ)
- 26 [&] These authors contributed equally to this work.

27 Abstract

Background: *Taenia solium* is a tapeworm that causes taeniosis in humans and cysticercosis in humans and pigs. Within Eastern and Southern Africa (ESA), there are many countries in which information on the presence of human taeniosis and cysticercosis is missing. This systematic review aimed to describe the current information available and gaps in the epidemiology of human *T. solium* infections in ESA.

Methods/Principle Findings: Scientific literature published between 1st January 2000 and 20th
June 2022 in international databases [MEDLINE (Ovid), Embase (Ovid), Global Health (Ovid),
Scopus (Elsevier), African Index Medicus (via WHO Global Index Medicus), and Open Grey] was
systematically reviewed for ESA following the PRISMA approach. The study area included 27

It is made available under a CC-BY 4.0 International license .

37 countries that make up the ESA region. Information on either taeniosis, cysticercosis or NCC was 38 available for 16 of 27 countries within the region. Most case reports for cysticercosis and NCC 39 were from South Africa, while Tanzania had the most aggregated cysticercosis reports. Eleven 40 countries reported on NCC with seven countries reporting data on NCC and epilepsy. Unconfirmed 41 human T. solium taeniosis cases were reported in nine countries while two countries (Madagascar 42 and Zambia) reported confirmed T. solium cases. The cysticercosis seroprevalence ranged between 43 0.99 - 40.8% on antigen (Ag) tests and between 1.7 - 45.3% on antibody (Ab) tests, while NCC-44 suggestive lesions on brain CT scans showed a prevalence range between 1.0 - 76%. The human 45 taeniosis prevalence based on microscopy ranged between 0.1 - 14.7%. Based on Copro Ag-ELISA studies conducted in Kenya, Rwanda, Tanzania, and Zambia, the highest prevalence of 46 47 19.7% was reported in Kenya

48 Conclusions: Despite the public health and economic impact of *T. solium* in ESA, there are still 49 large gaps in knowledge about the occurrence of the parasite, and the resulting One Health disease 50 complex, and monitoring of *T. solium* taeniosis and cysticercosis is mostly not in place

51 Author summary

Taenia solium is a tapeworm that causes three diseases, taeniosis in humans and cysticercosis in humans and pigs. Neurocysticercosis, which occurs when the central nervous system is involved has been associated with up to 57% of epilepsy cases in sub–Saharan Africa. Diagnosing neurocysticercosis among people with epilepsy is vital to prevent further morbidity and mortality from the disease as well as to reduce the negative socio-cultural beliefs associated with epilepsy. Within Eastern and Southern Africa, there are many countries in which information on the presence of human taeniosis, cysticercosis and neurocysticercosis is missing. This systematic

It is made available under a CC-BY 4.0 International license .

review aimed to describe the current information available and gaps in the epidemiology of human *T. solium* infections in Eastern and Southern Africa. We found that Information on either taeniosis, cysticercosis or NCC was available only for 16 of 27 countries within the region. We also found that most of the studies on *T. solium* taeniosis, cysticercosis and neurocysticercosis within the region have been done in Kenya, Madagascar, Mozambique, Rwanda, South Africa, Tanzania and Zambia. Understanding the epidemiology of *T. solium* infections is essential for monitoring, prevention and control of the disease complex in a One Health approach.

66 Introduction

67 Rationale

68 *Taenia solium* is a tapeworm that causes taeniosis in humans and cysticercosis in humans and pigs. 69 The life cycle of *T. solium* involves pigs as intermediate hosts (cysticercosis) while humans are 70 definitive hosts (taeniosis). Humans may also act as accidental intermediate hosts when larvae of 71 the parasite settle in muscles, subcutaneous or organ tissues causing human cysticercosis (HC). If they lodge in the central nervous system (CNS), including the brain and the spinal cord, the disease 72 73 is called neurocysticercosis (NCC) [1,2]. An individual may also have cysticerci in the CNS as 74 well as in other parts of the body, which is referred to as (neuro) cysticercosis. NCC may be 75 asymptomatic, but it can also cause various neurological signs/symptoms such as epileptic seizures 76 and epilepsy, chronic progressive headache, focal neurological deficits, signs and symptoms of 77 increased intracranial pressure and in rare cases, death [3–5]. Globally, NCC is estimated to be 78 responsible for 30% of cases of acquired epilepsy in endemic areas [6] and 22% in sub-Saharan

It is made available under a CC-BY 4.0 International license .

Africa [7]. However, this proportion depends on the infection pressure of *T. solium* and thus mayvary considerably.

81 T. solium is common in Asia and Latin America, especially in areas where pigs are reared under 82 free-ranging conditions, where pork is eaten and where hygiene is limited [8,9]. Hence, T. solium 83 is also likely endemic in most African countries [10]. Nevertheless, information on the endemicity 84 of T. solium is limited and there are many countries from which no published information is 85 available for either human or porcine cysticercosis. The purpose of this review was to gather all 86 published information on human T. solium infections (T. solium taeniosis/(neuro)cysticercosis 87 (TSTC)) in eastern and southern Africa (ESA) within the period 1st January 2000 and 20th June 88 2022 and describe the gaps in the epidemiology of TSTC in this area.

89 Methods

The systematic review was conducted following the Preferred Reporting Items for Systematic
Reviews and Meta-Analyses (PRISMA) guidelines [11] (S1 Table). The review was registered
with the International Prospective Register of Systematic Reviews (PROSPERO) (registration
number: CRD 42022343072).

94 Search strategy

All published articles were searched using the electronic databases, MEDLINE (Ovid), Embase
(Ovid), Global Health (Ovid), Scopus (Elsevier), African Index Medicus (via WHO Global Index
Medicus), and Open Grey. The reference lists of included studies were also scanned to identify
further eligible documents. The searches were conducted on 20th June 2022.

For all questions subject headings (where applicable) and text words describing *T. solium* taeniosis/cysticercosis/neurocysticercosis and prevalence, epidemiology, control, elimination, or eradication were searched. The search strategy was intended to obtain all available literature on human taeniosis, cysticercosis and neurocysticercosis in ESA conducted and published in the last 22 years. Grey literature was also searched for any relevant publications. The search terms for all databases can be found in (S1 File).

105 **Procedures**

106 **Data management**

107 Literature search results were uploaded to the bibliographic software Mendeley Desktop and 108 Covidence. Duplicates were removed based on author names and titles. Screening questions based 109 on the inclusion and exclusion criteria were developed and citation abstracts and full-text articles 110 were uploaded.

111 Selection process

The selection process was conducted in two stages. Firstly, titles and abstracts were screened independently by two reviewers (GZ and CM) and then full-text reports were obtained by the same reviewers for a detailed assessment with respect to the inclusion criteria. For records that referred to the same study, only one was selected.

It is made available under a CC-BY 4.0 International license .

116 **Data collection process**

- 117 The data were collected using standardized data collection forms in Microsoft Excel (S2 File).
- 118 Data extracted included demographic information, methodology, and all the relevant reported
- 119 outcomes. GZ carried out the data extraction with verification by CM.

120 Data items

Predefined tables summarizing individual cases included the year of diagnosis, age, gender, the country where cases were detected, diagnostic method, and location of the lesion (for cysticercosis). Tables summarizing aggregated cases or prevalence data included country, level of data collection (e.g. national or regional), timeframe, number of cases (or prevalence proportion), number of people tested, and diagnostic method.

126 Eligibility criteria

127 Inclusion criteria

128 We included all studies describing human TSTC in ESA, which was defined as the area covered 129 by the following countries/territories: Angola, Botswana, Burundi, Comoros, Djibouti, Eritrea, 130 Ethiopia, Kenya, Lesotho, Madagascar, Malawi, Mauritius, Mayotte, Mozambique, Namibia, 131 Reunion, Rwanda, Seychelles, Socotra, Somalia, Somaliland, South Africa, Eswatini (former 132 Kingdom of Swaziland), Tanzania, Uganda, Zambia and Zimbabwe. Included in this review were 133 studies that describe the epidemiology, disease occurrence, burden, prevalence, incidence, 134 prevention and control of TSTC in any of the twenty-seven countries of ESA. All observational 135 studies including case series and case reports and systematic reviews of such studies were included.

It is made available under a CC-BY 4.0 International license .

136 Only studies conducted between 1st January 2000 and 20th June 2022 were considered. No

137 language limits were imposed on the search, although studies in languages other than English were

138 only included if they could be adequately translated by using Google Translate.

139

140 Exclusion criteria

141 We excluded studies that did not concern *T. solium*, did not concern humans, did not report data

142 from within ESA and studies conducted before 1st January 2000.

143 **Risk of bias assessment**

The risk of bias in the studies was assessed using items recommended by the Agency for Health Care Research and Quality (AHRQ) [12]. This assessment covered baseline characteristics, inclusion-exclusion criteria, confounding and modifying variables, performance bias, attrition bias, detection bias and reporting bias. Each of these factors was rated and categorized as low, or high risk of bias. If the information available was insufficient, the risk of bias was considered 'unsure'.

150 Statistical analyses

No statistical pooling of results was conducted and the findings of the review are presented in a narrative synthesis with tables and figures to aid data presentation. Prevalence data if not already provided were calculated using the numerator and denominator of the study sample. Data were analyzed separately for taeniosis, cysticercosis, and neurocysticercosis. For the latter two, data were also presented separately for the general population and people with epilepsy.

It is made available under a CC-BY 4.0 International license .

156 Ethical considerations

157 Ethical recommendation was not required for this review.

158 **Results**

159 Study selection

160 Our search yielded 2114 records of which 1198 were duplicates. Through screening of titles and 161 abstracts, 776 (85%) records were excluded and for a further 8, no full text was found. Full-text 162 assessment with respect to the inclusion criteria was performed on 132 records of which 19 (14%) 163 were excluded (17 reported data described in other included studies, one did not concern T. solium, 164 and one reported results outside the scope of the study). One hundred and thirteen (113) studies 165 were finally included in the review. Of those, 49 were concerned with only (neuro) cysticercosis, 166 57 with taeniosis, and 7 with both cysticercosis and taeniosis. Of the 113 reports reviewed, 25 were 167 case reports, 82 were cross-sectional, 3 cohort and 3 case-control studies. The flow diagram for 168 the search is shown in (Fig 1). The reference for included studies is shown in (S2 Table).

169

170

171

172

It is made available under a CC-BY 4.0 International license .

174

175



176



178 Risk of bias

179 The risk of bias assessment revealed that most studies had a low risk of reporting, detection,

180 attrition and performance bias. However, 7 studies had a high risk of performance bias, 20 had a

It is made available under a CC-BY 4.0 International license .

- 181 high risk of bias due to inadequate inclusion and exclusion criteria, and 45 articles had a high risk
- 182 of bias due to confounding and modifying variables.

183 **Results of individual studies**

184 Human (neuro) cysticercosis

185 Information on human (neuro) cysticercosis was obtained from 62 sources. In total, 25 (40%) 186 records provided individual information on NCC and CC. Twelve (48%) of these records were on 187 NCC only, seven (28%) on CC and six (24%) on both NCC and CC. Thirty-seven (60%) records 188 reported aggregated information on NCC and CC. Ten (27%) of these records were on NCC only, 189 twenty-one (57%) on CC only, and six (16%) on both NCC and CC. Information on human 190 neuro(cysticercosis) was available for 14 of 27 ESA countries. No reports were available for 191 Angola, Botswana, Comoros, Djibouti, Eritrea, Ethiopia, Lesotho, Mayotte, Reunion, Seychelles, 192 Socotra, Somaliland and Somalia.

193 Human cysticercosis case reports

194 Individual human CC cases during the period 2000 to 2022 were reported in 13 hospital case 195 records from six of the ESA countries with, South Africa reporting four cases; Madagascar, 196 Malawi, Rwanda and Tanzania reporting two cases each and Mauritius reporting one case (Table 197 1). Individual NCC cases were reported in 18 hospital records from eight of the ESA countries 198 with South Africa reporting eight cases while Madagascar, Malawi and Zambia reported two cases 199 each. Mozambique, Eswatini, Tanzania and Zimbabwe reported one case each (Table 2). All 200 reports were based on individual patients presenting with various signs/symptoms, and 201 investigations leading to their diagnosis of either CC or NCC. Twelve (48%) of the 25 case reports

202	described only NCC diagnosed through the use of brain computed tomography (CT) scan (7) and
203	magnetic resonance imaging (MRI) (5), while seven (28%) presented with CC affecting other
204	organs i.e. bronchus, breast, abdomen, neck and subcutaneous tissue diagnosed through
205	histopathology and other investigations. Six (24%) case reports were on patients who presented
206	with both NCC and CC affecting other organs (disseminated CC). At a district hospital in Rwanda
207	ten cases of cutaneous CC were reported with the youngest being only 2 months old [13]. The
208	oldest reported case of NCC was from South Africa in an 86-year-old female with a pharyngeal
209	cyst, cutaneous cysts and active brain cysts [14]. The median age in years was 41 (17 -48 year).
210	Twenty-three (92%) of the case reports had information on the affected gender with 54% of the
211	reported cases being female and 46% being male.
212	
213	
214	
215	
216	
217	
218	
219	

It is made available under a CC-BY 4.0 International license .

221

223 Table 1. Individual human cysticercosis case reports as detected through histopathology of sample specimens from patients in

Country	Age (yrs.)	Gender	Sample	Location of pathogen	Comment	Reference
Madagascar	20	Male	Bronchial specimen	Bronchus	A 20-year-old male who presented with hemoptysis following bronchial cysticercosis.	[50]
	15	Female	Breast lamp	Breast	Breast cysticercosis in a 15-year-old girl.	[51]
Malawi	41	Female	Skin nodule	Subcutaneous all over the body*	A 41-year-old patient on ARVs with disseminated cysticercosis and more than 50 cysts in the brain.	[52]
	56	Female	Biopsy of neck mass	Left anterior and posterior triangles of the neck	A 56-year-old woman with a history of gradual but progressive growing left sided neck swelling for the past 25 years.	[53]
Mauritius	22	Female	Excision biopsy	Upper part of right breast	A 22-year-old woman with history of painless mobile swelling in the right side of the breast.	[54]
Rwanda	2 months - 65 yrs.	-	Anatomical department files	Subcutaneous nodules	A case report of 10 cases of cutaneous cysticercosis confirmed by histopathology in the anatomical pathology department at a district hospital (Nyamasheke) in eastern Rwanda.	[13]
	46	Female	Subcutaneous nodule	Subcutaneous nodules on trunk and extremities	A 46-year-old woman who presented with a two-week history of bilateral lower limb weakness, causing difficulty walking.	[55]
South Africa	42	Male	Biopsy of subcutaneous nodule	Extensive calcifications in the muscle and subcutaneous tissue. Active cysts within myocardium [#]	A 42-year-old man with a rare case of myocardial cysticercosis with multiple calcifications in the muscle and subcutaneous tissues.	[56]
	86	Female	Excised cyst from pharynx	Left posterior oropharynx	An 86-year-old woman with cysts in the left posterior oropharynx, brain and subcutaneous tissues on the neck and	[14]
		Subcutat nodule b	Subcutaneous nodule biopsy	Subcutaneous nodules on the back	back.	

224 eastern and southern Africa (2000 - 2022)

Country	Age (yrs.)	Gender	Sample	Location of pathogen	Comment	Reference
	49	Male	Excisional nodule biopsy	Subcutaneous nodules on the trunk and upper extremities	A 49-year-old man with generalized cysticercosis.	[57]
	8	Nr	Cyst from pleural effusion	Right lung	An 8-year-old child with pulmonary cysticercosis who presented with a large right pleural effusion.	[58]
Tanzania	4	Male	Subcutaneous sample biopsy	Sub mandibular and cervical region and abdomen	A 4-year-old HIV positive child with disseminated cysticercosis.	[59]
	48	Male	Subcutaneous nodule	Neck	A 48-year-old male driver with disseminated cysticercosis treated as malaria.	[60]

225

226 ARV, Antiretroviral medication; HIV, Human immunodeficiency virus; Yrs., Years.

227 *detected through musculoskeletal ultrasound of nodules

228 #detected through chest and soft tissue Rontgen radiation (X-rays)

231 Table 2. Individual human neurocysticercosis (NCC) case reports as detected by Brain CT in eastern and southern Africa (2000

- 2022)

Country	Age	Gender	Location of pathogen	Comment	Reference
Eswatini	40	Male	Multiple intra-parenchymal cysts, mural scoleces, calcified granulomas, right sided and parafalcine septated subdural collections, pathognomonic of racemose, stage I and stage IV NCC	A 40-year-old male with a history of chronic headache, unsteady gait and progressive deterioration in vision.	[69]
Madagascar	71	Male	Not described*	A 71-year-old man with dysarthria and memory problems.	[61]
	31	Male	Two ring enhancing lesions in the left capsulo-lenticular. [¶]	A 31-year-old man presenting with dysarthria, blurred vision and monoparesis of the right upper limb 7 weeks after starting anti-tuberculosis medication.	[62]
Malawi	58	Female	Multicystic lesion in the left temporal lobe ^{¶#}	A 58-year-old woman with coincidental racemose NCC who was being treated for HIV associated cryptococcal meningitis, clinically and microbiologically responding to antifungal therapy.	[63]
	41	Female	Multiple intra-parenchymal cyst in the brain	A 41-year-old patient on ARVs with disseminated cysticercosis and more than 50 cysts in brain.	[52]
Mozambique	18	Female	Multiple lesions with different stages ranging from cerebral ring-enhancing cysts with scolices to transitional or degenerative colloidal cysts to multiple parenchymal calcifications	An 18-year-old female admitted to Hospital Central de Quelimane, Zambezia province in Mozambique, 8 days after delivery with a misdiagnosis of eclampsia that turned out to be NCC.	[64]
South Africa	52	Female	Granuloma within bilateral frontal and temporal lobes; Multicystic racemose lesion in the right cerebellar hemisphere; active cyst within the left cerebellar hemisphere; hydrocephalus with blocked ventriculo- peritoneal shunt; Intraventricular cyst	A 52-year-old woman with multiple types of NCC pathology.	[65]

Country	Age	Gender	Location of pathogen	Comment	Reference
	(yrs.)				
	42	Male	Multiple active and calcified lesions in the brain.	A 42-year-old man with NCC and disseminated cysticercosis with myocardial involvement and with multiple calcifications in the muscle and subcutaneous regions.	[56]
	86	Female	Active brain cysts (not specific)	An 86-year-old woman with NCC and cysts in the left posterior oropharynx and subcutaneous tissues on the neck and back.	[14]
	49	Male	Bilateral multiple parenchymal non- enhancing cerebral cystic lesions	A 49-year-old man with generalized cysticercosis.	[57]
	46	Female	Spinal and vertebra region. A spinal lesion at T12, causing vertebral collapse ^{#\$†}	A 46-year-old HIV+ patient with NCC affecting the spine with severe pain with spasms and weakness in lower limbs.	[66]
	30	Female	Cyst in the occipital horn of the right lateral ventricle	A 30-year-old woman with acute unilateral hydrocephalus.	[67]
	8	-	A non-specific solitary sub centimeter calcified granuloma abutting the tentorium cerebellum	Letter to the editor of a case report on an 8-year-old child with pulmonary cysticercosis who presented to Red Cross children' s hospital, Cape Town, with a large right pleural effusion.	[58]
	32	Male	Brain with isolated cystic lesions with scolex and ring enhancing lesion in colloidal stage	A 32-year-old known PWE admitted with COVID-19 and developed recurrent epileptic seizures due to dying cysts after beginning of NCC treatment.	[68]
	42	Female	Not described	A 42-year-old known PWE admitted with COIVD-19 and NCC with poor response to treatment and in status epilepticus for a long time.	
Tanzania	48	Male	Multiple cystic lesions with dot sign diffusely distributed in the brain, neck, chest, including the heart, abdomen, and pelvis [®]	A 48-year-old driver with disseminated cysticercosis treated as malaria.	[60]
Zambia	48	Female	Numerous round points with T1 hypo intense, T2 hyper intense dot-form cores seen in all cerebral slices. [#]	A 48-year-old woman who presented with subcutaneous nodules on the face, recurrent headache and epileptic seizures in Chambishi Zambia.	[70]

	Country	Age	Gender	Location of pathogen	Comment	Reference			
		(yrs.)							
		60	Male	Thoracic intramedullary cystic lesions. Also cystic lesions in subcutaneous tissue ^{#†}	A 60-year-old man with progressive lower extremity weakness and numbness at University Teaching Hospital in Zambia.	[71]			
	Zimbabwe	35	Male	A Suprasellar cystic lesion compressing the optic nerve ^{#\$}	A 35-year-old man presenting with pain accompanied by mild vision loss in the right eye due to a cystic mass compressing the optic nerve.	[72]			
233									
234	Ag-ELISA, Antigen based enzyme linked immunosorbent assay; ARVs, Antiretroviral medication; CSF, Cerebrospinal fluid; CT, Computed tomography; EITB,								
235	Enzyme linked electroimmunotransfer blot; ELISA, Enzyme linked immunosorbent assay; HIV, Human immunodeficiency virus; MRI, Magnetic resonance								
236	imaging; NCC,	Neurocyst	ticercosis; Y	rs., Years.					
237	* Serology with	n Ag-ELIS	A and EITE	B also used for diagnosis					
238	¶CSF ELISA a	lso used fo	r diagnosis						
239	[#] MRI employe	d as detect	ion method						
240	CT of head, n	eck chest, a	abdomen, p	elvis employed as detection method					
241	^{\$} Histopatholog	y employe	d as detecti	on method					
242	[†] NCC affecting	g the spine	with no bra	in involvement					
243 244									

It is made available under a CC-BY 4.0 International license .

245 Aggregated human cysticercosis

246 The number of aggregated human CC reports (community-based, cross-sectional studies) for the 247 period 2000 to 2022 in ESA was 27 (Table 3). These were reports from 9 nine countries, the 248 majority of which were conducted in Tanzania (n=7) followed by Madagascar (n=6) and Zambia 249 (n=4). South Africa, Mozambique, Rwanda and Kenya had two reports each. For Burundi and 250 Uganda one report each was available. There was great variation in the aggregated number of 251 human CC cases across countries with over 2307 cases reported from Madagascar only. The 252 number of human CC cases identified in other ESA countries in documents published between 253 2000 and 2022 is shown in (Fig 2) below.



254

Fig 2. Aggregated human cysticercosis cases reported in community-based cross-sectional studies in eastern and southern Africa between the years 2000 and 2022.

257

258

Country	No.	Prevalence	method	Comment	Reference
	Tested	No. (%)			
Burundi	972	397 (40.8)	Ag-ELISA	A prevalence matched case-control design used in the Kiremba area, Burundi to evaluate the role of cysticercosis in PWE.	[73]
Kenya	2113	122 (5.8)	HP10-ELISA	A cross-sectional survey around the Lake Victoria region to evaluate human and animal zoonotic and non-zoonotic disease in a rural farming community.	[23]
	17	0 (0) ₽	Ag-ELISA	A cross-sectional study to determine the prevalence of cysticercosis and NCC	[18]
	12	1 (8.3) ₽		among PWE (17) and PWOE (12) in a pig keeping community in Western Kenya.	
Madagascar	73	6 (8.2) ₽	EITB	Screening of peace corps volunteers for cysticercosis and NCC before their return from Madagascar.	[74]
	1751	485 (27.7)	B158 / B60	A cross-sectional study to investigate the prevalence of active cysticercosis and associated risk factors in school children (3-16years) in seven cities in Madagascar.	[75]
			Ag-ELISA		
	2094	527 (25.2)	Ag-ELISA	An epidemiology study to evaluate cysticercosis seroprevalence in Madagascar in	[76]
	2048	269 (13.1)	rT24-ELISA	two groups. Group 1: All age groups (2-24) in Antananarivo, capital of Madagascar $(n = 2094)$.	
	1760	489 (27.8)	Ag-ELISA	Group 2: Children aged between 3 to 16 years old from seven different cities (n =	
	1799	242 (13.4)	rT24-ELISA	1799).	
	3619	76 (2.1)	Serology (not specified)	A retrospective study on 76 patients hospitalized for NCC in the pediatrics department of the Tamatave University Hospital Center to evaluate the epidemio- clinical aspect of NCC.	[77]
	237	35 (14.8)	IgG-Western blot	A cross-sectional descriptive study at the Regional Referral Hospital in Antsirabe to determine the seroprevalence of cysticercosis as well as its associated risk factors in patients from the region of Vakinankaratra with clinical suspicion.	[78]
	543	166 (30.6)	Ag-ELISA	A review of cysticercosis in Madagascar. Data for 2017 national survey in	[79]
		12 (2.2)	EITB	Infanadiana district.	
Mozambique	1723	267 (15.5)	Ag-ELISA	A cross-sectional study to estimate the prevalence of NCC in Angónia district, Tete province, Mozambique based on prevalence of human <i>T. solium</i> cysticercosis.	[80]

260 Table 3. Aggregated human cysticercosis cases identified in eastern and southern Africa (2000 - 2022).

Country	No.	Prevalence	method	Comment	Reference
-	Tested	No. (%)			
Mozambique	601	61 (10.2)	IgG-Western blot	A cross-sectional serological study among HIV infected people in Beira (Ponta Gea Health Centre (PGHC)), located in the central part of the Mozambique.	[81]
Rwanda	211	46 (21.8) ᢪ	Immunoblot CSF ELISA	A health-facility based study in the southern province of Rwanda to determine the prevalence of NCC among PWE.	[82]
	572	76 (13.3)	EITB	A cross-sectional study conducted in 680 children from a rural primary school in	[24]
	71	27 (38)	Ag-ELISA	Gakenke district, Northern province to assess the prevalence of <i>T. solium</i> taeniosis cysticercosis.	
South Africa	113	42 (37.2) ₽	IgG-Western blot	A descriptive prospective study to survey the prevalence of NCC in PWE in Lusikisiki, Eastern Cape province.	[19]
	281	89 (32.6)	Ab-ELISA	A cross-sectional study of NCC prevalence among PWE at St Elizabeth hospital	[83]
Tanzania		15 (7.9)	Ag-ELISA	Eastern Cape province.	
Tanzania	20	6 (30) 🖡	Ag-ELISA	A cross-sectional study to assess the prevalence of NCC among PWE in northern Tanzania.	[84]
	830	139 (16.7) 🖡	Ag-ELISA	A cross-sectional study to determine prevalence and risk factors of human <i>T. solium</i> infections in Mbeya Region of Tanzania.	[16]
		376 (45.3)	Ab-ELISA		
	218	6 (2.8) ₽	Ag-/ Ab-ELISA	A cross-sectional study among PWE in a door-to-door study in an established demographic surveillance site in Hai district North East Tanzania.	[85]
	404	10 (2.5)	Ag-/ Ab-ELISA	A cross-sectional study to estimate the prevalence of and factors associated with <i>T</i> . <i>solium</i> infections among HIV+ and HIV - individuals in Mbulu district, northern Tanzania.	[86]
	302	3 (0.99)	Ag-ELISA	A cross-sectional study to investigate the presence and prevalence of TSTC in PWE	[87]
		8 (2.65)	Ab-ELISA	in Dar es Salaam.	
	528	9 (1.7)	Ab-ELISA	Tanzania results from a multi-center case-control study using prevalent cases of ACE identified from HDSS in Tanzania, Kenya, Uganda and South Africa.	[88]
	1051	171 (16.3)	Ab-ELISA	A cross-sectional study conducted to determine the status and health burden of NCC in Mbulu district.	[89]
Uganda	84	2 (2.4)	Ab-/ Ag-ELISA	Uganda results from a multi centre case-control study using prevalent cases of ACE identified from HDSS in Tanzania, Kenya, Uganda and South Africa.	[90]

Country	No.	Prevalence	_ method	Comment	Reference
	Tested	No. (%)			
Zambia	708	41 (5.8)	Ag-ELISA	A community-based cross-sectional survey to determine the prevalence of human taeniosis and cysticercosis in Petauke district in Eastern province.	[20]
	867	(12.2–14.5)	Ag-ELISA	A prospective study conducted to ascertain the incidence of human cysticercosis in	[91]
		(33.5–38.5)	EITB	Katete district in Eastern province.	
	56	22 (39.3)	Ab-ELISA	A cross-sectional study to assess NCC prevalence among PWE in Katete district in	[41]
		13 (23.2)	Ag-ELISA	Eastern province.	
	9	9 (-)	Histopathology Full body dissection postmortem	A descriptive study looking at postmortem examinations of the past 12 months for cardiac cysticercosis and NCC in sudden and unexpected community deaths in Lusaka.	[15]

261

262 ACE, Active convulsive epilepsy; Ab-ELISA, Antibody based ELISA; Ag-ELISA, Antigen based ELISA; B158/B60, an antigen detecting ELISA; CSF,

263 Cerebrospinal fluid; CT, Computed tomography; EITB, Enzyme linked electroimmunotransfer blot; ELISA, Enzyme linked immunosorbent Assay; HDSS, Health

264 demographic surveillance systems; HIV, Human immunodeficiency virus; HP10-ELISA, an antigen detecting ELISA; NCC, Neurocysticercosis; No., Number;

265 PWE, People with epilepsy; PWOE, People without epilepsy; rT24-ELISA; A recombinant antibody detecting ELISA; TSTC, *Taenia solium* taeniosis cysticercosis.

266

267 Employed imaging in addition to serology.

It is made available under a CC-BY 4.0 International license .

To ascertain cases of aggregated human CC, various diagnostic methods were utilized in the included reports with eight (30%) studies employing a combination of antibody and antigen (Ab/Ag) based enzyme linked immunosorbent assay (ELISA), five (19%) employed Ag-ELISA, while three reports (11%) utilized a combination of Ag-ELISA and electroimmunotransfer blot assay (EITB). One report (4%) was based on postmortem findings of CC through full-body dissection [15]. The other methods used for CC diagnosis are shown in (Fig 3).

274



275



278 The prevalence of human CC in ESA in different study populations showed wide variation within

and between countries (Fig 4 and Table 3). The CC seroprevalence ranged between 0.99 - 40.8%

280 on Ag tests and between 1.7 - 45.3% on Ab tests. The highest point prevalence was reported in

281 Tanzania (45.3%) from a study conducted in the Mbozi area of Mbeya District [16].

It is made available under a CC-BY 4.0 International license .



282

Fig 4. The distribution of human cysticercosis seroprevalence in the included studies for
eastern and southern Africa.

For the studies conducted within Tanzania, this was an outlier value. High prevalence variation
was also observed within countries as seen in Madagascar, Rwanda, South Africa and Zambia (Fig
5). The lowest point prevalence (0.99%) was also reported from Tanzania. For details on sources
refer to Table 3.

It is made available under a CC-BY 4.0 International license .





Fig 5. Prevalence of human cysticercosis in eastern and southern Africa based on serology
data in reviewed studies with high variation observed in Rwanda, South Africa, Tanzania
and Zambia

294 Aggregated human neurocysticercosis

For reports presenting aggregated NCC cases (Table 4), 16 records were available. Four (25%) employed imaging (brain CT or MRI) only, whereas 11 (69%) incorporated both imaging and serological tests, and one (6%) was based on postmortem full-body dissection. The NCCsuggestive lesions on brain CT scans showed a prevalence range between 1.0 - 76% in different study populations in ESA (Fig 6 and Table 4).

300

301

302

It is made available under a CC-BY 4.0 International license .



304



307 One report utilized MRI to identify lesions suggestive of NCC in HIV+ children. In this report,

308 four children (8.5% of the total) were found to have NCC, with imaging suggesting the vesicular

309 (active) stage of NCC in all of them [17].

- 310
- 311
- 312

Country	No.	Prevalence	Comment			
	Tested	No. (%)				
Kenya	17	¶(0) 0	A cross-sectional study to determine the prevalence of cysticercosis and NCC among PWE	[18]		
	12	1 (8.3) ₽	(17) and PWOE (12) in a pig keeping community in Western Kenya.			
Madagascar	73	1 (1.4) ᢪ	reening of peace corps volunteers for cysticercosis and NCC before their return from adagascar.			
Malawi	98	1 (1)	Hospital study on patients presenting with a stroke in Blantyre hospital.	[92]		
Mozambique	151	77 (51) 🎙	A cross-sectional study to estimate the prevalence of NCC in Angónia district, Tete province, Mozambique based on prevalence of human <i>T. solium</i> cysticercosis.			
Namibia	117	96 (51.4)	An analytical cross-sectional study conducted among patients with a first seizure presenting to the casualty and medical in-patient wards of Oshakati hospital in northern Namibia.	[93]		
Rwanda	211	46 (21.8) ₽	A health-facility based study in the southern province of Rwanda to determine the prevalence of NCC among PWE.	[82]		
South Africa	113	69 (61.1)₿	A descriptive prospective study to survey the prevalence of NCC in PWE in Lusikisiki, Eastern Cape province.	[19]		
	92	34 (37) ₽	A cross-sectional study of NCC prevalence among PWE at St Elizabeth hospital Eastern Cape province.	[83]		
	75	14 (18.7)	A retrospective review of the brain CT findings in adults presenting with new-onset seizures in a tertiary hospital in Gauteng province.	[94]		
Tanzania	212	68 (32.1)	A cross-sectional study to assess the prevalence of NCC among PWE in northern Tanzania.	[84]		
	55	30 (54.6) ₽	A cross-sectional study to determine prevalence and risk factors of human <i>T. solium</i> infections in Mbeya Region of Tanzania.	[16]		
	200	8 (4) ᢪ	A cross-sectional study among PWE in a door-to-door study in an established demographic surveillance site in Hai district North East Tanzania.	[85]		
	25	19 (76) 🖡	A cross-sectional study conducted to determine the status and health burden of NCC in Mbulu district.	[89]		

Table 4. Aggregated human neurocysticercosis cases as detected by Brain CT scan in eastern and southern Africa (2000 - 2022)

Country	No.	Prevalence	nce Comment		
	Tested	No. (%)			
Zambia	49	20 (41) 🎙	A cross-sectional study to assess NCC prevalence among PWE in Katete district in Eastern province.	[41]	
	34	3 (8.8) *	Observational study of cognition and HIV in children's hospital Lusaka. Three of 34 HIV+	[17]	
	13	1 (7.7) *	participants and one of 13 controls were found to have neurocysticercosis.		
	9	9 (-) #	A descriptive study looking at postmortem examinations for the past 12 months for cardiac cysticercosis and NCC in sudden and unexpected community deaths in Lusaka.	[15]	

315

316 CC+, Cysticercosis positive; CT, Computed tomography; EITB, Enzyme linked electroimmunotransfer blot; HIV, Human immunodeficiency virus; MRI, Magnetic

317 resonance imaging; NCC, Neurocysticercosis; No., Number; PWE, People with epilepsy; PWOE, People without epilepsy.

- 318 *NCC detected through use of MRI
- 319 [#] NCC detected through full body dissection postmortem histopathology

320 ^P Employed serology in addition to imaging

It is made available under a CC-BY 4.0 International license .

322 Neurocysticercosis and epilepsy

323	Among the 16 records reporting aggregated NCC, nine (56%) reported data linking NCC and
324	epilepsy (Table 5). Three of these records were reported from South Africa whereas one record
325	each was reported from Kenya, Mozambique, Namibia, Rwanda, Tanzania and Zambia. Three
326	(33%) of these NCC epilepsy studies were community-based studies and six (67%) were
327	conducted in hospital/health facility settings. The prevalence of NCC among people with epilepsy
328	ranged between $0 - 61\%$. One study in Kenya in a <i>T. solium</i> endemic area found none of the
329	participants with epilepsy had serological evidence of cysticercosis and none had radiographic
330	findings consistent with NCC [18]. Whereas at St Elizabeth's Hospital, Lusikisiki, Eastern Cape,
331	61% of the patients presenting with epilepsy had NCC associated epilepsy, the prevalence being
332	highest in the 10 - 19-year-old age group [19]. For details on sources refer to Table 5.

341 Table 5. Human neurocysticercosis with epilepsy as detected by Brain CT scan in eastern and southern Africa (2000 - 2022).

Country	No	Prevalence	Comment	Reference
	Tested	No (%)		
Kenya	17	0 (0) ^ℙ	A cross-sectional study to determine the prevalence of cysticercosis and NCC among PWE in a pig keeping community in Western Kenya	[18]
Mozambique	151	77 (51) 『	A cross-sectional study to estimate the prevalence of NCC in Angónia district, Tete province, Mozambique based on prevalence of human <i>T. solium</i> cysticercosis.	[80]
Namibia	117	96 (51.4)	An analytical cross-sectional study conducted among patients with a first seizure presenting to the casualty and medical in-patient wards of Oshakati hospital in northern Namibia.	[93]
Rwanda	211	46 (21.8) ₽	A health-facility based study in the southern province of Rwanda to determine the prevalence of NCC among PWE.	[82]
South Africa	113	69 (61.1)	A descriptive prospective study to survey the prevalence of NCC in PWE in Lusikisiki, Eastern Cape province.	[19]
	92	34 (37) ₽	A cross-sectional study of NCC prevalence among PWE at St Elizabeth hospital Eastern Cape province.	[83]
	75	14 (18.7)	A retrospective review of the brain CT findings in adults presenting with new-onset seizures in a tertiary hospital in Gauteng province.	[94]
Tanzania	212	68 (32.1) ₽	A cross-sectional study to assess the prevalence of NCC among PWE in northern Tanzania.	[84]
Zambia	49	20 (41)	A cross-sectional study to assess NCC prevalence among PWE in Katete district in Eastern province.	[41]

342

343 CT, Computed tomography; NCC, Neurocysticercosis; PWE, People with epilepsy.

344 ^P Employed serology in addition to imaging

It is made available under a CC-BY 4.0 International license .

345 Human taeniosis

- 346 A total of 64 records were identified providing information on taeniosis cases in ESA for the period
- 347 2000 to 2022 (Table 6). These records were reported in 11 out of 27 ESA countries (Fig 7) with
- 348 Ethiopia reporting the most records (n=38) followed by Tanzania (n=6) and Zambia (n=5). Kenya
- 349 and Uganda had three records each while Angola, Mozambique, and South Africa had two records
- 350 each. Malawi, Madagascar and Rwanda had one record each. There were no cases reported from
- 351 Botswana, Burundi, Comoros, Djibouti, Eritrea, Lesotho, Mauritius, Mayotte, Namibia, Reunion,
- 352 Seychelles, Socotra, Somalia, Somaliland, Swaziland, and Zimbabwe.



It is made available under a CC-BY 4.0 International license .

Fig 7. The distribution of human taeniosis in eastern and southern Africa showing prevalence data on microscopy and *prevalence based on serology (rES33-immunoblot, Ag-ELISA), and Copro-Ag ELISA)

Several groups of people were studied for human taeniosis with 25 records reporting taeniosis cases in school children, 18 records on patients presenting at health facilities, 15 on consenting community volunteers, and six on food handlers. There was great variation in aggregated numbers of human taeniosis cases across countries with 595 cases reported from Ethiopia alone and even more (4834) reported from Mozambique. The number of human taeniosis cases identified for the other nine countries in ESA is shown in (Fig 8).



364 Fig 8. Taeniosis cases reported in eastern and southern Africa between 2000 – 2020

The largest number of cases were reported as *Taenia spp*. (Table 6) without specifying the type of species. In only two records from Madagascar and Zambia, *T. solium* was confirmed using PCR [20,21]. Microscopy only was employed for 53 studies, and copro Ag-ELISA only in three studies. Five studies used both copro Ag-ELISA and microscopy for taeniosis diagnosis. Three studies

It is made available under a CC-BY 4.0 International license .

used the rES33 EITB for the presence of a tapeworm antibodies in the body. Immunological data
(copro Ag-ELISA or rES33 EITB) on taeniosis was only available from Kenya, Rwanda,
Tanzania, and Zambia.

Within the ESA region, the prevalence of human taeniosis based on microscopy ranged between 0.1 - 14.7% with the highest prevalence reported in a study conducted among food handlers at Wolkite University in Ethiopia [22] (Figs 7 and 9). Among studies based on copro Ag-ELISA conducted in Kenya, Rwanda, Tanzania, and Zambia, the highest prevalence was reported in Kenya at 19.7% [23]. Rwanda had a taeniosis prevalence of 1.4% [24], Tanzania ranged between 0.5 to 5.2% [16,25] [16], [22] and Zambia ranged between 0.3 to 13.8% [20,26] (Fig 10). For details on sources refer to Table 6.



379

Fig 9. Prevalence of taeniosis in eastern and southern Africa based on microscopy data in
reviewed studies

It is made available under a CC-BY 4.0 International license .





384

southern Africa.

383 Fig 10. Prevalence of taeniosis based on copro-Ag ELISA in reviewed studies in eastern and

385			
386			
387			
388			
389			
390			
391			
392			
393			
394			

395 Table 6. Aggregated human taeniosis cases detected by stool microscopy and reported as *Taenia spp* in eastern and southern

Africa (2000 - 2022).

Country	No.	Prevalence	Comment	Reference
	Tested	No (%)		
Angola	1237	1 (0.1)	A cross-sectional study of 1,237 preschool children (0-5 year olds), 1,142 school-aged children	[95]
	1142	2 (0.2)	(6-15 year olds) and 960 women (15 year olds) conducted to understand the distribution of malnutrition anemia malaria schistosomiasis (intestinal and urinary) and geobelminths in	
	960	1 (0.1)	north-western province.	
	328	2. (0.6)	A cross-sectional study conducted among 328 children attending a primary school in Lubango.	[96]
Ethiopia	93	4 (4.3)	A cross-sectional study among food handlers in Awassa town.	[97]
	372	2 (0.5)	A cross-sectional study among HIV infected and non-infected patients in Jimma zone.	[98]
	1021	14 (1.4)	A cross-sectional community based parasitological survey conducted among population living around the Gilgel Gibe dam catchment areas.	[99]
	248	2 (0.8)	A cross-sectional study among HIV negative subjects in Bahir Dar city.	[100]
	459	13 (2.8)	A cross-sectional study to assess the magnitude and pattern of intestinal parasitism in lowland (2.8%) and highland (1.2%) dwellers in Gamo area, south Ethiopia.	[101]
	399	5 (1.2)		
	399	28 (7)	A cross-sectional study to assess prevalence of intestinal parasites infection and associated risk factors among school children in Dagi primary school.	[102]
	343	14 (4.1)	A cross-sectional study to determine the prevalence of intestinal parasitic infections in patients living with HIV/AIDS at Hawassa University referral hospital.	.[103]
	778	10 (1.3)	A cross-sectional study among 12 primary schools in Bahir Dar region.	[104]
	121	5 (4.1)	A cross-sectional study among prison inmates (4.1%) and tobacco farm workers (0.9%) in	[105]
	115	1 (0.9)	Northcentral Ethiopia.	
	385	6 (1.6)	A cross-sectional study among elementary school children in Sanja, northwest Ethiopia.	[106]
	415	1 (0.2)	A cross-sectional study among TB patients in Gondar hospital.	[107]
	272	1 (0.4)	A cross-sectional study among food handlers in food establishments at Hassawa University.	[108]
	260	1 (0.4)	A cross-sectional study among diarrhea children at Jimma health centre.	[109]

Country	No.	Prevalence	_ Comment	Reference
	Tested	No (%)		
Ethiopia	400	3 (0.8)	A cross-sectional study among children of selected primary schools in Chencha town south Ethiopia.	[110]
	307	4 (1.3)	A cross-sectional study among university students at Mekelle University.	[111]
	32,191	322 (1.0)	A retrospective observational study of all stool samples examined for parasites from patients presenting with diarrhea in the period 2007-2012 at Gambo Rural hospital.	[112]
	464	4 (0.9)	A cross-sectional study among patients attending at Anbesame health center, South Gondar.	[113]
	374	20 (5.3)	A cross-sectional study among school children in Dawro Zone, Southern Ethiopia.	[114]
	95	4 (4.2)	A cross-sectional study to determine the prevalence of intestinal parasite among HIV/AIDS patients attending ART clinic at Arba Minch hospital southern Ethiopia.	[115]
	345	14 (4.1)	A cross-sectional study among food handlers at Arba Minch University student's cafeteria, South Ethiopia.	[116]
	401	2 (0.5)	A cross-sectional study among preschool-aged children in Chuahit, Dembia district, Northwest Ethiopia	[117]
	173	4 (2.3)	A cross-sectional study among HIV infected individuals in Wolaita Sodo hospital.	[118]
	503	13 (2.6)	A cross-sectional parasitological and malacological survey among school children in Wolaita zone south Ethiopia.	[119]
	94	2 (2.1)	A cross-sectional study among food handlers at cafeteria of Jimma University specialized hospital.	[120]
	427	2 (0.5)	A cross-sectional study among patients seen at Adwa health centre North Ethiopia.	[121]
	279	2 (2.4)	A cross-sectional study among primary school children of Haike primary school Northeast Ethiopia.	[122]
	256	3 (1.2)	A cross-sectional study conducted at University of Gondar hospital to assess the prevalence of intestinal parasites among pulmonary tuberculosis suspected patients.	[123]
	213	5 (2.3)	A cross-sectional study among TB patients in Arba Minch facility.	[124]
	108	5 (4.6)	A cross-sectional study to compare prevalence of intestinal parasites and associated factors	[125]
	204	4 (2.0)	among adult pre-ART (4.6%) and on ART (2.0%) patients in Goncha Siso Enesie Woreda, East Gojjam, Northwest Ethiopia.	

Country	No.	Prevalence	Comment	Reference
	Tested	No (%)		
Ethiopia	280	8 (2.9)	A cross-sectional study to determine the prevalence of schistosomiasis and other intestinal helminthiasis among school children and free ranging vervet monkeys in Bochessa area, Ziway. Results for school children.	[126]
	247	1 (0.4)	A cross-sectional study to assess intestinal parasitic infections among under-five children attending in Debre Birhan referral hospital, Northeast Ethiopia.	[127]
	391	4 (1.0)	A cross-sectional study conducted among primary school children in Arbaminch Zuria district.	[128]
	273	6 (2.2)	A cross-sectional study among rural school children in Northwest Ethiopia.	[129]
	170	25 (14.7)	A cross-sectional study among food handlers at Wolkite University.	[22]
	406	12 (3)	A cross-sectional study among primary school children in Jawi town, northwest Ethiopia.	[130]
	200	3 (1.5)	A cross-sectional study among food handlers at Wollo University Northeastern Ethiopia.	[131]
	383	5 (1.3)	A cross-sectional study conducted among Gara Riketa primary school children at Hawassa Tula Sub-City, Southern Ethiopia.	[132]
	525	11 (2.1)	A cross-sectional study in three primary schools in Sululta after deworming.	[133]
Kenya	797	1 (0.13)	A cross-sectional study to determine a baseline prevalence of IPIs among children of five years and below at Webuye HDSS area in western Kenya.	[134]
	2113	416 (19.7) *	A cross-sectional survey around the Lake Victoria region to evaluate human and animal	[23]
		6 (0.3)	zoonotic and non-zoonotic disease in a rural farming community.	
	310	17 (5.5)	A cross-sectional study among school children in lodwa municipality Turkana county.	[135]
Madagascar	459	10 (2.2) #	A cross-sectional study aimed at assessing the prevalence of <i>T. Solium</i> infections and associated risk factors in twelve remote villages surrounding Ranomafana national park, Ifanadiana district.	[21]
Malawi	190	4 (2.1)	A cross-sectional survey among preschool children in Mangochi district.	[136]
Mozambique	269	1 (0.4)	A cross-sectional study among children living in urban environs of Maputo and attending the health clinic at xipamanine.	[137]
	83,331	4833 (5.8)	A cross-sectional study among school children across the country.	[138]
Rwanda	144	2 (1.4) *	A cross-sectional study conducted in 680 children from a rural primary school in Gakenke district, Northern province to assess the prevalence of <i>T. solium</i> taeniosis cysticercosis.	[24]

Country	No.	Prevalence	_ Comment	Reference
	Tested	No (%)		
South Africa	510	- (0-1.4)	A cross-sectional study conducted to estimate the prevalence of fecal helminths in a series of rural and urban African school pupils residing at a relatively high altitude above sea level.	[139]
	998	55 (5.6)	A cross-sectional study to determine the prevalence and risk factors of schistosomiasis and STH infection among preschool aged children aged 15 years in Ngwavuma area of Mkhanyakude district.	[140]
Tanzania	820	43 (5.2) *	A cross-sectional study conducted in Mbozi area of Mbeya district. Of the stool samples testing	[16]
		34 (4.1) &	positive for taeniosis by copro-Ag-ELISA, 34 (4.1%) were Ab positive against adult <i>T. solium</i> . Taenia equa were detected in $9(1.1\%)$ stool samples by routine coprology	
		9 (1.1)		
	404	5 (1.2) &	A cross-sectional study to estimate the prevalence of and factors associated with <i>T. solium</i> infections among HIV+ and HIV - individuals in Mbulu district, northern Tanzania	[86]
	1517	9 (0.6) *	Taeniosis estimates after 3 rounds of MDA for schistosomiasis in the two regions Mbozi (0.6%)	[141]
	1501	7 (0.5)	and Mbeya (0.5%).	
	1514	12 (0.8) *	Repeated MDA to school aged children in Mbozi (0.8%) and school aged children in Mbeya (0.5%), Tanzania.	[25]
	1500	8 (0.5)		
	302	5 (1.7) &	A cross-sectional study to investigate the presence and prevalence of TSTC in PWE in Dar es Salaam.	[87]
Uganda	162	1 (0.6)	A cross-sectional study among children attending Mwanamugimu nutrition unit of Mulago hospital, Kampala, Uganda.	[142]
	331	2 (0.6)	A cross-sectional study among slum dwellers without risk of flooding at least 2 km away from the Nakivubo wetland in Kampala.	[143]
	476	14 (2.9)	A cross-sectional study aimed at determining the prevalence and the outcomes (including anaemia) of intestinal parasitic infections and co-infections among children in Kiryandongo refugee camp.	[144]
Zambia	297	1 (0.3)	Prospective study conducted among HIV infected patients in Lusaka.	[145]
	403	4 (0.9)	A cross-sectional study to determine the prevalence and intensity of STH infections, intestinal protozoa and the presence of multiple infections in children attending preschool centres in Kafue district Zambia.	[146]

Country	No.	Prevalence	_ Comment	Reference
	Tested	No (%)		
	718	2 (0.3) #	A community-based cross-sectional survey to determine the prevalence of human taeniosis and	[20]
	712	45 (6.3)*	cysticercosis in Petauke district in Eastern province.	
	226	27 (11.9)*	A prospective study conducted to ascertain the incidence of human cysticercosis in Katete district in Eastern province.	[91]
	54	8 (14.8)*	A cross-sectional study to assess NCC prevalence among PWE in Katete district in Eastern province.	[41]

397

398 Ag-ELISA, Antigen based ELISA; EITB, Enzyme linked electrotransfer blot; ELISA, Enzyme Linked Immunosorbent Assay; HDSS, Health and demographic

399 surveillance system; HIV, Human immune deficiency virus; IPIs; Intestinal parasite infections; MDA, Mass drug administration; NCC, Neurocysticercosis; No.,

400 Number; PWE, People with epilepsy; STH; Soil transmitted helminths; TSTC, *Taenia solium* taeniosis cysticercosis.

401 *Detection by copro Ag-ELISA.

402 *"T. solium* species confirmed.

403 [&] Detection based on serology using EITB (rES33 immunoblot).

It is made available under a CC-BY 4.0 International license .

404 **Discussion**

405 This study aimed at collecting epidemiological data on T. solium infections in humans in the ESA 406 region for the period 2000 to 2022. For both cysticercosis and taeniosis, there was no data from 407 Comoros, Djibouti, Eritrea, Mayotte, Reunion, Seychelles, Socotra, Somalia and Somaliland that 408 could be retrieved. Somalia has for some time now experienced internal conflicts and this may 409 have a bearing on the diagnosis and reporting of human cysticercosis and taeniosis cases. For the 410 other six island states and territories, a lack of scientific interest in reporting cases has been cited 411 [27]. The same can be said about Botswana and Lesotho from which data on cysticercosis and 412 taeniosis was also lacking. Cysticercosis data was lacking in even more countries.

413 According to the World Bank IBRD-IDA report of 2022 [28] the population using safely managed 414 sanitation services for both Comoros and Djibouti only stood at 37% with no data for other island 415 states. People practicing open defecation were higher in Eritrea (67%) with Somalia reporting 416 (23%) and Djibouti (16%). On the Réunion island, pork is said to be the most popular meat with 417 a total annual pork consumption of about 20 000 tones. Half of this is produced on the island and 418 the rest is imported frozen from Europe [29]. Comoros on the other hand has a small livestock 419 sector with most of its meat including pork imported from Tanzania, Madagascar and France [30]. 420 Seventy per cent of the Somali human population subsists in pastoralism with sheep and goats 421 being the dominant animals. Camels are primarily raised for milk production and small ruminants 422 are for generating cash income for the family [31]. For Mayotte, Eritrea and Djibouti, information 423 on pig rearing could not be found as these countries focus mostly on cattle, sheep, goats, and 424 camels [32]. However, for Eritrea, pig breeding has recently begun to be promoted [33] and so is 425 the case for Seychelles whose 2011 census of agriculture showed that 2% (483/24770) of the

426 households were raising pigs in small holdings [34]. Therefore, while it is not possible to ascertain 427 the presence or absence of human *T. solium* infections in these countries, there is a possibility that 428 these infections do exist as pork consumption seems to be practiced and risk factors are present; 429 some countries even import pork from confirmed *T. solium* endemic countries. Therefore, the 430 numbers collected for this review represent an underestimation of the *T. solium* epidemiology for 431 human disease in ESA.

432 Regarding human cysticercosis cases, tapeworm carriers are the sole cause of infection in both 433 rural and urban areas of the ESA region [10,35]. Tapeworm carriers, if not treated, pose a risk to 434 themselves if they ingest infective eggs leading to cysticercosis but can also pose a risk to other 435 people in contact [21,36]. A study in Madagascar found at a household level, the overall 436 seroprevalence of IgG detected by Ab-ELISA and EITB within the same household in which a 437 tapeworm carrier has been detected was 46% and 34% respectively [21]. Individual human 438 cysticercosis cases were identified from a third of the 27 ESA countries. South Africa reported the 439 highest number of individual human cysticercosis cases followed by Madagascar. This could be 440 due to increased awareness of cysticercosis as a possible diagnosis in, for example, people with 441 epilepsy in these countries compared to other countries within the region. South Africa also has 442 more resources compared to the other countries within the region. For example, as early as 2003 443 South Africa already had 214 CT scans and 111 neurologists. This is an advantage in terms of diagnosis and management of particularly NCC that is not present in other countries within the 444 445 region [37]. With poor pig management, poor meat inspection, and poor sanitation those who do 446 not eat pork are equally at risk of infection with cysticercosis as those eating pork [38-40]. 447 Additionally, certain cultural practices within the region have also been cited as contributing 448 factors to cysticercosis. For example, in South Africa, it has been reported that some unqualified

It is made available under a CC-BY 4.0 International license .

449 traditional healers "baloi" used *Taenia* segments and added them to medicinal mixtures as 450 strengthening ingredients. There are also reports of *T. solium* segments being added to the beer of 451 unfaithful husbands or lovers as punishment [37].

452 NCC has been associated with up to 57% of epilepsy cases in sub–Saharan Africa to which the 453 region under consideration belongs [7,41]. This is higher than the global average of 30% of NCC 454 among PWE [6]. This association is higher than the global average of 30% [6]. For most countries 455 within ESA, the diagnosis of NCC is problematic due to the scarcity of neuroimaging, serology 456 and trained neurologists [42,43]. This diagnostic challenge is not only present in ESA but also in 457 Europe where a lack of awareness of NCC leads to underdiagnoses [44]. Only a quarter of the 458 countries within the ESA reported data on NCC and epilepsy. Epilepsy is one of the most common 459 neurological disorders in many parts of sub-Saharan Africa and NCC seems to be a major cause 460 of it in *T. solium* cysticercosis endemic areas. Diagnosing NCC among people with epilepsy is 461 vital to prevent further morbidity and mortality from the disease as well as to reduce the negative 462 socio-cultural beliefs associated with epilepsy in the region [42,45–47].

463 Regarding human T. solium taeniosis in ESA, only Kenya, Madagascar, Rwanda, Tanzania, and 464 Zambia conducted community-based studies and employed serological tests that specifically 465 aimed at determining the burden of infection. The other countries reported data on taeniosis as 466 incidental findings following microscopic examinations of stool samples conducted for soil-467 transmitted helminths. The reported human taeniosis cases were also reported on an aggregated 468 level without evidence of species determination. Thus, cases reported as either taeniosis or *taenia* 469 spp could not for instance be differentiated from cases due to T. saginata which is also widely 470 distributed in ESA [27]. Only two studies, one from Madagascar and another from Zambia were 471 able to confirm the *T. solium* species after collecting tapeworm proglottids [20,21].

It is made available under a CC-BY 4.0 International license .

This lack of species differentiation creates uncertainty regarding the epidemiology of *T. solium* infections in ESA. There is therefore a possible underestimation of the magnitude of *T. solium* taeniosis in the region. This is worsened by the fact that *T. solium* taeniosis is not even a notifiable disease in most of these countries. Application of molecular methods to differentiate *T. solium* species on stool examination is not widely practiced and it is difficult to differentiate *taenia* eggs based on microscopy [35,48]. More sensitive and specific diagnostic tools need to be used to identify the true prevalence and outline the epidemiology of *T. solium*.

479 Human taeniosis in the region is due to the high prevalence of porcine cysticercosis in ESA 480 countries which is ranked amongst the highest in the world [10]. Over the last decade, pig 481 production as a risk factor for human taeniosis in the rural communities of ESA countries has 482 increased mainly because pig raising is an attractive alternative to other animals as it does not 483 require grazing land compared to ruminants. Free-range pig keeping is popular, especially in rural 484 areas of the region as it is easy and cheap. This coupled with the general lack of slaughterhouses 485 and inspection of pork with a lack of sanitation facilities has contributed to the risk for people to 486 attract taeniosis and cysticercosis [3,10,49].

487 Study limitation

Our study is limited by the fact that search platforms were used only for online publications. Thus, unpublished data which could have contributed significantly to a more nuanced picture of the epidemiology of human *T. solium* infections in ESA could have been missed. Another limitation of our study is the difficulty in comparing epidemiological data from different study populations using different diagnostic tests used and the different diagnostic approaches.

It is made available under a CC-BY 4.0 International license .

493 **Conclusion**

- In this review on human *T. solium* taeniosis and (neuro) cysticercosis within ESA, we found wide variations in the prevailing prevalent estimates depending on the quality of the study and the diagnostic methods used. There remain large gaps with regards to *T. solium* infections within the ESA region with 11 countries without any information on human taeniosis and cysticercosis, and even more countries (18) without any information for cysticercosis.
- 499 Considering the public health and economic impact that T. solium infections have, it is important 500 to understand its epidemiology in humans. Human taeniosis, cysticercosis and NCC cases need to 501 be reported and notified to public health authorities. More community-based surveys as well as 502 ante-mortem and postmortem porcine surveys must be conducted for control and surveillance 503 purposes. Lack of awareness about the presence of T. solium among medical, community and 504 government authorities including inadequate technology for diagnosis of taeniosis, cysticercosis 505 and NCC, in both humans and pigs (cysticercosis) remains a challenge that needs to be addressed 506 for proper surveillance, prevention and control.

507 Acknowledgment

508 None.

509 Supplement

- 510 S1 File. Documentation of literature search. (DOCX)
- 511 S2 File. Data collection forms. (DOCX)

It is made available under a CC-BY 4.0 International license .

512 S1 Table. PRISMA 2020 checklist. (DOCX)

513 S2 Table. References retrieved through online international databases. (XLS)

514 Funding

- 515 This study was fully funded by the German Federal Ministry of Education and Research (BMBF)
- 516 under CYSTINET-Africa 2018-Mar-002 and 01KA1618. The funder had no role in the design of
- 517 the study and collection, analysis, and interpretation of data and in writing the manuscript.

518 Availability of data and material

519 All references found eligible in our review are included in the supplementary materials (S2 Table).

520 **Competing interests**

521 The authors declare that they have no competing interests.

522 Author contributions

523 GZ, CM and HS conducted the systematic review of the literature and extracted data. GZ analysed 524 the data and drafted the first version of the manuscript. AA, ASW, CM, DS, GZ, IGK, KEM, VS 525 contributed to the design of the study. All authors contributed to interpretation of data and writing 526 of the paper. All authors read and approved the final manuscript.

527

It is made available under a CC-BY 4.0 International license .

529 **References**

- J. Sotelo, O.H. Del Brutto, Review of neurocysticercosis, Neurosurg. Focus. 12 (2002) e1.
 https://doi.org/10.3171/foc.2002.12.6.2.
- 532 [2] K.D. Murrell, WHO/FAO/OIE guidelines for the surveillance, prevention and control of
 533 taeniosis/cysticercosis. Paris: World Health Organisation for Animal Health (OIE), 2005,
 534 2005. http://www.oie.int (accessed May 6, 2019).
- H. Carabin, A. Millogo, A. Cissé, S. Gabriël, I. Sahlu, P. Dorny, C. Bauer, Z. Tarnagda,
 L.D. Cowan, R. Ganaba, Prevalence of and Factors Associated with Human Cysticercosis
 in 60 Villages in Three Provinces of Burkina Faso, PLoS Negl. Trop. Dis. 9 (2015)
 e0004248. https://doi.org/10.1371/journal.pntd.0004248.
- A. Winkler, H. Richter, Landscape analysis : management of neurocysticercosis with an
 emphasis on low- and middle-income countries, World Heal. Organ. (2015) 65.
 www.neurokopfzentrum.med.tum.de/neurologie/42b.html (accessed February 22, 2018).
- 542 [5] D. Stelzle, V. Schmidt, L. Keller, B.J. Ngowi, W. Matuja, G. Escheu, P. Hauke, V.
 543 Richter, E. Ovuga, B. Pfausler, E. Schmutzhard, A. Amos, W. Harrison, J. Kaducu, A.S.
 544 Winkler, Characteristics of people with epilepsy and Neurocysticercosis in three eastern
 545 African countries–A pooled analysis, PLoS Negl. Trop. Dis. 16 (2022) e0010870.
 546 https://doi.org/10.1371/journal.pntd.0010870.
- 547 [6] P.C. Ndimubanzi, H. Carabin, C.M. Budke, H. Nguyen, Y.J. Qian, E. Rainwater, M.
 548 Dickey, S. Reynolds, J.A. Stoner, A systematic review of the frequency of
 549 neurocyticercosis with a focus on people with epilepsy, PLoS Negl. Trop. Dis. 4 (2010)
 550 e870. https://doi.org/10.1371/journal.pntd.0000870.
- L.F. Owolabi, B. Adamu, A.M. Jibo, S.D. Owolabi, A.I. Imam, I.D. Alhaji,
 Neurocysticercosis in people with epilepsy in Sub-Saharan Africa: A systematic review
 and meta-analysis of the prevalence and strength of association, SeizureEuropean J.
 Epilepsy. 76 (2020) 1–11. https://doi.org/10.1016/j.seizure.2020.01.005.
- 555 [8] E. Bruno, A. Bartoloni, L. Zammarchi, M. Strohmeyer, F. Bartalesi, J. Bustos, S. 556 Santivañez, H.H. Garcia, A. Nicoletti, M. Bonati, F. Severino, V. Confalonieri, C. 557 Pandolfini, Z. Bisoffi, D. Buonfrate, A. Angheben, M. Albonico, J. Muñoz, R. Pool, A. 558 Requena-Mendez, M. Roura, A. Hardon, C. Pell, P. Chiodini, J. Moreira, R. Sempértegui, 559 M. Anselmi, E. Gotuzzo, M.A. Mena, F. Torrico, D. Lozano, G.C. Rojas, T.H. Cabrera, 560 J.O. Morón, I.A. Cuellar, J.A. Suarez, G. Tognoni, C.L. Caro, Epilepsy and 561 Neurocysticercosis in Latin America: A Systematic Review and Meta-analysis, PLoS 562 Negl. Trop. Dis. 7 (2013). https://doi.org/10.1371/journal.pntd.0002480.
- 563 [9] D. Ng-Nguyen, R.J. Traub, V.A.T. Nguyen, K. Breen, M.A. Stevenson, Spatial
 564 distribution of Taenia solium exposure in humans and pigs in the Central Highlands of
 565 Vietnam, PLoS Negl. Trop. Dis. 12 (2018). https://doi.org/10.1371/journal.pntd.0006810.

- I.K. Phiri, H. Ngowi, S. Afonso, E. Matenga, M. Boa, S. Mukaratirwa, S. Githigia, M.
 Saimo, C. Sikasunge, N. Maingi, G.W. Lubega, A. Kassuku, L. Michael, S. Siziya, R.C.
 Krecek, E. Noormahomed, M. Vilhena, P. Dorny, A.L. Willingham, The emergence of
 Taenia solium cysticercosis in Eastern and Southern Africa as a serious agricultural
 problem and public health risk, in: Acta Trop., Elsevier, 2003: pp. 13–23.
 https://doi.org/10.1016/S0001-706X(03)00051-2.
- M.J. Page, J.E. McKenzie, P.M. Bossuyt, I. Boutron, T.C. Hoffmann, C.D. Mulrow, L.
 Shamseer, J.M. Tetzlaff, E.A. Akl, S.E. Brennan, R. Chou, J. Glanville, J.M. Grimshaw,
 A. Hróbjartsson, M.M. Lalu, T. Li, E.W. Loder, E. Mayo-Wilson, S. McDonald, L.A.
 McGuinness, L.A. Stewart, J. Thomas, A.C. Tricco, V.A. Welch, P. Whiting, D. Moher,
 The PRISMA 2020 statement: An updated guideline for reporting systematic reviews,
 BMJ. 372 (2021). https://doi.org/10.1136/bmj.n71.
- 579 [12] M. Viswanathan, A. MT, N. Berkman, S. Chang, L. Hartling, M. et al. McPheeters,
 580 Assessing the risk of bias of individual studies in systematic reviews of health care
 581 interventions methods guide chapter | AHRQ effective health care program, (2016).
 582 www.effectivehealthcare.ahrq.gov. (accessed September 7, 2018).
- 583 [13] M. Mugabe, D. Ruhangaza, E. Hakizimana, Cysticercosis: Report of 10 cases diagnosed at a district hospital in Rwanda, Mod. Pathol. 32 (2019).
 585 http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emexb&NEWS=N&AN 586 =631814441.
- 587 [14] S. Sobnach, S.A. Khosa, S. Pather, S. Longhurst, D. Kahn, P.J. Raubenheimer, First case
 588 report of pharyngeal cysticercosis, Trans. R. Soc. Trop. Med. Hyg. 103 (2009) 206–208.
 589 https://doi.org/10.1016/j.trstmh.2008.08.017.
- 590 [15] C. Himwaze, L.A. Mucheleng'anga, V. Telendiy, A. Hamukale, J. Tembo, N. Kapata, F.
 591 Ntoumi, A. Zumla, Cardiac cysticercosis and neurocysticercosis in sudden and unexpected
 592 community deaths in Lusaka, Zambia: a descriptive medico-legal post-mortem
 593 examination study, Int. J. Infect. Dis. 115 (2022) 195–200.
 594 https://doi.org/10.1016/j.ijid.2021.11.042.
- 595 [16] G. Mwanjali, C. Kihamia, D.V.C. Kakoko, F. Lekule, H. Ngowi, M.V. Johansen, S.M.
 596 Thamsborg, A.L. Willingham, Prevalence and Risk Factors Associated with Human
 597 Taenia Solium Infections in Mbozi District, Mbeya Region, Tanzania, PLoS Negl. Trop.
 598 Dis. 7 (2013) e2102. https://doi.org/10.1371/journal.pntd.0002102.
- A. Buda, O. Dean, H.R. Adams, S. Mwanza-Kabaghe, M.J. Potchen, E.G. Mbewe, P.P.
 Kabundula, S.M. Moghaddam, M. Mweemba, B. Matoka, M.M. Mathews, G.L. Birbeck,
 D.R. Bearden, Neurocysticercosis Among Zambian Children and Adolescents With
 Human Immunodeficiency Virus: A Geographic Information Systems Approach, Pediatr.
 Neurol. 102 (2020) 36–43. https://doi.org/10.1016/j.pediatrneurol.2019.07.017.
- M.M. Diaz, D. Sokhi, J. Noh, A.K. Ngugi, F.J. Minja, P. Reddi, E.M. Fèvre, A.-C.L.
 Meyer, Prevalence of Epilepsy, Human Cysticercosis, and Porcine Cysticercosis in
 Western Kenya, Am. J. Trop. Med. Hyg. 106 (2022) 1450–1455.

- 607 https://doi.org/10.4269/ajtmh.21-0594.
- 608 [19] G.S. Ocana, J.C.O. Sablon, I.O. Tamayo, L.A. Arena, L.M.S. Ocana, S. Govender,
 609 Neurocysticercosis in patients presenting with epilepsy at St. Elizabeth's Hospital,
 610 Lusikisiki, South African Med. J. 99 (2009) 588–591. https://doi.org/10.7196/SAMJ.3358.
- [20] K.E. Mwape, I.K. Phiri, N. Praet, J.B. Muma, G. Zulu, P. van den Bossche, R. de Deken,
 N. Speybroeck, P. Dorny, S. Gabriël, Taenia solium infections in a rural area of Eastern
 Zambia-A community based study, PLoS Negl. Trop. Dis. 6 (2012) 1–9.
 https://doi.org/10.1371/journal.pntd.0001594.
- A. Rahantamalala, R.L. Rakotoarison, E. Rakotomalala, M. Rakotondrazaka, J. Kiernan,
 P.M. Castle, L. Hakami, K. Choi, A.S. Rafalimanantsoa, A. Harimanana, P. Wright, S.G.
 Lapierre, M. Schoenhals, P.M. Small, L.A. Marcos, I. Vigan-Womas, Prevalence and
 factors associated with human Taenia solium taeniosis and cysticercosis in twelve remote
 villages of Ranomafana rainforest, Madagascar, PLoS Negl. Trop. Dis. 16 (2022)
 e0010265. https://doi.org/10.1371/journal.pntd.0010265.
- [22] T.A. Bafa, E.M. Sherif, A.H. Hantalo, G.G. Woldeamanuel, Magnitude of
 enteropathogens and associated factors among apparently healthy food handlers at
 Wolkite University Student's Cafeteria, Southern Ethiopia, BMC Res. Notes. 12 (2019)
 1–6. https://doi.org/10.1186/s13104-019-4599-z.
- E.M. Fèvre, W.A. de Glanville, L.F. Thomas, E.A.J. Cook, S. Kariuki, C.N. Wamae, An
 integrated study of human and animal infectious disease in the Lake Victoria crescent
 small-holder crop-livestock production system, Kenya, BMC Infect. Dis. 17 (2017) 1–14.
 https://doi.org/10.1186/s12879-017-2559-6.
- [24] L. Acosta Soto, L.A. Parker, M.J. Irisarri-Gutiérrez, J.A. Bustos, Y. Castillo, E. Perez, C.
 Muñoz-Antoli, J.G. Esteban, H.H. García, F.J. Bornay-Llinares, Evidence for
 Transmission of Taenia solium Taeniasis/Cysticercosis in a Rural Area of Northern
 Rwanda, Front. Vet. Sci. 8 (2021) 237. https://doi.org/10.3389/fvets.2021.645076.
- U.C. Braae, P. Magnussen, B. Ndawi, W. Harrison, F. Lekule, M.V. Johansen, Effect of
 repeated mass drug administration with praziquantel and track and treat of taeniosis cases
 on the prevalence of taeniosis in Taenia solium endemic rural communities of Tanzania,
 Acta Trop. 165 (2017) 246–251. https://doi.org/10.1016/j.actatropica.2015.10.012.
- K.E. Mwape, I.K. Phiri, N. Praet, P. Dorny, J.B. Muma, G. Zulu, N. Speybroeck, S.
 Gabriël, Study and ranking of determinants of taenia solium infections by classification tree models, Am. J. Trop. Med. Hyg. 92 (2015) 56–63. https://doi.org/10.4269/ajtmh.13-0593.
- [27] V. Dermauw, P. Dorny, U.C. Braae, B. Devleesschauwer, L.J. Robertson, A. Saratsis, L.F.
 Thomas, Epidemiology of Taenia saginata taeniosis/cysticercosis: A systematic review of
 the distribution in southern and eastern Africa, Parasites and Vectors. 11 (2018) 578.
 https://doi.org/10.1186/s13071-018-3163-3.

645 646	[28]	The World Bank IBRD-IDA, Countries Data, (2022). https://data.worldbank.org/country (accessed November 17, 2022).
647 648 649	[29]	L. Phillips, Successful meat production on Réunion Island, Farmer's Wkly. (2014). https://www.farmersweekly.co.za/animals/cattle/successful-meat-production-on-reunion- island/ (accessed September 27, 2022).
650 651	[30]	H.C. Metz, Comoros: A Country Study, Washingt. GPO Libr. Congr. (1994). http://countrystudies.us/comoros/ (accessed September 27, 2022).
652 653 654	[31]	A.A. Elmi, Livestock production in Somalia with special emphasis on camels, Nomad. People. 29 (1991) 87–103. https://www.jstor.org/stable/43123342 (accessed November 17, 2022).
655 656	[32]	EAFF 2022, East Africa Farmers Federation, (n.d.). https://www.eaffu.org/djibouti/ (accessed September 23, 2022).
657 658 659	[33]	Eritrea Ministry of Information, Livestock Population and Types, (2012). https://shabait.com/2012/12/24/livestock-population-and-types-of-eritrea/ (accessed September 27, 2022).
660 661	[34]	Mr Marc Naiken; Ms. Laura Ah Time, Seychelles Census of Agriculture, (2011). https://wedocs.unep.org/xmlui/handle/20.500.11822/9520 (accessed November 17, 2022).
662 663 664	[35]	K.E. Mwape, S. Gabriël, The Parasitological, Immunological, and Molecular Diagnosis of Human Taeniasis with Special Emphasis on Taenia solium Taeniasis, Curr. Trop. Med. Reports. 1 (2014) 173–180. https://doi.org/10.1007/s40475-014-0028-5.
665 666	[36]	H.H. Garcia, O.H. Del Brutto, Neurocysticercosis: Updated concepts about an old disease, Lancet Neurol. 4 (2005) 653–661. https://doi.org/10.1016/S1474-4422(05)70194-0.
667 668 669	[37]	N.A. Mafojane, C.C. Appleton, R.C. Krecek, L.M. Michael, A.L. Willingham, The current status of neurocysticercosis in Eastern and Southern Africa, in: Acta Trop., Elsevier, 2003: pp. 25–33. https://doi.org/10.1016/S0001-706X(03)00052-4.
670 671 672	[38]	H.J. Heinz, G.M. Macnab, Cysticercosis in the Bantu of southern Africa., S. Afr. J. Med. Sci. 30 (1965) 19–31. https://pubmed.ncbi.nlm.nih.gov/5854976/ (accessed May 15, 2022).
673 674 675	[39]	I.K. Phiri, P. Dorny, S. Gabriel, A.L. Willingham, N. Speybroeck, J. Vercruysse, The prevalence of porcine cysticercosis in Eastern and Southern provinces of Zambia, Vet. Parasitol. 108 (2002) 31–39. https://doi.org/10.1016/S0304-4017(02)00165-6.
676 677 678 679	[40]	F. Jansen, P. Dorny, S. Gabriël, V. Dermauw, M.V. Johansen, C. Trevisan, The survival and dispersal of Taenia eggs in the environment: what are the implications for transmission? A systematic review, Parasites and Vectors. 14 (2021) 1–16. https://doi.org/10.1186/s13071-021-04589-6.

680 681 682 683	[41]	K.E. Mwape, J. Blocher, J. Wiefek, K. Schmidt, P. Dorny, N. Praet, C. Chiluba, H. Schmidt, I.K. Phiri, A.S. Winkler, S. Gabriël, Prevalence of neurocysticercosis in people with epilepsy in the Eastern province of Zambia, PLoS Negl. Trop. Dis. 9 (2015) e0003972. https://doi.org/10.1371/journal.pntd.0003972.
684 685	[42]	A. Winkler, Sylvia, Epilepsy and Neurocysticercosis in Sub-Saharan Africa, Nov. Asp. Cysticercosis Neurocysticercosis. (2013). https://doi.org/10.5772/53289.
686 687 688	[43]	A. Millogo, A. Kongnyu Njamnshi, M. Kabwa-PierreLuabeya, Neurocysticercosis and epilepsy in sub-Saharan Africa, Brain Res. Bull. 145 (2019) 30–38. https://doi.org/10.1016/j.brainresbull.2018.08.011.
689 690 691 692 693 694 695 696	[44]	M. Laranjo-González, B. Devleesschauwer, C. Trevisan, A. Allepuz, S. Sotiraki, A. Abraham, M.B. Afonso, J. Blocher, L. Cardoso, J.M. Correia Da Costa, P. Dorny, S. Gabriël, J. Gomes, M.Á. Gómez-Morales, P. Jokelainen, M. Kaminski, B. Krt, P. Magnussen, L.J. Robertson, V. Schmidt, E. Schmutzhard, G.S.A. Smit, B. Šoba, C.R. Stensvold, J. Starič, K. Troell, A.V. Rataj, M. Vieira-Pinto, M. Vilhena, N.A. Wardrop, A.S. Winkler, V. Dermauw, Epidemiology of taeniosis/cysticercosis in Europe, a systematic review: Western Europe, Parasites and Vectors. 10 (2017) 1–14. https://doi.org/10.1186/s13071-017-2280-8.
697 698 699	[45]	R. Baskind, G.L. Birbeck, Epilepsy-associated stigma in sub-Saharan Africa: The social landscape of a disease, Epilepsy Behav. 7 (2005) 68–73. https://doi.org/10.1016/J.YEBEH.2005.04.009.
700 701 702 703	[46]	M. Atadzhanov, A. Haworth, E.N. Chomba, E.K. Mbewe, G.L. Birbeck, Epilepsy- associated stigma in Zambia: what factors predict greater felt stigma in a highly stigmatized population?, Epilepsy Behav. 19 (2010) 414–8. https://doi.org/10.1016/j.yebeh.2010.08.017.
704 705 706 707 708	[47]	M.T. Medina, R.L. Aguilar-Estrada, A. Alvarez, R.M. Durõn, L. Martínez, S. Dubõn, A.L. Estrada, C. Zúniga, D. Cartagena, A. Thompson, E. Ramirez, L. Banegas, J.R. Osorio, A. V. Delgado-Escueta, J.S. Collins, K.R. Holden, Reduction in rate of epilepsy from neurocysticercosis by community interventions: The Salamá, Honduras study, Epilepsia. 52 (2011) 1177–1185. https://doi.org/10.1111/j.1528-1167.2010.02945.x.
709 710 711 712	[48]	J.S. McCarthy, S. Lustigman, G.J. Yang, R.M. Barakat, H.H. García, B. Sripa, A.L. Willingham, R.K. Prichard, M.G. Basáñez, A research agenda for helminth diseases of humans: Diagnostics for control and elimination programmes, PLoS Negl. Trop. Dis. 6 (2012). https://doi.org/10.1371/journal.pntd.0001601.
713 714 715 716	[49]	S. Thys, K.E. Mwape, P. Lefèvre, P. Dorny, T. Marcotty, A.M. Phiri, I.K. Phiri, S. Gabriël, Why Latrines Are Not Used: Communities' Perceptions and Practices Regarding Latrines in a Taenia solium Endemic Rural Area in Eastern Zambia, PLoS Negl. Trop. Dis. 9 (2015) e0003570. https://doi.org/10.1371/journal.pntd.0003570.
717 718	[50]	F.R. Rabenjamina, J. Ranaivoravo, V. Raholimina, A. Ramialiharisoa, A case report of bronchial cysticercosis, Arch. Inst. Pasteur Madagascar. 66 (2000) 43–45.

719		https://europepmc.org/article/med/12463034 (accessed October 23, 2021).
720 721 722	[51]	H.N. Rakoto-Ratsimba, S.S.E.N. Rabesalama, H.J.C. Razafimahandry, A. Ranaivozanany, Case report of solitary breast cysticercosis in Madagascar, Med. Trop. 67 (2007) 179–180. https://pubmed.ncbi.nlm.nih.gov/17691439/ (accessed October 18, 2021).
723 724 725	[52]	T. Heller, C. Wallrauch, D. Kaminstein, S. Phiri, Case report: Cysticercosis: sonographic diagnosis of a treatable cause of epilepsy and skin nodules, Am. J. Trop. Med. Hyg. 97 (2017) 1827–1829. https://doi.org/10.4269/ajtmh.17-0257.
726 727	[53]	S.J. Uledi, A rare gigantic solitary cysticercosis pseudotumour of the neck, J. Surg. Case Reports. 2010 (2010) 5–5. https://doi.org/10.1093/jscr/2010.9.5.
728 729	[54]	S. Agnihotri, O.P. Talwar, S. Pudasaini, R. Baral, Cysticercosis of breast - A case report, Polish J. Pathol. 57 (2006) 53–54. https://doi.org/10.3348/jkrs.1995.32.5.835.
730 731 732 733 734 735 736 737	[55]	J. Tuan, L. Kailani, P. Ngabitsinze, S. Umuganwa, F. Munyaneza, E. Musoni, A.L. Canales, M. Nkeshimana, Disseminated cysticercosis in rwanda—case report of a patient presenting with difficulty with walking and skin nodules, Rwanda Med. J. 77 (2020) 1–4. https://www.google.com/search?q=Disseminated+cysticercosis+in+rwanda—case+report+of+a+patient+presenting+with+difficulty+with+walking+and+skin+nodules &oq=Disseminated+cysticercosis+in+rwanda—case+report+of+a+patient+presenting+with+difficulty+with+walking (accessed July 5, 2022).
738 739 740 741	[56]	M.B. Thomas, K.M. Thomas, A.A. Awotedu, E. Blanco-Blanco, M. Anwary, Cardiocysticercosis, SAMJ South African Med. J. 97 (2007) 504–506. https://go.gale.com/ps/i.do?p=AONE&sw=w&issn=02569574&v=2.1⁢=r&id=GALE% 7CA167881603&sid=googleScholar&linkaccess=fulltext (accessed October 24, 2021).
742 743 744	[57]	E. Agaba, D. Modi, O. Gunduz, Z. Modi, Subcutaneous nodules of cysticercosis as a sign of asymptomatic neurocysticercosis in an HIV positive patient, Rev. Soc. Bras. Med. Trop. 51 (2018) 861–863. https://doi.org/10.1590/0037-8682-0178-2018.
745 746 747	[58]	Z. Nowalaza, M. Zampoli, K. Pillay, S. Singh, H.J. Zar, Pulmonary cysticercosis in an urban South African child, Pediatr. Pulmonol. 56 (2021) 4060–4062. https://doi.org/10.1002/ppul.25682.
748 749 750	[59]	D.W. McCormick, J.M. Bacha, N.K. El-Mallawany, C.L. Kovarik, J.S. Slone, L.R. Campbell, Disseminated cysticercosis and Kaposi sarcoma in a child with HIV/AIDS: A case report, BMC Infect. Dis. 20 (2020) 1–5. https://doi.org/10.1186/s12879-020-05039-x.
751 752 753	[60]	F.R. Lyimo, A.M. Jusabani, H. Makungu, M. Mtolera, S. Surani, Thinking Outside Malaria: A Rare Case of Disseminated Cysticercosis With Cardiopulmonary Involvement From Urban Tanzania, Cureus. 13 (2021). https://doi.org/10.7759/cureus.12851.
754 755	[61]	P. Rajaonarison, S. Ralamboson, C. Andriamamonjy, R. Ramanampamonjy, R. Ce, F. Razafindratrimo, R. Villeneuve, A. Andriantsimahavandy, Diagnostic de la

756 757 758		neurocysticercose : à propos d'un cas, Arch. Inst. Pasteur Madagascar. 67 (2001) 53–56. https://www.researchgate.net/publication/237675160_Diagnostic_de_la_neurocysticercos e_a_propos_d'un_cas (accessed October 23, 2021).
759 760 761 762	[62]	R.A. Rakotoarivelo, R.L. Andrianasolo, D. Ranoharison, R.O.S. Rakoto, F.A. Sendrasoa, M.J. de Dieu Randria, Neurocysticercosis as an important differential of paradoxical response during antituberculosis therapy in HIV-negative patient, Asian Pacific J. Trop. Dis. 1 (2011) 333–334. https://doi.org/10.1016/S2222-1808(11)60077-7.
763 764 765 766 767	[63]	N. Kalata, J. Ellis, L. Benjamin, S. Kampondeni, P. Chiodini, T. Harrison, D.G. Lalloo, R.S. Heyderman, Neurological deterioration in a patient with HIV-associated cryptococcal meningitis initially improving on antifungal treatment: a case report of coincidental racemose neurocysticercosis, BMC Infect. Dis. 21 (2021) 1–5. https://doi.org/10.1186/s12879-021-06425-9.
768 769 770 771 772 773	[64]	V. Saldanha, G. Saldanha, R.P. Reys, C.A. Benson, E.V. Noormahomed, Neurocysticercosis in Child Bearing Women: An Overlooked Condition in Mozambique and a Potentially Missed Diagnosis in Women Presenting with Eclampsia., EC Microbiol. 14 (2018) 736–740. http://www.ncbi.nlm.nih.gov/pubmed/31681909%0Ahttp://www.pubmedcentral.nih.gov/a rticlerender.fcgi?artid=PMC6824723 (accessed July 5, 2022).
774 775	[65]	A.M. Vitali, R. Jithoo, Multiple forms of neurocysticercosis in a single patient, Br. J. Neurosurg. 18 (2004) 650–651. https://doi.org/10.1080/02688690400022847.
776 777 778	[66]	T. Motsepe, D. Ackerman, Spinal and vertebral neurocysticercosis in an HIV-positive female patient, South. African J. Epidemiol. Infect. 27 (2012) 133–136. https://doi.org/10.1080/10158782.2012.11441499.
779 780 781 782	[67]	K. Jivan, A. Mochan, G. Modi, Intraventricular neurocysticercosis causing acute unilateral hydrocephalus, African J. Psychiatry (South Africa). 13 (2010) 315–317. https://www.researchgate.net/publication/47460261_Intraventricular_neurocysticercosis_c ausing_acute_unilateral_hydrocephalus (accessed October 22, 2021).
783 784 785	[68]	H.F. Sibat, Neurocysticercosis, Epilepsy, COVID-19 and a Novel Hypothesis: Cases Series and Systematic Review, Clin. Schizophr. Relat. Psychoses. 15 (2021). https://doi.org/10.3371/CSRP.FH.121421.
786 787 788 789	[69]	B. Ndlovu, Combined surgical and medical treatment of racemose neurocysticercosis in Swaziland, Neuroradiology. 60 (2018) 109–110. https://www.embase.com/search/results?subaction=viewrecord&id=L621458617&from=e xport.
790	[70]	F. Mukonde, Abstracts, Epilepsia. 55 (2014) 4–246. https://doi.org/10.1111/epi.12675.
791 792	[71]	J. Manske, C. McVey, H. Chengazi, O. Siddiqi, D. Saylor, Intramedullary spinal neurocysticercosis (P5.9-031), Neurology. 92 (2019).

793 794 795	[72]	A. Musara, N. Soko, S. Shamu, Suprasellar cysticercosis cyst with optic nerve compression masquerading as an arachnoid cyst, Middle East Afr. J. Ophthalmol. 26 (2019) 114–116. https://doi.org/10.4103/meajo.MEAJO_142_18.
796 797 798	[73]	G. Nsengiyumva, M. Druet-Cabanac, B. Ramanankandrasana, B. Bouteille, L. Nsizabira, P.M. Preux, Cysticercosis as a major risk factor for epilepsy in Burundi, East Africa, Epilepsia. 44 (2003) 950–955. https://doi.org/10.1046/j.1528-1157.2003.55302.x.
799 800 801	[74]	P. Leutscher, A. Andriantsimahavandy, Cysticercosis in Peace Corps Volunteers in Madagascar, N. Engl. J. Med. 350 (2004) 311–312. https://doi.org/10.1056/nejm200401153500325.
802 803 804 805 806	[75]	J Carod, J. Razafimahefa, M. Randrianarison, R Ramahefarisoa, M. Rakotondrazaka, M. Andriantseheno, The burden of neurocysticercosis: A multi-purpose survey in Madagascar, Clin. Microbiol. Infect. 18 (2012) 52–53. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed10&NEWS=N&A N=70822228 (accessed July 8, 2022).
807 808 809 810 811	[76]	J Carod, M. Rakotondrazaka, R Ramahefarisoa, D. Menard, J Soares, P. Dorny, P. Wilkins, Updated epidemiology of cysticercosis in Madagascar, Trop. Med. Int. Heal. 16 (2011) 270. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed10&NEWS=N&A N=70589687 (accessed July 9, 2022).
812 813 814 815	[77]	M.L. Rakotomahefa Narison, A.B.A. Ratsimbazafy, H. Randrianjafinimpanana, N. Randrianaivo, T.N. Andriatahina, H.S. Raobijaona, Aspect épidémio-clinique de la neurocysticercose dans le service de pédiatrie au CHU de Tamatave- Madagascar, Arch. Pediatr. 21 (2014) 810. https://doi.org/10.1016/S0929-693X(14)72070-6.
816 817 818 819 820	[78]	N.J. Zafindraibe, J. Ralalarinivo, A.I. Rakotoniaina, M.N. Maeder, M.R. Andrianarivelo, B. Contamin, A. Michault, A. Rasamindrakotroka, Séroprévalence de la cysticercose et facteurs de risque associés chez un groupe de patients vus au centre hospitalier régional de référence d'antsirabe, madagascar, Pan Afr. Med. J. 28 (2017). https://doi.org/10.11604/pamj.2017.28.260.10463.
821 822	[79]	J.F. Carod, P. Dorny, Cysticercosis in Madagascar, J. Infect. Dev. Ctries. 14 (2020) 1031–1042. https://doi.org/10.3855/JIDC.13450.
823 824 825 826	[80]	Y.A. Assane, C. Trevisan, C.M. Schutte, E.V. Noormahomed, M.V. Johansen, P. Magnussen, Neurocysticercosis in a rural population with extensive pig production in Angónia district, Tete Province, Mozambique, Acta Trop. 165 (2017) 155–160. https://doi.org/10.1016/j.actatropica.2015.10.018.
827 828 829 830 831	[81]	E.V. Noormahomed, N. Nhacupe, C. Mascaró-Lazcano, M.N. Mauaie, T. Buene, C.A. Funzamo, C.A. Benson, A Cross-sectional Serological Study of Cysticercosis, Schistosomiasis, Toxocariasis and Echinococcosis in HIV-1 Infected People in Beira, Mozambique, PLoS Negl. Trop. Dis. 8 (2014) e3121. https://doi.org/10.1371/journal.pntd.0003121.

- R. Rottbeck, J.F. Nshimiyimana, P. Tugirimana, U.E. Düll, J. Sattler, J.C. Hategekimana,
 J. Hitayezu, I. Bruckmaier, M. Borchert, J.B. Gahutu, S. Dieckmann, G. Harms, F.P.
 Mockenhaupt, R. Ignatius, High Prevalence of Cysticercosis in People with Epilepsy in
 Southern Rwanda, PLoS Negl. Trop. Dis. 7 (2013).
 https://doi.org/10.1371/journal.pntd.0002558.
- [83] H. Foyaca-Sibat, L.D. Cowan, H. Carabin, I. Targonska, M.A. Anwary, G. SerranoOcaña, R.C. Krecek, A.L. Willingham, Accuracy of serological testing for the diagnosis
 of prevalent neurocysticercosis in outpatients with epilepsy, Eastern Cape Province, South
 Africa, PLoS Negl. Trop. Dis. 3 (2009) e562.
- 841 https://doi.org/10.1371/journal.pntd.0000562.
- [84] A.S. Winkler, J. Blocher, H. Auer, T. Gotwald, W. Matuja, E. Schmutzhard, Epilepsy and neurocysticercosis in rural Tanzania - An imaging study, Epilepsia. 50 (2009) 987–993.
 https://doi.org/10.1111/j.1528-1167.2008.01867.x.
- [85] E. Hunter, K. Burton, A. Iqbal, D. Birchall, M. Jackson, J. Rogathe, A. Jusabani, W. Gray,
 E. Aris, G. Kamuyu, P.P. Wilkins, C.R. Newton, R. Walker, Cysticercosis and epilepsy in
 rural Tanzania: A community-based case-control and imaging study, Trop. Med. Int. Heal.
 20 (2015) 1171–1179. https://doi.org/10.1111/tmi.12529.
- [86] V. Schmidt, C. Kositz, K.H. Herbinger, H. Carabin, B. Ngowi, E. Naman, P.P. Wilkins, J.
 Noh, W. Matuja, A.S. Winkler, Association between Taenia solium infection and
 HIV/AIDS in northern Tanzania: A matched cross sectional-study, Infect. Dis. Poverty. 5
 (2016). https://doi.org/10.1186/s40249-016-0209-7.
- [87] V. Schmidt, M.C. O'Hara, B. Ngowi, K.H. Herbinger, J. Noh, P.P. Wilkins, V. Richter, C.
 Kositz, W. Matuja, A.S. Winkler, Taenia solium cysticercosis and taeniasis in urban
 settings: Epidemiological evidence from a health-center based study among people with
 epilepsy in Dar es Salaam, Tanzania, PLoS Negl. Trop. Dis. 13 (2019) e0007751.
 https://doi.org/10.1371/journal.pntd.0007751.
- [88] J.O. Mageto, C.I. Muleke, C.R. Newton, G. Samuel, Multiple parasitic infections as risk
 factors of active convulsive epilepsy., Sch. J. Appl. Med. Sci. 3 (2015) 178–184.
 http://saspublisher.com/wp-content/uploads/2015/01/SJAMS-31C178-184.pdf.
- 861 [89] B.J. Mwang'Onde, M.J. Chacha, G. Nkwengulila, The status and health burden of
 862 neurocysticercosis in Mbulu district, northern Tanzania, BMC Res. Notes. 11 (2018) 1–5.
 863 https://doi.org/10.1186/s13104-018-3999-9.
- [90] G. Kamuyu, C. Bottomley, J. Mageto, B. Lowe, P.P. Wilkins, J.C. Noh, T.B. Nutman,
 A.K. Ngugi, R. Odhiambo, R.G. Wagner, A. Kakooza-Mwesige, S. Owusu-Agyei, K. AeNgibise, H. Masanja, F.H.A. Osier, P. Odermatt, C.R. Newton, Exposure to Multiple
 Parasites Is Associated with the Prevalence of Active Convulsive Epilepsy in Sub-Saharan
 Africa, PLoS Negl. Trop. Dis. 8 (2014) e2908.
- 869 https://doi.org/10.1371/journal.pntd.0002908.
- 870

871 872 873	[91]	K.E. Mwape, I.K. Phiri, N. Praet, N. Speybroeck, J.B. Muma, P. Dorny, S. Gabriël, The Incidence of Human Cysticercosis in a Rural Community of Eastern Zambia, PLoS Negl. Trop. Dis. 7 (2013) e2142. https://doi.org/10.1371/journal.pntd.0002142.
874 875 876 877	[92]	J.J.J. Kumwenda, G. Mateyu, S. Kampondeni, A.P. Van Dam, L. Van Lieshout, E.E.E. Zijlstra, A. van Dam, L. van Lieshout, E.E.E. Zijlstra, Differential diagnosis of stroke in a setting of high HIV prevalence in Blantyre, Malawi, Stroke. 17 (2005) 960–964. https://doi.org/10.1161/01.STR.0000162585.97216.ef.
878 879 880	[93]	I.L. Segamwenge, N.P. Kioko, C. Mukulu, O. Jacob, W. Humphrey, J. Augustinus, Neurocysticercosis among patients with first time seizure in Northern Namibia, Pan Afr. Med. J. 24 (2016). https://doi.org/10.11604/pamj.2016.24.127.8908.
881 882 883	[94]	S.H. Mabaso, D. Bhana-Nathoo, S. Lucas, An audit of CT brain findings in adults with new-onset seizures in a resource restricted setting in South Africa, South African J. Radiol. 26 (2022) 7. https://doi.org/10.4102/SAJR.V26I1.2294.
884 885 886 887 888	[95]	J.C. Sousa-Figueiredo, D. Gamboa, J.M. Pedro, C. Fançony, A.J. Langa, R.J.S. Magalhães, J.R. Stothard, S.V. Nery, Epidemiology of malaria, schistosomiasis, geohelminths, anemia and malnutrition in the context of a demographic surveillance system in northern Angola, PLoS One. 7 (2012) e33189. https://doi.org/10.1371/journal.pone.0033189.
889 890 891	[96]	D. Oliveira, F.S. Ferreira, J. Atouguia, F. Fortes, A. Guerra, S. Centeno-Lima, Infection by intestinal parasites, stunting and anemia in school-aged children from southern Angola, PLoS One. 10 (2015) e0137327. https://doi.org/10.1371/journal.pone.0137327.
892 893 894	[97]	S.T. Mariam, B. Roma, S. Sorsa, S. Worku, L. Erosie, Assessment of sanitary and hygienic status of catering establishments of Awassa Town, Ethiop. J. Heal. Dev. 14 (2000). https://doi.org/10.4314/ejhd.v14i1.9934.
895 896 897	[98]	M. Awole, S. Gebre-Selassie, T. Kassa, G. Kibru, Prevalence of intestinal parasites in HIV-infected adult patients in Southwestern Ethiopia, Ethiop. J. Heal. Dev. 17 (2003). https://doi.org/10.4314/ejhd.v17i1.9783.
898 899 900 901	[99]	Z. Mekonnen, S. Suleman, A. Biruksew, T. Tefera, L. Chelkeba, Intestinal polyparasitism with special emphasis to soil-transmitted helminths among residents around Gilgel Gibe Dam, Southwest Ethiopia: a community based survey, BMC Public Health. 16 (2016) 1–7. https://doi.org/10.1186/s12889-016-3859-2.
902 903 904 905	[100]	A. Alemu, Y. Shiferaw, G. Getnet, A. Yalew, Z. Addis, Opportunistic and other intestinal parasites among HIV/AIDS patients attending Gambi higher clinic in Bahir Dar city, North West Ethiopia, Asian Pac. J. Trop. Med. 4 (2011) 661–665. https://doi.org/10.1016/S1995-7645(11)60168-5.
906 907 908	[101]	T. Wegayehu, T. Tsalla, B. Seifu, T. Teklu, Prevalence of intestinal parasitic infections among highland and lowland dwellers in Gamo area, South Ethiopia, BMC Public Health. 13 (2013) 1–7. https://doi.org/10.1186/1471-2458-13-151.

909 910 911	[102]	M. Alamir, W. Awoke, A. Feleke, Intestinal parasites infection and associated factors among school children in Dagi primary school, Amhara National Regional State, Ethiopia Health (Irvine. Calif). 05 (2013) 1697–1701. https://doi.org/10.4236/health.2013.510228.	a,
912 913 914	[103]	S. Fekadu, K. Taye, W. Teshome, S. Asnake, Prevalence of parasitic infections in HIV- positive patients in southern Ethiopia: A cross-sectional study, J. Infect. Dev. Ctries. 7 (2013) 868–872. https://doi.org/10.3855/jidc.2906.	
915 916 917	[104]	B. Abera, G. Alem, M. Yimer, Z. Herrador, Epidemiology of soil-transmitted helminths, Schistosoma mansoni, and haematocrit values among schoolchildren in Ethiopia, J. Infec Dev. Ctries. 7 (2013) 253–260. https://doi.org/10.3855/jidc.2539.	t.
918 919 920	[105]	H. Mamo, Intestinal parasitic infections among prison inmates and tobacco farm workers in Shewa Robit, north-central Ethiopia, PLoS One. 9 (2014) e99559. https://doi.org/10.1371/journal.pone.0099559.	
921 922 923	[106]	L. Worku, D. Damte, M. Endris, H. Tesfa, M. Aemero, Schistosoma mansoni infection and associated determinant factors among school children in Sanja Town, northwest Ethiopia, J. Parasitol. Res. 2014 (2014). https://doi.org/10.1155/2014/792536.	
924 925 926 927 928	[107]	M. Alemayehu, W. Birhan, Y. Belyhun, M. Sahle, B. Tessema, Prevalence of smear positive tuberculosis, intestinal parasites and their co-infection among tuberculosis suspects in Gondar University Hospital and Gondar Poly Clinic, North West Ethiopia, J. Microb. Biochem. Technol. 6 (2014) 179–184. https://doi.org/10.4172/1948-5948.1000140.	
929 930 931 932 933 934 935 936	[108]	M. Desta, D. Asrat, W. Yimtubezinash, N. Demiss, Prevalence of intestinal parasites and Salmonella and Shigella among food handlers at food service establishments in the main campus and Health Sciences College of Hawassa University, Hawassa, Ethiopia, Ethiop. J. Heal. Dev. 28 (2014) 29–34. http://www.etpha.org/publications/the-ethiopian-journal-of-health-development.html%0Ahttp://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&AGE=fulltext&D=cagh&AN=20153041959%0Ahttp://wa4py6yj8t.search.serialssolution.com/?url_ver=Z39.88-2004&rft_val_fmt= (accessed October 18, 2021).	р IS
937 938 939 940	[109]	G. Beyene, H. Tasew, Prevalence of intestinal parasite, Shigella and Salmonella species among diarrheal children in Jimma health center, Jimma southwest Ethiopia: A cross sectional study, Ann. Clin. Microbiol. Antimicrob. 13 (2014) 10. https://doi.org/10.1186/1476-0711-13-10.	
941 942 943	[110]	A. Abossie, M. Seid, Assessment of the prevalence of intestinal parasitosis and associated risk factors among primary school children in Chencha town, Southern Ethiopia, BMC Public Health. 14 (2014). https://doi.org/10.1186/1471-2458-14-166.	ł
944 945 946 947	[111]	A. Gebreyesus, K. Adane, L. Negash, T. Asmelash, S. Belay, M. Alemu, M. Saravanan, Prevalence of Salmonella typhi and intestinal parasites among food handlers in Mekelle University student cafeteria, Mekelle, Ethiopia, Food Control. 44 (2014) 45–48. https://doi.org/10.1016/j.foodcont.2014.03.040.	
		5	56

948 949 950 951 952	[112]	J.M. Ramos, N. Rodríguez-Valero, G. Tisiano, H. Fano, T. Yohannes, A. Gosa, E. Fruttero, F. Reyes, M. Górgolas, Different profile of intestinal protozoa and helminthic infections among patients with diarrhoea according to age attending a rural hospital in southern Ethiopia, Trop. Biomed. 31 (2014) 392–397. https://pubmed.ncbi.nlm.nih.gov/25134911/ (accessed October 27, 2021).
953 954 955	[113]	M.B. Shiferaw, A.D. Mengistu, Helminthiasis: Hookworm infection remains a public health problem in Dera District, South Gondar, Ethiopia, PLoS One. 10 (2015) e0144588. https://doi.org/10.1371/journal.pone.0144588.
956 957 958 959	[114]	A. Bereket, T. Zewdneh, Prevalence of intestinal helminthiasis and associated risk factors among schoolchildren in Dawro Zone, Southern Ethiopia, J. Biol. Agric. Healthc. 5 (2015) 76–82. https://www.iiste.org/Journals/index.php/JBAH/article/view/23238 (accessed October 27, 2021).
960 961 962 963 964 965 966 967	[115]	G. Tigist, E. Fantahun, E. Hailu, H. Kibe, O. Fesseha, W. Godana, W.G. Tsadik, Prevalence of Intestinal Parasitosis Among HIV / AIDS Patients Attending ART Clinic of Arbaminch Hospital., Researchgate.Net. (2015) 1–6. https://www.researchgate.net/profile/Wanzahun- Godana/publication/283351146_PREVALENCE_OF_INTESTINAL_PARASITOSIS_A MONG_HIVAIDS_PATIENTS_ATTENDING_ART_CLINIC_OF_ARBAMINCH_HO SPITAL/links/5636572908aeb786b703f0ae/PREVALENCE-OF-INTESTINAL- PARASITOSIS-AMONG-HIV (accessed October 22, 2021).
968 969 970 971	[116]	M. Mama, G. Alemu, Prevalence and factors associated with intestinal parasitic infections among food handlers of Southern Ethiopia: Cross sectional study Infectious Disease epidemiology, BMC Public Health. 16 (2016) 1–7. https://doi.org/10.1186/s12889-016-2790-x.
972 973 974 975	[117]	A. Alemu, Y. Tegegne, D. Damte, M. Melku, Schistosoma mansoni and soil-transmitted helminths among preschool-aged children in Chuahit, Dembia district, Northwest Ethiopia: Prevalence, intensity of infection and associated risk factors, BMC Public Health. 16 (2016). https://doi.org/10.1186/s12889-016-2864-9.
976 977 978 979	[118]	A. Arota Amado, F. Wadilo Wada, F. Solomon Bisetegn, Y. Abreham Leka, Non- opportunistic intestinal parasitic infections among HIV-infected individuals at Wolaita Sodo Hospital, South Ethiopia, J. Coast. Life Med. 4 (2016) 353–357. https://doi.org/10.12980/jclm.4.2016j6-57.
980 981 982 983	[119]	B. Alemayehu, Z. Tomass, F. Wadilo, D. Leja, S. Liang, B. Erko, Epidemiology of intestinal helminthiasis among school children with emphasis on Schistosoma mansoni infection in Wolaita zone, Southern Ethiopia, BMC Public Health. 17 (2017). https://doi.org/10.1186/s12889-017-4499-x.
984 985 986	[120]	H. Girma, G. Beyene, Z. Mekonnen, Prevalence of intestinal parasites among food handlers at cafeteria of Jimma University Specialized Hospital, Southwest Ethiopia, Asian Pacific J. Trop. Dis. 7 (2017) 467–471. https://doi.org/10.12980/apjtd.7.2017D7-20.

- [121] M. Alemu, B. Kinfe, D. Tadesse, W. Mulu, T. Hailu, E. Yizengaw, Intestinal parasitosis
 and anaemia among patients in a Health Center, North Ethiopia, BMC Res. Notes. 10
 (2017) 1–7. https://doi.org/10.1186/s13104-017-2957-2.
- [122] D.G. Feleke, S. Arega, M. Tekleweini, K. Kindie, A. Gedefie, Schistosoma mansoni and other helminthes infections at Haike primary school children, North-East, Ethiopia: A cross-sectional study, BMC Res. Notes. 10 (2017) 1–6. https://doi.org/10.1186/s13104-017-2942-9.
- 994 [123] Y. Tegegne, T. Wondmagegn, L. Worku, A. Jejaw Zeleke, Prevalence of Intestinal
 995 Parasites and Associated Factors among Pulmonary Tuberculosis Suspected Patients
 996 Attending University of Gondar Hospital, Gondar, Northwest Ethiopia, J. Parasitol. Res.
 997 2018 (2018). https://doi.org/10.1155/2018/9372145.
- 998 [124] G. Alemu, M. Mama, Intestinal helminth co-infection and associated factors among
 999 tuberculosis patients in Arba Minch, Ethiopia, BMC Infect. Dis. 17 (2017) 1–9.
 1000 https://doi.org/10.1186/s12879-017-2195-1.
- [125] T. Asmare, W. Awoke, G. Alem, Prevalence of Intestinal Parasites and Associated Factors among AdultPre-ART and ART Patients in Goncha Siso Enesie Woreda, East Gojjam,Northwest Ethiopia, 2014, J. AIDS Clin. Res. 8 (2017).
 https://doi.org/10.4172/2155-6113.1000734.
- [126] D. Teklemariam, M. Legesse, A. Degarege, S. Liang, B. Erko, Schistosoma mansoni and other intestinal parasitic infections in schoolchildren and vervet monkeys in Lake Ziway area, Ethiopia, BMC Res. Notes. 11 (2018) 1–6. https://doi.org/10.1186/s13104-018-3248-2.
- 1009 [127] T. Zemene, M.B. Shiferaw, Prevalence of intestinal parasitic infections in children under 1010 the age of 5 years attending the Debre Birhan referral hospital, North Shoa, Ethiopia, 1011 BMC Res. Notes. 11 (2018) 1–6. https://doi.org/10.1186/s13104-018-3166-3.
- [128] G. Alemu, Z. Aschalew, E. Zerihun, Burden of intestinal helminths and associated factors three years after initiation of mass drug administration in Arbaminch Zuria district, Southern Ethiopia, BMC Infect. Dis. 18 (2018). https://doi.org/10.1186/s12879-018-3330-3.
- 1016 [129] M. Alemu, A. Anley, K. Tedla, Magnitude of Intestinal Parasitosis and Associated Factors
 1017 in Rural School Children, Northwest Ethiopia, Ethiop. J. Health Sci. 29 (2019) 923–928.
 1018 https://doi.org/10.4314/ejhs.v29i1.14.
- [130] B. Sitotaw, H. Mekuriaw, D. Damtie, Prevalence of intestinal parasitic infections and associated risk factors among Jawi primary school children, Jawi town, north-west Ethiopia, BMC Infect. Dis. 19 (2019). https://doi.org/10.1186/s12879-019-3971-x.
- 1022
- 1023

- [131] E. Kebede, A. Seid, S. Akele, Prevalence and associated risk factors of intestinal parasitic infections among asymptomatic food handlers in Wollo University student's cafeteria, Northeastern Ethiopia, BMC Res. Notes. 12 (2019) 1–6. https://doi.org/10.1186/s13104-019-4182-7.
- [132] T. Eyamo, M. Girma, T. Alemayehu, Z. Bedewi, Soil-Transmitted Helminths And Other
 Intestinal Parasites Among Schoolchildren In Southern Ethiopia, Res. Rep. Trop. Med.
 Volume 10 (2019) 137–143. https://doi.org/10.2147/rrtm.s210200.
- [133] N. Mohamed, A. Muse, M. Wordofa, D. Abera, A. Mesfin, M. Wolde, K. Desta, A.
 Tsegaye, B. Taye, Increased prevalence of cestode infection associated with history of deworming among primary school children in Ethiopia, Am. J. Trop. Med. Hyg. 101
 (2019) 641–649. https://doi.org/10.4269/ajtmh.19-0284.
- [134] A.A. Obala, C.J. Simiyu, D.O. Odhiambo, V. Nanyu, P. Chege, R. Downing, E. Mwaliko,
 A.W. Mwangi, D. Menya, D. Chelagat, H.D.N. Nyamogoba, P.O. Ayuo, W.P. O'Meara,
 M. Twagirumukiza, D. Vandenbroek, B.B.O. Otsyula, J. De Maeseneer, Webuye health
 and demographic surveillance systems baseline survey of soil-transmitted helminths and
 intestinal protozoa among children up to five years, J. Trop. Med. 2013 (2013).
 https://doi.org/10.1155/2013/734562.
- [135] B.N. Lowoko, B. Ngugi, A. Malai, J.K. Mutai, Factors associated with intestinal parasites among school going children in Lodwar Municipality, Turkana County, Kenya, East Afr. Med. J. 94 (2018) 649–663. https://doi.org/10.4314/eamj.v94i8.
- [136] T.P.W. Jones, J.D. Hart, K. Kalua, R.L. Bailey, A prevalence survey of enteral parasites in preschool children in the Mangochi District of Malawi, BMC Infect. Dis. 19 (2019).
 https://doi.org/10.1186/s12879-019-4439-8.
- [137] E. V. Noormahomed, J.G. Pividal, S. Azzouz, C. Mascaró, M. Delgado-Rodríguez, A.
 Osuna, Seroprevalence of anti-cysticercus antibodies among the children living in the
 urban environs of Maputo, Mozambique, Ann. Trop. Med. Parasitol. 97 (2003) 31–35.
 https://doi.org/10.1179/000349803125002742.
- [138] G. Augusto, R. Nalá, V. Casmo, A. Sabonete, L. Mapaco, J. Monteiro, Geographic
 distribution and prevalence of schistosomiasis and soil-transmitted helminths among
 schoolchildren in mozambique, Am. J. Trop. Med. Hyg. 81 (2009) 799–803.
 https://doi.org/10.4269/ajtmh.2009.08-0344.
- [139] A.R.P. Walker, L.A. Dini, B.F. Walker, J.A. Frean, Helminthiasis in African children in a relatively low risk region in south Africa: implications for treatment?, S Afr J Epidemiol Infect South. 15 (2000). https://www.cabdirect.org/cabdirect/abstract/20013078709
 (accessed October 22, 2021).
- [140] H. Sacolo-Gwebu, M. Chimbari, C. Kalinda, Prevalence and risk factors of
 schistosomiasis and soil-transmitted helminthiases among preschool aged children (1-5
 years) in rural KwaZulu-Natal, South Africa: A cross-sectional study, Infect. Dis. Poverty.
 8 (2019) 1–12. https://doi.org/10.1186/s40249-019-0561-5.

1063 1064 1065 1066	[141]	U.C. Braae, P. Magnussen, W. Harrison, B. Ndawi, F. Lekule, M.V. Johansen, Effect of National Schistosomiasis Control Programme on Taenia solium taeniosis and porcine cysticercosis in rural communities of Tanzania, Parasite Epidemiol. Control. 1 (2016) 245–251. https://doi.org/10.1016/j.parepi.2016.08.004.
1067 1068 1069 1070 1071	[142]	E. Buzigi, Prevalence of Intestinal Parasites , and its Association with Severe Acute Malnutrition Related Diarrhoea, J. Biol. Agric. Healthc. 5 (2015). https://www.researchgate.net/publication/271763982_Prevalence_of_Intestinal_Parasites_ and_its_Association_with_Severe_Acute_Malnutrition_Related_Diarrhoea (accessed October 18, 2021).
1072 1073 1074 1075 1076	[143]	S. Fuhrimann, M.S. Winkler, N.B. Kabatereine, E.M. Tukahebwa, A.A. Halage, E. Rutebemberwa, K. Medlicott, C. Schindler, J. Utzinger, G. Cissé, Risk of Intestinal Parasitic Infections in People with Different Exposures to Wastewater and Fecal Sludge in Kampala, Uganda: A Cross-Sectional Study, PLoS Negl. Trop. Dis. 10 (2016) e0004469. https://doi.org/10.1371/journal.pntd.0004469.
1077 1078 1079 1080	[144]	P. Oboth, Y. Gavamukulya, B.J. Barugahare, Prevalence and clinical outcomes of Plasmodium falciparum and intestinal parasitic infections among children in Kiryandongo refugee camp, mid-Western Uganda: A cross sectional study, BMC Infect. Dis. 19 (2019) 1–8. https://doi.org/10.1186/s12879-019-3939-x.
1081 1082 1083 1084	[145]	K. Modjarrad, I. Zulu, D.T. Redden, L. Njobvu, D.O. Freedman, S.H. Vermund, Prevalence and predictors of intestinal helminth infections among human immunodeficiency virus type 1-infected adults in an urban African setting, Am. J. Trop. Med. Hyg. 73 (2005) 777–782. https://doi.org/10.4269/ajtmh.2005.73.777.
1085 1086 1087	[146]	J. Siwila, I.G.K. Phiri, H.L. Enemark, M. Nchito, A. Olsen, Intestinal helminths and protozoa in children in pre-schools in Kafue district, Zambia, Trans. R. Soc. Trop. Med. Hyg. 104 (2010) 122–128. https://doi.org/10.1016/j.trstmh.2009.07.024.
1088		
1089		
1090		
1091		
1092		
1093		