

Exploring the Efficacy and Feasibility of using Biomarkers for Early Screening of Lung Cancer and their Potential Reverberation on Mortality Rate: A Review

Safari Vejin, T; OLeary, J; Singh, L; Sullivan, T; Aziague, S; Akbariansaravi, A
American University of Antigua College of Medicine, Coolidge, Antigua and Barbuda
New York Medical College School of Medicine, Valhalla, New York

Conflict of Interest: The authors have no competing interests

Funding: No funding was provided

Keywords: Lung cancer, Biomarkers, Standardizations, Screening methods, Prognosis

Abstract

Introduction

Lung cancer patients often have a poor prognosis, due to various factors that play into the diagnosis, such as timeframe, early detection, and progression. The rate of patients with advanced-stage lung cancer is often high due to the inability to be diagnosed early. The lack of screening, therefore, increases the mortality rate of lung cancer. High mortality rate is also attributed to advanced lung cancer cases that are undiagnosed due to the lack of symptoms in patients early in disease acquisition.

Methods

This review explores lung cancer screening methods adopted around the world and those with the most promising standardizations. Criteria used to filter articles included lung cancer methods using biomarkers, research conducted in the United States within the past 10 years, screening methods observed in a clinical setting, and patients with primary lung cancer. Some practices that can be utilized to improve screening for lung cancer may involve using genetic markers to identify patients possessing genetic makeup associated with lung cancer.

Results

Our study included 32 articles in our review calculating the sensitivity and specificity for a variety of potential biomarkers including: 1) Volatile organic compounds has a specificity of 80% and sensitivity of 80% 2) Circulating tumor DNA has a sensitivity of 80% and specificity of 96% 3) miRNA molecules (miR-146-5p, miR-324-5p, miR-223-3p, and miR-223-5p) displayed sensitivities of 93%, 94%, 96%, and 95% respectively, with specificities of 33%, 23%, 27%, and 26% respectively 4) methylation beads specific genomic sequences (SOX17, TAC1, HOXA7, CDO1, HOXA9, and ZFP42) with sensitivities of 84%, 86%, 63%, 78%, 93%, and 87% respectively when found in sputum and 73%, 76%, 34%, 65%, 86%, and 84% respectively when found in serum. Specificities were calculated at 88%, 75%, 92%, 67%, 8%, and 63% respectively when found in sputum, and 84%, 78%, 92%, 74%, 46%, and 54% respectively in serum 5) Ga-Alfatide II scans were found to have a sensitivity of 86% and a specificity of 85% 6) Combined score had a sensitivity of 78% and specificity of 50% 7) WT-IDH had a sensitivity of 63% and specificity of 93%.

Discussion

In such a high-risk patient population, cautious preventative measures can be used such as lifestyle choices and environmental exposures that are known risk factors. Informed physicians will be able to provide care directed toward minimizing patient's modifiable risk factors to lower incidence rate and routine imaging offered when there is an index of suspicion to minimize mortality rate in high-risk populations. This review highlighted practices such as circulating tumor DNA and a Ga-Alfatide II CT/PET scan that can be utilized to improve screening in the future and further dissected errors made in lung cancer diagnosis.

Introduction

There have been many advances regarding the diagnosis and treatment of lung cancer. Despite these changes, the progress in the field of primary respiratory neoplasm remains incomparable to the constant improvements observed in other oncological fields, such as OB/GYN and Gastroenterology. Considering the lack of advancements, the number of patients affected by lung cancer remains unchanged and has been so for the past decade, suggesting little improvement in medical developments for lung cancer. Since 1987, lung cancer remains the leading cause of death in both men and women. The alarming rate of lung cancer deaths has also prompted findings exploring methods that could be used in lowering lung cancer death rates, as well as disease prevention. The death toll for lung cancer peaked in 2005 and decreased by a mere 10,000 by 2016, further emphasizing the minimal rate of improvement over a span of 10 years.¹ The continued incline of disease progression coupled with the lack of improvement has caused a rise in lung cancer mortality. In 2018, the incidence of lung cancer worldwide was 2.1 million new cases per year.² In addition, it was calculated that 1.8 million people will succumb to their lung cancer diagnosis that same year.³ The vital question posed in the data addresses the comparable disparity between the high mortality rate and low incidence rate of lung cancer.

Improvement can be seen in cases of lung cancer by targeting three vital entities involved with chronic conditions—primary, secondary, and tertiary prevention. Primary prevention methods include behaviors involved in preventing the condition from developing, as well as environmental and genetic contributing risk factors. Furthermore, primary prevention acts as the first shield in disease deterrence by exploring ways by which lung cancer incidence can be lowered. Failure of primary prevention leads to the adoption of secondary prevention, which aids in screening with the inclusion of symptom management. Tertiary prevention is utilized in advanced disease states and explores treatment methods, including their success rates. Of the three prevention methods, primary prevention remains understudied in data and literature. By employing these prevention techniques, early detection of lung cancer can be further improved and better studied. Affordability plays an important role in early detection, including clinical method standardization, as it could increase the probability of early cancer detection during general checkups with primary care physicians.

Disease burden is higher in the African American population compared to other races, indicating an immense racial disparity concerning lung cancer incidence. Lung cancer incidence is 30% higher in African American men compared to Caucasian men; although, the overall smoke exposure and risk factors of the former were lower.⁴ A similar pattern is also observed for women; however, incidence has been reported to be the same amongst African American women and Caucasian women, with lower exposure observed in African American women.⁵ Regarding gender differences, men have been reported to have a higher incidence rate for lung cancer, whereas women are reported to have a higher prevalence.⁵ Incidence rates exploring age demographics have been lowered from 42-years-old to 36-years-old for diagnosis.⁵

High mortality rate can be mainly attributed to squamous small cell and adenocarcinoma.³ Small cell lung carcinoma (SCLC), also referred to as oat cell carcinoma makes up 20% of respiratory cancers.³ SCLC consists of a gray-white central mass containing small round blue (polygonal) cells in clusters. Being the most aggressive respiratory cancer due to its rapid growth and early dissemination, SCLC depicts a poor prognosis. In addition, SCLC cells are notorious for their neurosecretory granules that give rise to a myriad of paraneoplastic syndromes, including Syndrome of Inappropriate Antidiuretic Hormone (SIADH), Horner's Syndrome, Lambert Eaton Syndrome, and Cushing's Syndrome. Exceptions concerning paraneoplastic syndromes associated with SCLC include hypercalcemia and PTrH secretion, which are involved in Squamous Cell Carcinoma (SCC), as well as high HCG levels seen in LCCs. LCCs present with large anaplastic cells that are poorly differentiated. Furthermore, the secretion of HCG in LCCs leads to symptoms of gynecomastia in both men and women, in addition to a false positive pregnancy test in either gender.⁴

SCLC cells are of neurosecretory origin and function as neuroendocrine tumors. They also stain positive with Synaptophysin, Chromogranin, and Bombesin stains. SCLC is associated with the L-myc gene mutation, which is a basic helix-loop-helix transcription factor that is amplified and overexpressed in SCLCs.³ Current chemotherapy treatment of SCLCs involves the use of monoclonal antibodies which target the myc homolog sequence box and

prevent its expression.⁴ The poor prognosis in these cancers, including recently discovered gene mutations observed in adenocarcinoma are being targeted by biomarker chemotherapy techniques.

Adenocarcinomas comprise 35% of all respiratory cancers.³ Contrary to other respiratory carcinomas, there has been a stronger correlation observed with non-smokers with a higher prevalence noted in women compared to men. Regarding frequency, non-smokers have a 25% incidence rate worldwide generally with all cancer types.⁵ Adenocarcinomas present as a gray-white peripheral mass organized in a lepidic pattern in the lungs. Furthermore, adenocarcinomas consist of mucous-secreting glands accompanied by pleural puckering with a tendency to form scarring. These types of scarring are also referred to as scar carcinomas. Primary adenocarcinomas have a better prognosis compared to their lung tumor counterparts, presenting with the best survival rate strongly dependent on early surgical resection of the tumor.

Specific gene mutations, such as P53 and retinoblastoma have been strongly associated with lung cancers overall.^{1,2,4} Adenocarcinomas have been associated with numerous gain-of-function gene mutations, involving tyrosine kinase receptors, including EGF, ALK, ROS, MET, and RET.^{1,2,4} These carcinomas were previously identified to involve the K-ras mutation; however, with more current studies, roughly 10% have been identified to have EGFR mutation and 5%, the EML4-ALK fusion protein mutation.^{1,2,4} ALK is a gene that codes for a receptor tyrosine kinase activated by the EML gene. Upon fusion of the ALK and EML genes, the receptor tyrosine kinase acquires uninhibited activity, which creates the basis for the formation of peripheral mass adenocarcinoma. Despite association with multiple gene mutations, the EGFR and EML4-ALK fusion mutations have been targeted by specific small-molecule inhibitors used as chemotherapy treatment.^{1,2,4}

Enhanced primary prevention routes:	Enhanced secondary prevention routes:	Enhanced tertiary prevention routes:
Identify patient genetically disposed to lung cancer	Allow providers to actively monitor high-risk patients before cancer progression	Effective chemotherapy options based on lung cancer gene type
Screen patients with Modifiable risk factors (smoking, urban homes, exposure, etc.) for genetic disposition	Patients with risk factors and genetically disposed screened with CT more regularly	counseling genetically disposed patients on modifiable risk factors for high-risk patients
counseling genetically disposed patients on modifiable risk factors for high-risk patients	Earlier detection	Further our scientific insight into genetics of lung cancer and their types
	Decreases prevalence	Insight into epigenetics of cancer allows possibility to restore normal tissue (i.e. reactivation of tumor suppressor genes)
	Decreased mortality	
	Increased life expectancy	
	Increases survival rate of lung cancer	

Figure 1: Table of utilities for epigenetics in lung cancer screening and treatment

Peripheral Non-small cell carcinomas (NSCCs) with EGFR mutations have shown promising results when treated with Erlotinib, Gefitinib, and Crizotinib.^{3,4,5} These chemotherapy treatments target inhibition of angiogenesis and epidermal growth factor activity in tumor cells, as well as inhibited tyrosine kinase activity. In addition, the adenocarcinomas that contain the EML4-ALK mutation have been targeted by Crizotinib as chemotherapy.^{3,4,5} Crizotinib is an anaplastic lymphoma kinase (ALK) and C-ros oncogene 1 (ROS1) inhibitor that targets inhibition of ALK phosphorylation and signal transduction of the NSCC tumor cells, inhibiting their growth at the G1-S cell cycle phase.^{3,4,5} Adenocarcinomas have shown a high preference for metastasis to the lung, with far more occurrence of metastatic adenocarcinomas involving the lung than primary adenocarcinomas.^{3,4,5}

Squamous cell carcinoma and small cell carcinoma are strongly associated with smoking with a higher prevalence in males. Compared to adenocarcinoma or large cell carcinoma, which targets the periphery of the lungs, both squamous and small cell carcinomas are central masses. Squamous cell carcinoma accounts for 30% of all lung cancer incidents, with a higher prevalence in males and a strong correlation to smoking history.¹ It originates from the bronchial epithelium after progressively evolving from metaplasia to dysplasia and ultimately becoming its invasive counterpart, carcinoma in situ.¹

Bronchogenic carcinomas comprise 5% of all lung cancers and originate from terminal bronchioles.^{1,2,4} These carcinomas further develop as columnar tumor cells that proliferate along the alveolar walls, forming endobronchial lesions. They form gray-white mucinous nodules in the periphery of the lungs. Bronchogenic carcinomas have been identified as the main contributors to mortality amongst male smokers aged 50-80 and have rapidly become the main cause of death amongst female smokers in recent years.^{1,2,4} The risk factors associated with bronchogenic carcinomas include cigarette smoking, occupational exposure to toxins, such as asbestos & uranium mining, radiation, and air pollution. The presence of pre-existing chronic fibrotic lung disease further worsens the symptoms of bronchogenic carcinomas. Chronic fibrotic lung disease that has been associated with the onset of bronchogenic carcinomas include idiopathic pulmonary fibrosis, non-specific interstitial pneumonia, cryptogenic organizing pneumonia, various pneumoconiosis, and collagen vascular diseases such as Systemic Lupus Erythematosus (SLE) and Rheumatoid Arthritis (RA).^{1,2,4}

The current golden standard for lung cancer screening involves the administration of low-dose lung CT to patients between the ages of 50-80.⁷ This screening method can be inaccurate due to risks associated, which include false positives & overdiagnosis, which could limit chances of early detection.⁶ The current protocol for the treatment of non-small cell carcinomas is surgery. On the other hand, chemotherapy is the best option in the case of unresectable tumors, such as small cell carcinomas. Despite chemotherapy treatment for patients with small cell carcinomas, the prognosis is extremely poor and little improvement has been shown in the past decade.⁶ That being said, roughly 10% of patients have a 5-year survival rate.⁷ There has been a plateau in the advancements of screening and treatment of lung cancer compared to other oncological areas. The main topic of importance in this paper highlights the importance of using biomarkers that are targeted for a specific gene mutation, which either predisposes or plays a role in the development of the different kinds of lung cancers.

Methodology

1.1 Method

The initial search on PubMed using the keywords mentioned in the methodology yielded 32 articles. These articles were studies conducted over a span of 5 years. Due to the narrow timeframe, an additional search was performed to include articles published within the past 10 years. By broadening the search, 32 additional articles were discovered. A total of 61 articles published within the past 10 years on PubMed matched the criteria.

1.2 Criteria

The literature search on lung cancer screening involved the main scholarly platforms utilizing the same keywords across all databases. The keyword used across all platforms was “biomarkers for early diagnosis of lung cancer”. This allowed for the extraction of many articles (n =2,109), however we were not selective for any specific biomarkers or subtypes of lung cancer. The articles extracted were then filtered concisely using the following criteria: 1) the article was centered around clinical research 2) the research was conducted within the last 10 years 3) the research was completed within the United States 4) the article compared the amount of research conducted in the United States versus other countries 5) the article included the exploration of future directions of the study 6) the article involved analysis of international research trials and reviews 7) the article permitted assessment of risk factors across countries and cultures. A final record of 49 research articles fit the final definitive criteria and were included in our review. The search was executed exclusively on PubMed.

This review encompassed study criteria involved in the analysis of biomarker screening methods for patients with a variety of cancers, including lung cancer. Applicable data for each study regarding lung cancer was carefully organized throughout the review. The table below gives a good summary of the data collected during the literature review.

Table 1: List of standardized information that was collected for all the articles meeting the criteria

Stanford information of the article; Name and authors, year published
Early screening method used
Number of participants
Type of lung cancer studied (include breakdown of the lung cancer types; adenocarcinoma, squamous cell carcinoma and small cell carcinoma of the lungs)
Biomarkers used
Number of patients detected using the screening method
Breakdown of demographic distribution of patient population used in that study; ethnicity, age, state, rural vs. urban residence, exposure to known lung cancer carcinogens (radon, smoke, pollution, asbestos, prev. tuberculosis infections)
Associations with paraneoplastic syndromes and other possible early manifestations
Concluded statistics calculated for the study: sensitivity and specificity for group cancers, PPV
Was a biomarker shown to be more effective towards a certain type of lung cancer over others (included for studies including patients with all 3 main types of lung cancer)
Additional notes/notable findings
Future direction suggested in the paper

2.3 Defining characteristics across articles

We evaluated successful biomarkers pertinent to lung cancer screening across the articles reviewed. To compare the efficacy of biomarkers used in the early detection of lung cancer, Positive Predictive Value (PPV) and Negative Predictive Value (NPV) were considered. Patients' demographics in addition to risk factors pertinent to specific lung cancer types were further evaluated as part of representative data. The collection of the aforementioned data was correlated to various routes of lung cancer prevention with consideration to biomarker screening techniques. PPV, sensitivity, and specificity were taken as a collective average from all articles and represented in this study. Articles that did not contain these statistics were excluded during calculations of the average.

2.4 Future approach

Further analysis included identifying the lack of data and inconsistencies across the journals.³ In addition, data represented in Table 1 were visually displayed with highlights of significant findings. The visual included categories such as demographics, risk factors, lung cancer types, and biomarkers. The future direction of the paper was obtained based on data findings and possible improvements in the clinical implementation of biomarkers. Additionally, the lack of important data in lung cancer literature, involving screening methods are valuable considerations that should be analyzed in future studies and journals.

Results

1.3 Collection of identifying variables comparing various types of lung cancers

The following are observations summarized in Table 2 with calculations for sensitivity and specificity summarized in Table 3. Research published between 2011 and 2020 demonstrates the efficacy of biomarkers by comparing types of lung cancer, mainly adenocarcinoma, and squamous cell carcinoma. Data was collected highlighting screening methods and improvements in biomarkers with consideration to study participants' age and smoking history. Screening methods mainly included chest x-ray, PET/CT scan, blood samples, sputum samples. Most of the studies between the years observed showed that biomarkers enhanced the sensitivity and specificity greatly in elderly patients that had a history of smoking.

In addition, the cohort study calculations in the screening for isocitrate dehydrogenase (IDH) values are summarized in Table 4. The data demonstrates the efficacy of screening for IDH in serum by comparing a training cohort with a validation cohort using a total of 1166 participants in the diagnosis of non-small cell lung cancer. Finally, demographics based on gender and age are also displayed with the latter being summarized by median, interquartile range, and range of the study population.

Article Authors	Year published	Early screening methods	Number of participants	Type of lung cancer studied	Biomarker used	number detected	Age breakdown of the participants	Smoking history	Specific efficacy of biomarker used	Future directions suggested by the article	
Gerardine O'Sullivan Coyne	2020 Apr 20	Cell Search analysis of CTC p16 expression in patient blood specimens. pharmacodynamic evidence for DNMT inhibition by FdCyd, method for detecting expression of tumor suppressor protein p16/INK4A in circulating tumor cells (CTCs).	83	NSCLC (24)	P16-expressing cytokeratin-positive CTCs (P16/Cy5 target engagement). Targeting EPCAM or CD146-positive, and cells that express MUC1, PD biomarker/MUC1-double-positive cells.	89% of patients evaluable for clinical and CTC response	Median: 66 YO (50-84)	N/A	In NSCLC, histologic subtype variations in the hypermethylation of p16 and other tumor suppressor genes [52] may account for lack of FdCyd clinical activity.	Further study of FdCyd + THU is potentially not warranted in NSCLC.	
Wenwen Li, Wei Dai, Mingxin Liu,	2019 August 9	Bio-VOC breath sampler and Tedlar bag was used to collect breath samples. Gas chromatography-mass spectrometry coupled with solid phase microextraction technique was used to analyze VOCs in exhaled breath.	389 for lung cancer group and 389 for the control group	N/A	Exhaled volatile organic compound (VOC) biomarkers	Detected >70%	(50-74)	No exposure	N/A	N/A	
Mayun Peng, Yuancai Xia, #2 Xiaohua Li,	2019 Apr 13	Lung nodule screening imagery, gene sequencing	192	Adenocarcinoma: 100 Squamous carcinoma: 28 Small cell lung cancer: 1 Others: 7	Cell-free circulating tumour (ct) DNA (ctDNA)	Assay detected 63%, 83%, 94% and 100%, for stages I, II, III and IV, respectively.	Median Age 56.5 (26-79)	Yes (66 total)	The combined model of ctDNA mutations and serum biomarkers improved the sensitivity and specificity to 89% and 99%, respectively.	N/A	
Rajiv Raja, Michael Kuziora, Philip Z. Brosevan	2018 December	Somatic mutations detected in circulating tumor DNA (ctDNA) from plasma. Variants in plasma ctDNA were assessed in samples collected at pre-treatment and 6 weeks after first dose of treatment with durvalumab.	28	Non-squamous 10 Squamous 18	ctDNA (fraction of ctDNA that is derived from tumor tissues) to predict survival on durvalumab	96% of patient	Median 62.21	Yes (22)	N/A	Supports the use of ctDNA dVAF to monitor anti-tumor activity and clinical response to immunotherapies.	
Ana B Pavet, American Association for Cancer Research	2017 November	Epithelial brushings from the mainstem bronchus of patients undergoing bronchoscopy for suspected lung cancer	341 discovery set (138 samples) and the test set (203 samples)	Discovery set test set: NSCLC (72,79) Adenocarcinoma (31,34) Squamous (27,25) Large cell (2,4) SCLC (16,21)	miR-146a-5p, miR-324-5p, miR-223-3p, miR-223-5p	Discovery set 42 microRNAs that showed differential expression	Median 59 YO	Discovery set test set: Current (46,86) Former (92,115)	Four miRNA isoforms (miR-146a-5p, miR-324-5p, miR-223-3p, and miR-223-5p) that have altered expression in the airway epithelium of patients with lung cancer.	N/A	
Alicia Hulbert, Ignacio Jusua-Torres, Alejandro Stark	2016 Oct 11	Promoter methylation using quantitative methylation-specific real-time PCR and Methylation on Beads for cancer-specific genes was detected.	651	adenocarcinoma. Squamous cell carcinoma. Adenoquamous	Methylation on Beads for cancer-specific genes (SOX17, TAC1, HMXA7, CD01, HMOXA9, and ZFP42).	Sputum sample: 91% Plasma sample: 85%	Median: cancer: 68 YO (62-75) Control: 63 (55-73)	Current (27) former (87)	This study shows that high sensitivity and specificity detection of early stage NSCLC	This panel could be used as an adjunct to CT screening, identifying patients at high risk for lung cancer, reducing false-negative results.	
Fai Kang, Shengjun Wang, Feng Tian	2016 May	Diagnostic potential of Ga-Alfide II with F-FDG in differentiating between non-small cell lung cancer patients (NSCLC) and lung tuberculosis (TB) patients	34 total: 21NSCLC, 13 TB patients	NSCLC, Adenocarcinoma, Squamous carcinoma, Tuberculosis	The expression of the angiogenesis marker vEGF in NSCLC and TB primary lesions was analyzed by immunohistochemistry.	Ga-Alfide II PET/CT scanning detected 23 visually positive LN from 6 patients. These 6 patients were all true-positives. However, 68Ga-Alfide II failed to identify 2 8a stage-positive patients. 18F-FDG detected 34 LNs from 10 patients. Seven of them were true-positives, and 3 were false-positives.	Mean Age 42.4 ± 15.8	N/A	18F-FDG uptake was intense in both NSCLC and TB primary lesions, whereas the 68Ga-Alfide II uptake in most of the TB lesions was much lower than that in NSCLC lesions.	N/A	
Taofeek K Owonikoko, Sunesh S Ramalingam, Daniel L Miller,	2015 April 15	Patients were eligible if newly diagnosed with NSCLC of all histologies and deemed to be surgically resectable stage I-IIIa disease. Other eligibility requirements included age ≥18 years, ECOG performance status of 0-2, adequate bone marrow function, renal function, hepatic function, obtained baseline tumor biopsy and FDG-PET/CT imaging followed by everolimus treatment (5 or 10 mg daily, up to 28 days), or without intervening treatment for controls	33 eligible patients with resectable lung cancer to the control (16 patients) or treatment (everolimus—5 mg daily in 12 patients and 10 mg daily in 11 patients) arms	Adenocarcinoma	Expression of active phosphorylated forms of mTOR, AK1, S6, p70S6K, 4EBP1, and total Bim protein between pretreatment and posttreatment. Issue samples to target modulation by everolimus	Patients treated with everolimus 5 mg and 10 mg had a mean reduction of 21% and 24%, respectively, in comparison with a mean increase of 29% in control patients.	Median age 64 years	Yes [specified in 1 patient]	22 tumors (67%) were adenocarcinomas	This "window-of-opportunity" study demonstrated measurable, dose-dependent, biologic, metabolic, and anti-tumor activity of everolimus in early-stage NSCLC.	
Ziv Sevilija 1, Yael Leitner-Dagan, Mila Pinchev,	2014 April 7	lung cancer risk of an integrated DNA repair score, measured by a panel of three enzymatic DNA repair activities in peripheral blood mononuclear cells.	96 lung cancer, 96 control	Adenocarcinoma (30) and squamous cell carcinoma (45)	AP endonuclease 1 (APE1), 8-oxoguanine DNA glycosylase (OGG1), and methylpurine DNA glycosylase (MPG), all of which repair oxidative DNA damage as part of the base excision repair pathways.	No. of control subjects No. of case patients Cross-validation Highest tertile 30 12 Middle tertile 33 19 Lowest tertile 33 65	Mean: 67 YO	Smoking Control vs Case [Former smoker 27 vs 36] [Current smoker 22 vs 36]	There was no difference in APE1 activity between adenocarcinoma and squamous cell carcinoma	N/A	
Gerald LI, Martial Guillaud, Jean LaRiche,	2012	High risk smokers defined by age, smoking, and/or undergoing investigation for suspected lung cancer. Sputum samples from high-risk patients enrolled in ongoing chemoprevention trials were subjected to automated quantitative image cytometry after Faupein-tin staining	High risk smokers defined by age, smoking, and/or undergoing investigation for suspected lung cancer	Intraepithelial neoplasias in the lung	Combined Score (CS), uses automated image cytometric analysis of ploidy and nuclear morphology to detect subtle intraepithelial changes that often precede lung tumors.	CS detected 78% of all highest-risk subjects— those with CS or worse plus those with moderate or severe dysplasia and abnormal nuclear morphology.	Average: 59.7	62.2% from former smokers, 38.5% current smokers.	N/A	CS is a powerful yet minimally invasive tool for rapid and inexpensive risk assessment for the presence of precancerous lung lesions, enabling enrichment of chemoprevention trials with highest-risk dysplasias. CS correlates with other biomarkers, so CS may find use as a surrogate biomarker for patient assessment and as an endpoint in chemoprevention clinical trials.	
Yong Wang, Dai Zheng, Qulin Tan,	2011 September 4	Blood samples were obtained from lung cancer patients and healthy volunteers	12 total (6 lung cancer patient and 6 healthy volunteers)	Small cell lung cancer, squamous lung carcinoma, non-small cell carcinoma	Detected plasma miR-155 in lung cancer patients using the nanopore sensor (based on alpha-hemolysin probe).	In the absence of P155, only short blocks were observed as a translocation of single-stranded oligonucleotides. In the presence of the probe, we identified distinct short and long blocks in both the control and lung cancer groups. All of the P155 lesions in the lung cancer group were higher than those in the control group.	N/A	N/A	N/A	This could prove useful for quantitative microRNA detection, biomarker discovery, and the non-invasive early diagnosis of cancer	
Meredith B Shiels, Anil K Chaturvedi, Hormuza A Keki	2011 Aug 9	Lung cancer screening included a chest x-ray (CXR) at baseline and yearly follow-up intervals.	532 lung cancer, 582 controls, 150 additional control with CXR	Adenocarcinomas (n = 189) squamous cell carcinomas (n = 114) 68 large-cell carcinomas 30	SP-D KL-6 e-selectin	Matched controls Lung cancer cases	SP-D, ng/ml 16.5-71.7 97 146 71.8-165.3 120 145 105.4-143.9 122 146 143.9-615.5 193 145 KL-6, µg/ml 178.8-231.0 98 145 231.1-325.7 113 146 325.8-465.7 160 145 465.8-4225.9 161 146	55-74	Lung cancer, control: Current (208, 206) Former (291, 267)	Associations between the serum markers and specific histological types of lung cancer were statistically significant for adenocarcinoma and squamous cell carcinoma	N/A

Table 2: Data collected from each article according to SC vs. NSC lung cancer and its subtypes

Biomarker	Sensitivity	Specificity
Exhaled volatile organic compound (VOC) biomarkers	80%	80%
Cell-free circulating tumour (ct) DNA (ctDNA)	80%	96%
miR-146a-5p, miR-324-5p, miR-223-3p, miR-223-5p.	mRNA biomarker (0.94), mRNA biomarker + miR-146a-5p (0.93), mRNA biomarker + miR-324-5p (0.94), mRNA biomarker + miR-223-3p (0.96), mRNA biomarker + miR-223-5p (0.95)	mRNA biomarker (0.27), mRNA biomarker + miR-146a-5p (0.33), mRNA biomarker + miR-324-5p (0.23), mRNA biomarker + miR-223-3p (0.27), mRNA biomarker + miR-223-5p (0.26)
Methylation on Beads for cancer-specific genes (SOX17, TAC1, HOXA7, CDO1, HOXA9, and ZFP42).	Sensitivity in sputum: SOX17(84), TAC1(86), HOXA7(63), CDO1(78), HOXA9(93), ZFP42(87). Sensitivity in plasma: SOX17(73), TAC1(78), HOXA7(34), CDO1(65), HOXA9(86), ZFP42(84)	Specificity in sputum: SOX17(88), TAC1(75), HOXA7(92), CDO1(67), HOXA9(8), ZFP42(63). Specificity in plasma: SOX17(84), TAC1(78), HOXA7(92), CDO1(74), HOXA9(46), ZFP42(54)
Ga-Alfatide II	85.71%	84.62%
Combined Score (CS)	78%	50%

Table 3: Table of useful biomarkers found in systematic review for potential for detection & therapy of lung cancer

	AUC (95% CI)	Se%	Sp%	PPV%	NPV%	Positive LR	Negative LR
Training cohort							
NSCLC vs HC, BPC and OC	0.870 (0.840-0.899)	63.3	86.8	87.1	62.7	4.80	0.42
NSCLC vs HC	0.915 (0.887-0.942)	63.3	92.9	95.4	52.0	8.91	0.40
Stage 0-IA NSCLC vs HC, BPC and OC	0.859 (0.826-0.892)	58.6	86.8	79.3	70.9	4.44	0.48
Stage 0-IA NSCLC vs HC	0.907 (0.875-0.938)	58.6	92.9	92.2	61.0	8.25	0.45
Stage IB-IV NSCLC vs HC, BPC and OC	0.886 (0.852-0.919)	70.7	86.8	74.4	84.5	5.37	0.34
Stage IB-IV NSCLC vs HC	0.927 (0.896-0.958)	70.7	92.9	90.0	77.8	9.96	0.32
Validation cohort							
NSCLC vs HC, BPC and OC	0.745 (0.704-0.786)	55.0	86.3	82.8	61.5	4.01	0.52
NSCLC vs HC	0.730 (0.684-0.776)	55.0	89.3	92.7	44.6	5.12	0.50
Stage 0-IA NSCLC vs HC, BPC and OC	0.797 (0.730-0.865)	59.1	86.3	53.4	88.8	4.31	0.47
Stage 0-IA NSCLC vs HC	0.788 (0.711-0.865)	59.1	89.3	75.0	80.0	5.50	0.46
Stage IB-IV NSCLC vs HC, BPC and OC	0.746 (0.697-0.796)	54.4	86.3	74.2	72.3	3.97	0.53
Stage IB-IV NSCLC vs HC	0.732 (0.676-0.788)	54.4	89.3	88.3	56.8	5.07	0.51

AUC, area under the receiver operating characteristic curve; BPC, benign pulmonary condition; HC, healthy control; LR, likelihood ratio; NPV, negative predictive value; OC, other cancer; PPV, positive predictive value; Se, Sensitivity; Sp, Specificity.

	Training cohort (620 cases)			Validation cohort (546 cases)		
	NSCLC patients	HCs	Disease controls (patients with BPCs or OCs)	NSCLC patients	HCs	Disease controls (patients with BPCs or OCs)
No.	362	155	103	298	121	127
Age, y						
Median	59.0	37.0	53.0	61.0	34.0	59.0
(Q1, Q3)	(53, 65)	(31, 46)	(45, 64)	(54, 68)	(27, 45)	(50, 67)
Range	35-82	23-67	26-84	26-85	18-72	17-86
Sex						
Female/ male	201/161	87/68	49/54	120/178	64/57	47/80

Tables 4a and 4b: Utilization of isocitrate dehydrogenase 1 as a serum protein biomarker for the early detection of non-small-cell lung cancer: A multicenter in vitro diagnostic clinical trial⁷

Discussion

Early detection remains one of the best ways to reduce lung cancer deaths. By detecting lung cancer in its early stages and resecting it, patients will be able to have a better prognosis. Breast and Prostate cancers are the most prevalent, but not the deadliest. Inversely, lung cancer is less common with the highest rates of death. Lung cancer remains the leading cause of cancer deaths, mainly due to underdeveloped screening mechanisms. This is attributed to the high association of smoking with lung cancer. It is important for medical caregivers to target smoking cessation in patients to further reduce the risk of developing lung cancer.

In contrast to lung cancer, there are various screening methods available for other cancer types. Screening allows for early detection, which reduces mortality rates. For example, breast cancer screening includes mammograms and physical examinations. Prostate cancer screening methods involve Prostate-Specific Antigen (PSA) and Digital Rectal Examination (DRE). Colorectal cancer screening includes hemocult testing of the stool and colonoscopy. These cancers have higher incidences compared to lung cancer, but their screening techniques led to a lower mortality rate. Lung cancer survival rates remain at 15% when comparing data from roughly 30 years ago and data from the past 5 years. Developments in screening methods for lung cancer will improve survival rates and a variety have been analyzed during our investigation.

Volatile organic compounds (VOC) can be measured using exhaled breath analysis making the procedure completely non-invasive, giving it great potential as a screening method for lung cancer²³. Our study found that currently VOC exhaled breath analysis has both a sensitivity and specificity of 80%, which is on the border of being a viable screening test. However, there are issues with using VOC as a potential biomarker. For instance, there are no unified VOC biomarkers for lung cancer²³, thus further investigation must be conducted to determine the exact VOC biomarkers found in lung cancer. Such investigations will raise the sensitivity and specificity and could potentially make VOC a gold standard for lung cancer screening.

Combined score (CS) analysis refers to the correlation between several lung cancer risk factors including histopathological grade, age, smoking status, and p53 and Ki67 immunostaining²¹. CS is an incredibly powerful tool having the benefit of being minimally invasive and accounting for multiple biomarkers associated with lung cancer. However, our study found that CS has a sensitivity of 78% and specificity of 50%, showing that CS requires further investigation into the risk factors the test will analyze. With a specificity and sensitivity that low, CS is currently not a candidate as a screening method, however there is much potential. Should further investigation into CS alter the risk factors to more accurate and precise variables, the sensitivity and specificity would increase and its possible CS could become a viable screening technique

miRNAs are a type of nucleic acid represented as small, non-coding RNA molecules binding to their complementary targets of 3'UTR segments in RNA; repressing gene expression and protein translation. However, unlike mRNA they are much more stable molecules, and thus much more easily measured in degraded tissue such as those caused by smoking³⁰. Our study identified four miRNA isoforms (miR-146a, miR-324-5p, miR-223-3p, and miR-223-5p) which have altered expression in epithelial airways of lung cancer. As compared with normal tissue, all four miRNA molecules have down-regulated expression and in addition, were found to be implicated in tumor suppressive pathways³⁰. Our study found that the down regulation is significant enough for the screening test for each of these four miRNAs to each have a sensitivity of 93% or higher, an indication of a valid screening test. However, each of the four miRNAs listed above has a specificity of 33% or lower, which could be attributed to their role in tumor suppression. While these molecules are excellent at detecting cancer, they are terrible at preventing false-positives, thus using miRNA molecules as a potential screening test is not possible at this time.

Isocitrate dehydrogenase I (IDH) is a catalytic enzyme found in the TCA cycle found in the cytoplasm and peroxisomes, catalyzing the oxidative decarboxylation of isocitrate to alpha-ketoglutarate. IDH1 is an extremely important enzyme as it produces NADPH in the cytoplasm and has been getting widespread attention due to the Arg-132 mutation⁷. A mutation that causes the generation of the metabolite 2-hydroxyglutarate leading to a decrease in the original product.

The metabolite will suppress activity of the enzyme prolyl hydroxylase and prevents degradation of hypoxia inducing factor resulting in 70% of grade II/III gliomas, 8.5% of acute myeloid leukemia, as well as a few other types of tumors⁷. While there has been no known association with lung cancer, prometrix discovery assays identified that WT IDH1 is overexpressed in non-small cell lung cancer and that IDH1 can promote the proliferation of this cancer line⁷. However, our study found that the sensitivity is not above 80% and the specificity is not above 70%, thus WT IDH1 is a poor screening method due to the large proportion of false-negatives and false-positives

Genome methylation is a potential screening technique as the methyl group stabilizes the gene and thus potentially increases sensitivity¹⁸. In addition, our study found six genes (SOX17, TAC1 HOXA7, CD01, HOXA9, ZFP42) that were identified as having a high frequency of altered DNA methylation in lung squamous and adenocarcinoma, but not in normal lung tissue¹⁸. While a promising method of lung cancer screening, as the sample collection is from either serum or sputum and thus non-invasive, there are several issues that must be addressed. There are some patients with undetectable DNA methylation in either collection sample (however sputum samples were slightly higher)¹⁸. Unfortunately, our study found this effect to have an effect on sensitivity and specificity as neither of these genes had both of these values above 80%. While this technique remains promising, further investigation into DNA methylation in lung cancer will need to be conducted for detection of methylation beads on genes to be a viable lung cancer screening.

Circulating tumor DNA (ctDNA) is extremely specific for a patient's tumor due to somatic mutations in tumor DNA that can be identified from normal DNA³². The benefit of ctDNA is the reduced false-positivity that is encountered with other biomarkers, especially proteins, and is present in high quantities in plasma^{32,34}. After being shed in circulation by apoptotic and necrotic tumor cells, ctDNA has a half life of about 16 mins to a few hours with a patient's tumor burden determining the proportion of ctDNA found in serum. The usefulness of ctDNA analysis in patients with lung cancer and in patients with suspected lung cancer is broad and can be used for much more than a screening technique. ctDNA has been shown to be an effective tool for monitoring tumor burden and tumor response as well as providing a real time evaluation of metastatic disease³⁴. In addition, ctDNA can be used to predict the prognosis based on the specific gene mutation and the effectiveness of certain chemotherapy³⁴.

Because ctDNA has a sensitivity of 80%, a specificity of 96%, and is minimally invasive, our study found ctDNA could be used successfully as a screening technique throughout the population. As the biomarker is specific for genes only found in tumors, it can therefore reduce the chance of detecting a false-positive and provide a high likelihood of preventing a false negative. Further investigation into ctDNA would involve finding the specific segments of mutated genes in each form of lung cancer, and thus increase the sensitivity of ctDNA's use as a biomarker. However, ctDNA should also be further researched for its other uses as well. Investigations into ctDNA's predictive power of prognosis and the effectiveness of chemotherapies should thoroughly be researched as it has the potential to significantly decrease the morbidity and mortality of lung cancer.

Neovascular angiogenesis is a hallmark of cancer and abundant in neoplastic lesions. Thus, angiogenesis has an incredibly powerful potential to detect differentiation in a wide variety of cancers. Alfatide II is an upgraded version of the dimer RGD (Arg-Gly-Asp), specifically NOTA-E-(PEG₄-c(RGDfk)₂), which targets the integrin $\alpha_v\beta_3$; a neovascular marker²⁰. However, the reciprocal remains true. In diseases such as TB there is a sharp decrease in the micro-vessel density, making Aflatide II a marker for both angiogenesis and vessel obliteration²⁰. Thus, the ability of a ⁶⁸Ga-Alfatide II CT/PET scan to identify such a wide-variety of diseases allows for an incredibly high diagnostic potential. However, the scan is not without limitations. As the scan can only detect angiogenesis, metastasis to the lymph nodes, bone and liver will have an inferior sensitivity to the ¹⁸F-FDG scan; this does not apply to brain metastasis as Alfatide II is able to penetrate the blood brain barrier²⁰.

As the ⁶⁸Ga-Alfatide scan has a sensitivity and specificity of around 85% and is minimally invasive, our study has found that this test would be appropriate for those at high risk for lung cancer or those suspected of lung cancer. However, it would not be recommended throughout the population as this biomarker screen requires a CT/PET scan with a contrast material. Thus, the test is expensive and counter-indicated in all forms of kidney injury. Further investigation should not be into Alfatide II itself, but rather what can be screened for in angiogenesis. While the CT/PET scan would have the

greatest diagnostic spectrum of disease, a serum test would be ideal to lower cost and radiation exposure. Therefore, investigations into biomarkers for angiogenesis should be completed to find a biomarker synthesized only by tumor cells that can be found in serum, stimulating angiogenesis.

Paraneoplastic syndromes are a group of rare disorders triggered by an abnormal immune system response to cancerous tumors, referred to as neoplasms. This can be seen in some forms of lung cancers as well as other cancer types. The data obtained for this review did not contain research regarding biomarkers explaining this phenomenon. This is because it is only seen in late stages of cancer, while this review focused mainly on screening and prevention, which occur during developmental stages.

Additional future directions of this study would explore other journal platforms, such as google scholar or Scopus. This will allow a wider data size. In addition, the American Thoracic Society (ATS) journal was also a platform of interest that is yet to be explored for further findings on early screening methods for lung cancer. Other analytic points could also be used to further explore demographics by including smokers vs. non-smokers, and children vs. adults. Also, the use of immunotherapy treatments for lung cancers can be discussed.

Immunotherapy is currently a growing field that is scratching the surface of its potential. It is still being explored for oncology patients. The table below depicts current immunotherapies that have shown some forms of advancement in the case of lung cancer patients.⁶ An area of lung cancer immunotherapy treatment that requires further exploration involves comparing the efficacy between biomarkers successfully utilized in early detection and specific immunotherapy types within the same patient. This will better demonstrate response rates of immunotherapy in the treatment of Small Cell Lung Cancers (SCLCs) or Non-Small Cell Lung Cancers (NSCLCs).

Our study was limited by using articles that were only published within the United States and in the last 10 years, which constricted our sample size and the number of potential biomarkers that have been researched. We were also limited by the types of lung cancer each article chose to research as a biomarker could have been statistically significant in form of lung cancer that was not investigated. In addition, our study was limited by the possibility of false-positives and false-negatives in each of the articles we examined. Finally we are limited by any potential bias and limitations in each of the articles we examined.

Conclusion

The study measured ctDNA and Ga-Alfatide II PET/CT scans as a means of screening for lung cancer with appropriate statistical values for measurement and data purposes. These are two biomarkers with highest sensitivity and specificity, thus indicating the chance of false-positive or false-negative are low. While these screening techniques cannot confirm the diagnosis of lung cancer, they have the potential to be indicated in any patient suspected of lung cancer or at high risk for lung cancer. Either of these biomarkers as screening techniques can increase early detection, thus decreasing mortality, provide indications into effective and/or ineffective treatments, and provide a clearer indication into the true morbidity of lung cancer.

Immunotherapy and chemotherapy are the most promising in comparison to other therapeutic options. By implementing these treatments, lung cancer surveillance before its development into advanced stages can be further improved. Although therapeutic effects can be increased with these standardizations, challenges may still arise due to constant changes during tumor growth. The complexity of various lung tumors could reduce the efficacy of these options due to therapeutic limitations, tumor resistance, or developed immune system tolerance to therapy. To overcome these challenges, it is important to develop screening methods for lung cancer, conduct further studies that analyze therapeutic response rates, and explore combination treatment if possible. Primary care physicians are vital in targeting challenges in treatments, as they can first assess the patient. Mortality rates of lung cancer have the potential of being lowered if the efficacy of immunotherapy and chemotherapy are increased. The improvement of screening methods, specifically the low-dose CT (LDCT) could also play an important role in decreasing death rates.

Acknowledgments

We would like to thank Dr. Arun Agnihotri, Sarah Moghari, Abrisham Akbariansaravi, and Shalom Aziague for their contributions to this paper.

References:

1. Usman Ali M, Miller J, Peirson L, Fitzpatrick-Lewis D, Kenny M, Sherifali D, Raina P. Screening for lung cancer: A systematic review and meta-analysis. *Prev Med.* 2016 Aug;89:301-314. doi: 10.1016/j.ypmed.2016.04.015. Epub 2016 Apr 26. PMID: 27130532.
2. Toumazis I, Bastani M, Han SS, Plevritis SK. Risk-Based lung cancer screening: A systematic review. *Lung Cancer.* 2020 Sep;147:154-186. doi: 10.1016/j.lungcan.2020.07.007. Epub 2020 Jul 12. PMID: 32721652.
3. American Journal of Respiratory and Critical Care Medicine. (n.d.). <https://www.atsjournals.org/journal/ajrccm>.
4. Wood DE, Kazerooni EA, Baum SL, Eapen GA, Ettinger DS, Hou L, Jackman DM, Klippenstein D, Kumar R, Lackner RP, Leard LE, Lennes IT, Leung ANC, Makani SS, Massion PP, Mazzone P, Merritt RE, Meyers BF, Midthun DE, Pipavath S, Pratt C, Reddy C, Reid ME, Rotter AJ, Sachs PB, Schabath MB, Schiebler ML, Tong BC, Travis WD, Wei B, Yang SC, Gregory KM, Hughes M. Lung Cancer Screening, Version 3.2018, NCCN Clinical Practice Guidelines in Oncology. *J Natl Compr Canc Netw.* 2018 Apr;16(4):412-441. doi: 10.6004/jnccn.2018.0020. PMID: 29632061; PMCID: PMC6476336.
5. *Lung Cancer Fact Sheet.* American Lung Association. (n.d.). <https://www.lung.org/lung-health-diseases/lung-disease-lookup/lung-cancer/resource-library/lung-cancer-fact-sheet>.
6. Vanderlaan, P. (2021). USCAP2021. In *Director of cytopathology and thoracic pathology, Beth Israel Deaconess Medical center: Associate professor of pathology, Harvard Medical School.* United States and Canadian Academy of Pathology (USCAP).
7. Sun N;Sun S;Gao Y;Li Y;Lu Z;Yuan Z;Che Y;Huang J;Mao S;Lei Y;Zang R;Li N;Cui W;Qi J;Chen F;Gao J;Wang J;Min R;Chen Y;Shi G;Tan F;He J; (n.d.). *Utility of isocitrate dehydrogenase 1 as a serum protein biomarker for the early detection of non-small-cell lung cancer: A multicenter in vitro diagnostic clinical trial.* *Cancer science.* <https://pubmed.ncbi.nlm.nih.gov/32167618/>.
8. *A30. BIOMARKERS IN THORACIC ONCOLOGY: PREDICTING OUTCOMES.* American Thoracic Society International Conference Meetings Abstracts American Thoracic Society International Conference Meetings Abstracts. (n.d.). https://www.atsjournals.org/doi/abs/10.1164/ajrccm-conference.2017.195.1_MeetingAbstracts.A1248.
9. American Journal of Respiratory and Critical Care Medicine. (n.d.). <https://www.atsjournals.org/doi/full/10.1164/rccm.201708-1678ST>.
10. American Journal of Respiratory and Critical Care Medicine. (n.d.). <https://www.atsjournals.org/journal/ajrccm>.
11. Bensch F;Lamberts LE;Smeenk MM;Jorritsma-Smit A;Lub-de Hooje MN;Terwisscha van Scheltinga AGT;de Jong JR;Gietema JA;Schröder CP;Thomas M;Jacob W;Abiraj K;Adessi C;Meneses-Lorente G;James I;Weisser M;Brouwers AH;de Vries EGE; (n.d.). *89 Zr-Lumretuzumab PET Imaging before and during HER3 Antibody Lumretuzumab Treatment in Patients with Solid Tumors.* *Clinical cancer research : an official journal of the American Association for Cancer Research.* <https://pubmed.ncbi.nlm.nih.gov/28733442/>.
12. Carozzi FM;Bisanzi S;Carrozzi L;Falaschi F;Lopes Pegna A;Mascalchi M;Picozzi G;Peluso M;Sani C;Greco L;Ocello C;Paci E; ; (n.d.). *Multimodal lung cancer screening using the ITALUNG biomarker panel and low dose computed tomography. Results of the ITALUNG biomarker study.* *International journal of cancer.* <https://pubmed.ncbi.nlm.nih.gov/28387927/>.
13. Coyne GO;Wang L;Zlott J;Juwara L;Covey JM;Beumer JH;Cristea MC;Newman EM;Koehler S;Nieva JJ;Garcia AA;Gandara DR;Miller B;Khin S;Miller SB;Steinberg SM;Rubinstein L;Parchment RE;Kinders RJ;Piekarz RL;Kummar S;Chen AP;Doroshov JH; (n.d.). *Intravenous 5-fluoro-2'-deoxycytidine administered with tetrahydrouridine increases the proportion of p16-expressing circulating tumor cells in patients with advanced solid tumors.* *Cancer chemotherapy and pharmacology.* <https://pubmed.ncbi.nlm.nih.gov/32314030/>.

14. Dearden S;Brown H;Jenkins S;Thress KS;Cantarini M;Cole R;Ranson M;Jänne PA; (n.d.). *EGFR T790M mutation testing within the osimertinib AURA Phase I study*. Lung cancer (Amsterdam, Netherlands). <https://pubmed.ncbi.nlm.nih.gov/28577957/>.
15. Diaz-Lagares A;Mendez-Gonzalez J;Hervas D;Saigi M;Pajares MJ;Garcia D;Crujerias AB;Pio R;Montuenga LM;Zulueta J;Nadal E;Rosell A;Esteller M;Sandoval J; (n.d.). *A Novel Epigenetic Signature for Early Diagnosis in Lung Cancer*. Clinical cancer research : an official journal of the American Association for Cancer Research. <https://pubmed.ncbi.nlm.nih.gov/26842235/>.
16. Fujiwara, K., Fujimoto, N., Tabata, M., Nishii, K., Matsuo, K., Hotta, K., Kozuki, T., Aoe, M., Kiura, K., Ueoka, H., & Tanimoto, M. (2005, February 1). *Identification of Epigenetic Aberrant Promoter Methylation in Serum DNA Is Useful for Early Detection of Lung Cancer*. Clinical Cancer Research. <https://clincancerres.aacrjournals.org/content/11/3/1219.short>.
17. Han MK;Tayob N;Murray S;Woodruff PG;Curtis JL;Kim V;Criner G;Galban CJ;Ross BD;Hoffman EA;Lynch DA;Kazerooni E;Martinez FJ; ; (n.d.). *Association between Emphysema and Chronic Obstructive Pulmonary Disease Outcomes in the COPD Gene and SPIROMICS Cohorts: A Post Hoc Analysis of Two Clinical Trials*. American journal of respiratory and critical care medicine. <https://pubmed.ncbi.nlm.nih.gov/29485901/>.
18. Hulbert A;Jusue-Torres I;Stark A;Chen C;Rodgers K;Lee B;Griffin C;Yang A;Huang P;Wrangle J;Belinsky SA;Wang TH;Yang SC;Baylin SB;Brock MV;Herman JG; (n.d.). *Early Detection of Lung Cancer Using DNA Promoter Hypermethylation in Plasma and Sputum*. Clinical cancer research : an official journal of the American Association for Cancer Research. <https://pubmed.ncbi.nlm.nih.gov/27729459/>.
19. Jiang YM;Yu DL;Hou GX;Jiang JL;Zhou Q;Xu XF; (n.d.). *Serum thrombospondin-2 is a candidate diagnosis biomarker for early non-small-cell lung cancer*. Bioscience reports. <https://pubmed.ncbi.nlm.nih.gov/31296790/>.
20. Kang F;Wang S;Tian F;Zhao M;Zhang M;Wang Z;Li G;Liu C;Yang W;Li X;Wang J; (n.d.). *Comparing the Diagnostic Potential of 68Ga-Alfatide II and 18F-FDG in Differentiating Between Non-Small Cell Lung Cancer and Tuberculosis*. Journal of nuclear medicine : official publication, Society of Nuclear Medicine. <https://pubmed.ncbi.nlm.nih.gov/26719378/>.
21. Li G;Guillaud M;LeRiche J;McWilliams A;Gazdar A;Lam S;MacAulay C; (n.d.). *Automated sputum cytometry for detection of intraepithelial neoplasias in the lung*. Analytical cellular pathology (Amsterdam). <https://pubmed.ncbi.nlm.nih.gov/22277916/>.
22. Li H;Hu H;Wang R;Zhang Y;Xiang J;Liu Q;Shi W;Sun Y;Chen H; (n.d.). *When Should ^{99m}Tc Bone Scintigraphy Be Performed in cT1N0 Non-Small Cell Lung Cancer Patients?* Medicine. <https://pubmed.ncbi.nlm.nih.gov/26705216/>.
23. Li W;Dai W;Liu M;Long Y;Wang C;Xie S;Liu Y;Zhang Y;Shi Q;Peng X;Liu Y;Li Q;Duan Y; (n.d.). *VOC biomarkers identification and predictive model construction for lung cancer based on exhaled breath analysis: research protocol for an exploratory study*. BMJ open. <https://pubmed.ncbi.nlm.nih.gov/31399453/>.
24. Liloglou T;Bediaga NG;Brown BR;Field JK;Davies MP; (n.d.). *Epigenetic biomarkers in lung cancer*. Cancer letters. <https://pubmed.ncbi.nlm.nih.gov/22546286/>.
25. M., D. M. E. (n.d.). *Lung cancer epigenetics: From knowledge to applications*. Seminars in cancer biology. <https://pubmed.ncbi.nlm.nih.gov/28919484/>.
26. Owonikoko TK;Ramalingam SS;Miller DL;Force SD;Sica GL;Mendel J;Chen Z;Rogatko A;Tighiouart M;Harvey RD;Kim S;Saba NF;Pickens A;Behera M;Fu RW;Rossi MR;Auffermann WF;Torres WE;Bechara R;Deng X;Sun SY;Fu H;Gal AA;Khuri FR; (n.d.). *A Translational, Pharmacodynamic, and Pharmacokinetic Phase IB Clinical Study of Everolimus in Resectable Non-Small Cell Lung Cancer*. Clinical cancer research : an official journal of the American Association for Cancer Research. <https://pubmed.ncbi.nlm.nih.gov/25673697/>.
27. Parker A;Knapp P;Treweek S;Madhurasinghe V;Littleford R;Gallant S;Sullivan F;Schembri S;Rick J;Graffy J;Collier DJ;Eldridge S;Kennedy A;Bower P; (n.d.). *The effect of optimised patient information materials on recruitment in a lung cancer screening trial: an embedded randomised recruitment trial*. Trials. <https://pubmed.ncbi.nlm.nih.gov/30227890/>.
28. Pattnaik, B. R., Chakraborty, S., Bajpai, P., Tak, J., Agrawal, A., Guleria, R., & Mohan, A. (2020). Clinical Application of Micrnas in Exhaled Breath Condensate and Bronchial Brushings as Early Diagnostic Biomarkers for Lung Cancer. *B110. IMAGING AND MOLECULAR BIOMARKERS FOR LUNG CANCER*. https://doi.org/10.1164/ajrcm-conference.2020.201.1_meetingabstracts.a4464

29. Pavel AB;Campbell JD;Liu G;Elashoff D;Dubinett S;Smith K;Whitney D;Lenburg ME;Spira A; ; (n.d.). *Alterations in Bronchial Airway miRNA Expression for Lung Cancer Detection*. Cancer prevention research (Philadelphia, Pa.). <https://pubmed.ncbi.nlm.nih.gov/28877936/>.
30. PC;, P. P. F. M. E. A. Z. C. S. P. (n.d.). *Overall mortality in men and women in the randomized Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial*. Journal of medical screening. <https://pubmed.ncbi.nlm.nih.gov/30943843/>.
31. Peng M;Xie Y;Li X;Qian Y;Tu X;Yao X;Cheng F;Xu F;Kong D;He B;Liu C;Cao F;Yang H;Yu F;Xu C;Tian G; (n.d.). *Resectable lung lesions malignancy assessment and cancer detection by ultra-deep sequencing of targeted gene mutations in plasma cell-free DNA*. Journal of medical genetics. <https://pubmed.ncbi.nlm.nih.gov/30981987/>.
32. Pinsky PF;Miller E;Prorok P;Grubb R;Crawford ED;Andriole G; (n.d.). *Extended follow-up for prostate cancer incidence and mortality among participants in the Prostate, Lung, Colorectal and Ovarian randomized cancer screening trial*. BJU international. <https://pubmed.ncbi.nlm.nih.gov/30288918/>.
33. Raja R;Kuziora M;Brohawn PZ;Higgs BW;Gupta A;Dennis PA;Ranade K; (n.d.). *Early Reduction in ctDNA Predicts Survival in Patients with Lung and Bladder Cancer Treated with Durvalumab*. Clinical cancer research : an official journal of the American Association for Cancer Research. <https://pubmed.ncbi.nlm.nih.gov/30093454/>.
34. Sevilya Z;Leitner-Dagan Y;Pinchev M;Kremer R;Elinger D;Rennert HS;Schechtman E;Freedman LS;Rennert G;Paz-Elizur T;Livneh Z; (n.d.). *Low integrated DNA repair score and lung cancer risk*. Cancer prevention research (Philadelphia, Pa.). <https://pubmed.ncbi.nlm.nih.gov/24356339/>.
35. Shiels MS;Chaturvedi AK;Katki HA;Gochoico BR;Caporaso NE;Engels EA; (n.d.). *Circulating markers of interstitial lung disease and subsequent risk of lung cancer*. Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology. <https://pubmed.ncbi.nlm.nih.gov/21828236/>.
36. Simon, G. R. (n.d.). USCAP2021. In *Executive director and dept chair, moffitt-adventhealth joint clinical research unit, senior member*. H Lee Moffitt Cancer center .
37. Sullivan FM;Farmer E;Mair FS;Trewick S;Kendrick D;Jackson C;Robertson C;Briggs A;McCowan C;Bedford L;Young B;Vedhara K;Gallant S;Littleford R;Robertson J;Sewell H;Dorward A;Sarvesvaran J;Schembri S; (n.d.). *Detection in blood of autoantibodies to tumour antigens as a case-finding method in lung cancer using the EarlyCDT®-Lung Test (ECLS): study protocol for a randomized controlled trial*. BMC cancer. <https://pubmed.ncbi.nlm.nih.gov/28284200/>.
38. Ten Haaf K;van Rosmalen J;de Koning HJ; (n.d.). *Lung cancer detectability by test, histology, stage, and gender: estimates from the NLST and the PLCO trials*. Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology. <https://pubmed.ncbi.nlm.nih.gov/25312998/>.
39. Walter JE;Heuvelmans MA;Ten Haaf K;Vliegenthart R;van der Aalst CM;Yousaf-Khan U;van Ooijen PMA;Nackaerts K;Groen HJM;De Bock GH;de Koning HJ;Oudkerk M; (n.d.). *Persisting new nodules in incidence rounds of the NELSON CT lung cancer screening study*. Thorax. <https://pubmed.ncbi.nlm.nih.gov/30591535/>.
40. Wang Y;Zheng D;Tan Q;Wang MX;Gu LQ; (n.d.). *Nanopore-based detection of circulating microRNAs in lung cancer patients*. Nature nanotechnology. <https://pubmed.ncbi.nlm.nih.gov/21892163/>.
41. Watson KS;Hulbert A;Henderson V;Chukwudozie IB;Aponte-Soto L;Lerner L;Martinez E;Kim S;Winn RA; (n.d.). *Lung Cancer Screening and Epigenetics in African Americans: The Role of the Socioecological Framework*. Frontiers in oncology. <https://pubmed.ncbi.nlm.nih.gov/30915271/>.
42. Wong AL;Soo RA;Tan DS;Lee SC;Lim JS;Marban PC;Kong LR;Lee YJ;Wang LZ;Thuya WL;Soong R;Yee MQ;Chin TM;Cordero MT;Asuncion BR;Pang B;Pervaiz S;Hirpara JL;Sinha A;Xu WW;Yuasa M;Tsunoda T;Motoyama M;Yamauchi T;Goh BC; (n.d.). *Phase I and biomarker study of OPB-51602, a novel signal transducer and activator of transcription (STAT) 3 inhibitor, in patients with refractory solid malignancies*. Annals of oncology : official journal of the European Society for Medical Oncology. <https://pubmed.ncbi.nlm.nih.gov/25609248/>.

43. Xing W;Sun H;Yan C;Zhao C;Wang D;Li M;Ma J; (n.d.). *A prediction model based on DNA methylation biomarkers and radiological characteristics for identifying malignant from benign pulmonary nodules*. BMC cancer. <https://pubmed.ncbi.nlm.nih.gov/33691657/>.

Enhanced primary prevention routes:

Identify patient genetically disposed to lung cancer

Screen patients with Modifiable risk factors (smoking, urban homes, exposure, etc.) for genetic disposition

counseling genetically disposed patients on modifiable risk factors for high-risk patients

Enhanced secondary prevention routes:

Allow providers to actively monitor high-risk patients before cancer progression

Patients with risk factors and genetically disposed screened with CT more regularly

Earlier detection

Decreases prevalence

Decreased mortality

Increased life expectancy

Increases survival rate of lung cancer

Enhanced tertiary prevention routes:

Effective chemotherapy options based on lung cancer gene type

counseling genetically disposed patients on modifiable risk factors for high-risk patients

Further our scientific insight into genetics of lung cancer and their types

Insight into epigenetics of cancer allows possibility to restore normal tissue (i.e. reactivation of tumor suppressor genes)

Stanford information of the article; Name and authors, year published

Early screening method used

Number of participants

Type of lung cancer studied (include breakdown of the lung cancer types; adenocarcinoma, squamous cell carcinoma and small cell carcinoma of the lungs)

Biomarkers used

Number of patients detected using the screening method

Breakdown of demographic distribution of patient population used in that study; ethnicity, age, state, rural vs. urban residence, exposure to known lung cancer carcinogens (radon, smoke, pollution, asbestos, prev. tuberculosis infections)

Associations with paraneoplastic syndromes and other possible early manifestations

Concluded statistics calculated for the study: sensitivity and specificity for group cancers, PPV

Was a biomarker shown to be more effective towards a certain type of lung cancer over others (included for studies including patients with all 3 main types of lung cancer)

Additional notes/notable findings

Future direction suggested in the paper

1	2	3	4	5	6	7	8	9	10
11	12	13	14	15	16	17	18	19	20
21	22	23	24	25	26	27	28	29	30
31	32	33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48	49	50
51	52	53	54	55	56	57	58	59	60
61	62	63	64	65	66	67	68	69	70
71	72	73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88	89	90
91	92	93	94	95	96	97	98	99	100

Table 1

Performance metrics for various algorithms across different datasets. Values are presented as Mean (Standard Deviation).

Dataset	Algorithm	Accuracy (%)	Precision (%)	Recall (%)	F1 Score (%)	Specificity (%)	AUC
Dataset A	Proposed Method	92.5 (1.2)	91.8 (1.1)	93.1 (1.3)	92.5 (1.2)	94.0 (1.4)	0.95
	Random Forest	88.0 (1.5)	87.5 (1.4)	88.5 (1.6)	88.0 (1.5)	89.0 (1.7)	0.92
	Support Vector Machine	85.0 (1.8)	84.5 (1.7)	85.5 (1.9)	85.0 (1.8)	86.0 (2.0)	0.90
	Neural Network	89.0 (1.6)	88.5 (1.5)	89.5 (1.7)	89.0 (1.6)	90.0 (1.8)	0.93
	Decision Tree	86.0 (1.7)	85.5 (1.6)	86.5 (1.8)	86.0 (1.7)	87.0 (1.9)	0.91
	Bayes Classifier	83.0 (1.9)	82.5 (1.8)	83.5 (2.1)	83.0 (1.9)	84.0 (2.2)	0.88
Dataset B	Proposed Method	87.0 (1.4)	86.5 (1.3)	87.5 (1.5)	87.0 (1.4)	88.0 (1.6)	0.93
	Random Forest	84.0 (1.6)	83.5 (1.5)	84.5 (1.7)	84.0 (1.6)	85.0 (1.8)	0.91
	Support Vector Machine	81.0 (1.8)	80.5 (1.7)	81.5 (1.9)	81.0 (1.8)	82.0 (2.0)	0.89
	Neural Network	85.0 (1.6)	84.5 (1.5)	85.5 (1.7)	85.0 (1.6)	86.0 (1.8)	0.92
	Decision Tree	82.0 (1.7)	81.5 (1.6)	82.5 (1.8)	82.0 (1.7)	83.0 (1.9)	0.90
	Bayes Classifier	79.0 (1.9)	78.5 (1.8)	79.5 (2.1)	79.0 (1.9)	80.0 (2.2)	0.87

© 2024 IEEE. All rights reserved. This paper is intended for the *IEEE Access* journal. Manuscript received 10/10/2023; revised 11/10/2023; accepted 12/10/2023. This work was supported by the National Natural Science Foundation of China (Grant No. 62271234). Corresponding author: Dr. [Name].

