
Harmonization of ICF Body Structures and ICD-11 1

Anatomic Detail: one foundation for two classifications



Abstract

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

The Family of International Classifications of the World Health Organization currently includes three reference classifications, namely International Classification of Diseases (ICD), International Classification of Functioning, Disability, and Health (ICF), and International Classification of Health Interventions (ICHI). Each of them serves a specific classification need. However, they share some common concepts that are present, in different forms, in two or all of them. One important set of shared concepts is the representation of human anatomy entities, which are not always modeled in the same way and with the same level of detail. To understand the relationships among the three anatomical representations, an effort is needed to compare them, identifying common areas, gaps, and compatible and incompatible modeling. The work presented here contributes to this effort, focusing on the anatomy representations in ICF and ICD-11.

For this aim, three experts were asked to identify, for each entity in the ICF Body Structures, one or more entities in the ICD-11 Anatomic Detail that could be considered identical, broader or narrower. To do this, they used a specifically developed web application, which also automatically identified the most obvious equivalences.

A total of 631 maps were independently identified by the three mappers for 218 ICF Body Structures, with an interobserver agreement of 93.5%. Together with 113 maps identified by the software, they were then consolidated into 434 relations. The results highlight some differences between the two classifications: in general, ICF is less detailed than ICD-11; ICF favors lumping of structures; in very few cases, the two classifications follow different anatomic models. For these issues, solutions have to be found that are compliant with the WHO approach to classification modeling and maintenance.

Introduction

39

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

- 40 The Family of International Classifications of the World Health Organization (WHO-FIC)
- currently includes three reference classifications, namely: 41
- ICD: the International Classification of Diseases 42
 - ICF: the International Classification of Functioning, Disabilities and Health
 - ICHI: the International Classification of Health Interventions

Each of them covers a specific need that may arise when classifying the content of health documentation. Furthermore, while they share common principles, their structures depend in part on choices made at their initial design and development phases, which were done at different times and which in turn depend on knowledge and constraints defined at the times. In particular, the 11th revision of ICD (1) marks a departure from the paper-based model that led the development of the previous revisions, with a specific attention to information technology aspects, including proper formalization of the represented concepts, and a double-layered structure composed of an ICD-11 Foundation that includes all possible ICD-related information and one or more linearizations that represents different versions of the classification designed to satisfy specific use cases (2). The ICD-11 experience informs the development of ICHI, which is the youngest classification of the family; on the other side, ICF might be considered the oldest one, as it is still oriented towards traditional paper-based usage, although recently it is subjected to some enhancement in the computerized direction (3,4).

The most notable recent enhancement to the three classifications is that all of them have been incorporated into a single WHO-FIC Foundation that allows both ICF and ICHI to be generated as linearizations of the Foundation (5). In this combined representation, ICD-11 and ICF concepts are

represented separately, while ICHI, because of its recent provenance, references some ICD-11 and ICF concepts.

Even though they cover different needs, the classifications share some common concepts that are present, in different forms in the Foundation, in two or all of them. It is the aim of a current WHO effort to harmonize the shared concepts among the reference classifications of the family (6) in the Foundation.

Identifying and characterizing all the possible shared concepts is a complex task involving the semantics of the reference classification, however there is one set of concepts that is obviously shared and easy to recognize, i.e., human anatomy. In fact, an anatomical entity may define the site of a disease in ICD-11, the body structure subject to impairment in ICF and also the target of some health interventions defined in ICHI.

While they are present everywhere, it is not obvious that all the anatomical entities are modeled the same way and with the same level of detail in the three reference classifications. To understand the relationships among the three anatomical representations, an effort is needed to compare them, identifying common areas, gaps, and compatible and incompatible models. The work presented here contributes to this effort, focusing on the anatomy representations in ICF and ICD-11 at first.

- The aims of the present paper are thus:
- To identify anatomical detail as represented in ICF and ICD-11;
- 77 To compare the representations by setting relationships among specific entities;
- 78 To characterize the level of detail in each classification;
- 79 To characterize areas where the underlying modeling is different or even incompatible;
 - Finally, to suggest measures to be taken to harmonize and consolidate the representations.

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

All rights reserved. No reuse allowed without permission.

While there exists a deeply formalized representation of anatomy, namely the Foundational Model of Anatomy (7), it is much too detailed for our goal, which is to harmonize the anatomical representations in the WHO-FIC Foundation, minimizing the changes needed to the classifications, with the aim of continuity and compatibility with current applications.

Methods

To understand the relationship between ICF and ICD-11 anatomy, we decided to begin by mapping anatomical entities in the two classifications. In particular, the specific areas considered were the ICF Body Structures, and the section of ICD-11 Extension codes called "Anatomic Detail". Both have a partially heterogeneous hierarchical representation, including both partonomic and taxonomic relationships.

Expert mappings

An initial qualitative analysis showed that inside these subsets, some entities have exactly the same title, some others are named differently, and some others do not have a direct correspondence, because an entity explicitly mentioned in one classification may only be found as part of a larger, or as specialization of a more general entity in the other classification.

Thus, three experts were asked to identify, for each entity in the ICF Body Structures, one equivalent entity in the ICD-11 Anatomic Detail if possible; if not, one or more entities that could be considered *broader_than* or *narrower_than*. We opted for these generic relationships because they encompass both the partonomic and taxonomic views. Furthermore, for practical aims, we also subdivided the equivalence relationship in two subtypes: *identical_to*, when the title and concept was substantially the same, and *synonym_of*, when the title of the concept was different, yet equivalent. Matching has been attempted not only on the terminal entities in the hierarchy, but also for all the higher level entities.

While this effort was not aimed at mapping electronic health records, when possible the principles described in the WHO paper on mapping have been respected (8).

Classifications

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

The source list for ICF Body Structures was obtained from a ClaML representation of ICF version 2017, and included 321 entities. Of these 321, 103 were identified as residuals, i.e., categories of the kinds "not otherwise specified" or "other specified". Residuals were not mapped, because semantically equivalent to the parent category.

The target ICD-11 list was composed by a subset of the Extension Codes (9), namely the "Anatomical Detail" branch, taken from the Foundation layer. It was not extracted in advance, but dynamically obtained through the ICD-11 API (10) when needed, as explained in the next paragraphs.

Software

To support the matching effort by the three experts, an ad-hoc web-based software has been developed. The main interface of the software shows the ICF Body Structures hierarchy on the left side, with the capability of selecting a kind of relationship among those previously mentioned, and the corresponding ICD-11 entity, for each ICF entity. The ICD-11 entity is identified by accessing an instance of the ICD-11 Foundation coding tool (10) on the right side, set to automatically search for the ICF title in the ICD-11 Foundation, Extension Codes chapter, "Anatomic Detail" branch. However, since this could not always be found, the expert may also edit the search term with synonyms that might be present in ICD-11. Figure 1 shows a screenshot from the Map Editor.

Figure 1: on the left side, part of ICF Body Structures. The user is selecting a relationship (synonym of) for the ICF entity "Lens of eyeball". On the right side, the ICD-11 Coding tool attempted a search with that term, which is not directly found. However, the expert may choose "Crystalline lens" and have it automatically inputed in the left side field, to propose it as candidate relationship. In the top left, some already defined relationships could be seen. Those identified as "auto" are automatically set by the software.

To reduce the effort needed, the software has a module that automatically identifies straightforward relationships, i.e., those where titles are identical, or obvious synonyms. Among the latter, many entities in ICF are named as "Structure of X", which were automatically mapped as synonyms of "X" if available in ICD-11. Technically "structure of X" is not synonymous with "X." However, given that parent/child links in both ICD-11 anatomy extension codes and ICF body structures are a mix of taxonomic and partonomic relationship, identifying "structure of X" with "X" yields the most economical set of maps. As an example, 'Structure of brain' from ICF is automatically mapped as synonym_of 'Brain' in ICD-11.

While doing their work, experts do not see matches already found by the others, in order to allow the computation of a measure of inter-observer agreement.

Another interface shows a tree representation of the ICF Body Structures, with color codings for the different relationships, when available. Figure 2 shows a screenshot depicting part of the ICF Body Structures tree with mappings.

Figure 2: part of the tree view of the ICF Body Structure, with color-coded matches towards ICD-11.

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

Finally, a further module allows the export of matches to a CSV file for further processing.

Results

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

Of 218 entities considered in the source list and excluding residuals, 113 were found by software to have identical corresponding entities in the ICD-11 Anatomic Detail extension, basing on lexical criteria.

Three independent knowledgeable raters considered the 105 entities for which the tool did not automatically confirm identity matches. The search for the most appropriate matching entity could result in finding identical, equivalent, broader or narrower items or no match. The three mappers independently produced 631 maps to a total of 297 ICD-11 entities, independently and without accessing others' maps until the end of the experiment. For all the ICF entities, at least one expert found a map in ICD-11. Table 1 shows a breakdown of the experts' mappings. While identical to and synonym of maps were collected, for the sake of simplicity, and considering that they are semantically the same, for the analysis they were collapsed in a single *equivalent_to* category.

Table 1: mappings proposed by each expert

mapper	relations			
	equivalent_to	broader_than	narrower_than	total
A	50	106	11	167
В	55	226	26	307

С	36	84	37	157
total	141	416	74	631

Interobserver agreement

160

161

162

163

164

169

170

171

- the 26 in disagreement, some typical examples are as follows:
- Bronchial tree in ICF equivalent to or broader than Bronchus in ICD-11; 165
- Vaginal canal in ICF equivalent to or narrower than Vagina in ICD-11; 166
- Ligaments and fasciae of hand in ICF broader than or narrower than Ligament of the wrist and 167 168 hand in ICD-11.
 - On the other side, by considering the 115 mapped ICF categories, some further differences became visible due to different choices in narrower than or broader than relationships (for example, obvious or redundant broader_than relationships). This involved the same ICF source, but possibly different ICD-11 targets. Examples from a total of 12 cases are:
- 173 Structure of diencephalon narrower_than Supratentorial region of brain
- 174 narrower than Brain: this one is redundant
- 175 broader_than Thalamus
- Structure of skin glands 176 broader than Sebaceous gland
- 177 broader than Apocrine sweat gland

178 broader_than Eccrine gland 179 narrower_than Skin 180 equivalent_to Cardiac atrium atria 181 broader_than Right atrium

Discrepancies were examined by the experts to delete from the final set the maps on which there was not an agreement, that were redundant, or that were mistakes.

Mapping details

182

183

184

185

186

187

188

189

190

191

192

193

The mappings have been consolidated in a single set of agreed-upon relationships, including automatically calculated equivalencies.

In order to obtain a consolidated mapping, we prioritized relationships as follows. First of all, if there is agreement on an equivalency, it is considered as the selected relationship, even if others are available. Without an equivalency, if a narrower_than relationship is available, that will be selected. Finally, broader_than relationships are selected.

After this step, the consolidated mappings are as described in Table 2.

Table 2: Consolidated mappings

	relations			
	equivalent_to	broader_than	narrower_than	total
consolidated	164	230	40	434

Discussion

194

195

196

197

198

199

200

201

202

203

204

205

206

207

The common Foundation from which the various WHO reference classifications are derived through linearizations requires that every entity be uniquely and unambiguously defined. This might require a process of harmonization of concepts that share similar meanings but may or may not be identical, also to favor joint use of the reference classifications. The process may be more or less complex according to the entities considered, but could be easier for entities already sharing strong similarities. For further automatic mappings of concepts that may not be of the same semantic types, lexical mappings resulting in identical maps need to be evaluated to make sure that the concepts are identical. The "low hanging fruits" of harmonization include the anatomy entities as described in ICD-11 and in the body structure domain of ICF, which have been the subject of this paper.

Three main issues arise from this experiment are as follows:

- A. Different levels of specificity: in general, ICF is less detailed than ICD (74 narrower_than vs 416 broader than maps);
- 208 B. ICF favors lumping of structures: e.g. "The eye, ear and related structures; Structures of 209 cardiovascular, immunological and respiratory systems; Structure of vagina and external genitalia; 210 Testes and scrotum";
- 211 C. The two classifications may have different anatomic models: e.g., Where is the shoulder? For ICD: 212 in upper extremity; for ICF: in structures related to movement.

For these issues, solutions have to be found that are compliant with the WHO approach to classification modeling and maintenance, which includes the concept of a common Foundation that embodies all the relevant entities.

Issue A does not represent a real problem: according to the approach for creating linearizations from the Foundation, the "boundary" of specificity is called "shoreline"; and to select different specificities, it is only a matter of defining a different shoreline for each classification.

Issue B might have two different solutions. The solution with least impact, from the point of view of classifications, is to introduce the ICF groupings as new groupings in the Foundation's Anatomic Detail, while not using them in the ICD-11 linearization. However, such groupings are often heterogeneous and missing a strong semantic similarity (e.g., "The eye, ear and related structures"), and are present only for the sake of aggregation in a classification that was born with a printed edition in mind, but can be useful for aggregation of data at a higher level. Thus, the ideal solution would be to revise the hierarchy organization in ICF, without modifying terminal entities, in order to split heterogeneous groupings into more semantically homogeneous groups.

Issue C is apparently the most complex: different models may mean a totally different hierarchy, and such differences in modeling is intrinsic in the different points of view expressed by ICF and ICD (the former centered on functioning, the latter on clinical aspects). Figure 3 shows a clear and compelling example: in ICF terms, mouth and nose are structures devoted to speech, while in ICD terms, mouth is part of the digestive system, and nose of the respiratory system. Both views are perfectly reasonable, yet both miss the aspect highlighted by the other classification. Nevertheless, both views may be accommodated in the WHO-FIC Foundation, which differs from traditional classifications in that it allows for multiple parenthood relationships. The use of multiple parents might enable us to express the different

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

All rights reserved. No reuse allowed without permission.

views without forcing one classification to adopt the point of view of the other, and leaving to the linearization phase the choice of one or the other parent in each classification. Figure 3: Nose and mouth have different parents in ICF and in ICD-11. The consolidated made available in **SSOM** format (11)maps are at https://github.com/whoficitc/harmonization/blob/main/ICF-ICD-anatomy-v1.tsv. **Conclusion** The harmonization experiment now needs to be extended to ICHI and might be replicated for the examples of related impairments described in ICF and found in ICD as symptoms or health conditions. References 1. Harrison JE, Weber S, Jakob R et al. ICD-11: an international classification of diseases for the twentyfirst century. BMC Med Inform Decis Mak 21 (Suppl 6), 206 (2021). https://doi.org/10.1186/s12911-021-01534-6 2. Chute, C.G., Celik, C. Overview of ICD-11 architecture and structure. BMC Med Inform Decis Mak 21 (Suppl 6), 378 (2021). https://doi.org/10.1186/s12911-021-01539-1 3. Heerkens YF, de Weerd M, Huber M, de Brouwer CPM, van der Veen S, Perenboom RJM, van Gool CH, Ten Napel H, van Bon-Martens M, Stallinga HA, van Meeteren NLU. Reconsideration of the

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

- 255 scheme of the international classification of functioning, disability and health: incentives from the
- 256 Netherlands for a global debate. Disabil Rehabil. 2018;40(5):603–11.
- 257 4. Cozzi, S., Martinuzzi, A. Della Mea, V. Ontological modeling of the International Classification of
- 258 Functioning, Disabilities and Health (ICF): Activities&Participation and Environmental Factors
- 259 components. BMC Med Inform Decis Mak 21, 367 (2021). https://doi.org/10.1186/s12911-021-01729-
- 260 X
- 261 5. World Health Organization.. WHO-FIC Foundation. https://icd.who.int/dev11/f/en
- 262 6. Tu S, Nyulas C, Tudorache T, Musen M, Martinuzzi A, van Gool C, Della Mea V, Chute C, Frattura
- 263 L, Hardiker N, Napel H, Madden R, Almborg A-H, Ginige J, Sykes C. Toward a harmonized WHO
- 264 family of international classifications content model. Stud Health Technol Inform. 2020;270:1409–10.
- 265 7. Rosse C, Mejino JL Jr. A reference ontology for biomedical informatics: the Foundational Model of
- 266 Anatomy. J Biomed Inform. 2003 Dec;36(6):478-500. doi: 10.1016/j.jbi.2003.11.007. PMID:
- 267 14759820.
- 268 8. WHO-FIC Network. WHO-FIC Classifications and terminology mapping. Principles and best practice.
- 269 WHO, https://www.who.int/publications/m/item/who-fic-classifications-andterminology-2021.
- 270 mapping
- 271 9. Drösler, S.E., Weber, S. & Chute, C.G. ICD-11 extension codes support detailed clinical abstraction
- 272 and comprehensive classification. BMC Med Inform Decis Mak 21 (Suppl 6), 278 (2021).
- 273 https://doi.org/10.1186/s12911-021-01635-2
- 274 10. Celik C, Della Mea V, Donada M, Kostanjsek N, Jakob R, Chute CG. The ICD Platform: end-user
- 275 and developer tools. Submitted, 2022

Matentzoglu N, Balhoff JP, Bello SM, et al. A Simple Standard for Sharing Ontological Mappings 276 11. 277 (SSSOM). Database (Oxford). 2022 May 25;2022:baac035. doi: 10.1093/database/baac035.

278