

## Daily exposure to PM<sub>2.5</sub> and 1.5 million deaths: A time-stratified case-crossover analysis in the Mexico City Metropolitan Area

Iván Gutiérrez-Avila<sup>1\*</sup>, Horacio Riojas-Rodríguez<sup>2</sup>, Elena Colicino<sup>1</sup>, Johnathan Rush<sup>1</sup>, Marcela Tamayo-Ortiz<sup>3</sup>, Víctor Hugo Borja-Aburto<sup>4</sup>, Allan C. Just<sup>1,5</sup>

<sup>1</sup> Department of Environmental Medicine and Public Health, Icahn School of Medicine at Mount Sinai, New York, NY, USA

<sup>2</sup> Dirección de Salud Ambiental, Instituto Nacional de Salud Pública, Cuernavaca Morelos, México

<sup>3</sup> Instituto Mexicano del Seguro Social. Unidad de Investigación en Salud Ocupacional, México City, México

<sup>4</sup> Instituto Mexicano del Seguro Social, México City, México

<sup>5</sup> Institute for Exposomic Research, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

### Abstract

**Background.** Satellite-based PM<sub>2.5</sub> predictions are being used to advance exposure science and air-pollution epidemiology in developed countries; including emerging evidence about the impacts of PM<sub>2.5</sub> on acute health outcomes beyond the cardiovascular and respiratory systems, and the potential modifying effects from individual-level factors in these associations. Research on these topics is lacking in Latin America.

**Methods.** We used a time-stratified case-crossover study design with 1,479,950 non-accidental deaths from Mexico City Metropolitan Area for the period of 2004-2019. Daily 1x1 km PM<sub>2.5</sub> (median=23.4 µg/m<sup>3</sup>; IQR=13.6 µg/m<sup>3</sup>) estimates from our satellite-based regional model were employed for exposure assessment at the sub-municipality level. Associations between PM<sub>2.5</sub> with broad-category (organ-system) and cause-specific mortality outcomes were estimated with distributed lag conditional logistic models. We also fit models stratifying by potential individual-level effect modifiers including; age, sex, and individual SES-related characteristics namely: education, health insurance coverage, and job categories.

**Results.** PM<sub>2.5</sub> exposure was associated with higher total non-accidental, cardiovascular, cerebrovascular, respiratory, and digestive mortality. A 10-µg/m<sup>3</sup> PM<sub>2.5</sub> higher cumulative exposure over one week (lag<sub>06</sub>) was associated with higher cause-specific mortality outcomes including hypertensive disease [2.28% (95%CI: 0.26%–4.33%)], acute ischemic heart disease [1.61% (95%CI: 0.59%–2.64%)], other forms of heart disease [2.39% (95%CI: -0.35%–5.20%)], hemorrhagic stroke [3.63% (95%CI: 0.79%–6.55%)], influenza and pneumonia [4.91% (95%CI: 2.84%–7.02%)], chronic respiratory disease [2.49% (95%CI: 0.71%–4.31%)], diseases of the liver [1.85% (95%CI: 0.31%–3.41%)], and renal failure [3.48% (95%CI: 0.79%–6.24%)]. No differences in effect size of associations were observed between SES strata.

**Conclusions.** Exposure to PM<sub>2.5</sub> was associated with mortality outcomes beyond the cardiovascular and respiratory systems, including specific death-causes from the digestive and genitourinary systems, with no indications of effect modification by individual SES-related characteristics.

**Keywords:** PM<sub>2.5</sub>, cause-specific mortality, short-term exposure, case-crossover study

### Correspondence

\*Iván Gutiérrez-Avila, Department of Environmental Medicine and Public Health, Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, Box 1057, New York, NY 10029, USA. Email: [ivan\\_2c@hotmail.com](mailto:ivan_2c@hotmail.com)

## Background

Human exposure to fine particulate matter (PM<sub>2.5</sub>) is a well-documented risk factor for cardiovascular and respiratory mortality and morbidity. Nonetheless, exposure to PM<sub>2.5</sub> can affect nearly every organ system through different mechanisms of damage, including oxidative stress, systemic inflammation, and immune dysregulation (Keswani Anjeni et al., 2022). Recent epidemiologic studies have reported associations between short- and long-term exposure to PM<sub>2.5</sub> with mental, behavioral, nervous, cerebrovascular, metabolic, digestive, and genitourinary mortality and morbidity, mostly in the US and Asia (Bowe et al., 2019; Li et al., 2018; Wang et al., 2020; Yu et al., 2019). Little evidence on acute mortality outcomes beyond the cardiovascular and respiratory spectrum have been reported in countries from low- and middle-income regions, including Latin America. Research focused on associations from under-studied causes of death is important to inform concentration-response functions that may be used in health impact assessments, including those quantifying the benefits of improving air quality. In the public-policy framework, omission of some health impacts may lead to underestimation of the benefits from interventions aimed to reduce the burden of disease from PM<sub>2.5</sub> relative to their costs (World Health Organization, 2013).

The development of highly spatially-resolved PM<sub>2.5</sub> models (1x1-km PM<sub>2.5</sub> predictions) based on satellite data, has facilitated the assessment of human exposure at fine spatial scales, even in regions with low to moderate PM<sub>2.5</sub> monitoring, such as Latin America (Gouveia et al., 2021). Also, the use of satellite-based PM<sub>2.5</sub> models' predictions, as well as the combinations of different methods to predict PM<sub>2.5</sub> for epidemiologic research, may reduce exposure measurement error, compared to the use of city-wide averages from small ground monitoring networks (Alexeeff et al., 2015; Samoli et al., 2020; Wei et al., 2022). Highly spatially-resolved PM<sub>2.5</sub> predictions for exposure assessment along with individual-level health records, allows exploration of effect modification by the inclusion of individual (or area-level shared) characteristics which are difficult to consider when using aggregated data, like in traditional city-wide time-series studies (Wei et al., 2019; Yitshak-Sade et al., 2019). Age, sex, socio-economic status (SES), and race, are among the factors that seem to modify vulnerability to PM<sub>2.5</sub> exposure in high-income countries (Franklin et al., 2006; Heo et al., 2022; Renzi et al., 2021; Wang et al., 2020). Such characteristics are of interest for identification of vulnerable populations, science-based policy and targeted public-health interventions (Xu et al., 2020). However, little has been reported about the potential modifying effects of these factors in middle-income countries like Mexico, where their distribution may differ compared to other regions.

In this study, we analyzed all ~1.5 million adult mortality records from the Mexico City Metropolitan Area for the period from 2004 to 2019, and estimated the daily percent increase in non-accidental mortality for broad-category and cause-specific mortality outcomes; as well as sex and age-group (adults and the elderly) categories, and SES-related indicators, associated with a  $10\text{-}\mu\text{g}/\text{m}^3$  higher  $\text{PM}_{2.5}$  concentration.

## Methods

### Mortality data

We obtained mortality records from the National Institute of Statistics and Geography of Mexico (INEGI) for the period spanning February 1<sup>st</sup> 2004 to December 31<sup>st</sup>, 2019. We excluded the month of January 2004 in order to keep only case-days and control-days with complete exposure histories of  $\text{PM}_{2.5}$  and temperature, i.e. up to one week (lag 6) before the date of event. All mortality records included information on date of death, geographic identifiers for place of residence, underlying cause of death classified according to the International Classification of Diseases Tenth Revision (ICD-10), sex, age, education, healthcare affiliation, and job category.

We restricted our analysis to non-accidental causes of death excluding most ICD-10 codes from the blocks V–Y (i.e. various accidents and side effects of treatments), and deaths in people  $\geq 18$  years-old. ICD-10 codes X6–Y0 were retained based on the evidence that  $\text{PM}_{2.5}$  can trigger intentional self-harm and aggressive behavior (Berman et al., 2019; Davoudi et al., 2021). We organized individual ICD-10 coded death records into mutually exclusive broad-category (organ-system) and cause-specific mortality outcomes. The selected ICD-10 codes included in our research were chosen to facilitate comparison with recently-published studies focused on under-studied associations between short-term exposure to  $\text{PM}_{2.5}$  with mortality and morbidity outcomes (Li et al., 2018; Yu et al., 2019).

To address the potential modifying effects of age, sex and SES on the  $\text{PM}_{2.5}$ -mortality relationship, we performed stratified analyses by age-group (adults: 18-64 years-old, and elderly: +65 years-old) and sex (males and females) for all broad-category mortality outcomes. For all non-accidental mortality, we also performed stratified analyses by age-group (adults and the elderly), degree of education (basic and more than basic education level), type of healthcare affiliation (having health social security affiliation or not), and employment status based on job categories. We used health social security affiliation rather than health insurance alone, since it is related to formal employment and other individual social benefits that differ from specific healthcare programs aimed to provide health services to the most vulnerable

populations in Mexico. Employment status was defined from the original job category variable in the mortality records that included the categories "Does not work" and "Looking for a job", which were combined into one single category "Not employed". However, this categorization does not rule out the possibility that part of the category "Does not work" can also include retired people, with a higher SES compared with those who are actually unemployed. The rest of the job categories were combined into the category "Employed". We also explored education-specific and job-specific associations. For the latter, given that the Mexican classification of occupations was updated in 2012, and specific occupations coded before 2012 were redistributed into new job categories in the most recent version, it was not possible to homogenize all job categories for the whole period of study. Thus, analyses of specific job categories were restricted to the period from 2013 to 2019. We compared if the effect size of PM<sub>2.5</sub> exposure differed by age group, and also by education, healthcare affiliation, and employment status, in the broad-category mortality outcomes, and total non-external mortality by following the methods described by Altman and Bland, 2003.

### **Environmental data**

We utilized daily PM<sub>2.5</sub> predictions with spatial resolution of 1×1 km from our recently developed model based on extreme gradient boosting (XGBoost), and inverse-distance weighting (IDW) that uses aerosol optical depth data, meteorology, and land-use variables to predict daily mean PM<sub>2.5</sub> for the Mexico City Metropolitan Area (Gutiérrez-Avila et al., 2022). Daily mean air temperature with the same 1x1 km resolution was obtained from our satellite-based land surface temperature model for Central Mexico (Gutiérrez-Avila et al., 2021). We restricted our analyses to the spatial domain with available predictions for both PM<sub>2.5</sub> and temperature exposures over the Mexico City Metropolitan Area. The study area was composed of 667 sub-municipal geographic areas defined as "localities" by the Instituto Nacional de Estadística y Geografía. Localities correspond to the third level of subnational division in Mexico, after states and municipalities (Instituto Nacional de Geografía y Estadística, 2010). The total population in the Mexico City Metropolitan Area in 2010 was 20,116,842 (Secretaría de Desarrollo Agrario, Territorial y Urbano, 2018). For the same year, the population living in the localities included in our study region was 19,711,516 inhabitants, ~96% of the total Mexico City Metropolitan Area. Urban localities had a population of 19,346,527 (median of 8,523 inhabitants; range of 910 to 1,815,786) inhabitants, and rural localities had a total population of 364,989 (median of 582 inhabitants; range 1 to 5,135) inhabitants. We assigned exposures to PM<sub>2.5</sub> and temperature for all mortality records at the locality level. For 255 urban localities with census-provided polygons (median land area of 2.9 km<sup>2</sup>; range 0.2 to 129.4 km<sup>2</sup>), we estimated daily exposures using population-weighted aggregation with population density from the Gridded Population of the World (GPWv4) ~1-km raster cells (Center for International Earth Science

Information Network-CIESIN-Columbia University, 2016), which utilizes data from the 2010 Mexican Census, and the R package *exactextractr* (Baston, 2020). For 412 rural localities, the census only assigned points (rather than enclosing polygons), and for this subset of rural localities we assigned exposure to PM<sub>2.5</sub> and temperature using the 1x1 km grid cell containing the corresponding census-assigned points.

### Statistical Analysis

We estimated the association between short-term exposure to PM<sub>2.5</sub> with mortality using a time-stratified case-crossover design (Levy et al., 2001). Case days were defined as the date of death, and control days (3 to 4 days per month) were chosen on the same day of week as the case day within the same month and year. This approach controls for potential confounding effects by day of week, seasonal patterns and long-term time trends (Janes et al., 2005). Time-invariant covariates are controlled by design (within-person matching), and are not considered to be confounders; thus, daily variations in exposure to PM<sub>2.5</sub> and temperature on the case day are compared to exposures on control days. All models were adjusted for non-linear effects of temperature with quadratic b-splines (4 degrees of freedom), and equally-spaced knots in the space of this predictor (Gasparrini, 2021). The odds ratios for all dependent variables associated with short-term exposure to PM<sub>2.5</sub> were estimated with linear terms in stratified Cox proportional hazards models, equivalent to conditional logistic regression. We explored non-linearities in the concentration-response relationships between PM<sub>2.5</sub> with all broad-category mortality outcomes by comparing the fit of generalized additive models (GAMs) including parsimonious non-linear (penalized thin-plate splines) PM<sub>2.5</sub> terms (lag<sub>0</sub> to lag<sub>6</sub>) with the fit of linear PM<sub>2.5</sub> terms using likelihood ratio tests. To address the delayed effects of exposure to PM<sub>2.5</sub> on all mortality outcomes we included distributed lag terms up to 6 days before the case day (seven terms for lags 0 to 6) to estimate mortality risks (Gasparrini et al., 2010). For ease of interpretation, all odds ratios were converted to percentage increase per 10 µg/m<sup>3</sup> higher PM<sub>2.5</sub>.

All analyses were performed in R version 4.2.1 (R Core Team, 2020) with packages: *data.table* (Dowle et al., 2019), *survival* (Therneau and Lumley, 2015), *dlm* (Gasparrini, 2011), and *mgcv* (Wood SN, 2015). Because the records used in all analyses were publicly available, the PI previously received a determination of exempt human research: 45 CFR 46. 101(b) (Category 4) from the Institutional Review Board of the Icahn School of Medicine at Mount Sinai.

### Results

The total number of non-accidental deaths ( $\geq 18$  years-old) analyzed from 2004 to 2019 in the Mexico City Metropolitan Area was 1,479,950. Table 1 describes the demographic characteristics of the study population.

**Table 1. Demographic characteristics of the analyzed deaths in people >18 years-old in the Mexico City Metropolitan Area from 2004-2019**

<b>Characteristic</b>	<b>Frequency</b>
All non-accidental deaths	1,479,950 (100%)
<b>Age group</b>	
Elderly 65+	927,795 (62.7%)
Adults 18-64	552,155 (37.3%)
<b>Sex</b>	
Men	751,804 (50.8%)
Women	728,146 (49.2%)
<b>Education</b>	
Primary School	731,468 (49.4%)
Mid-High School	340,172 (23%)
No education	191,426 (12.9%)
College	153,100 (10.3%)
Unspecified	63,784 (4.3%)
<b>Insurance</b>	
Health social security affiliation	928,678 (62.8%)
No health social security affiliation	496,491 (33.5%)
Unspecified	54,781 (3.7%)
<b>Job type</b>	
Does not apply before 2013	738,294 (49.9%)
Not employed	477,573 (32.3%)
Sales workers	53,918 (3.6%)
Unspecified	48,332 (3.3%)
Craft and related trades workers	47,834 (3.2%)
Professionals and technicians	44,423 (3%)
Plant and machine operators	21,727 (1.5%)
Agricultural, forestry and fishery	12,926 (0.9%)
Personal services	11,337 (0.8%)
Clerical support	9,581 (0.6%)
Elementary occupations	9,475 (0.6%)
Managers	4,530 (0.3%)
<b>PM<sub>2.5</sub></b>	
Mean (SD)	24.6 (10.8)
Median [Min, Max]	23.4 [0.200, 202]
<b>Tmean</b>	
Mean (SD)	16.6 (2.58)
Median [Min, Max]	16.8 [3.30, 24.1]

There were slightly more deaths in men than women, and deaths in the elderly ( $\geq 65$  years-old) accounted for ~63% of all deaths. Around 50% of the decedents attended only elementary school, 63% had health social security affiliation, and 32% were not working at the time of death. The five most common job categories in the mortality records for the period 2013-2019 were: not-employed or not working at the time of death, sales, unspecified occupations, craft and related trades, and professionals and technicians. Over the study period, the mean population-weighted exposure to PM<sub>2.5</sub> and temperature in the study region were 24.6 $\mu\text{g}/\text{m}^3$ , and 16.6° C, respectively.

Table 2 shows the total number of organ-system and cause-specific deaths ( $\geq 18$  years-old) analyzed over the study period. Circulatory system diseases were the most prevalent death causes, accounting for almost 30% of the total, followed by digestive (12%) and respiratory (10%) diseases.

**Table 2. Underlying cause-specific mortality (>18 years-old) in the Mexico City Metropolitan Area from 2004-2019**

<b>Disease</b>	<b>ICD code</b>	<b>Frequency</b>
All non-accidental deaths		1,479,950
<b>Disease group</b>		
Circulatory System	I00-I99	427,420
Digestive system	K00-K93	172,906
Respiratory system	J00-J99	142,866
Genitourinary	N00-N99	57,372
Nervous system	G00-G99	22,735
Intentional self-harm	X60-X84	10,021
Mental and behavioral	F00-F99	9,577
<b>Specific death cause</b>		
Acute ischemic heart disease	I20-I22, I24	212,201
Diseases of the liver	K70-K77	92,562
Chronic respiratory disease	J40-J47	67,062
Hypertensive diseases	I10-I15	53,675
Influenza and Pneumonia	J09-J18	52,972
Renal failure	N17-N19	30,787
Other forms of heart disease	I30-I52	29,516
Stroke hemorrhagic	I60-I62	28,412
Chronic ischemic heart disease	I25	21,313
Disorders of gallbladder, biliary tract, and pancreas	K80-K87	16,266
Diseases of esophagus, stomach and duodenum	K20-K31	12,322
Stroke ischemic	I63	10,684
Suicides	X6-X84	9,954
Pulmonary heart disease	I26-I28	6,792
Diseases of the arteries	I70-I79	6,299
Chronic rheumatic heart diseases	I05-I09	4,314
Extrapyramidal and movement disorders	G20-G26	3,531

Among the specific death causes, acute ischemic heart disease and hypertensive disease were the leading causes from the circulatory system. Diseases of the liver were the leading death causes from the digestive system, and chronic respiratory disease, influenza and pneumonia were the most common causes of death from the respiratory system (Table 2).

Results from our assessment of linearity in the concentration-response functions between short-term exposure to PM<sub>2.5</sub> with all broad-category mortality outcomes showed no significant differences in model's fit when including non-linear PM<sub>2.5</sub> exposures; therefore all our further results are based on linear concentration-response functions. Figure 1 shows associations between PM<sub>2.5</sub> exposure with the broad-category death causes analyzed over the study period. For consistency with previous studies, two-days (lag<sub>0</sub> + lag<sub>1</sub>) and one-week, (sum from lag<sub>0</sub> to lag<sub>6</sub>) cumulative associations with PM<sub>2.5</sub> are shown. Among the broad-category death causes, two-days exposure to PM<sub>2.5</sub> (lag<sub>01</sub>) was associated with total higher non-accidental mortality [adults = 0.72% (95% CI: 0.26%–1.19%), elderly = 1.04% (95% CI: 0.68%–1.40%)], with sex-specific associations for men of 0.84% (95% CI: 0.44%–1.24%) and women of 1.00% (95% CI: 0.60%–1.41%), cardiovascular mortality [adults = 1.03% (95% CI: -0.3%–2.10%), elderly = 1.01% (95% CI: 0.41%–1.62%)], cerebrovascular mortality [adults = 2.84% (95% CI: 0.52%–5.21%)], respiratory mortality [adults = 2.51% (95% CI: 0.63%–4.44%), elderly = 2.02% (95% CI: 1.00%–3.04%)] and digestive mortality [elderly = 1.16% (95% CI: -0.01%–2.35%)]. Cumulative associations over one week (lag<sub>06</sub>) were in general larger compared with associations for lag<sub>01</sub> for most mortality outcomes; with the largest associations observed for cerebrovascular [adults = 3.65% (95% CI: 0.49%–6.91%)], respiratory [adults = 2.61% (95% CI: 0.04%–5.25%), elderly = 3.46% (95% CI: 2.05%–4.90%)], and genitourinary [adults = 3.35% (95% CI: 0.11%–6.69%), elderly = 2.30% (95% CI: -0.17%–4.83%)] mortality. Our test to identify differences in effect sizes by age and sex in all broad-categories of death did not suggest effect modification from these variables in the associations between PM<sub>2.5</sub> and mortality. Results for specific single lag (lag<sub>0</sub> ... lag<sub>6</sub>) and cumulative associations (lag<sub>01</sub> ... lag<sub>06</sub>) for all broad-categories of mortality and PM<sub>2.5</sub> are reported in Table S1.






Figure 1. Cumulative percentage increase (%) and 95% CI for broad-category death causes by age group (adults  $\geq 18$ -64 years-old and elderly  $\geq 65$  years-old) per  $10 \mu\text{g}/\text{m}^3$  increase of  $\text{PM}_{2.5}$  over two days ( $\text{lag}_{01}$ ) and one week ( $\text{lag}_{06}$ ) in the Mexico City Metropolitan Area for 2004-2019.

Associations between  $\text{PM}_{2.5}$  exposure and cause-specific mortality for  $\text{lag}_{01}$  and  $\text{lag}_{06}$  are shown in Figure 2. Cumulative exposure to  $\text{PM}_{2.5}$  over two days ( $\text{lag}_{01}$ ) was associated with mortality from hypertensive disease [1.14% (95%CI: -0.33%–2.62%)], acute ischemic heart disease [1.21% (95%CI: 0.47%–1.96%)], influenza and pneumonia [2.97% (95%CI: 1.48%–4.48%)], and chronic respiratory disease [1.51% (95%CI: 0.23%–2.81%)]. Associations for cumulative exposure to  $\text{PM}_{2.5}$  over one week ( $\text{lag}_{06}$ ) were observed for hypertensive disease [2.28% (95%CI: 0.26%–4.33%)], acute ischemic heart disease [1.61% (95%CI: 0.59%–2.64%)], other forms of heart disease [2.39% (95%CI: -0.35%–5.20%)], hemorrhagic stroke [3.63% (95%CI: 0.79%–6.55%)], influenza and pneumonia [4.91% (95%CI: 2.84%–7.02%)], chronic respiratory disease [2.49% (95%CI: 0.71%–4.31%)], diseases of the liver [1.85% (95%CI: 0.31%–3.41%)], and renal failure [3.48% (95%CI: 0.79%–6.24%)].




Figure 2. Cumulative percentage increase (%) and 95% CI for cause-specific mortality by age group (adults  $\geq 18$  years-old and elderly  $\geq 65$  years-old) per  $10 \mu\text{g}/\text{m}^3$  increase of  $\text{PM}_{2.5}$  over two ( $\text{lag}_{01}$ ) days and one week ( $\text{lag}_{06}$ ) in the Mexico City Metropolitan Area for 2004-2019.

Cumulative associations from distributed lags can capture the impact of multiple days after initial exposure to  $\text{PM}_{2.5}$  on mortality. In general, consistent associations between short-term exposure to  $\text{PM}_{2.5}$  with all-cause (i.e. total non-accidental) or broad-category death causes (mostly from cardiovascular and respiratory diseases) have been observed during the first week of exposure across study sites. Therefore, communicating results for  $\text{lag}_{01}$  and  $\text{lag}_{06}$ , has become the standard, as the evidence shows immediate ( $\text{lag}_{01}$ ) and more prolonged ( $\text{lag}_{06}$ ) effects on cardiovascular, and respiratory mortality, respectively. That said, the onset and duration of the mechanisms of damage leading to death can vary for other health outcomes, with mortality from  $\text{PM}_{2.5}$  exposure occurring with a delay of a few days (World Health Organization. Regional Office for Europe, 2021). Little has been documented about the timing of when the largest single lag effects of  $\text{PM}_{2.5}$  on cause-specific mortality occur, and these “peaks” in mortality can be missed when reporting associations only for  $\text{lag}_{01}$ , or  $\text{lag}_{06}$ , and this information can provide meaningful information for health professionals. Figure 3 shows the largest single lag associations for two cause-specific mortality outcomes (hemorrhagic stroke and influenza and pneumonia); and to complement this information, results for all single lag and cumulative associations for all cause-specific mortality outcomes are reported in Table S2. Same-day exposure to  $\text{PM}_{2.5}$  ( $\text{lag}_0$ ) showed the largest single lag associations with diseases of the arteries [4.62% (95%CI: 0.73%–8.67%)], hemorrhagic stroke [2.98% (95%CI: 1.09%–4.90%)], hypertensive disease [1.34% (95%CI: 0.02%–2.68%)], and acute ischemic heart disease [0.64% (95%CI: -0.02%–1.31%)].  $\text{Lag}_1$  showed the largest association with diseases of the esophagus, stomach and duodenum [3.40% (95%CI: 0.26%–6.64%)], and chronic respiratory disease [1.72% (95%CI: 0.44%–3.02%)].  $\text{Lag}_4$  had the largest effects for chronic

heart disease [3.74% (95%CI: 1.37%–6.16%)], and ischemic stroke [3.08% (95%CI: -0.22%–6.50%)], and lag<sub>6</sub> for influenza and pneumonia [2.19% (95%CI: 0.88%–3.52%)].




Figure 3. Single lag associations as percent increase (%) and 95% CI for hemorrhagic-stroke and influenza and pneumonia per 10  $\mu\text{g}/\text{m}^3$  increase of  $\text{PM}_{2.5}$

Figure 4 shows the associations between exposure to  $\text{PM}_{2.5}$  with non-accidental mortality in adults ( $\geq 18$ -64 years-old) and the elderly ( $\geq 65$  years-old), stratified by education level (having or not basic education), health insurance affiliation (having or not health social security affiliation), and employment status (either being working or not when the person died). Associations for lag<sub>06</sub> in those with no education were 2.70% (95%CI: 0.18%–5.28%), and 2.33% (95%CI: 1.16%–3.52%) for adults and the elderly, respectively. While for those having at least basic education the associations were 1.64% (95%CI: 0.97%–2.31%) for adults, and 1.55% (95%CI: 1.00%–2.10%)], for the elderly. For the same exposure window, mortality risk in adults without health social security affiliation was 2.19% (95%CI: 1.22%–3.16%), and 1.75% (95%CI: 0.82%–2.68%) for the elderly. Adults with health insurance showed a 1.22% (95%CI: 0.35%–2.09%) mortality risk, and the elderly 1.74% (95%CI: 1.15%–2.34%). Results by employment status for the period from 2013-2019, showed that associations for adults not working at the time of death were 1.68% (95%CI: 0.36%–3.02%), and 1.83% (95%CI: 1.03%–2.63%) for the

elderly; compared to those who were working with mortality risks of 0.90% (95% CI: -0.48%–2.29%) and 1.70% (95% CI: 0.23%–3.20%), for adults and the elderly, respectively. Our test of heterogeneity in effect size between SES related strata, does not suggest effect modification between SES strata. Figure S1, shows job-specific associations with PM<sub>2.5</sub> exposure for the same period of time.




Figure 4. Cumulative percentage increase (%) and 95% CI for non-accidental mortality by education, health insurance affiliation, and employment status (adults 18-64 years-old, elderly  $\geq 65$  years-old) per 10  $\mu\text{g}/\text{m}^3$  increase in PM<sub>2.5</sub> over two (lag<sub>01</sub>) days and one week (lag<sub>06</sub>) in the Mexico City Metropolitan Area for 2004-2019. Associations for employment status are for the period from 2013-2019.

## Discussion

This study presents a detailed analysis on the associations between short-term exposure to PM<sub>2.5</sub> with total non-external, broad-catergory (organ-system) and cause-specific mortality outcomes; and the role that age, sex, and potential SES-related effect modifiers play in such associations in the Mexico City Metropolitan Area. Our research involved the use of state-of-the-art methods to estimate daily exposure to PM<sub>2.5</sub>, using highly spatially-resolved satellite-based PM<sub>2.5</sub> predictions. With our PM<sub>2.5</sub> predictions, we were able to assign daily exposure to PM<sub>2.5</sub> to all case and control days at the sub-municipality level; leveraging the smallest geographic identifier for place of residence included in the Mexican mortality records. To our knowledge, this is the most comprehensive analysis in Central Mexico in terms of geographic extent, amount of mortality outcomes analyzed, and assessment of potential effect modifiers in these associations using individual mortality records.

Our results support previous evidence on the associations between exposure to PM<sub>2.5</sub> with increased risks of non-accidental, cardiovascular, cerebrovascular and respiratory mortality in the Mexico City region (Borja-Aburto et al., 1998; Gutiérrez-Avila et al., 2018). We also found associations between PM<sub>2.5</sub> exposure with causes of death not previously reported in the Mexico City Metropolitan Area, such as digestive and genitourinary outcomes. Our results for non-accidental, cardiovascular and respiratory mortality in adults, the three more consistent broad-category mortality outcomes linked to PM<sub>2.5</sub> exposure for lag<sub>01</sub>, were larger than the associations reported by Li et al. (2018) for non-accidental [0.25% (95% CI: 0.11%–0.38%)], circulatory system [0.39% (95% CI: 0.21%–0.58%)], and respiratory [0.43%; 95% CI: 0.05%–0.78%] related mortality, and also larger than the results from Xu et al. (2020) for all-cause [0.13% (95% CI: 0.01%–0.27%)], cardiovascular [0.02% (95% CI: -0.17%–0.21%)], and respiratory [0.81% (95% CI: 0.39%–0.96%)] mortality for Beijing. Our results for the same health outcomes were similar to those reported in the meta-analyses of Atkinson et al. (2014) for all-cause [1.04% (95% CI: 0.52%–1.56%)], cardiovascular [0.84% (95% CI: 0.41%–1.28%)], and respiratory [1.51% (95% CI: 1.01%–2.01%)] mortality (Atkinson et al., 2014). Orellano et al. (2020) also reported similar associations for all-cause [0.65% (95% CI: 0.44%–0.86%)] and cardiovascular [0.92% (95% CI: 0.61%–1.23%)] mortality, but their associations for respiratory [0.73% (95% CI: 0.29%–1.16%)] and cerebrovascular [0.72% (95% CI: 0.12%–1.32%)] mortality were lower compared to our results (Orellano et al., 2020). However, it is possible that the selection of heterogeneous effect estimates from different lags structures that were combined in the meta-analyses of Atkinson et al. (2014), and Orellano et al. (2020) could have induced some biases. Differential toxicity related to PM<sub>2.5</sub> composition, variation in the underlying health status of the population, and distribution of potential effect modifiers between study regions can also play a role in the differences reported in the literature. Also, it has been suggested that at higher concentrations of PM<sub>2.5</sub> like those observed in Asian cities, the slopes in the concentration-response function between PM<sub>2.5</sub> and mortality flattens out (supralinear exposure-response curve) for cardiorespiratory outcomes, which might also explain the differences in effect estimates between study regions (Liu et al., 2019). Future research efforts exploring the shape of the concentration response functions for mortality outcomes other than cardiorespiratory outcomes could aid to understand the differences in mortality risks across study regions.

Among the cause-specific mortality associations with PM<sub>2.5</sub>, those related to the circulatory system (e.g. diseases of the arteries, hypertensive disease, acute ischemic heart disease, and stroke) were the most clear, with the largest associations on the same day of exposure (lag<sub>0</sub> in Table S2). Li et al. (2018) also reported consistent associations between PM<sub>2.5</sub> with mortality outcomes from the circulatory and respiratory systems in Beijing. However, we found larger associations compared to those reported by Li et al. (2018) for ischemic heart disease (0.46%; 95% CI: 0.19%–0.72%), influenza and pneumonia

(0.35%; 95% CI: -0.32%–1.03%), and chronic respiratory disease (0.62%; 95% CI: 0.08%–1.16%). Li et al. (2018) also reported associations between PM<sub>2.5</sub> and sub-specific causes of ischemic heart disease such as: acute ischemic heart disease (ICD-10 codes: I20-I22, I24), acute myocardial infarction (ICD-10 codes: I21-I22), and myocardial infarction (ICD-10 codes: I21-I23). We did not explore such subtypes of ischemic heart disease, as Li et al. (2018) did, because the grouping of such health outcomes involves the overlap of several ICD-10 codes; while we only focused on reporting associations for mutually exclusive health outcomes. We found larger associations than those reported by Xu et al. (2020) for ischemic heart disease (-0.06%; 95% CI: -0.33%–0.22%), and chronic respiratory disease (0.96%; 95% CI: 0.35%–1.57%) also in Beijing. Our results for specific cardiovascular and cerebrovascular outcomes were larger than those reported by Chen et al. (2018) for 30 Chinese counties (Chen et al., 2018). Intentional-self harm mortality was not associated with PM<sub>2.5</sub> exposure in our study, which is consistent with results of Astudillo-García et al. (2019) from a time-series study focused on suicides and exposure to different air pollutants including PM<sub>2.5</sub> in Mexico City (Astudillo-García et al., 2019) .

Contrary to the results from Li et al. (2018), we did not find associations between short-term exposure to PM<sub>2.5</sub> with acute mortality outcomes from the nervous system (e.g. extrapyramidal and movement disorders), however we observed positive associations between cumulative exposure to PM<sub>2.5</sub> with digestive (diseases of the liver) and genitourinary (renal failure) mortality. Exposure to PM<sub>2.5</sub> has been associated with increased serum levels of hepatic enzymes, such as  $\gamma$ -glutamyltranspeptidase (GTP), and alanine transaminase (ALT), which are biomarkers of liver damage. PM<sub>2.5</sub> can induce systemic oxidative stress and inflammation when deposited in the airways, and when swallowed PM<sub>2.5</sub> are removed from the airways by mucociliary clearance leading to gastrointestinal exposure with the largest internal dose and adverse effects observed in the liver (Kim et al., 2014; Tavera Busso et al., 2020). On the other hand, recent evidence has shown that PM<sub>2.5</sub> is related to renal injury and increases the risk of nephropathy, as well as increased risk of first hospital admission from kidney and total urinary system diseases (Lee et al., 2022; Rasking et al., 2022). PM<sub>2.5</sub> can unbalance the kidney function by accumulation in the kidney tissue, endothelial dysfunction, abnormal renin-angiotensin system, and immune complex deposition. The mechanisms of damage from PM<sub>2.5</sub> to the kidney involve inflammation, oxidative stress, apoptosis, DNA damage, and autophagy (Xu et al., 2022).

The most recent version of the EPA's Integrated Science Assessment (ISA) for Particulate Matter (2019) has defined the associations between short-term exposure to PM<sub>2.5</sub> with cardiovascular and respiratory mortality as causal and likely causal, respectively; but less evidence is available for making conclusions about other causes of death (EPA, 2019). Although our results suggest that short-term exposure to PM<sub>2.5</sub>

may trigger cause-specific mortality beyond the circulatory and respiratory systems, more epidemiological evidence is needed to understand the links between PM<sub>2.5</sub> exposure with the different specific death causes evaluated in our study. Emerging evidence about the adverse effects of PM<sub>2.5</sub> exposure on multiple prevalent but rarely studied causes of hospital admissions (Wei et al., 2019) adds to the body of epidemiologic evidence that supports our findings, and it opens the possibility to replicate such results in different locations with ongoing or future health studies.

Information on the timing when health effects occur after initial exposure to PM<sub>2.5</sub>, or any other air pollutant, is of importance for public health administrators and pollution control managers. For instance, identifying the lag days over which health effects are observed can inform the development of air quality standards, risk communication tools (e.g. air quality indices), quantification of health impacts, mitigation strategies (e.g. planning and allocation of clinical resources), and the design of emission control measures. Our results are consistent with the epidemiologic evidence showing that short-term exposure to PM<sub>2.5</sub> has an immediate effect on non-accidental and cardiovascular mortality, and a more prolonged effect on respiratory mortality (EPA, 2019). However, when considering specific death causes (Table S2) within the broad-category mortality outcomes, there seems to be more variability about the timing when the largest (single and cumulative) associations are observed. For instance, chronic ischemic heart disease and ischemic stroke showed the largest associations with PM<sub>2.5</sub> on lag<sub>4</sub>, while the rest of the death causes from the circulatory system showed the largest association on lag<sub>0</sub> and lag<sub>1</sub>. For respiratory mortality, our results are within the range of observed associations, with chronic respiratory disease showing the largest association with PM<sub>2.5</sub> on lag<sub>1</sub>, while the association with influenza and pneumonia remained high at lag<sub>6</sub>. On the other hand, it is known that the strength of the relationship between exposure to PM<sub>2.5</sub> and health effects varies depending on the exposure duration i.e., short- or long-term exposure, but less is known about the strength of this relationship for sub-daily exposures e.g. hourly peak exposures. Few epidemiologic studies have focused on the associations between sub-daily exposures to PM<sub>2.5</sub> with morbidity and mortality outcomes (Lin et al., 2017; Madsen et al., 2012). Results from such investigations are inconclusive on whether sub-daily exposures pose a greater risk of mortality, or not, compared to daily averaged exposures; being possible that the sub-daily metrics of PM<sub>2.5</sub> employed for exposure assessment in previous studies had a limited spatiotemporal variability (EPA, 2019). In our research, we only focused on exposures to daily mean PM<sub>2.5</sub> concentrations assuming that the same averaging time could have the same relevance for all mortality outcomes. However, we can not rule out the possibility that there might be averaging times (sub-daily, or sub-chronic exposures) that could be of greater relevance for the different death causes analyzed in our research.

It is known that subgroups of the population with lower SES indicators may have higher prevalence of health disorders and comorbidities, lower living standards, and higher exposure to air pollutants (e.g. living closer to a highway or lack of green spaces) (Shavers, 2007; Xu et al., 2020). We used three indicators of SES, namely education, health social security affiliation, and employment status to assess their roles as potential effect modifiers in the  $PM_{2.5}$ -mortality relationship. Although in general point estimates for the lower SES related strata (lower education level, not having health social security affiliation, and not being working at the time of death) seemed larger than those in the higher SES indicators; we did not find evidence of effect modification in the  $PM_{2.5}$ -mortality associations (analyzed on the log scale) between SES strata (Altman and Bland, 2003). In Mexico there is limited evidence about the role of education in the association between short-term exposure to  $PM_{2.5}$  and mortality, however O'Neill et al. (2008) did not observe evidence of heterogeneity in the association between short-term exposure to  $PM_{10}$  and mortality across education levels (O'Neill et al., 2008). Health insurance access can modify the susceptibility to detrimental effects of environmental stressors, including  $PM_{2.5}$ , by providing access to medications, supplements and overall preventive services (Woolhandler and Himmelstein, 2017). Our results did not show significant effect modification between strata of health social security affiliation. Also, our results for employment status did not suggest greater risk of mortality for those in the not-employed strata compared with people in the employed category. The evidence on the effects of working status on health suggests that unemployment is related to poor health (e.g. greater risk of mental illness, physical complaints, and an increased risk for coronary heart diseases) and early mortality (Kim and von dem Knesebeck, 2015). Although our models were further stratified by age group to reduce its potential confounding effects, our results should be carefully interpreted, since there was not a well-defined measure of unemployment excluding all the economically inactive (Clemens et al., 2015). Also, we did not have information on the health status of those in the “non-working” group before death, so it is not possible to determine if a deteriorating health status led to a “non-working” condition (healthy worker effect). When evaluating the role of specific job types in the association between short-term exposure to  $PM_{2.5}$  and mortality, the lack of consistency in the codification of specific job categories throughout the period of study led us to evaluate these associations for a shorter time period (2013-2019), likely reducing power to identify associations by individual job categories (Figure S1). Two major problems with the use of death certificates for classifying employment include ambiguity about whether unemployed persons are actually retired or if a prior occupation is reported for retired decedents and thus may not represent their recent exposures.



Other limitations in our study could be related to the inaccurate classification of death causes for some diseases. In general, inaccurate classification of death causes in Mexico is considered low (Naghavi et al., 2010). However, errors might exist in the classification of some mortality outcomes, including those from the circulatory system, which have been recognized among the most affected by the use of “garbage codes”, i.e. those that are not underlying causes of death such as heart failure, but codes that define poorly-specified diagnoses not clearly identifying a death cause. For Mexico City, we have previously reported that the use of garbage codes tends to bias effect estimates of cardiovascular outcomes to the null, and that proportional redistribution of garbage codes into actual death causes may help to reduce such bias (Gutiérrez-Avila et al., 2018). Also, there are contributions to exposure measurement error that accumulate from the combination of prediction model error and the use of locality of residence to represent all relevant time-activity, which may bias effect estimates (albeit less than the use of a single city-wide exposure time series). Finally, the study of effect modification by cause-specific mortality, age group, sex, and SES indicators can be sensitive to analytic methods. Although our stratified analyses have been the standard in case-crossover and time-series studies, the sparse numbers in some of the strata could have reduced statistical power to detect associations.

## **Conclusion**

Our findings support the growing evidence that short-term exposure to PM<sub>2.5</sub> may trigger cause-specific mortality beyond cardiovascular and respiratory outcomes, including mortality from diseases of the digestive and genitourinary systems. This evidence can be used to inform and update local environmental health policies, and to more comprehensively assess the benefits of interventions aimed to reduce PM<sub>2.5</sub> pollution in the Mexico City Metropolitan Area. Our findings suggest that short-term exposure to PM<sub>2.5</sub> increases the risk of acute mortality in the Mexico City Metropolitan Area, regardless of age groups, and SES characteristics.

### **Funding information**

This work was supported by grants from the National Institutes of Health (NIH): R01ES031295 and P30ES023515.

### **Ethical Approval**

Because the records used in all analyses were already publicly available, the senior author (ACJ) received a determination of exempt human research [45 CFR 46. 101(b) (Category 4)] from the Institutional Review Board of the Icahn School of Medicine at Mount Sinai.

### **Competing interests**

The authors declare that they have no conflicts of interest regarding this study.

.

## References

- Alexeeff, S.E., Schwartz, J., Kloog, I., Chudnovsky, A., Koutrakis, P., Coull, B.A., 2015. Consequences of kriging and land use regression for PM<sub>2.5</sub> predictions in epidemiologic analyses: insights into spatial variability using high-resolution satellite data. *J. Expo. Sci. Environ. Epidemiol.* 25, 138–144.
- Altman, D.G., Bland, J.M., 2003. Interaction revisited: the difference between two estimates. *BMJ* 326, 219.
- Astudillo-García, C.I., Rodríguez-Villamizar, L.A., Cortez-Lugo, M., Cruz-De la Cruz, J.C., Fernández-Niño, J.A., 2019. Air Pollution and Suicide in Mexico City: A Time Series Analysis, 2000–2016. *Int. J. Environ. Res. Public Health* 16, 2971.
- Atkinson, R.W., Kang, S., Anderson, H.R., Mills, I.C., Walton, H.A., 2014. Epidemiological time series studies of PM<sub>2.5</sub> and daily mortality and hospital admissions: a systematic review and meta-analysis. *Thorax* 69, 660–665.
- Baston, D., 2020. exactextractr: Fast extraction from raster datasets using polygons.
- Berman, J.D., Burkhardt, J., Bayham, J., Carter, E., Wilson, A., 2019. Acute Air Pollution Exposure and the Risk of Violent Behavior in the United States. *Epidemiology* 30, 799–806.
- Borja-Aburto, V.H., Castillejos, M., Gold, D.R., Bierzwinski, S., Loomis, D., 1998. Mortality and Ambient Fine Particles in Southwest Mexico City, 1993-1995. *Environmental Health Perspectives*. <https://doi.org/10.2307/3434129>
- Bowe, B., Xie, Y., Yan, Y., Al-Aly, Z., 2019. Burden of Cause-Specific Mortality Associated With PM<sub>2.5</sub> Air Pollution in the United States. *JAMA Netw Open* 2, e1915834.
- Center for International Earth Science Information Network-CIESIN-Columbia University, 2016. Gridded population of the world, version 4 (GPWv4): population density.
- Chen, C., Zhu, P., Lan, L., Zhou, L., Liu, R., Sun, Q., Ban, J., Wang, W., Xu, D., Li, T., 2018. Short-term exposures to PM<sub>2.5</sub> and cause-specific mortality of cardiovascular health in China. *Environ. Res.* 161, 188–194.
- Clemens, T., Popham, F., Boyle, P., 2015. What is the effect of unemployment on all-cause mortality? A cohort study using propensity score matching. *Eur. J. Public Health* 25, 115–121.
- Davoudi, M., Barjasteh-Askari, F., Amini, H., Lester, D., Mahvi, A.H., Ghavami, V., Rezvani Ghalhari, M., 2021. Association of suicide with short-term exposure to air pollution at different lag times: A systematic review and meta-analysis. *Sci. Total Environ.* 771, 144882.
- Dowle, Srinivasan, Gorecki, 2019. data.table: Extension of “data.frame”. R Package Version 1.12.8.
- EPA, 2019. Integrated science assessment (ISA) for particulate matter (final report, Dec 2019). US Environ. Prot. Agency.
- Franklin, M., Zeka, A., Schwartz, J., 2006. Association between PM<sub>2.5</sub> and all-cause and specific-cause mortality in 27 US communities. *J. Expo. Sci. Environ. Epidemiol.* 17, 279.
- Gasparrini, A., 2021. Distributed lag linear and non-linear models for time series data. London School of Hygiene & Tropical Medicine, UK.
- Gasparrini, A., 2011. Distributed Lag Linear and Non-Linear Models in R: The Package dlnm. *J. Stat. Softw.* 43, 1–20.
- Gasparrini, A., Armstrong, B., Kenward, M.G., 2010. Distributed lag non-linear models. *Stat. Med.* 29, 2224–2234.
- Gouveia, N., Kephart, J.L., Dronova, I., McClure, L., Granados, J.T., Betancourt, R.M., O’Ryan, A.C., Texcalac-Sangrador, J.L., Martinez-Folgar, K., Rodriguez, D., Diez-Roux, A.V., 2021. Ambient fine particulate matter in Latin American cities: Levels, population exposure, and associated urban factors. *Sci. Total Environ.* 772, 145035.
- Gutiérrez-Avila, I., Arfer, K.B., Carrión, D., Rush, J., Kloog, I., Naeger, A.R., Grutter, M., Páramo-Figueroa, V.H., Riojas-Rodríguez, H., Just, A.C., 2022. Prediction of daily mean and one-hour maximum PM<sub>2.5</sub> concentrations and applications in Central Mexico using satellite-based machine-

- learning models. *J. Expo. Sci. Environ. Epidemiol.* <https://doi.org/10.1038/s41370-022-00471-4>
- Gutiérrez-Avila, I., Arfer, K.B., Wong, S., Rush, J., Kloog, I., Just, A.C., 2021. A spatiotemporal reconstruction of daily ambient temperature using satellite data in the Megalopolis of Central Mexico from 2003 to 2019. *Int. J. Climatol.* 41, 4095–4111.
- Gutiérrez-Avila, I., Rojas-Bracho, L., Riojas-Rodríguez, H., Kloog, I., Just, A.C., Rothenberg, S.J., 2018. Cardiovascular and Cerebrovascular Mortality Associated With Acute Exposure to PM 2.5 in Mexico City. *Stroke.* <https://doi.org/10.1161/strokeaha.118.021034>
- Heo, S., Son, J.-Y., Lim, C.C., Fong, K.C., Choi, H.M., Hernandez-Ramirez, R.U., Nyhan, K., Dhillon, P.K., Kapoor, S., Prabhakaran, D., Spiegelman, D., Bell, M.L., 2022. Effect modification by sex for associations of fine particulate matter (PM<sub>2.5</sub>) with cardiovascular mortality, hospitalization, and emergency room visits: systematic review and meta-analysis. *Environmental Research Letters.* <https://doi.org/10.1088/1748-9326/ac6cfb>
- Instituto Nacional de Geografía y Estadística, 2010. *Manual de Cartografía Censal.*
- Janes, H., Sheppard, L., Lumley, T., 2005. Case-crossover analyses of air pollution exposure data: referent selection strategies and their implications for bias. *Epidemiology* 16, 717–726.
- Keswani Anjeni, Akselrod Hana, Anenberg Susan C., 2022. Health and Clinical Impacts of Air Pollution and Linkages with Climate Change. *NEJM Evidence* 1, EVIDra2200068.
- Kim, J.W., Park, S., Lim, C.W., Lee, K., Kim, B., 2014. The role of air pollutants in initiating liver disease. *Toxicol. Res.* 30, 65–70.
- Kim, T.J., von dem Knesebeck, O., 2015. Is an insecure job better for health than having no job at all? A systematic review of studies investigating the health-related risks of both job insecurity and unemployment. *BMC Public Health.* <https://doi.org/10.1186/s12889-015-2313-1>
- Lee, W., Wu, X., Heo, S., Fong, K.C., Son, J.-Y., Benjamin Sabath, M., Braun, D., Park, J.Y., Kim, Y.C., Lee, J.P., Schwartz, J., Kim, H., Dominici, F., Bell, M., 2022. Associations between long term air pollution exposure and first hospital admission for kidney and total urinary system diseases in the US Medicare population: nationwide longitudinal cohort study. *BMJ Medicine* 1. <https://doi.org/10.1136/bmjmed-2021-000009>
- Levy, D., Lumley, T., Sheppard, L., Kaufman, J., Checkoway, H., 2001. Referent selection in case-crossover analyses of acute health effects of air pollution. *Epidemiology* 12, 186–192.
- Lin, H., Liu, T., Xiao, J., Zeng, W., Guo, L., Li, X., Xu, Y., Zhang, Y., Chang, J.J., Vaughn, M.G., Qian, Z.M., Ma, W., 2017. Hourly peak PM<sub>2.5</sub> concentration associated with increased cardiovascular mortality in Guangzhou, China. *J. Expo. Sci. Environ. Epidemiol.* 27, 333–338.
- Li, T., Yan, M., Sun, Q., Anderson, G.B., 2018. Mortality risks from a spectrum of causes associated with wide-ranging exposure to fine particulate matter: A case-crossover study in Beijing, China. *Environ. Int.* 111, 52–59.
- Liu, C., Chen, R., Sera, F., Vicedo-Cabrera, A.M., Guo, Y., Tong, S., Coelho, M.S.Z.S., Saldiva, P.H.N., Lavigne, E., Matus, P., Valdes Ortega, N., Osorio Garcia, S., Pascal, M., Stafoggia, M., Scortichini, M., Hashizume, M., Honda, Y., Hurtado-Díaz, M., Cruz, J., Nunes, B., Teixeira, J.P., Kim, H., Tobias, A., Íñiguez, C., Forsberg, B., Åström, C., Ragettli, M.S., Guo, Y.-L., Chen, B.-Y., Bell, M.L., Wright, C.Y., Scovronick, N., Garland, R.M., Milojevic, A., Kyselý, J., Urban, A., Orru, H., Indermitte, E., Jaakkola, J.J.K., Rytí, N.R.I., Katsouyanni, K., Analitis, A., Zanobetti, A., Schwartz, J., Chen, J., Wu, T., Cohen, A., Gasparrini, A., Kan, H., 2019. Ambient Particulate Air Pollution and Daily Mortality in 652 Cities. *N. Engl. J. Med.* 381, 705–715.
- Madsen, C., Rosland, P., Hoff, D.A., Nystad, W., Nafstad, P., Naess, O.E., 2012. The short-term effect of 24-h average and peak air pollution on mortality in Oslo, Norway. *Eur. J. Epidemiol.* 27, 717–727.
- Naghavi, M., Makela, S., Foreman, K., O'Brien, J., Pourmalek, F., Lozano, R., 2010. Algorithms for enhancing public health utility of national causes-of-death data. *Popul. Health Metr.* 8, 9.
- O'Neill, M.S., Bell, M.L., Ranjit, N., Cifuentes, L.A., Loomis, D., Gouveia, N., Borja-Aburto, V.H., 2008. Air pollution and mortality in Latin America: the role of education. *Epidemiology* 19, 810–819.

- Orellano, P., Reynoso, J., Quaranta, N., Bardach, A., Ciapponi, A., 2020. Short-term exposure to particulate matter (PM<sub>10</sub> and PM<sub>2.5</sub>), nitrogen dioxide (NO<sub>2</sub>), and ozone (O<sub>3</sub>) and all-cause and cause-specific mortality: Systematic review and meta-analysis. *Environment International*. <https://doi.org/10.1016/j.envint.2020.105876>
- Rasking, L., Vanbrabant, K., Bové, H., Plusquin, M., De Vusser, K., Roels, H.A., Nawrot, T.S., 2022. Adverse Effects of fine particulate matter on human kidney functioning: a systematic review. *Environ. Health* 21, 24.
- R Core Team, 2020. R: A Language and Environment for Statistical Computing.
- Renzi, M., Marchetti, S., De' Donato, F., Pappagallo, M., Scortichini, M., Davoli, M., Frova, L., Michelozzi, P., Stafoggia, M., 2021. Acute Effects of Particulate Matter on All-Cause Mortality in Urban, Rural, and Suburban Areas, Italy. *Int. J. Environ. Res. Public Health* 18. <https://doi.org/10.3390/ijerph182412895>
- Samoli, E., Butland, B.K., Rodopoulou, S., Atkinson, R.W., Barratt, B., Beevers, S.D., Beddows, A., Dimakopoulou, K., Schwartz, J.D., Yazdi, M.D., Katsouyanni, K., 2020. The impact of measurement error in modeled ambient particles exposures on health effect estimates in multilevel analysis: A simulation study. *Environ Epidemiol* 4, e094.
- Secretaría de Desarrollo Agrario, Territorial y Urbano, 2018. Delimitación de las zonas metropolitanas de México 2015.
- Shavers, V.L., 2007. Measurement of socioeconomic status in health disparities research. *J. Natl. Med. Assoc.* 99, 1013–1023.
- Tavera Busso, I., Mateos, A.C., González Peroni, A., Graziani, N.S., Carreras, H.A., 2020. Hepatic alterations associated with fine particulate matter exposure. *Toxicol. Res.* 36, 139–148.
- Therneau, Lumley, 2015. Package “survival.” [brieger.esalq.usp.br](http://brieger.esalq.usp.br).
- Wang, B., Eum, K.-D., Kazemiparkouhi, F., Li, C., Manjourides, J., Pavlu, V., Suh, H., 2020. The impact of long-term PM exposure on specific causes of death: exposure-response curves and effect modification among 53 million U.S. Medicare beneficiaries. *Environ. Health* 19, 20.
- Wei, Y., Qiu, X., Yazdi, M.D., Shtein, A., Shi, L., Yang, J., Peralta, A.A., Coull, B.A., Schwartz, J.D., 2022. The impact of exposure measurement error on the estimated concentration-response relationship between long-term exposure to PM<sub>2.5</sub> and mortality. *Environ. Health Perspect.* 130, 77006.
- Wei, Y., Wang, Y., Di, Q., Choirat, C., Wang, Y., Koutrakis, P., Zanobetti, A., Dominici, F., Schwartz, J.D., 2019. Short term exposure to fine particulate matter and hospital admission risks and costs in the Medicare population: time stratified, case crossover study. *BMJ* 367, 16258.
- Wood SN, 2015. Package “mgcv.” R package version.
- Woolhandler, S., Himmelstein, D.U., 2017. The Relationship of Health Insurance and Mortality: Is Lack of Insurance Deadly? *Ann. Intern. Med.* 167, 424–431.
- World Health Organization, 2013. Health Risks of Air Pollution in Europe HRAPIE Project: Recommendations for concentration–response functions for cost–benefit analysis of particulate matter, ozone and nitrogen dioxide.
- World Health Organization. Regional Office for Europe, 2021. Review of evidence on health aspects of air pollution: REVIHAAP project: technical report (No. WHO/EURO:2013-4101-43860-61757). World Health Organization. Regional Office for Europe.
- Xu, M., Sbihi, H., Pan, X., Brauer, M., 2020. Modifiers of the effect of short-term variation in PM<sub>2.5</sub> on mortality in Beijing, China. *Environmental Research*. <https://doi.org/10.1016/j.envres.2019.109066>
- Xu, W., Wang, S., Jiang, L., Sun, X., Wang, N., Liu, X., Yao, X., Qiu, T., Zhang, C., Li, J., Deng, H., Yang, G., 2022. The influence of PM<sub>2.5</sub> exposure on kidney diseases. *Hum. Exp. Toxicol.* 41, 9603271211069982.
- Yitshak-Sade, M., James, P., Kloog, I., Hart, J.E., Schwartz, J.D., Laden, F., Lane, K.J., Fabian, M.P., Fong, K.C., Zanobetti, A., 2019. Neighborhood Greenness Attenuates the Adverse Effect of PM<sub>2.5</sub> on Cardiovascular Mortality in Neighborhoods of Lower Socioeconomic Status. *Int. J. Environ. Res.*

Public Health 16. <https://doi.org/10.3390/ijerph16050814>

Yu, Y., Yao, S., Dong, H., Wang, L., Wang, C., Ji, X., Ji, M., Yao, X., Zhang, Z., 2019. Association between short-term exposure to particulate matter air pollution and cause-specific mortality in Changzhou, China. *Environ. Res.* 170, 7–15.