

Influence of soil-transmitted helminth infections on physical activity, physical fitness, and cardiovascular disease risk in primary schoolchildren in Gqeberha, South Africa

Siphesihle Nqweniso^{1*}, Cheryl Walter¹, Rosa du Randt¹, Larissa Adams¹, Johanna Beckmann², Jean T. Coulibaly^{3,4,5,6}, Jan Degen², Danielle Dolley¹, Stefanie Gall², Nandi Joubert^{2,3,4}, Kurt Z. Long^{3,4}, Ivan Müller², Madeleine Nienaber¹, Uwe Pühse², Harald Seelig², Peter Steinmann^{3,4}, Jürg Utzinger^{3,4}, Markus Gerber² & **Christin Lang**²

¹ Department of Human Movement Science, Nelson Mandela University, Gqeberha, South Africa; ² Department of Sport, Exercise and Health, University of Basel, Basel, Switzerland; ³ Swiss Tropical and Public Health Institute, Allschwil, Switzerland; ⁴ University of Basel, Basel, Switzerland ; ⁵ Centre Suisse de Recherches Scientifiques en Côte d'Ivoire, Abidjan, Côte d'Ivoire; ⁶ Unité de Formation et de Recherche Biosciences, Université Félix Houphouët-Boigny, Abidjan, Côte d'Ivoire.

Abstract

Objective: School-aged children in low- and middle-income countries carry the highest burden of intestinal helminth infections, such as soil-transmitted helminths (STH). STH infections have been associated with negative consequences for child physical and cognitive development and wellbeing. Thus, the aim of this study was to determine the association of STH infections in schoolchildren from Gqeberha, focusing on physical activity, physical fitness, and clustered cardiovascular disease (CVD) risk score.

Methods: This cross-sectional study involved 680 schoolchildren (356 girls and 324 boys; mean age 8.19 years, SD±1.4) from disadvantaged communities in Gqeberha (formerly, Port Elizabeth), South Africa. Stool samples were collected and examined for STH infections using the Kato-Katz method. Physical activity (accelerometer) and physical fitness (grip strength, 20 m shuttle run) were measured using standard procedures. Furthermore, anthropometry, blood pressure, as well as glycated haemoglobin and lipid profile from capillary blood samples were assessed. We employed one-way ANOVAs to identify the associations of STH infections in terms of species and infection intensity with physical activity, physical fitness, and clustered CVD risk score.

Results: We found a low STH infection prevalence (7.2%) in our study, with participants infected with at least one intestinal helminth species. In comparison to their non-infected peers, children infected with STH had lower mean grip strength scores, but higher mean VO₂max estimation and higher levels of MVPA (p<.001). When considering type and intensity of infection, a positive association of *A. lumbricoides* infection and MVPA was found. In contrast, light *T. trichiura*-infected children had significantly lower grip strength scores compared to non-infected children. VO₂max and MVPA were positively associated with light *T. trichiura* infection. No significant association between the clustered CVD risk score and infection with any STH species was evident.

Conclusions: STH-infected children had lower grip strength scores than their non-infected peers, yet, achieved higher VO₂max and MVPA scores. Our study highlights that the type and intensity of STH infection is relevant in understanding the disease burden of STH infections on children's health. The findings of our study must be interpreted cautiously due to the low infection rate, and more research is needed in samples with higher prevalence rates or case-control designs.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Introduction

More than 1.5 billion of the global population are infected with soil-transmitted helminths (STH), with most of these infections occurring in children living in low- and middle-income countries (LMICs) (1-3). The most prevalent helminth species are *Ascaris lumbricoides* (*A. lumbricoides*), *Trichuris trichiura* (*T. trichiura*), and hookworms, and infection with more than one worm species is not uncommon (1). According to the World Health Organization (WHO) (4), school-aged children in LMICs carry the highest burden of STH infections. The consequences of STH infections during childhood are manifold, including impaired growth, malnutrition, and delayed intellectual development and cognition (1, 5).

In recent years, a growing body of research has investigated the effects of children's STH infection on physical fitness (6-10) and physical activity levels (11). Physical activity is vital to young children's healthy physical and cognitive development. Later in life, regular physical activity becomes increasingly important in preventing non-communicable disease (NCD) risk factors (12). Conversely, high physical activity levels may pose a risk factor for STH infection because it increases exposure to contaminated soil when playing and running outside regularly, especially in communities with poor water, sanitation, and hygiene (WASH). STH infection itself may then lead to increased morbidity and lower levels of physical fitness. Studies investigating the impact of STH infection on children's physical fitness have reported lower grip strength scores (9, 10) and lower maximum oxygen uptake (VO₂max) (8). In contrast, Hürlimann et al. (13) observed higher VO₂max and standing broad jump scores in children infected with hookworm and *Schistosoma mansoni* (*S. mansoni*). The authors attributed these findings to infected children's general higher activity patterns. In line with this notion, a study conducted in marginalised communities in Gqeberha (formerly Port Elizabeth), South Africa, revealed that infected children self-reported higher physical activity levels than their non-infected peers (11). Especially water activities place children at an increased risk for schistosomiasis (13). However, other cross-sectional studies conducted in Côte d'Ivoire (7) and South Africa (10) did not find an association between STH infection and VO₂max. The inconsistent findings may be explained by the fact that previous studies focused primarily on the prevalence of STH infection alone, without considering the type and intensity of infection. An important factor in determining the morbidity of STH infections is infection intensity, which refers to the number of worms found in an individual (4, 14). In most individuals, light infections are not overtly symptomatic, but the symptoms progress as the infection gets more severe. Infection intensity is measured in eggs per gram (EPG) and categorised as light, moderate, or heavy (4). Moderately and heavily infected individuals experience more health consequences than lightly-infected individuals (15). Furthermore, hookworm infection is mainly associated with

iron deficiency anaemia resulting from chronic gastrointestinal bleeding. Anaemia can also be a symptom of moderate to heavy infections with *T. trichiura*. In contrast, infections with *A. lumbricoides* do not exhibit these symptoms, suggesting that the type of STH infection may also be relevant to the observed impact on VO₂max and overall disease burden (15).

While in LMICs, children continue to suffer from infectious diseases, such as STH, their parents are more likely to suffer from NCDs. There has been a long-standing notion that NCDs, such as cardiovascular diseases (CVDs), type 2 diabetes, hypertension, and obesity, are the diseases of western high-income countries (16). However, current epidemiological data indicate that NCDs, and CVD risk factors in particular, now disproportionately affect people in LMICs, where more than three-quarters of global NCD deaths occur (16). To what extent childhood STH infection may function as a potential driver or prevention factor for CVD risk factors is still understudied. Previous research has focused on adult populations (17, 18) or animals (19). The studies conducted on humans have shown an association between *S. mansoni* and reduced risk of diabetes and CVD risk (17, 18). Based on two systematic reviews among adult populations, helminth infections may offer a protective mechanism by modulating the immune system of their hosts, thereby preventing diabetes and improving insulin sensitivity (20, 21).

Taken together, CVD risk factors are characterised by multiple overlapping lifestyles and clinical risk factors (e.g. obesity, diabetes or impaired glucose tolerance, physical inactivity, hypertension), which the WHO Global Health Observatory data have identified as common and preventable risk factors that underlie most CVDs. Childhood physical activity is a preventive factor for various CVDs later in life. Yet, outdoor physical activities may provide a potential risk factor for STH infection among school-aged children. To date, little is known about the relationship between STH infections, habitual physical activity levels, physical fitness, and CVD risk factors in primary schoolchildren living in marginalised areas. Therefore, the objective of this study is threefold: first, to describe the prevalence, type, and intensity of STH infections in four schools in Gqeberha, South Africa. Second, to identify whether STH-infected and non-infected children differ regarding their physical activity, physical fitness levels, and clustered CVD risk score. Third, determine whether STH infection type and intensity are associated with habitual physical activity, physical fitness levels, and clustered CVD risk score.

Methods

Study design and location

This study presents cross-sectional baseline data from a larger cluster-randomised controlled trial (*KaziAfy*). The *KaziAfy* study aims to assess the effectiveness of physical activity and multi-micronutrient supplementation interventions on primary schoolchildren living in South Africa, Tanzania, and Côte d'Ivoire (22). In the South African cohort of the study, baseline data collection was conducted from February to April 2019 in Gqeberha, Eastern Cape province. The study was conducted at four primary schools in socioeconomically disadvantaged communities inhabited by predominantly coloured (mixed-race) and black African children. The selected schools are categorised as quintile three schools. In South Africa, quintile three schools are situated in underprivileged communities and are non-fee-paying schools (23).

Participants and procedure

Based on a priori power analysis (calculations based on G*power 3.1) (22), the final target sample comprised 1320 children per study site. Detailed information about sampling and selection of schools, study participants, management of helminth infections, and referrals can be found in the published study protocol (22). In brief, participants were recruited from four peri-urban primary schools in Gqeberha, South Africa. The inclusion criteria included schools that; (i) were quintile three public schools, (ii) had facilities available to implement physical education classes, and (iii) did not participate in another research project or clinical trial. Children were included in the study if; (i) they were in grades 1–4 (aged 6-12 years) at baseline, (ii) their parent/guardian had signed the informed consent form, and (iii) they were not participating in other research projects at the same time as our study, and they did not have any adverse medical condition preventing participation in physical activity. All children submitted an informed consent form signed by their parent/guardian, giving verbal assent before being assessed. Children's participation was voluntary, and it was explained that withdrawal from the study was possible at any point without further obligation. Data collection was conducted by biokineticists, nurses and trained research assistants during school hours over two days.

Ethical considerations

Approval of the study was obtained from the responsible ethics committees at the Nelson Mandela University in Gqeberha, South Africa (reference number: H19-HEA-HMS-001) and

the 'Ethikkommission Nordwest- und Zentralschweiz' in Switzerland (EKNZ; reference number: Req-2018-00608). Permission was also sought from the Eastern Cape Department of Education and Department of Health, South Africa. The KaziAfya intervention study was registered on August 9, 2018, with ISRCTN (<http://www.isrctn.com/ISRCTN29534081>) and was conducted following the study protocol (22), the principles of the Declaration of Helsinki and the guidelines of Good Clinical Practice (GCP) issued by the International Conference of Harmonisation (ICH). The study was guided by principles of ethics, including voluntary participation, confidentiality, and anonymity.




Figure 1: Participant flow chart

Measures

All procedures used in this study were based on standardised, validated and quality-controlled standard operating procedures (22).

Anthropometric measurements

Body height was assessed using a Seca Stadiometer (Surgical SA; Johannesburg, South Africa), measured to the nearest 0.1 cm. With shoes off, children stood against the stadiometer, shoulders relaxed, back erect, and heels touching the stadiometer. Body weight

(to nearest 0.1 kg) and percentage body fat (BF%) were assessed using bioelectrical impedance analysis (BIA) via an electronic scale (Tanita MC-580; Tanita Corp., Tokyo, Japan). Children were required to fast for 3 hours before data collection. With minimal clothing, children stood barefoot on the metal plates of the scale, allowing for optimal contact with the plates. After inputting height measurement on the Tanita scale, body mass index (BMI) could be read from the scale's monitor.

Socioeconomic status (SES)

We assessed household SES using a parental survey, where parents/guardians had to answer questions related to several items about asset ownership (e.g. number of bedrooms, people per household). The items included and exact scoring to estimate SES have been described elsewhere (24). Items possessed by >95% or <5% of the sample were excluded, with the remaining items used to generate a wealth index via principal component analysis (PCA). The use of PCA to generate a wealth index has been used and validated previously (25). A high score reflected high SES; quintiles were built based on the wealth index, with quintiles 1 and 5 representing lowest/poorest and highest/wealthiest, respectively.

Soil-transmitted helminth (STH) infection

A pre-labelled (unique ID and name) stool sample container was handed out to participants on the morning of data collection. The participants were instructed to return with the container filled with their morning stool the next day (at least 15 g). Sample collection was done in the morning, and containers were transported to a laboratory for diagnostic work-up within the same day. For the detection of STH infection (*A. lumbricoides*, *T. Trichiura* and hookworms), the Kato-Katz technique was used (26). For each sample, a duplicate 41.7 mg thick smear was prepared and read under a microscope by experienced laboratory technicians from Nelson Mandela University. The number of helminth eggs was counted and recorded for each species separately. According to WHO guidelines, EPG was used to calculate STH infection intensity (light, moderate, heavy) (4).

Physical fitness

Two tests from the Eurofit physical fitness test battery were used (27), namely (i) grip strength test and (ii) 20 m shuttle run test. The grip strength test was used to assess upper body strength, and the 20 m shuttle run test was used to assess cardiorespiratory fitness (28). Using the Saehan hydraulic hand dynamometer (MSD Europe BVBA; Tisselt, Belgium), children were tested on both right and left hands after adjusting the grip span. The test was conducted with the child seated upright, shoulders relaxed, and elbow bent at a 90-degree angle alongside the body. With alternating hands, six trials were conducted (with 30-sec rest

in between). The final score was determined by calculating an average of all the trials (to the nearest 1 kg). The 20 m shuttle run test was conducted in the schoolyard on flat ground, adhering to a standard test protocol (28, 29). Children were instructed to run to a pre-recorded signal with an initial running speed of 8.5 km/h that increased by 0.5 km/h after every minute. The test was terminated when a child failed to follow the pace in two consecutive 20 m intervals. The last completed lap and corresponding speed were recorded and converted to maximal oxygen uptake (VO₂ max) using the speed and age of each participant (28).

Moderate-to-vigorous physical activity (MVPA)

Physical activity was assessed with an accelerometer device (ActiGraph®wGT3X-BT; Pensacola, USA). Children were instructed to wear the device around the hip for seven consecutive days and to take it off only when in contact with water. Devices collected data from 6 a.m. to midnight, and data were deemed valid if children wore the device for at least 8 hours on four weekdays and one weekend day (30), and the accelerometer device (ActiGraph) had been validated for children (31). Moderate and vigorous physical activity were determined using raw data and the ActiLife® software version 6.13.2, with raw data set to 10-sec epochs (32). Children-specific cut-off points were used to determine MVPA (33).

Cardiovascular risk factors

An Omron M3® digital blood pressure monitor (Omron Healthcare Europe; Hoofddorp, The Netherlands) was used to measure blood pressure three times after the child had been resting for 5 minutes, with a 1-minute rest between measures. An average of the last two blood pressure measurements was used as the final reading. For the assessment of blood lipid profiles (high-density lipoprotein (HDL), triglycerides and total cholesterol to HDL ratio) and glycated haemoglobin (HbA1c) level, capillary blood was analysed using the Afinion test (Alere Technology, Abbot; Wädenswil, Switzerland). The finger prick technique was used to collect drops of blood, with children having fasted for 3 hours before the test.

Statistical analysis

Data were double-entered, validated using EpiData version 3.1 (EpiData Association; Odense, Denmark) and merged into a single file. Descriptive statistics are reported as mean (M) and standard deviation (SD) or number (n) and percentage (%). One-way ANOVAs were used to determine whether grip strength, VO₂max, MVPA and clustered CVD risk score differed between infected and non-infected children. Additional subgroup analyses were carried out at one school with the highest prevalence. Similarly, one-way ANOVAs were

used to determine whether grip strength, VO₂max, MVPA, and clustered CVD risk scores differed by STH infection type and intensity. When the p-value revealed significant differences between some of the tested variables, a Tukey's Honest Significant Difference (HSD) test was used to assess the significant differences between pairs of group means. Possible group differences in children's sex, age and SES are controlled for in the models. The VO₂max calculation already considers age (28). To obtain a clustered cardiovascular risk score, the individual risk factors were z-standardized and added up using the following formula to calculate the clustered risk score: (systolic + diastolic blood pressure/2) + body fat + ratio of total cholesterol to HDL + triglycerides + HbA1c. Previous publications based on studies conducted in Europe, the United States of America (34, 35), and South Africa, Gqeberha (36, 37) have used this formula. SES was determined by calculating a wealth index generated through a PCA, with higher scores reflecting higher SES. Details about the wealth index calculation have been described elsewhere (24). All statistical analyses were carried out using SPSS Version 27 (IBM Corporation, Armonk, NY, USA) for Windows. For all statistical analyses, we use $p < 0.05$ to denote significance.

Results

Sample characteristics

A total of 1369 children provided informed consent, with 65 dropping out before baseline assessment due to relocation or moving to a different school. Of the 1304 remaining children, 624 were not included in the data analysis as they had missing information for at least one of the following: age, sex, stool sample, grip strength, 20m shuttle run, accelerometer or clustered CVD score (Figure 1). Subsequent analyses were based on 680 children with complete data (356 girls, 324 boys), a mean age of 8.19 years ($SD \pm 1.4$). There were significant age differences between the four schools ($p < 0.001$); thus, age was included as a covariate in all subsequent analyses. No sex and SES differences were observed. Table 1 shows the descriptive statistics of all measured variables for the sample ($n=680$). The mean grip strength, VO₂max estimate and MVPA scores for the study cohort were 11.40 kg, 47.56 ml/kg/min and 80.09 min/day, respectively.

Table 1. Characteristics of study participants from disadvantaged communities in Gqeberha, South Africa (N = 680)

Parameters	N	Total	School 1	School 2	School 3	School 4	Statistics		
		(n = 230)	(n = 223)	(n = 144)	(n = 83)	F	Group		
		M (SD)	M (SD)	M (SD)	M (SD)	M (SD)		p	η^2
Age and anthropometry									
Age (years)	680	8.19 (1.42)	8.38 (1.48)	8.24 (1.45)	8.19 (1.42)	7.59 (0.99)	6.44	< 0.001	0.028
Height (cm)	680	123.98 (9.08)	126.48 (8.94)	122.53 (9.25)	123.55 (9.40)	121.67 (6.72)	9.98	< 0.001	0.042
Weight (kg)	680	24.72 (6.13)	27.25 (6.66)	23.45 (5.67)	23.38 (5.24)	23.42 (4.76)	21.57	< 0.001	0.087
BMI (kg/m ²)	680	15.89 (2.35)	16.87 (2.71)	15.43 (2.07)	15.16 (1.78)	15.69 (1.95)	23.10	< 0.001	0.093
Socio-economic status	680	3.07 (1.44)	3.13 (1.49)	3.03 (1.44)	3.17 (1.39)	2.88 (1.36)	0.73	0.535	0.005
STH infection									
	N	N (%)	N (%)	N (%)	N (%)	N (%)	χ^2	p	
<i>Ascaris lumbricoides</i> (%)	680	40 (5.8)	0 (0.0)	39 (5.7)	1 (0.1)	0 (0.0)	80.45	< 0.001	0.344
<i>Trichuris trichiura</i> (%)	680	48 (7.0)	0 (0.0)	47 (6.9)	1 (0.1)	0 (0.0)	98.99	< 0.001	0.381
Hookworm (%)	680	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.1)	7.22	0.065	0.103
Single infection	680	9 (1.3)	0 (0.0)	8 (1.2)	0 (0.0)	1 (0.1)			
Co-infection	680	40 (5.8)	0 (0.0)	39 (5.7)	1 (0.1)	0 (0.0)			
Physical fitness									
		M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	F	p	η^2
Grip strength (kg)	680	11.40 (4.87)	10.94 (4.23)	10.50 (3.48)	13.86 (7.12)	10.83 (3.45)	16.96	< 0.001	0.070
VO ₂ max (ml/kg/min)	680	47.56 (3.48)	46.79 (3.14)	48.84 (3.78)	46.07 (2.80)	48.88 (2.93)	30.09	< 0.001	0.118
MVPA (min/day)	680	80.09 (26.39)	73.75 (24.82)	84.14 (27.05)	82.34 (26.21)	82.86 (26.32)	7.01	< 0.001	0.030
Clustered CVD risk	680	-0.04 (2.68)	0.16 (2.75)	-0.14 (2.66)	-0.51 (2.56)	0.45 (2.68)	2.98	0.031	0.013

Note. STH = Soil-transmitted helminth infection, VO₂max = Cardiorespiratory fitness, MVPA = Moderate-to-vigorous physical activity, CVD = Cardiovascular disease, η^2 = partial eta squared: 0.01 = small, 0.06 = medium, 0.14 = large effect size.

Prevalence, type, and intensity of STH infection

Forty-nine out of 680 children (7.2%) were infected with at least one intestinal helminth species. Infection prevalence of *A. lumbricoides* and *T. trichiura* were 5.8% and 7.0%, respectively. Only one child was infected with hookworm, and no separate analyses were conducted for this child. Out of the 49 infected children, co-infection was found in 40 (81.6%) children who harboured both *A. lumbricoides* and *T. trichiura*. Stratification by school revealed that school 2 had the highest prevalence of both *A. lumbricoides* and *T. trichiura* (Table 1). Children infected with *A. lumbricoides* presented mostly with light infection intensity. In contrast, those infected with *T. trichiura* showed mostly heavy infection intensity, with no moderate infection intensity observed in *T. trichiura* infection (data not shown).

MVPA, physical fitness, and CVD risk among infected and non-infected children

As shown in Table 2a, children infected with STH had significantly lower mean grip strength than their non-infected peers ($F(1,679) = 5.44$, $p = 0.020$, $\eta^2 = 0.008$). Mean VO₂max estimate was found to be significantly higher in infected compared to non-infected children ($F(1,679) = 18.42$, $p < 0.001$, $\eta^2 = 0.026$). Similar results were found for MVPA, with infected children achieving higher scores compared to non-infected children ($F(1,679) = 19.54$, $p < 0.001$, $\eta^2 = 0.028$). No significant differences between infected and non-infected children in respect of the clustered CVD risk score could be established ($F(1,679) = 0.368$, $p > 0.05$, $\eta^2 = 0.001$).

Due to infection occurring in one of the four schools under investigation, further analyses were carried out, focusing on the high infection area school (school 2, $n = 223$). Table 2b indicates that STH-infected children showed significantly lower mean grip strength and higher VO₂max compared to non-infected children ($F(1,222) = 4.65$, $p = 0.032$, $\eta^2 = 0.021$) and ($F(1,222) = 4.59$, $p = 0.033$, $\eta^2 = 0.020$), respectively. Mean MVPA scores were significantly higher for infected children compared to their non-infected peers ($F(1,222) = 13.07$, $p < 0.001$, $\eta^2 = 0.056$). These results remained significant even after adjusting for covariates (age, BMI). No significant differences were found between infected and non-infected children regarding the clustered CVD risk score ($F(1,222) = 0.354$, $p > 0.05$, $\eta^2 = 0.002$).

Table 2a. Physical fitness, physical activity (MVPA), and clustered CVD risk in STH-infected and non-infected children (n = 680)

Measures	Non-infected	STH infected	Statistics		
	(n = 631)	(n = 49)	F	p	η^2
	M (SD)	M (SD)			
Physical fitness					
Grip strength (kg)	11.55 (4.97)	9.57 (2.97)	5.44	0.020	0.008
VO ₂ max (ml/kg/min)	47.39 (3.36)	49.78 (4.16)	18.42	<0.001	0.026
MVPA (min/day)	78.79 (26.18)	96.82 (23.34)	19.54	<0.001	0.028
Clustered CVD risk	-0.03 (2.71)	-0.20 (2.32)	0.368	0.544	0.001

Note. STH = Soil-transmitted helminth infection, VO₂max = Cardiorespiratory fitness, MVPA = Moderate-to-vigorous physical activity, CVD = Cardiovascular disease, η^2 = Partial eta squared: 0.01 = small, 0.06 = medium, 0.14 = large effect size.

Table 2b. Sub analyses: Physical fitness, physical activity (MVPA), and clustered CVD risk in STH-infected and non-infected children from high infection prevalence area (school 2: n = 223)

Measures	Non-infected	STH-infected	Statistics		
	(n= 176)	(n= 47)	F	p	η^2
	M (SD)	M (SD)			
Physical fitness					
Grip strength (kg)	10.77 (3.55)	9.50 (3.01)	4.65	0.032	0.021
VO ₂ max (ml/kg/min)	48.55 (3.64)	49.93 (4.13)	4.59	0.033	0.020
MVPA (min/day)	80.81 (26.96)	96.59 (23.78)	13.07	<0.001	0.056
Clustered CVD risk	-0.16 (2.76)	-0.08 (2.25)	0.354	0.552	0.002

Note. STH = Soil-transmitted helminth infection, VO₂max = Cardiorespiratory fitness, MVPA = Moderate-to-vigorous physical activity, CVD = Cardiovascular disease, η^2 = Partial eta squared: 0.01 = small, 0.06 = medium, 0.14 = large effect size.

Association of infection type and intensity with MVPA, physical fitness, and CVD risk

When considering STH infection type and intensity (Table 3), infection with *A. lumbricoides* showed no significant overall effect with regard to infection intensity and grip

strength and VO₂max ($F(3,220) = 0.59$, $p = 0.621$, $\eta^2 = 0.008$) and ($F(3,220) = 0.83$, $p = 0.480$, $\eta^2 = 0.011$), respectively. There was a significant overall effect with regards to infection intensity and MVPA ($F(3,220) = 3.42$, $p = 0.018$, $\eta^2 = 0.045$). A post hoc Tukey test showed no significant differences between infection intensities ($p > 0.05$). Infection intensity did not show any significant overall effects on clustered CVD risk scores.

T. trichiura infection intensity had a significant overall effect on grip strength ($F(3,220) = 5.14$, $p = 0.007$, $\eta^2 = 0.045$). A post hoc Tukey test indicated a significant difference between non and light infection ($p=0.007$). There was a significant overall effect for infection intensity and VO₂max, in that VO₂max increases from non to light infection yet, decreases again for heavy infections ($F(3,220) = 4.82$, $p = 0.009$, $\eta^2 = 0.042$). Again, a post hoc test showed a significant difference between non and light infection ($p=0.011$). Similar to VO₂max, MVPA and infection intensity with *T. trichiura* showed a significant overall effect ($F(3,220) = 6.79$, $p = 0.001$, $\eta^2 = 0.058$), with post hoc test showing a significant difference between non and heavy infection ($p=0.004$). We did not establish significant differences between the clustered CVD risk score and *T. trichiura* infection intensity ($p > 0.05$).

Discussion

The present study investigated the prevalence and intensity of STH infection and its association with selected health parameters in primary schoolchildren from disadvantaged communities in Gqeberha, South Africa. A key finding was that the prevalence of STH infection was highly dependent on the geographic location of schools. Overall, STH-infected children had lower grip strength scores than their non-infected peers yet, achieved higher VO₂max and MVPA scores. Sub-analyses result only showed significantly higher MVPA scores in infected children compared to their non-infected peers. Sub-analyses with *A. lumbricoides* infection intensity revealed no differences in grip strength and VO₂max but higher MVPA among infected children, independent of infection intensity. In contrast, for *T. trichiura* infection, lightly-infected children had significantly lower grip strength and achieved higher VO₂max estimates. While heavily-infected children had significantly higher MVPA levels than their light and non-infected peers. There was an unexpected reversal in this pattern for grip strength, with non-infected and *T. trichiura* heavily-infected children having higher grip strength scores compared to their lightly-infected peers; however, this may be due to the low number of lightly-infected children ($n = 8$). In contrast, clustered CVD risk score was not associated with STH infection intensity.

Table 3. Physical fitness, physical activity (MVPA), and clustered CVD risk in non-infected versus STH-infected children, separately for type (*A. lumbricoides* and *T. trichiura*) and intensity (light vs. moderate vs. heavy as determined by EPG) (school 2: n = 223)

<i>Ascaris lumbricoides</i>							
	Non-infected (n = 184)	Light (n = 16)	Moderate (n = 11)	Heavy (n = 12)	Statistics		
Parameters	M (SD)	M (SD)	M (SD)	M (SD)	F	p	η^2
Grip strength (kg)	10.60 (3.56)	10.13 (3.16)	10.64 (2.26)	9.31 (3.29)	0.59	0.621	0.008
VO ₂ max (ml/kg/min)	48.70 (3.74)	50.03 (3.94)	48.36 (4.55)	49.59 (3.59)	0.83	0.480	0.011
MVPA (min/day)	81.57 (26.94)	96.39 (21.66)	100.97 (30.95)	90.69 (21.85)	3.42	0.018	0.045
Clustered CVD risk	-0.21 (2.73)	0.17 (2.67)	-0.76 (2.18)	0.80 (1.82)	0.81	0.489	0.011

<i>Trichuris trichiura</i>							
	Not infected (n = 176)	Light (n = 8)	Moderate (n = 0)	Heavy (n = 39)	Statistics		
Parameters	M (SD)	M (SD)	M (SD)	M (SD)	F	p	η^2
Grip strength (kg)	10.77 (3.55)	7.00 (1.91)	--	10.02 (2.95)	5.14	0.007	0.045
VO ₂ max (ml/kg/min)	48.53 (3.64)	52.42 (4.19)	--	49.42 (3.98)	4.82	0.009	0.042
MVPA (min/day)	80.75 (26.90)	99.83 (22.14)	--	95.93 (24.33)	6.79	0.001	0.058
Clustered CVD risk	-0.17 (2.76)	-0.97 (1.69)	--	0.10 (2.32)	0.56	0.570	0.005

Note. STH = Soil-transmitted helminth infection, VO₂max = Cardiorespiratory fitness, MVPA = Moderate-to-vigorous physical activity, CVD = Cardiovascular disease, EPG = Eggs per gram, η^2 = Partial eta squared: 0.01 = small, 0.06 = medium, 0.14 = large effect size.

The present study had three research aims, each of which will now be discussed. Firstly, we wanted to report on the prevalence of STH infection. Overall, our study found a low prevalence of *T. trichiura* and/or *A. lumbricoides* infection (7.2%), with only one child infected with hookworms. The STH infection prevalence is much lower than what has been reported in previous studies in Gqeberha (31.0%) (10) and KwaZulu-Natal (1.2 – 26%) (38, 39). The low prevalence can be attributed to a national school deworming programme launched in 2016 by the South African Department of Education for schools in disadvantaged communities (40). Müller and colleagues (41) also reported shrinking risk profiles following three rounds of albendazole deworming of STH-infected children from communities located in close proximity to the current study setting (March 2015, October 2015 & May 2016). Even though the general prevalence in our sample population was found to be low, one school showed substantially higher STH infection compared to the other three schools. This observation is in line with findings from the DASH study that also reported high variations between schools located within similar geographical areas (9). Their study found that two schools situated in the same geographical area as the present study had high infection rates. This geographical area is one of the old apartheid group areas set aside for coloured people during the period of forced removals in the 1960s (42, 43), and it is characterised by high population density and inadequate water and sanitation facilities to cope with the high density of people living in the area (9). The present study revealed that 81.6% of the infected children were co-infected with *A. lumbricoides* and *T. trichiura*. This is higher than the findings of previous studies in China and South Africa (8, 9, 39), where 30-51% of their participants harboured both *A. lumbricoides* and *T. trichiura*, which is common due to the mechanism of infection (14).

Secondly, we aimed to explore whether STH-infected and non-infected children differed with regard to their physical activity, physical fitness levels, and clustered CVD risk score. These results are reported for both the whole group and sub-analyses. Consistent with existing literature, STH-infected children in the whole sample and sub-analyses had a lower grip strength score than their non-infected peers (8, 9). Interestingly, STH-infected children had significantly higher VO₂max than non-infected children. These findings contradict previous research conducted in China and South Africa, which found lower mean VO₂max estimates in STH-infected children (8, 9), but corroborate findings from Gerber and colleagues (11) where self-reported physical activity was associated with a higher infection rate in children from a similar area. A possible reason for the high VO₂max scores in the current study is that infected children have significantly lower BMI than non-infected children, allowing them to complete more laps on the 20 m shuttle run test. High BMI in children has been associated with poor performance in weight-bearing activities such as the 20 m shuttle

run test (44). This was also found to be the case in a study by Smith and colleagues (45), who reported that overweight/obese children completed fewer laps in the 20 m shuttle run test and had poor VO₂max scores. Similar to the VO₂max results, STH-infected children in the whole group and sub-analyses had significantly higher MVPA scores. These observations may be explained by the fact that physically active children are more likely to play outside (sometimes barefoot), which may place them at a higher risk of contracting infection, especially in poor water and sanitation areas. A previous study in KwaZulu-Natal found a positive association between schistosomiasis and physical fitness. The authors attributed this to a selective increase in exposure to infected water in more active children (46). Although a physically active lifestyle from a young age onwards is a key component in preventing NCDs later in life, physical activity guidelines need to address risk factors in disadvantaged areas. The importance of regular deworming in schoolchildren has been underscored by the WHO, as it ensures the reduction in morbidity caused by the infection and the occurrence of severe complications (4). On the other hand, previous research on adult populations suggests a protective mechanism of STH infection against metabolic outcomes. For instance, *S. mansoni* infection was positively associated with blood lipids (lower mean low-density lipoprotein, total cholesterol, and triglycerides) and diastolic blood pressure (17, 18). However, in our study, we could not establish such a relationship. However, we recommend that future studies investigate the possible link between STH infection and CVD risk factors in children.

Thirdly, we aimed to identify whether STH type and infection intensity differed regarding physical activity, physical fitness, and clustered CVD risk score. To our knowledge, previous studies have not considered the type and intensity of infection when investigating the effects of STH infections on children's health, especially physical activity, physical fitness and clustered CVD risk factors. We did not find any significant effect with regard to *A. lumbricoides* infection intensity and grip strength, and VO₂max. A study conducted in China where *A. lumbricoides* infected children had lower grip strength scores and showed increases in grip strength following one month of deworming treatment with albendazole (47). *A. lumbricoides* infected children had significantly higher MVPA compared to their non-infected counterparts; however, Tukey's HSD post hoc test did not show any significant differences between infection intensity. In our findings, light *T. trichiura* infection intensity resulted in significantly lower grip strength and higher VO₂max, which does not corroborate previous studies that reported a decrease in VO₂max of infected children (9). However, a study in China showed no clear relationship between infection intensity and physical fitness (8). We also found significantly higher MVPA in *T. trichiura*-infected children, with the post hoc test indicating significant differences between non and highly-infected

children. Our findings show a trend towards an inverse U-shaped relationship for *T. trichiura* infection, in that lightly-infected children achieved higher VO₂max and MVPA levels compared to their heavily-infected and non-infected peers. However, we reported low light *T. trichiura* infection (n=8) compared to heavy intensity (n=39), indicating an imbalanced portion of children with different infection intensities. This means we do not have sufficient infected children to confirm these results, but we can only assume this inverse relationship, and our data shows some support for this relationship. Due to our low infection prevalence and low numbers of light infection intensity, more research in samples with higher infectious rates or with case-control design is needed. Light STH infections usually have little to no long-term health consequences, whereas moderate and high infection intensity results in a high disease burden (1). Consequently, a light infection may pose an advantage in weight-bearing activities due to lower BMI, but this is outweighed by increasing infection intensity and hence, an increase in disease burden. Low BMI in *T. trichiura*-infected children has been reported in previous studies (8, 9) due to hindered nutrient uptake, and light infection may also contribute to this, especially with recurring infection (1, 14). Previous studies investigating the effects of STH infections on nutrition status and physical fitness reported that being anaemic and stunted had a negative impact on children's physical fitness (48). Anaemia is a common symptom of hookworm infection, and it can also be a symptom of moderate to heavy infection with *T. trichiura*, while *A. lumbricoides* do not exhibit these symptoms (15). Since hookworm has the most detrimental effects on children's physical fitness due to iron deficiency, anaemia and malnutrition (14, 48), and as we only observed one case of hookworm infection, we cannot make conclusions on the fitness benefits seen in STH-infected children.

Limitations and strengths

Due to the study's cross-sectional nature, we could not report inferences about causality and its direction. The overall sample size was small, and STH infection prevalence was low in our population. Our findings cannot be generalised to the entire South African population as children from only one province, and social strata were included in the investigation. We only collected one stool sample, whereas WHO recommends the collection of three stool samples per participant for accurate detection of infection. Furthermore, our diagnostic method has limitations regarding its ability to pick up low-intensity infection (15). Therefore, a concentration test with higher sensitivity is recommended for future studies. Apart from the limitations identified in the study, some strengths can be noted, such as the study's novelty in investigating the type and intensity of infection among children in Gqeberha, South Africa. Our results highlight the importance of focusing on the type and intensity of STH infections to

determine disease morbidity, as prevalence alone cannot display a complete picture of the disease burden. Most of the prevalence of infection observed in one area and the differences amongst infected and non-infected children regarding VO₂max and MVPA, which have not been reported elsewhere, warrant more extensive follow-up studies.

Conclusions

Our data suggest that STH infections hamper children's upper body strength. We also highlighted that there are STH infection hotspot areas in Gqeberha, which warrants regular deworming programmes and WASH interventions. Our findings revealed no clear relationship between STH infection and CVD risk factors. Hence, future research should seek a deeper understanding of the mechanism of how helminths mediate metabolic pathways, especially in children. Our study highlights that the type and intensity of STH infection are relevant in understanding the disease burden of STH infections on children's health. We hope that more empirical evidence will emerge from future studies with larger samples, from areas where the authorities perform no systematic deworming and where infection rates are likely to be higher, and/or with case-control designs with a balanced portion of children with different infection intensity.

Author Contributions

Conceptualization: SN, CL, JTC. Methodology: SN and CL. Formal analysis: SN and CL. Validation: MG and CL. Investigation: SN, LA, JB, JD, DD, SG, NJ, IM, MN. Visualization, SN. Supervision, CW, RduR, MG and CL. Project administration, SN, JB, CL, KZL, and MG. Funding acquisition: CW, KZL, UP, PS, JU MG and CL. Writing—original draft preparation, SN. All authors have read, contributed, and agreed to the final version of the manuscript.

Funding

The study is funded by the Fondation Botnar (Basel, Switzerland; project number 6071 'Physical activity and multi-micronutrient supplementation'), covering research expenses, staff salaries, study equipment and laboratory analyses. In-kind contributions are provided by all involved parties. The multi-micronutrient supplementation and the placebo products are sponsored by DSM Nutritional Products Ltd. (Basel, Switzerland). The funders played no role in the study design, data collection and data analysis, data interpretation, and preparation of the manuscript or decision to publish. The corresponding author had the responsibility for the decision to submit for publication.

Informed consent statement

Informed consent was obtained from all subjects involved in the study.

Data availability statement

All data analysed during this study will be included in the published articles.

Acknowledgments

The authors are grateful to the school authorities, school staff, participating children, and their parents/guardians for providing consent and assent to participate in the study. A special thanks to Zaahira Ismail for her administrative work and contribution to data collection. We are also thankful and appreciative to all the postgraduate students who participated and dedicated their time to data collection and data entry.

Conflicts of interest

We declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. Bethony J, Brooker S, Albonico M, Geiger SM, Loukas A, Diemert D, et al. Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm. *Lancet*. 2006; 367(9521): 1521-1532. doi: 10.1016/s0140-6736(06)68653-4
2. Bharti B, Bharti S, Khurana S. Worm Infestation: Diagnosis, Treatment and Prevention. *Indian J Pediatr*. 2018; 85(11): 1017-1024. doi: 10.1007/s12098-017-2505-z
3. WHO. Soil-transmitted helminth infections Geneva: World Health Organization; 2022 [Available from: <https://www.who.int/news-room/fact-sheets/detail/soil-transmitted-helminth-infections>].
4. WHO. Helminth control in school-age children: a guide for managers of control programmes. Geneva: World Health Organization; 2011. Available from: <https://apps.who.int/iris/handle/10665/44671>.
5. Hotez PJ, Molyneux DH, Fenwick A, Kumaresan J, Sachs SE, Sachs JD, et al. Control of neglected tropical diseases. *N Engl J Med*. 2007; 357(10): 1018-1027. doi: 10.1056/NEJMra064142
6. Stephenson LS, Latham MC, Adams EJ, Kinoti SN, Pertet A. Physical fitness, growth and appetite of Kenyan school boys with hookworm, *Trichuris trichiura* and *Ascaris lumbricoides* infections are improved four months after a single dose of albendazole. *J Nutr*. 1993; 123(6): 1036-1046. doi: 10.1093/jn/123.6.1036
7. Müller I, Coulibaly JT, Fürst T, Knopp S, Hattendorf J, Krauth SJ, et al. Effect of schistosomiasis and soil-transmitted helminth infections on physical fitness of school children in Côte d'Ivoire. *PLoS Negl Trop Dis*. 2011; 5(7): e1239. doi: 10.1371/journal.pntd.0001239
8. Yap P, Du ZW, Chen R, Zhang LP, Wu FW, Wang J, et al. Soil-transmitted helminth infections and physical fitness in school-aged Bulang children in southwest China: results from a cross-sectional survey. *Parasit Vectors*. 2012; 5: 50. doi: 10.1186/1756-3305-5-50
9. Müller I, Yap P, Steinmann P, Damons BP, Schindler C, Seelig H, et al. Intestinal parasites, growth and physical fitness of schoolchildren in poor neighbourhoods of Port Elizabeth, South Africa: a cross-sectional survey. *Parasit Vectors*. 2016; 9(1): 488. doi: 10.1186/s13071-016-1761-5
10. Gall S, Müller I, Walter C, Seelig H, Steenkamp L, Pühse U, et al. Associations between selective attention and soil-transmitted helminth infections, socioeconomic status, and physical fitness in disadvantaged children in Port Elizabeth, South Africa: An observational study. *PLoS Negl Trop Dis*. 2017; 11(5): e0005573. doi: 10.1371/journal.pntd.0005573

11. Gerber M, Müller I, Walter C, du Randt R, Adams L, Gall S, et al. Physical activity and dual disease burden among South African primary schoolchildren from disadvantaged neighbourhoods. *Prev Med*. 2018; 112: 104-110. doi: 10.1016/j.ypmed.2018.04.001
12. Blair SN, Cheng Y, Holder JS. Is physical activity or physical fitness more important in defining health benefits? *Med Sci Sports Exerc*. 2001; 33(6 Suppl): S379-399; discussion S419-320. doi: 10.1097/00005768-200106001-00007
13. Hürlimann E, Hounbedji CA, N'Dri PB, Bänninger D, Coulibaly JT, Yap P, et al. Effect of deworming on school-aged children's physical fitness, cognition and clinical parameters in a malaria-helminth co-endemic area of Côte d'Ivoire. *BMC Infect Dis*. 2014; 14: 411. doi: 10.1186/1471-2334-14-411
14. Hotez PJ, Bundy DA, Beegle K, Brooker S, Drake L, de Silva N, et al. *Helminth infections: soil-transmitted helminth infections and schistosomiasis*. In: Jamison D, Breman J, Measham A, Alleyne G, Claeson M, Evans D, et al., editors. *Disease Control Priorities in Developing Countries*. 2nd ed. Washington (DC): The International Bank for Reconstruction and Development / The World Bank; 2006.
15. Harhay MO, Horton J, Olliaro PL. Epidemiology and control of human gastrointestinal parasites in children. *Expert Rev Anti Infect Ther*. 2010; 8(2): 219-234. doi: 10.1586/eri.09.119
16. Ndubuisi NE. Noncommunicable Diseases Prevention In Low- and Middle-Income Countries: An Overview of Health in All Policies (HiAP). *Inquiry*. 2021; 58: 46958020927885. doi: 10.1177/0046958020927885
17. Wolde M, Berhe N, Medhin G, Chala F, van Die I, Tsegaye A. Inverse Associations of *Schistosoma mansoni* Infection and Metabolic Syndromes in Humans: A Cross-Sectional Study in Northeast Ethiopia. *Microbiol Insights*. 2019; 12: 1178636119849934. doi: 10.1177/1178636119849934
18. Sanya RE, Webb EL, Zziwa C, Kizindo R, Sewankambo M, Tumusiime J, et al. The Effect of Helminth Infections and Their Treatment on Metabolic Outcomes: Results of a Cluster-Randomized Trial. *Clin Infect Dis*. 2020; 71(3): 601-613. doi: 10.1093/cid/ciz859
19. Cooke A, Tonks P, Jones FM, O'Shea H, Hutchings P, Fulford AJ, et al. Infection with *Schistosoma mansoni* prevents insulin dependent diabetes mellitus in non-obese diabetic mice. *Parasite Immunol*. 1999; 21(4): 169-176. doi: 10.1046/j.1365-3024.1999.00213.x
20. Berbudi A, Ajendra J, Wardani AP, Hoerauf A, Hübner MP. Parasitic helminths and their beneficial impact on type 1 and type 2 diabetes. *Diabetes Metab Res Rev*. 2016; 32(3): 238-250. doi: 10.1002/dmrr.2673

21. Tracey EF, McDermott RA, McDonald MI. Do worms protect against the metabolic syndrome? A systematic review and meta-analysis. *Diabetes Res Clin Pract.* 2016; 120: 209-220. doi: 10.1016/j.diabres.2016.08.014
22. Gerber M, Ayekoé SA, Beckmann J, Bonfoh B, Coulibaly JT, Daouda D, et al. Effects of school-based physical activity and multi-micronutrient supplementation intervention on growth, health and well-being of schoolchildren in three African countries: the KaziAfyra cluster randomised controlled trial protocol with a 2x 2 factorial design. *Trials.* 2020; 21(1): 22. doi: 10.1186/s13063-019-3883-5
23. Hall K, Giese S. *Addressing quality through school fees and school funding.* In: Pendlebury S, Lake L, Smith C, editors. South African Child Gauge. Cape Town: Children's Institute, University of Cape Town; 2009. p. 35-40.
24. Gerber M, Lang C, Beckmann J, Degen J, du Randt R, Gall S, et al. Associations Between Household Socioeconomic Status, Car Ownership, Physical Activity, and Cardiorespiratory Fitness in South African Primary Schoolchildren Living in Marginalized Communities. *J Phys Act Health.* 2021; 18(8): 883-894. doi: 10.1123/jpah.2020-0839
25. Filmer D, Pritchett LH. Estimating wealth effects without expenditure data—or tears: an application to educational enrollments in states of India. *Demography.* 2001; 38(1): 115-132. doi: <https://doi.org/10.1353%2Fdem.2001.0003>
26. Katz N, Chaves A, Pellegrino J. A simple device for quantitative stool thick-smear technique in Schistosomiasis mansoni. *Rev Inst Med Trop Sao Paulo.* 1972; 14(6): 397-400.
27. Council of Europe. Eurofit: Handbook for the Eurofit Tests of Physical Fitness. Strasbourg: Council of Europe; 1993. Available from: <https://books.google.co.za/books?id=oOUfMwEACAAJ>.
28. Léger LA, Mercier D, Gadoury C, Lambert J. The multistage 20 metre shuttle run test for aerobic fitness. *J Sports Sci.* 1988; 6(2): 93-101. doi: 10.1080/02640418808729800
29. Léger L, Lambert J, Goulet A, Rowan C, Dinelle Y. Aerobic capacity of 6 to 17-year-old Quebecois--20 meter shuttle run test with 1 minute stages. *Can J Appl Sport Sci.* 1984; 9(2): 64.
30. Aadland E, Andersen LB, Skrede T, Ekelund U, Anderssen SA, Resaland GK. Reproducibility of objectively measured physical activity and sedentary time over two seasons in children; comparing a day-by-day and a week-by-week approach. *PLoS One.* 2017; 12(12): e0189304. doi: 10.1371/journal.pone.0189304
31. Crouter SE, Horton M, Bassett DR, Jr. Validity of ActiGraph child-specific equations during various physical activities. *Med Sci Sports Exerc.* 2013; 45(7): 1403-1409. doi: 10.1249/MSS.0b013e318285f03b

32. Freedson PS, Melanson E, Sirard J. Calibration of the Computer Science and Applications, Inc. accelerometer. *Medicine and science in sports and exercise*. 1998; 30(5): 777-781.
33. Evenson KR, Catellier DJ, Gill K, Ondrak KS, McMurray RG. Calibration of two objective measures of physical activity for children. *J Sports Sci*. 2008; 26(14): 1557-1565. doi: 10.1080/02640410802334196
34. Andersen LB, Harro M, Sardinha LB, Froberg K, Ekelund U, Brage S, et al. Physical activity and clustered cardiovascular risk in children: a cross-sectional study (The European Youth Heart Study). *The Lancet*. 2006; 368(9532): 299-304. doi: 10.1016/s0140-6736(06)69075-2
35. Stavnsbo M, Resaland GK, Anderssen SA, Steene-Johannessen J, Domazet SL, Skrede T, et al. Reference values for cardiometabolic risk scores in children and adolescents: Suggesting a common standard. *Atherosclerosis*. 2018; 278: 299-306. doi: 10.1016/j.atherosclerosis.2018.10.003
36. Müller I, Walter C, Du Randt R, Aerts A, Adams L, Degen J, et al. Association between physical activity, cardiorespiratory fitness and clustered cardiovascular risk in South African children from disadvantaged communities: results from a cross-sectional study. *BMJ Open Sport Exerc Med*. 2020; 6(1): e000823. doi: 10.1136/bmjsem-2020-000823
37. Nqweniso S, Walter C, du Randt R, Adams L, Beckmann J, Degen J, et al. Physical Activity, Cardiorespiratory Fitness and Clustered Cardiovascular Risk in South African Primary Schoolchildren from Disadvantaged Communities: A Cross-Sectional Study. *Int J Environ Res Public Health*. 2021; 18(4). doi: 10.3390/ijerph18042080
38. Sacolo-Gwebu H, Chimbari M, Kalinda C. Prevalence and risk factors of schistosomiasis and soil-transmitted helminthiasis among preschool aged children (1-5 years) in rural KwaZulu-Natal, South Africa: a cross-sectional study. *Infect Dis Poverty*. 2019; 8(1): 47. doi: 10.1186/s40249-019-0561-5
39. Zulu SG, Kjetland EF, Gundersen SG, Taylor M. Prevalence and intensity of neglected tropical diseases (schistosomiasis and soil-transmitted helminths) amongst rural female pupils in Ugu district, KwaZulu-Natal, South Africa. *S Afr J Infect Dis*. 2020; 35(1): 123. doi: 10.4102/sajid.v35i1.123
40. South African Government. Launch of National School Deworming Programme Johannesburg, South Africa: South African Government; 2016 [Available from: <https://www.gov.za/speeches/minister-angie-motshekga-launch-national-school-deworming-programme-16-feb-2016-0000>].
41. Müller I, Gall S, Beylveled L, Gerber M, Pühse U, Du Randt R, et al. Shrinking risk profiles after deworming of children in Port Elizabeth, South Africa, with special

- reference to *Ascaris lumbricoides* and *Trichuris trichiura*. *Geospat Health*. 2017; 12(2): 601. doi: 10.4081/gh.2017.601
42. Agherdien Y, George AC, Hendricks S, Du Pre RH. *South End as we knew it*. Port Elizabeth: Western Research Group; 1997.
 43. Du Plessis E, Southern Africa Development R, and Training Institute. *One community many communities : a Nelson Mandela Bay Heritage and History Project (NMBHHP) : forced removals in South End, North End, Salisbury Park, Fairview, Willowdene and Korsten*. Port Elizabeth: Southern Africa Development, Research and Training Institute; 2013.
 44. Moran CA, Peccin MS, Bombig MT, Pereira SA, Dal Corso S. Performance and reproducibility on shuttle run test between obese and non-obese children: a cross-sectional study. *BMC Pediatr*. 2017; 17(1): 68. doi: 10.1186/s12887-017-0825-9
 45. Smith D, Adams L, du Randt R, Degen J, Gall S, Joubert N, et al. Physical fitness and nutritional anthropometric status of children from disadvantaged communities in the Nelson Mandela Bay region. *S Afr J Sports Med*. 2020; 32(1): 1-8. doi: 10.17159/2078-516X/2020/v32i1a8158
 46. Kvalsvig JD. The effects of schistosomiasis on spontaneous play activity in black schoolchildren in the endemic areas. An ethological study. *S Afr Med J*. 1981; 60(2): 61-64.
 47. Yap P, Wu FW, Du ZW, Hattendorf J, Chen R, Jiang JY, et al. Effect of deworming on physical fitness of school-aged children in Yunnan, China: a double-blind, randomized, placebo-controlled trial. *PLoS Negl Trop Dis*. 2014; 8(7): e2983. doi: 10.1371/journal.pntd.0002983
 48. Bustinduy AL, Thomas CL, Fiutem JJ, Parraga IM, Mungai PL, Muchiri EM, et al. Measuring fitness of Kenyan children with polyparasitic infections using the 20-meter shuttle run test as a morbidity metric. *PLoS Negl Trop Dis*. 2011; 5(7): e1213. doi: 10.1371/journal.pntd.0001213