

***Title: Relationship between cognition and frailty-A cross-sectional study among elders in the tribal areas of Mysuru***

Vanmathi A<sup>1</sup>, Praveen Kulkarni<sup>2</sup>, Dennis Chauhan<sup>3</sup>, Sunil Kumar D<sup>4</sup>, Prathiba Periera<sup>5</sup>, Kishor M<sup>6</sup>, Sayana basheer<sup>7</sup>

1-Post graduate, Department of community medicine, JSS Medical College, Mysuru-570015, Karnataka, India, *email id-vanmathianbu09@gmail.com*

2-Associate dean, JSS Academy of Higher Education and Research and Vice Principal, JSS Medical College, Mysuru-570015, Karnataka, India *email id: prakulfi@gmail.com*

3-Director, Department of health and community-based program, Vivekananda memorial hospital, Sargur, -571121, Karnataka, India, *email id:dennis@svym.org.in*

4-Head of the department, Department of community medicine, JSS Medical College, Mysuru-570015, Karnataka, India, *email id: sunilkumard@jssuni.edu.in*

5-Professor, Department of geriatric medicine, JSS Medical college hospital, Mysuru-570015, Karnataka, India, *email id: ppatta08@gmail.com*

6- Associate professor, Department of Psychiatry, JSS Medical college hospital, Mysuru-570015, Karnataka, India, *email id: rkishormd@gmail.com*

7- Postgraduate, Department of community medicine, JSS Medical College, Mysuru-570015, Karnataka, India, *email id: [dr.sayanabasheer@gmail.com](mailto:dr.sayanabasheer@gmail.com)*

**Corresponding author: Vanmathi A**

**email id- [vanmathianbu09@gmail.com](mailto:vanmathianbu09@gmail.com)**

**Contributorship**

VA, PK, SK, PP, KM Contributed equally to the conception and design of the study, VA, DC and, SB contributed to the acquisition of data. VA and PK contributed to analysing and interpreting data and drafting the article. PP and KM critically revised the draft for important intellectual content. All the authors approved the final version and were accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved

**Abstract**

Background: Cognition declines as age advances. Frailty is a pre-disability state. cognitive impairment and frailty lower the quality of life of elders. studies done in this background were mostly in urban or rural areas. Hence the current study was conducted among elders in

**NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.**  
the tribal areas to find the relationship between cognition and frailty. Materials and methods:

It was a cross-sectional study done for a period of one year among 316 elders >60 years. Based on the reported prevalence of cognitive dysfunction among the elderly to be 25% (Rakesh M Patel et al), with 5% absolute allowable error, 5% of alpha error, and 10% non-response rate) residing in the tribal areas of Mysuru. WHO- 30 cluster sampling was used. MOCA-B was used for assessing cognition scale and the TFI scale was used for assessing frailty. Results: The prevalence of cognitive impairment and frailty was 95.3 % and 61.4% respectively. Factors like Gender, education, type of family, physical, financial and emotional dependency, and preference of treatment were associated with cognition. There was also a significant association between cognition and frailty. Conclusion: There was a higher prevalence of cognitive impairment among frail elders. Thus, early diagnosis and appropriate management are necessary for healthy ageing

## Introduction

Ageing gracefully is an art <sup>[1]</sup>.

According to WHO, one in six individuals in the world will be 60 years of age or older by the year 2030. By 2050, 80 per cent of elderly citizens will reside in developing and middle-income nations <sup>[2]</sup> In India there were about 104 million elders according to 2011 census <sup>[3]</sup>

Ageing is a biological process caused by many cellular and molecular damages resulting in the deterioration of physical and mental abilities leading to increased susceptibility to diseases and eventually death. <sup>[4]</sup> While some changes in older people's health are inherited, the majority are caused by people's living conditions, neighbourhoods, and communities, as well as by their characteristics such as their sex, race, or financial position. <sup>[2]</sup>

Tribals constitute about 8.6% of the total Indian population. People in the tribal region are the most neglected and vulnerable population with a high degree of malnutrition, morbidity and mortality. The tribal elders play the role of culture transformer and this transformation predisposes them to psychosocial issues such as depression, anxiety, stress, and cognitive deficiencies <sup>[5],[6]</sup>

Cognitive function is one aspect of overall brain function and is an essential component for carrying out daily tasks [7]. Cognitive dysfunction in the elderly is characterised by loss of memory, Asking the same question repeatedly or telling the same tale repeatedly, Losing sight of familiar faces and surroundings, and Experiencing troubles in judgement, such as being unable to decide what to do in an emergency, Modifications in behaviour or mood, eyesight issues, Having trouble organising and completing tasks, like following a recipe or keeping track of monthly bills Furthermore, it has substantial social effects, resulting in a loss of autonomy and independence, as well as increasing demand for permanent caregivers and healthcare help. [8]

Poor nutrition, Midlife obesity, hyperlipidemia, midlife hypertension, diabetes, smoking, alcohol use, sleep disturbances, and depression are among the modifiable risk factors for cognitive dysfunction. While other factors, such as physical activity, education, and social engagement, reduce the risk of cognitive dysfunction. The development of cognitive impairment lowers the quality of life, leading to severe painful physical and emotional stress. [9],[10]Frailty is a pre-disability state American Geriatrics Society describes frailty as *"a state of increased vulnerability to stressors due to age-related declines in physiologic reserve across neuromuscular, metabolic, and immune systems"* [11]. Frailty May lead to several adverse health outcomes such as falls, disability, hospitalization [12] etc, recently it has been linked to cognition also

Most of the reported studies in this field have been conducted in hospitals or on older people living in rural or urban areas. Despite the fact that few studies have been conducted in tribal regions, they are related to general health conditions among the elderly. Taking this into consideration, we propose the current study to find the relationship between nutrition and cognitive functioning of the elderly living in tribal areas of Mysuru.

## **Method and materials:**

A cross-sectional study was done for a period of 6 months among elders >60 years residing in the tribal areas of Mysuru for a period of 6 months. A total of 316 elders (Based on the reported prevalence of cognitive dysfunction among the elderly to be 25% (Rakesh M Patel et al) [13], with 5% absolute allowable error and 5% of alpha error and 10% non-response rate) who were willing to participate in the study were included after obtaining necessary consent. In case, the elderly was illiterate/ not able to comprehend,

consent was obtained in presence of an impartial witness. Elders who had visual and auditory deficiencies, those who were not able to speak and diagnosed with mental illness were excluded

WHO- 30 clusters sampling technique <sup>[14]</sup> was used. The line listing of the population of all the tribal hamlets was done by collecting data from local panchayats and the cumulative total was calculated. The cumulative population was divided by sample size (316) to get the sampling interval. In each cluster total of 11 elders were interviewed (316/30). The first cluster was identified from the list where the sampling interval exists and the subsequent cluster was identified by adding the sampling interval to the population of the previous cluster. During the survey, the centre of the hamlet was identified and one road was randomly selected. A household survey was conducted on the elders satisfying the inclusion and exclusion criteria who were residing in the house. If a household had more than one elderly, only one was chosen randomly irrespective of gender. If a house was locked, an elderly from the subsequent house was interviewed. If a hamlet didn't have enough samples (i.e.<11 elders), elders from other hamlets which had a higher elderly population were chosen. Ethical committee approval was obtained before data collection.

The data collection had three parts. a) sociodemographic profile, b) cognitive assessment using MOCA -B scale <sup>[15]</sup>. c) Tilburg frailty indicator <sup>[16]</sup>(TFI) for assessing frailty

The data thus obtained was coded and entered into MS excel 2019 spreadsheet and analysed using SPSS version 25 (licensed to JSSAHER). Data were expressed using descriptive statistics like frequency, percentage and proportion. The Chi-square test/Fisher's Exact Test was used to find the association between various socio-demographic variables and cognition and also with frailty. A correlation between frailty and cognition was found using the Pearson correlation. A p-value of < 0.05 was considered statistically significant.

## **Results:**

### **Sociodemographic profile:**

In the study comprising 316 elderlies, most of the participants were females (60.8%,192) And the majority belonged to the age group of 60-65 years (56.6%, 179) and 25.3% (80) were in the age group of 66-70 years. Most of the elderly interviewed were illiterates (82.3%,260) and very few were literates (17.7%,56). only 46.8%,148 were currently working whereas 53.2%, (168) were not involved in any occupation. Almost 99.1%,313 were Hindus. More than half of the elders were married (58.2%,184),36.4 %,115 were widowers 16,5.1% were unmarried. 70.9%, (224) belonged to the nuclear family and27.8%,88 were in a joint family,

Among the study participants 63%,199 stated they were not dependent on others for their day-to-day activities and 4,1.3% were fully dependent on others, 55.4% (175) participants were independent financially while 15,4.7% were completely dependent on others When questioned about their emotional dependency 101,32 % stated that they were fully involved in family decisions whereas 153,48.4% were partially involved and 62,19.6% stated that they were never involved

Out of 316 elders interviewed 39(12.3%) had comorbidities. More than half of the study population (59.5%,188) preferred allopathy as their first preference,34.5%,109 preferred home remedies, 55.7% (176) had addictive habits and 32.3%, 102 said that they have sleep disturbances (Table 1)

**Table 1: socio-demographic profile of the study population**

<b>Variables</b>	<b>Categories</b>	<b>Frequency</b>	<b>Percent (%)</b>
<b>Gender</b>	Male	124	39.2
	Female	192	60.8
<b>Age</b>	60-65	179	56.6
	66-70	80	25.3
	71-75	34	10.8

	>75	23	7.3
<b>Education</b>	Literate	56	17.7
	Illiterate	260	82.3
<b>Currently working</b>	Yes	148	46.8
	No	168	53.2
<b>Religion</b>	Hindu	313	99.1
	Muslim	2	0.6
	Christian	1	0.3
<b>Marital status</b>	Married	184	58.2
	Widow/Widower	115	36.4
	Separated	1	0.3
	Unmarried	16	5.1
<b>Type of family</b>	Nuclear	224	70.9
	Three generation	4	1.3
	Joint/extended	88	27.8
<b>Living status</b>	Spouse and children	45	14.2
	Spouse	68	21.5

	Children	162	51.3
	Alone	41	13
<b>Physical dependency</b>	Independent	199	63
	Partially dependent	113	35.7
	Fully dependent	4	1.3
<b>Financial dependency</b>	Independent	175	55.4
	Partially dependent	126	39.9
	Fully dependent	15	4.7
<b>Emotional dependency</b>	Always	101	32
	Sometimes	153	48.4
	Never	62	19.6
<b>Co-morbidities</b>	Present	39	12.3
	absent	277	87.7
<b>Treatment modalities</b>	Allopathy	188	59.5
	Indigenous	1	0.3
	Traditional	8	2.5
	Home remedies	109	34.5

	Magico-religious	10	3.2
<b>Addictive habits</b>	Present	176	55.7
	Absent	140	44.3
<b>Sleep disturbances</b>	Present	102	32.3
	Not present	177	56
	Sometimes	37	11.7

### **Cognition:**

301,95.3% had cognitive impairment whereas 15, 4.7% were found to be normal (Fig 1), when graded (according to the moca B scale)13%,42 elders had severe cognitive impairment,54.1%,171 had moderate and 28.2%,89 had mild cognitive impairment (Fig 2)

#### **Fig 1: Prevalence of cognitive impairment among the study population**

#### **Fig 2: Distribution of study population according to the severity of cognitive impairment**

Gender, education, type of family, physical, financial and emotional dependency, and preference of treatment were associated with a cognitive impairment which was statistically significant (Table 2)

**Table 2: Factors associated with cognitive impairment**

<b>Variables</b>	<b>Categories</b>	<b>Cognitive impairment present</b>	<b>Cognitive impairment absent</b>	<b>Chi-Square value</b>	<b>P value</b>



		<b>(301) (n, %)</b>	<b>(15) (n, %)</b>		
<b>Gender</b>	Male	113(37.5)	11(73.3)	7.677	<b>0.006*</b>
	Female	188(62.5)	4(26.7)		
<b>Age</b>	60-65	167(55.5)	12(80)	2.560	0.409
	66-70	78(25.9)	2(13.3)		
	71-75	33(11)	1(6.7)		
	>75	23(7.6)	0		
<b>Education</b>	Literate	46(15.3)	10(66.7)	25.87 3	<b>&lt;0.001*</b>
	Illiterate	255(84.7)	5(33.3)		
<b>Currently working</b>	Yes	138(45.8)	10(66.7)	2.487	0.115
	No	163(54.2)	5(33.2)		
<b>Religion</b>	Hindu	299(99.3)	14(93.3)	6.725	0.136
	Muslim	1(0.3)	1(6.7)		
	Christian	1(0.3)	0		
<b>Marital status</b>	Married	17(57.1)	12(80.0)	3.713	0.346
	Widow/Widower	112(37.2)	3(20.0)		
	Separated	1(0.3)	0		

	Unmarried	16(5.3)	0		
<b>Type of family</b>	Nuclear	218(71.8)	6(40.0)	9.688	<b>0.022*</b>
	Three generation	3(1.0)	1(6.7)		
	Joint/extended	80(26.6)	8(53.3)		
<b>Living status</b>	Spouse and children	42(14.0)	3(20.0)	2.720	0.450
	Spouse	64(21.3)	4(26.7)		
	Children	154(51.2)	8(53.3)		
	Alone	41(13.6)	0		
<b>Physical dependency</b>	Independent	185(61.5)	14(93.3)	6.716	<b>0.03*</b>
	Partially dependent	112(37.2)	1(6.7)		
	Fully dependent	4(1.3)	0		
<b>Financial dependency</b>	Independent	161(53.5)	14(93.3)	8.926	<b>0.011*</b>
	Partially dependent	125(41.5)	1(6.7)		
	Fully dependent	15(5.0)	0		

<b>Emotional dependenc y</b>	Always	90(29.9)	11(73.3)	11.49 1	<b>0.002*</b>
	Sometimes	149(49.5)	4(26.7)		
	Never	62(20.6)	0		
<b>Co- morbidity</b>	Present	36(12.0)	3(20.0)	0.854	0.410
	absent	265(88.0)	12(80.0)		
<b>Treatment modalities</b>	Allopathy	179(59.5)	9(60)	9.193	<b>0.05*</b>
	Indigenous	0	1(6.7)		
	Traditional	7(2.3)	1(6.7)		
	Home remedies	105(34.9)	4(26.7)		
	Magico- religious	10(3.3)	0		
<b>Addictive habits</b>	Present	168(55.8)	8(53.3)	0.036	1.000
	Absent	133(44.2)	7(46.7)		
<b>Sleep disturbanc es</b>	Present	95(31.6)	7(46.7)	3.792	0.102
	Not present	172(57.1)	5(33.3)		
	Sometimes	34(11.3)	3(20.0)		

n=Frequency,%=Percent,\*-Statistically significant p <0.05

**Frailty:**

61.4% (194) of the study population were frail according to the TFI scale whereas 38.6% (122) were non-frail (Fig 3).

**Fig 3: Distribution of study population based on frailty**

Factors like age, education, marital status, type of family, living status, Physical and emotional dependency and also the presence of comorbidities were significantly associated with frailty. (Table 3)

**Table 3: Factors associated with frailty**

<b>Variables</b>	<b>Categories</b>	<b>Frail (194), (n, %)</b>	<b>Non-frail (122), (n, %)</b>	<b>Chi- Square value</b>	<b>P value</b>
<b>Gender</b>	Male	70(36.1)	54(44.3)	2.102	0.147
	Female	124(63.9)	68(55.7)		
<b>Age</b>	60-65	100(51.5)	79(64.8)	9.13	0.028*
	66-70	60(30.9)	20(16.4)		
	71-75	19(9.8)	15(12.3)		
	>75	15(7.7)	8(6.6)		
<b>Education</b>	Literate	27(13.9)	29(23.8)	4.987	0.026*
	Illiterate	167(86.1)	93(76.2)		
<b>Currently working</b>	Yes	93(47.9)	55(45.1)	0.245	0.645
	No	101(52.1)	67(54.9)		
<b>Religion</b>	Hindu	192(99.0)	121(99.2)	2.848	0.241

	Muslim	2(1)	0		
	Christian	0	1(0.8)		
<b>Marital status</b>	Married	101(52.1)	83(68)	13.778	0.002*
	Widow/Widower	85(43.8)	30(24.6)		
	Separated	1(0.8)	0		
	Unmarried	7(3.6)	9(7.4)		
<b>Type of family</b>	Nuclear	154(79.5)	70(57.4)	18.225	<0.001*
	Three generation	1(0.5)	3(2.5)		
	Joint/extended	39(20.1)	49(40.2)		
<b>Living status</b>	Spouse and children	32(16.5)	13(10.7)	11.496	0.009*
	Spouse	33(17)	35(28.7)		
	Children	97(50)	65(53.3)		
	Alone	32(16.5)	9(7.4)		
<b>Physical dependency</b>	Independent	111(57.2)	88(72.1)	8.621	0.013*
	Partially dependent	79(40.7)	34(27.9)		

	Fully dependent	4(2.1)	0		
<b>Financial dependency</b>	Independent	106(54.6)	69(56.6)	4.278	0.118
	Partially dependent	75(38.7)	51(41.8)		
	Fully dependent	13(6.7)	2(1.6)		
<b>Emotional dependency</b>	Always	49(25.3)	52(42.6)	12.958	0.001*
	Sometimes	98(50.5)	55(45.1)		
	Never	47(24.2)	15(12.3)		
<b>Co-morbidities</b>	Present	32(16.5)	7(5.7)	8.011	0.005*
	absent	162(83.5)	115(94.3)		
<b>Treatment modalities</b>	Allopathy	123(63.4)	65(53.3)	6.745	0.166
	Indigenous	0	1(0.8)		
	Traditional	3(1.5)	5(4.1)		
	Home remedies	61(31.4)	48(39.3)		
	Magico-religious	7(3.6)	3(2.5)		
<b>Addictive</b>	Present	116(59.8)	60(55.7)	3.419	0.064

<b>habits</b>	Absent	78(40.2)	62(50.8)		
<b>Sleep disturbances</b>	Present	66(34)	36(29.5)	1.179	0.55
	Not present	104(53.6)	73(59.8)		
	Sometimes	24(12.4)	13(10.7)		

n=Frequency,%=Percent,\*-Statistically significant p <0.05

### Frailty and cognition:

Frailty and cognitive were found to be statistically significant(p<0.05) (table 4).

**Table 4: Association between frailty and cognition**

<b>Variables</b>	<b>Categories</b>	<b>Frail (194), (n, %)</b>	<b>Non-frail (122), (n, %)</b>	<b>Chi-Square value</b>	<b>P value</b>
<b>Cognitive impairment</b>	<b>Present</b>	189(62.8)	112(37.2)	5.231	0.02
	<b>Absent</b>	5(33.3)	10(66.7)		

n=Frequency,%=Percent,\*-Statistically significant p <0.05

when Pearson correlation was applied it was found that MoCA-B scores and TFI scores are negatively correlated which explained that as cognition increases frailty score decreased but the results were not statistically significant (Fig 4)

**Fig 4: scatter plot which shows the correlation between MoCA-B scores and TFI scores**

### Discussion:

Healthy ageing is a significant challenge in this 21<sup>st</sup> century. The elderly who lives in tribal regions will serve as a link between the traditional tribal way of life and the modernization initiatives being carried out by the government and non-governmental

organisations. Being elderly increases their vulnerability to physical and mental health issues [17] A study was conducted on 952 tribal people in the Vellore districts of South India by S. Sathiyarayanan et al. The majority in that survey (82.2%) were illiterate and female (59.8%), which was in accordance with the results of this study [18].

A study using the Bims scale (brief interview for mental state) done among elders in Chengalpattu, Tamilnadu by *Sujitha P et al* found that nearly 44% had mild-moderate cognitive impairment, and 36% had severe cognitive impairment. Only 20% of the respondents had intact cognitive levels [19] where as in our study when graded (according to the moca B scale) 13%, 42 elders had severe cognitive impairment, 54.1%, 171 had moderate and 28.2%, 89 had mild cognitive impairment.

In the current study, the prevalence of frailty was 61.4%, and female elderly were frailer than male elderly. Similar results were obtained in a study by R P Thakur et al. [20], where it was found that their frailty was 83.29% more among female elders than male elders. This might be a result of the study's participants being predominantly female.

The concomitant prevalence of frailty and cognitive changes can greatly predict mortality in the elderly [21] According to the study conducted by *Isabela Thaís Machado de Jesus et al* among 247 elders in the city of Sao Carlos, 87.4% (216) had cognitive impairment, and 84 (38.88%) had some level of frailty (mild, moderate or severe). Whereas in our study out of 301 elders who had cognitive impairment 189 (62.8) was frail. [22] Another study done among 84 elders on a haemodialysis unit by *Jyotish Chalil Gopinathan et al* Cognitive impairment was present in 89.7% of patients and significantly associated with frailty ( $P < 0.001$ , Fisher's exact test) [23] The variations may be due different scales used for assessing cognition and frailty. The limitation of the study includes a smaller sample size and was not able to cover a larger geographical area.

## Conclusion

Thus, the study showed that the tribal elders had a high prevalence of cognitive impairment and the prevalence of cognitive impairment was relatively high among frail people. Although frailty and cognitive decline are part of ageing, its onset and progression can be delayed by appropriate management. Elders diagnosed with cognitive dysfunction should be counselled and regularly monitored. Elderly who are frail should receive counselling on lifestyle improvements including exercise, appropriate nutrition intake, and frequent health checks,



among other things, for healthy ageing. All older people with pre-frailty should receive dietary and pharmaceutical supplements of micro and macronutrients. Geriatric clinics may be established at the primary care level to ensure appropriate care and service for the elderly, Additionally, there aren't enough medical professionals from tribal communities who would be more inclined to work in these places. Medical outreach camps and mobile health clinics can also contribute to bringing health services to remote populations.

## Acknowledgement


We authors were very thankful to the health facilitators of Vivekananda memorial hospital, sargur, Mysuru for their contribution for conducting the study. we were also grateful to the department of community medicine, JSS medical college for their cooperation

## References

1. Harris M. Is there a role for micronutrients in the prevention of cognitive declines associated with ageing. *Vitam Trace Elem.* 2012;1(2):1000e105
2. Ageing and health [Internet]. Who. int. 2022. Available from: <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health>
3. Ageing and health India [Internet]. Who. int. 2022. Available from: [https://www.who.int/india/health-topics/ageing#:~:text=According%20to%20Census%202011%2C%20India,60%2B\)%2C%20females%20outnumber%20males](https://www.who.int/india/health-topics/ageing#:~:text=According%20to%20Census%202011%2C%20India,60%2B)%2C%20females%20outnumber%20males)
4. Kadri A. *Iapsms Textbook Of Community Medicine.* 1st Ed. [S.L.]: Jaypee Brothers Medical P; 2019
5. Santhosam MA, Samuel U. A study on the health status of elderly Irular tribal women in Kancheepuram District. *IOSR Journal of Humanities and Social Science.* 2013;7(2):1-4.
6. Das S, Bose K. Adult tribal malnutrition in India: an anthropometric and socio-demographic review. *AnthropologicAl review.* 2015 Mar 30;78(1):47-65.
7. Cognitive health and older adults [Internet]. National Institute of ageing. Available from: <https://www.nia.nih.gov/health/cognitive-health-and-older-adults>
8. Jutkowitz E, Gozalo P, Trivedi A, Mitchell L, Gaugler JE. The effect of physical and cognitive impairments on caregiving. *Medical care.* 2020 Jul;58(7):601.


9. Sengupta P, Benjamin AI, Singh Y, Grover A. Prevalence and correlates of cognitive impairment in a north Indian elderly population. *WHO South-East Asia journal of public health*. 2014;3(2):135-43
10. Han F, Luo C, Lv D, Tian L, Qu C. Risk Factors Affecting Cognitive Impairment of the Elderly Aged 65 and Over: A Cross-Sectional Study. *Frontiers in Aging Neuroscience*. 2022;14.
11. Gómez-Gómez ME, Zapico SC. Frailty, Cognitive Decline, Neurodegenerative Diseases and Nutrition Interventions. *Int J Mol Sci*. 2019;20(11):2842. Published 2019 Jun 11. doi:10.3390/ijms20112842
12. Dasgupta A, Bandyopadhyay S, Bandyopadhyay L, Roy S, Paul B, Mandal S. How frail are our elderly? An assessment with Tilburg frailty indicator (TFI) in a rural elderly population of West Bengal. *J Fam Med Prim Care*. 2019 Jul;8(7):2242–8.
13. Patel RM, Singh US. Prevalence study of cognitive impairment and its associated sociodemographic variables using mini-mental status examination among elderly population residing in field practice areas of a medical college. *Indian journal of community medicine: official publication of Indian Association of Preventive & Social Medicine*. 2018 Apr;43(2):113.
14. World Health Organization. Training for mid-level managers (MLM): Module 7: the EPI coverage survey.
15. Julayanont P;Tangwongchai S;Hemrungronj S;Tunvirachaisakul C;Phanthumchinda K;Hongswat J;Suwichanarakul P;Thanasirorat S;Nasreddine ZS; The Montreal Cognitive Assessment-BASIC: A screening tool for mild cognitive impairment in illiterate and low-educated elderly adults [Internet]. *Journal of the American Geriatrics Society*. U.S. National Library of Medicine; Available from: <https://pubmed.ncbi.nlm.nih.gov/26648041/>
16. Gobbens RJ, Boersma P, Uchmanowicz I, Santiago LM. The Tilburg Frailty Indicator (TFI): new evidence for its validity. *Clinical interventions in aging*. 2020;15:265.
17. Arlappa N, Balakrishna N, Brahmam GN, Vijayaraghavan K. Nutritional status of the tribal elderly in India. *Journal of Nutrition for the Elderly*. 2006 Dec 13;25(2):23-39.
18. Sathiyarayanan S, Muthunarayanan L, Devaparthasarathy TA. Changing perspectives in tribal health: Rising prevalence of lifestyle diseases among tribal population in India. *Indian Journal of Community Medicine: Official Publication of Indian Association of Preventive & Social Medicine*. 2019 Oct;44(4):342.

19. Pandian S, Swetha NB, Umadevi R, Grace G A, Gopalakrishnan S. A cross sectional study on assessment of cognitive dysfunction and behavioural risk factors among senior citizens living in old age homes in Chengalpattu District, Tamilnadu. *National Journal of Community Medicine*. 2022;13(07):446–52.
20. Thakur R, Banerjee A, Nikumb V. Health Problems Among the Elderly: A CrossSectional Study. *Ann Med Health Sci Res*. 2013;3(1):19–25
21. Brigola AG, Rossetti ES, Santos BR, Neri AL, Zazzetta MS, Inouye K, Pavarini SC. Relationship between cognition and frailty in elderly: A systematic review. *Dementia & Neuropsychologia*. 2015 Apr;9:110-9.
22. Jesus IT, Orlando FD, Zazzetta MS. Frailty and cognitive performance of elderly in the context of social vulnerability. *Dementia & neuropsychologia*. 2018 Apr;12:173-80.
23. Gopinathan JC, Hafeeq B, Aziz F, Narayanan S, Aboobacker IN, Uvais NA. The Prevalence of Frailty and its Association with Cognitive Dysfunction among Elderly Patients on Maintenance Hemodialysis: A Cross-Sectional Study from South India. *Saudi Journal of Kidney Diseases and Transplantation*. 2020 Jul 1;31(4):767.




medRxiv preprint doi: <https://doi.org/10.1101/2023.01.31.23285235>; this version posted February 1, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

**Fig 2: Distribution of study population according to the severity of cognitive impairment**




medRxiv preprint doi: <https://doi.org/10.1101/2023.01.31.23285235>; this version posted February 1, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).



medRxiv preprint doi: <https://doi.org/10.1101/2023.01.31.23285235>; this version posted February 1, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](#).

Figure



Figure