- 1 Sexually transmitted infection knowledge levels, socio-demographic characteristics and
- 2 sexual behaviour among men who have sex with men: results from a cross-sectional
- 3 survey in Nairobi, Kenya.

4 Delvin Kwamboka Nyasani<sup>1,2</sup>, Meshack Onyambu<sup>1</sup>, Laura Lusike Lunani<sup>2</sup>, Geoffrey

#### 5 Oino Ombati<sup>1</sup>, Gaundensia Nzembi Mutua<sup>3</sup>, Matt Price<sup>3</sup>, <sup>4</sup>, Justus .O.Osero<sup>1</sup>

1 Kenyatta University Department of Community Health and Epidemiology, Nairobi, Kenya
2 KAVI Institute Of Clinical Research, University Of Nairobi, Nairobi, Kenya. 3. IAVI, New
York, United States of America. 4 Department of Epidemiology and Biostatistics, University
of California, San Francisco, California, United States of America.

10

#### 12 Abstract

#### 13 Background:

14 High rates of sexually transmitted infections (STIs) among men who have sex with men

15 (MSM) have been reported, there is little research on their STI knowledge. Our study sought

16 to characterize knowledge and awareness of STIs among MSM in Nairobi, Kenya.

#### 17 Methods

We mobilized MSM aged ≥18 years from Nairobi into a cross-sectional study. To determine
their understanding about STIs, a pre-tested structured questionnaire was administered.
Knowledge score was generated by summing up the number of responses answered correctly

by a participant. We dichotomized scores as "low" and "high", by splitting the group at <12

and  $\geq 12$  which was the mean.

#### 23 **Results**

A total of 404 participants were interviewed between the month of March and August 2020.

- 25 The mean age was 25.2 (SD=6.4) years. Majority were single (80.4%) and Christians
- 26 (84.2%). All participants had some formal education ranging from primary to tertiary; the

<sup>11</sup> Corresponding author, email: <u>dnyasani@kaviuon.org</u> or dnyasani01@yahoo.com

majority (92.3%) had secondary education or more. Most (64.0%) were employed and their
monthly income ranged from <50->150 USD. Almost all (98.5%) were Kenyans.

Of the 404 (90.6%) self-identified as male and (47.5%) reported to be exclusively top partners. Many (39.9%) reported being both bottom and top, while those reporting to be bottom partners were, (12.6%). The last 12 months, (55.4%) of the participants reported having sex with men only and (88.6%) reported to have had more than one sexual partner.

33 Participants scored an average of 12.2, out of 29 SD 4.5. The multivariable modelling revealed that participants aged  $\geq 25$  years were more likely to have a higher knowledge score 34 35 compared with the participants aged 18-24 years (aOR=0.973, CI: 0.616-1.538). Regarding education and occupation, participants who had tertiary education and those who were 36 employed were more likely to have a higher knowledge score compared with the participants 37 who had primary education and were not employed ( aOR=2.627, CI:1.142-6.043) 38 39 (aOR=0.922, CI:0.401-2.117). Participants who were earning (USD >150) were more likely to have a higher knowledge score compared to the ones who were not earning (aOR=2.520, 40 CI: 0.900-7.055). Further bisexual men were more likely to have a higher knowledge score 41 compared with the participants who were having sex with men only (aOR= 1.550, CI: 1026-42 2.342) 43

#### 44 Conclusion

45 Participant's knowledge level regarding STIs was low. We recommend health care workers46 to continue educating patients about STIs.

47

49

- 50
- 51

#### 52 Introduction

#### 53 Background

Over 1 million sexually transmitted infections (STIs) are acquired daily globally; common among them are chlamydia, gonorrhoea, syphilis and trichomoniasis. Many of these infections are frequently mild or asymptomatic (1) In low- and middle-income countries (LMICs), symptomatic STIs are often treated by syndromic management which frequently misses asymptomatic STIs that therefore remain untreated ad potentially infectious (2). Regardless of symptoms, STIs can cause serious morbidity including cancer, infertility, and enhanced HIV transmission (3,4).

61

62 Prevalence rates of STIs vary from area to area, nation to nation and between urban and rural 63 populations. The infections are more prevalent in Asia and sub-Saharan Africa (SSA), but 64 SSA is estimated to contribute a greater proportion of the cases of STIs every year, 65 highlighting a need for intervention (5).

66

The impact of STI epidemics is of particular importance among key, higher-risk populations who include: young people and adolescents, men who have sex with men (MSM), people in prisons, sex workers and people who inject drugs. These populations are often vulnerable to both HIV and other STIs not only because of their high-risk sexual and drug use behaviours, but also by structural barriers such as low access to quality health services, stigma and discrimination(6).

The risk may even be even greater among MSM because unprotected receptive anal sex; is a
greater risk factor for STI and HIV acquisition than unprotected vaginal sex (7).

In developed countries such as United States, England and Netherlands the data as also
shown that MSM bear a much higher burden of new cases of STI and HIV than their
heterosexual male counterparts (8–11)

The situation is similar in Africa, where increasing attention has been spent describing the epidemiology of STIs and HIV among MSM in recent years. In Uganda, a study conducted among 300 MSM, found that they had a higher HIV prevalence 13.7% compared to the general adult male population at 4.5% (12).

In Kenya, a study conducted in Nairobi among 563 male sex workers and other MSM found that the prevalence of testing positive for one or more STIs combined (Syphilis, gonorrhoea or chlamydia) among male sex workers was higher (15.0%) compared to those who were not sex workers (5.3%) (13). Also according to (14) among the estimated 22,000 MSM in Kenya, the HIV prevalence was 18.2% compared to the national general population at 5.9%.

87

According to (15) comprehensive HIV and AIDS knowledge among sexually active men had reduced the odds of them contracting STIs. Being that the HIV prevalence among MSM in Kenya is estimated to be three times higher than that of the general population, we sought to characterize knowledge and awareness of STIs among MSM in Nairobi City County, Kenya.

- 93
- 94
- 95 96
- 97
- 98
- 50
- 99

#### 100 Materials and Methods

#### 101 Study design

102 This was a cross-sectional study among sexually active MSM within Nairobi, Kenya, 103 conducted between the months of March-August 2020. We interviewed participants to 104 determine their understanding of STIs and assessed for association with socio-demographic 105 characteristic and sexual behaviour.

#### 106 Study participants

107 The study was designed to enrol men aged ≥18 years and who have sex with men from 108 places where MSM are known to either solicit their clients or where sexual activity was 109 common including " cruising" areas along streets, or bars with or without lodging (16). 110 They had to be willing and able to give informed consent to participate. MSM who were 111 obviously intoxicated were excluded from participation.

112

#### 113 **Participant recruitment**

Mobilization was done by MSM community mobilizers. We identified MSM community mobilizers from the Nairobi-based Sex Workers Outreach Programme (SWOP) with whom we had partnered for a recent study (17). The community mobilizers gave a brief description of the research activities to potential participants. Then using a systematic random sampling procedure they selected participants with the desired characteristics. A systematic random sampling is a sampling technique that the researcher selects participants to be included in the sample based on a systematic rule, using a fixed interval.

We had twenty one days to enrol 422 participants. We had purposively selected 5 subcounties that were densely populated with MSM(16). To get the percentage of the participants to be accessed from each sub-county, the researcher took the expected number in a specific sub-county divided by the total participants from all the 5 sub-counties and multiplied by 100.Then to get the numbers to be recruited from each sub county, it was the

- 126 percentage of the participants to be accessed from each sub-county multiplied by the required
- number of participants (n=422) (Table 1).
- 128Table 1: The Number of MSM in the selected Sub-Counties, at selected MSM geographic
- areas In Nairobi, Kenya and the sampled population.
- 130

Sub-County	Estimated number of MSM	Percentage of the	Number of MSM
_	available on all selected MSM	total MSM to be	required for the
	geographic areas	accessed	study ((n=422)
Starehe	2000 (40 MSM geographic areas)	55%	232
Westlands	1000 (27 MSM geographic areas)	27%	114
Dagoretti	360 (20 MSM geographic areas)	10%	42
South			
Ruaraka	180 (15 MSM geographic areas)	5%	21
Kasarani	100 (10 MSM geographic areas)	3%	13
Total	3640	100	422

#### 131

132

To get the first participant from each sub-county on each day of mobilization, we had 2 133 sealed papers written "Yes and No". The mobilizers issued the first two participants the 134 sealed papers if for example, the second participant selected "Yes", then they were issued 135 with a unique identification card. Then they continued mobilizing the second participant until 136 the required number was attained. If the sampled participant had declined to participate, the 137 next exiting participant was sampled. The sampled participants were then referred to the 138 researcher and the research assistants for interviews at a selected venue. The selected venue 139 was a safe and private place identified by the mobilizers where the interviews could be 140 conducted; without attracting attention to the participants. At the entrance to the selected 141 venue, a peer navigator confirmed participant identity by confirming that they had a unique 142 identification card and that they were not intoxicated. Thereafter informed consent was 143 obtained from the participants before the questionnaire was administered. Mobilization from 144 the MSM geographic areas was done on 4 selected peak days (i.e., Wednesday, Friday, 145 Saturday and Sunday) as identified by the key population implementing partners during the 146

key population size estimate mapping(16). During that time also the peak time identified
was 18.00hrs-22.00hrs and that is what we adopted. Peak day refers to a day when the
number of MSM present at the MSM geographic areas was more than usual.

- 150
- 151 **Procedure**

Participants were assigned a unique study identification number. Data was collected about socio-demographics and sexual behaviour including: age, marital status, religion, education level, occupation, level of income, nationality, gender identity, roles during sexual activity, sex with whom in the past 12 months, number of sexual partners in the past 12 months, type of sex, if received gifts/money in exchange for sex, whether gave gifts/money in exchange for sex and history of having ever contracted any STI.

To determine their understanding about STIs participants were then given a pre-tested 158 structured questionnaire (Supplemental material). The questionnaire had four open ended 159 questions and one short answer question. Interviewers were instructed not to prompt 160 participants but only tally all correct responses provided. If a participant gave an incorrect 161 answer it was ignored but if it was correct and not among the options given it was captured as 162 other and later analysed. The lead researcher in the field did quality control checks on the 163 questionnaires before the participant was reimbursed, allowing the team to avoid missing data 164 or clarify ambiguities. 165

166 Study Variables

#### 167 **Dependent variable**

Knowledge score was the dependent variable. In assessing knowledge levels the participants were asked the following questions; which diseases are spread via sexual intercourse? What the signs and symptoms of STIs were? How one could be infected with STIs? Can someone be infected with STIs via anal sexual intercourse? And Through which media could one get infected with STIs? (Additional details are shown in the supplemental material). Interviewerswere instructed not to prompt participants but only tally all correct responses provided.

174 A knowledge score was generated by summing up the number of responses answered 175 correctly by a participant; to give a score ranging from 0 to 29. For the purposes of analysis, 176 we dichotomized scores as "low" and "high", by splitting the group at <12 and  $\geq$ 12 177 (additional details are shown in Appendix 1).

#### **178** Independent variables

Independent variables included in this analysis included: age, marital status(single never 179 180 married, married and single ever married), religion (Christians and non-Christians), education (primary, secondary and post-secondary), occupation (unemployed and employed), level of 181 income (No income, <5000, 5001-10,000, 10001-15000 and >15000), nationality (Kenyans 182 and non-Kenyans), gender identity (male, transgender woman, intersex and non-conforming), 183 roles during sexual activity (top, bottom or versatile), sex with whom in the past 12 months 184 (men or both men and women), number of sexual partners in the past 12 months (one or more 185 than one), type of sex (vaginal, anal, oral, mutual masturbation), received gifts/money in 186 exchange for sex (yes, all the time, yes, sometimes, No), gave gifts/money in exchange for 187 sex (yes, all the time, yes, sometimes, No), and ever contracted STI (Yes, No) (see Table 1 188 and 2). 189

#### 190 Data analysis

Data was entered into statistical package for social sciences programme (IBM-SPSS) version 25. Quantitative data from the study questionnaires was coded. Single data entry was done into SPPS database after quality control checks had been done. Further cleaning was carried out after data entry using frequency distributions and cross tabulations until no more errors were detected. Participant's characteristics are presented by use of frequencies and percentages for categorical variables. Pearson's Chi-square and Fishers exact test where appropriate was done to characterize any statistical significant associations between STI knowledge, demographics and reported sexual behaviour. Variables with a  $p \le 0.05$  and age being a priori variable were included in a multivariable logistic regression model to characterize factors associated with higher or lower participants STI knowledge.

#### 201 Ethical consideration and study approval number

Approval of the study was sought from Kenyatta University Board of Post Graduate studies.

203 Ethical Clearance was given by Kenyatta University ethics review committee;

(PKU/1071/11121). Permission to conduct the study was sought from the National Council
for Science, Technology and Innovations (NACOSTI), and from Nairobi County and subcounty facilities. Confidentiality and anonymity of the information given by the participants
was protected by ensuring that the names of the participants were not indicated in the data
collection tools. Participants were reimbursed in the local currency equivalent of \$2 USD in
compensation for their time.

- 210
- 211
- 212
- 213
- 214
- 215
- 216
- 217
- 218
- 219
- 220
- 221

#### 222 **Results**

223 Demographics

A total of 404 participants were interviewed between the month of March and August 2020. They had a mean age of 25.2 (SD=6.4) years. The majority were single never married (80.4%; 325) and Christians (84.2%; 340). All participants had some formal education ranging from primary to tertiary level; the majority (92.3%; 373) had secondary education or more. Most of the participants (64.0%; 259) were employed and their level of monthly income ranged from \$<50->150 USD. Almost all the participants (98.5%; 398) were Kenyans. (Table 2)

231

232

233

234

235

# Table 2: Socio-demographic characteristics of study participants and their association with participant's knowledge score.

	Total	Knowledge score		
	(n=404)	<12	≥12	
Characteristics		(n=215)	(n=189)	P Value
	n(%)	n(%)	n(%)	
Mean age=25.20, SD=6.	41			
Age bracket				
18-24	241(59.7)	127(52.7)	114(47.3)	
≥25	163(40.3)	73(44.8)	90(55.2)	
				0.129
Marital Status				
Single never married	325(80.4)	162(49.8)	163(50.2)	
Married	54(13.4)	27(50.0)	27(50.0)	0.859
Single ever married	25(6.2)	11(44.0)	14(56.0)	
Religion				
Christian(Catholic &	340(84.2)	174(51.2)	166(48.8)	0.135
protestants)				
Non-Christian (Muslim,	64(15.8)	26(40.6)	38(59.4)	
Atheists, Hindu)				
Education level				
Primary	31(7.7)	19(61.3)	12(38.7)	0.001
Secondary	226(55.9)	126(55.8)	100(44.2)	
Post-secondary	147(36.4)	55(37.4)	92(62.6)	

238	Occupation				
	Unemployment/Student	145(36.0)	85(58.6)	60(41.4)	0.007
239	Employment/Self-	259(64.0)	115(44.4)	144(55.6)	
	employed				
240	Level of income				
2/1	None	120(29.7)	75(62.5)	45(37.5)	
241	<5000	54(13.4)	28(51.9)	26(48.1)	0.005
212	5001-10000	87(21.5)	38(43.7)	49(56.3)	
242	10001-15000	71(17.6)	33(46.5)	38(53.5)	
243	>15000	72(17.8)	26(36.1)	46(63.9)	
2.13	Nationality				
244	Kenyans	398(98.5)	196(49.2)	202(50.8)	0.446
	Non-Kenyans	6(1.5)	4(66.7)	2(33.3)	
245	5 <b>* KEY: by Chi-square test or Fisher's Exact Test where cell size was small</b>				

246

247

248

- 249
- 250

#### 251 Sexual behaviour

Most of the participants (90.6%; 366) self-identified as male and almost half (47.5%; 192) of them reported to be exclusively insertive ("Top") partners. Many (39.9%; 161) reported being versatile (both bottom and top), while those reporting to be receptive ("Bottom") partners were, (12.6%; 51). In the last 12 months, (55.4%; 224) of the participants reported having sex with men only and (44.6 %; 180) reported being bisexual. Also during that period, majority of the participants (88.6%; 358) reported to have had more than one sexual partner (Table 2).

Majority of the respondents (93.6%; 378) reported to have had anal sex, (42.1%; 170) had
vaginal sex, (28.5%; 115) reported oral and (19.6%; 79) reported mutual masturbation.
Almost half (47.5%; 192) of the participants, reported receiving gifts/money in exchange for

- sex, sometimes. Majority of the participants (95.3%; 385), engaged in transactional sex.
- Participants who reported to have ever contracted STI were (43.3%; 175). (Table 3)

### 264 Table 3: Sexual behavioural characteristics of study participants and their association

- 265 with participant's knowledge score
- 266

		Total	Knowledge score		
267		(n=404)	<12	≥12	
	Characteristics		(n=215)	(n=189)	P Value
268		n(%)	n(%)	n(%)	
	Gender	· · ·	• • •		
269	Male	366(90.6)	176(48.1)	190(51.9)	
	Transgender women	13(3.2)	8(61.5)	5(38.5)	0.364
270	Intersex	17(4.2)	11(64.7)	6(35.3)	
	Non-conforming	8(2.0)	5(62.5)	3(37.5)	
271	Roles during sexual act	ivity			
	Top(Insertive)	192(47.5)	94(49.0)	98(51.0)	0.209
272	Bottom(Receptive)	51(12.6)	31(60.8)	20(39.2)	
272	Versatile (Both	161(39.9)	75(46.6)	86(53.4)	
2/3	insertive & receptive)				
274	Sex with whom in the past 12 months				
2/4	Men	224(55.4)	123(54.9)	101(45.1)	
275	Both men and women	180(44.6)	77(42.8)	103(57.2)	0.017
275	Number of sexual partr	ers in the <b>p</b>	oast 12 mon	ths	
276	Only 1	46(11.4)	30(65.2)	16(34.8)	
270	>1	358(88.6)	170(47.5)	188(52.5)	0.584
277	Type of sex				
_,,	Vaginal	170(42.1)	69(72.4)	101(27.6)	0.003
278	Anal	378(93.6)	185(48.9)	193(51.1)	0.423
-	Oral	115(28.5)	50(43.5)	65(56.5)	0.152
279	Mutual masturbation	79(19.6)	34(43.0)	45(57.0)	0.212
	Received gifts/money in				
280	Yes, all the time	38(9.4)	14(36.8)	24(63.2)	0.163
	Yes, sometimes	192(47.5)	93(48.4)	99(51.6)	
281	No	174(43.1)	93(53.4)	81(46.6)	
	Gave gifts/money in exchange for sex				
282	Yes, all the time	16(4.0)	5(31.3)	11(68.8)	0.337
	Yes, sometimes	139(34.4)	71(51.1)	68(48.9)	
283	No	249(61.6)	124(49.8)	125(50.2)	
	Ever contracted STI				0.271
284	Yes	175(43.3)	81(46.3)	94(53.7)	
-	No	229(56.7)	119(52.0)	110(48.0)	
285	*Key: by Chi-square te	est or Fisher	's Exact Te	st where cell	size was small

#### 287 STI knowledge

- Out of a total of 29 possible correct answers the highest participant scored 25, we observed
- an average score of 12.2 correct, SD 4.5 (additional details shown in Appendix 1).As
- indicated on Table 4 below, majority of the participants were aware of gonorrhoea (92.8 %;
- 291 375), syphilis (88.1%; 356), HIV (65.6%; 265), genital warts (33.9%; 137), Herpes Simplex (
- 292 22.8%; 92), chancroid (20.5%;83), hepatitis B (17.3%;70), chlamydia (13.6%;55) and
- trichomoniasis (5.2%;21).
- 294 Regarding signs and symptoms of STIs, most of the participants were cognizant of penile
- discharge (66.8%; 270), burning sensation during urination (60.4%; 244), genital ulcer/sores
- 296 (39.6%; 160), anal discharge (21.0%; 85), swelling in the groin region (23.0%; 93), anal
- ulcer/sore (25.0 %; 101), anal pain (21.3%; 86), and scrotal swelling (14.1%; 57).
- Also almost all the participants knew that STIs were transmitted through unprotected sexual intercourse (97.5%; 394) i.e. vaginal, anal, oral and mutual masturbation. Also about the anal sexual intercourse risk to STIs, majority of the participants (83.7%; 338) were aware that they could be infected with STIs through unprotected anal sexual intercourse. About three quarters of the participants reported correctly that they could be infected with STIs through semen (78.0%; 315) and through blood (46.0%; 186), vaginal fluids (32.7%; 132), contaminated sharp objects (17.3%; 70) and anal fluid (29.2%; 118).
- 305
- 306
- 307
- 308
- 309
- 310
- 311
- 312

	Proportion correct
Sexually transmitted infections	
Gonorrhoea	92.8
Syphilis	88.1
HIV	65.6
Genital warts	33.9
Herpes Simplex	22.8
Chancroid	20.5
Hepatitis B	17.3
Chlamydia	13.6
Trichomoniasis	5.2
Symptomatology of STIs	
Penis discharge	66.8
Burning pain during urination	60.4
Genital ulcer/sore	39.6
Anal ulcer/sores	25.0
Swelling in groin region	23.0
Anal pain	21.3
Anal discharge	21.0
Scrotal swelling	14.1
Transmission mechanism	
Unprotected sexual intercourse	97.5
Contaminated sharp	
objects/wounds	5.9
Blood transfusion	3.0
Unhygienic conditions	2.2
Don't know	1.0
Anal intercourse exposure to STI	83.7
Transmission Media	
Semen	78.0
Blood	46.0
Vaginal fluids	32.7
Anal fluids	29.2
Contaminated sharp objects	17.3
Clothing	7.2
Food	1.7
N/B- Supporting information sho	ws how the questioned were
asked	

### 314 Table 4: Proportion of participants answering STI knowledge items correctly

## 320 Participant characteristics independently associated with participant's knowledge about

321 STIs.

The multivariable modelling revealed that participants who were aged  $\geq 25$  years were more 322 likely to have a higher knowledge score compared with the participants who were aged 18-24 323 years (adjusted odds ratio aOR=0.973, 95% CI 0.616-1.538). Regarding education, 324 participants who had tertiary education were three times more likely to have a higher 325 knowledge score compared with the participants who had primary education (aOR=2.627, 326 95% CI 1.142-6.043). Regarding occupation participants who were employed had a higher 327 328 knowledge score compared to the ones who were not employed (aOR=0.922, 95% CI 0.401-2.117). Under level of income, participants earning Kshs 5000-10,000 were three times likely 329 to have a higher knowledge score compared to the ones who were not earning (aOR 2.332, 330 331 95% CI 0.990-6.263) . Participants who were earning Kshs >15000 (USD >150) were also three times more likely to have a higher knowledge score compared to the ones who were not 332 earning (aOR=2.520, 95% CI 0.900-7.055). Further bisexual men were more likely to have a 333 higher knowledge score compared with the participants who were having sex with men only ( 334 aOR= 1.550, 95% CI 1026-2.342) (Table 5). 335

336

337

338

339

- 340
- 341

342

#### 344 Table 5: Participant characteristics independently associated with participant's

#### **Unadjusted Odds Adjusted Odds** Ratio (95% CI Ratio (95% CI Variable **P-Value** 0.129 Age bracket Reference 18-24yrs >25 yrs 1.373 (0.922-2.047) 0.973 (0.616-1.538) Highest reported education level **Reference- "Up to completed Primary**" 0.001 "Up to completed Secondary 1.257 (0.582-2.711) 1.284 (0.575-2.865) "Up to completed Post-secondary 2.648 (1.195-5.872) 2.627 (1.142-6.043) **Occupation Reference (Not** employed/student 0.007 Employed 1.774 (1.176-2.677) 0.922 (0.401-2.117) Level of income Reference (No 0.005 income). < 5000 1.548 (0.809-2.962) 1.857 (0.725-4.759) 5000-10000 2.149 (1.225-3.771) 2.491 (0.990-6.263) 10001-15000 1.919 (1.059-3.480) 1.940 (0.746-5.042) >15000 2.949 (1.608-5.408) 2.520 (0.900-7.055) Sex with whom Reference (Men) 0.017 **Bisexual** 1.629 (1.097-2.419) 1.550 (1026-2.342)

#### 345 **knowledge about STIs.**

346

347

348

349

350

351

352

353

354

355

356

357

358

Page **16** of **25** 

#### 359 Discussion

360 STI knowledge and awareness is a key element towards decreasing the incidence of STIs 361 among populations at high risk of infection. However, findings from this study among MSM 362 in Nairobi, Kenya, demonstrated low STI knowledge.

Participants in this study who reported HIV as one of the STIs were less compared to the one 363 who reported Gonorrhoea and Syphilis. It's expected that all the participants should be aware 364 of it because HIV has no cure and it has caused devastating effects globally(18). This 365 disconnect among some of the participants could be because patients on antiretroviral drugs 366 look healthy. This has changed the perception of HIV/AIDS from a very fatal to a chronic 367 and potentially manageable disease. Also the availability and administration of antiretroviral 368 therapy has significantly reduced mortality and morbidity linked with HIV and AIDS(19). 369 This disconnect of HIV as one of the STIs among some of the participants is worthy worrying 370 371 about; because the progress that has been made to curb the spread of HIV might be futile. HCWs need to continue emphasizing to STI patients and other people in the society that HIV 372 373 is one of the STIs and it has no cure. Being infected by other STIs such as Herpes Simplex 374 Virus type 2 (HSV-2) and Syphilis infections puts them at a higher risk of being infected by HIV (3,4). 375

Others have found similar results; a study conducted among sixty racially –and ethnically 376 diverse MSM and transgender women had also reported that the participants had low 377 knowledge about STIs (20). Also another study conducted among 50 at risk MSM at Boston 378 USA had found that the MSM lacked knowledge about the signs and symptoms of STDs 379 (21). However, worth noting is that the measures of STI knowledge used on the above past 380 studies varied greatly from those used in our survey. There is need for the participants to be 381 health educated about STIs as some studies have revealed that patients who are enlightened 382 have higher probability of seeking health care services compared to the once who are not 383

384 (22). By them seeking treatment it can help stop the transmission of STI to others. To take
385 care of the asymptomatic STIs; quarterly STI screening should be encouraged among MSM
386 who are at risk of STIs.

In this study some of our participants reported of having oral sexual intercourse. Therefore, there is need for at least annual screening of extra-genital sites (rectum and pharynx) among sexually active MSM(23); so that, the extra-genital STIs cannot be missed out.

This study did not reveal statistical significant association between knowledge of the MSM and their age. Others have found similar results too; a study that was conducted among MSM in Ireland reported that participants who were aged 18-24 years of age had lower knowledge about STIs compared with the older MSM (24). Low knowledge about STIs among the youth could be occasioned by government policies and laws that criminalize key population behaviours and by education and health systems that pay no attention to or reject them(6).

Participants who had tertiary education were three times more likely to have a higher knowledge score compared with the participants who had primary education. Consistent with our findings is a study which was conducted in Melaka Malasyia; they found that participants who were studying in a degree program were two times more likely to be knowledgeable about STIs than the one who were not (25)

Further this study revealed that sexual behaviour of the study participants in terms of sex with whom, bisexual men had higher odds of knowledge about STIs compared to gay men. These findings are contrary to (26,27) who reported that bisexual men had limited availability of culturally sensitive education materials or health information that is specifically targeted to their needs. However, their settings are different from our Kenyan context.

Also majority of the participants had more than one sexual partner. Contrary to our findings,
a study which was conducted among MSM in UK revealed that men who were
knowledgeable about STI had more sexual partners. The authors of that study suggested that,

in addition to knowledge, behaviours were determined by a complex range of psychological 409 and eco-social factors (28). Still consistent with the UK study is a study which was conducted 410 among 772 subjects in Estonia. They found that higher knowledge scores were not associated 411 with lower HIV infection rates (29). This affirms that while knowledge is a key component of 412 behaviour change models, by itself it is often insufficient to effect behaviour change. Models 413 such as the Information-Behavioural Skills Model suggest that to take steps to reduce risk of 414 415 STI infection, individuals would need not only knowledge about STIs, but also motivation to prevent it and the skills necessary to implement risk reduction actions, such as getting 416 417 regularly tested for STIs (30).

#### 418 Limitations

Our study has several limitations. Our sample represents those men who seek sex at selected
geographic areas in Nairobi, and may not be representative of MSM, but otherwise do not
visit these areas. However, our aim was to enrol very high at risk individuals.

422 Our data are cross sectional, and as such we cannot infer causality between our independent423 variables and our dependent variable, knowledge of STIs.

#### 424 Implications to programmes

Individual health education and psychological approaches help an individual to acquire 425 knowledge and skills to enact new behaviours to reduce risk (31). In this study participants 426 had low knowledge about STIs and majority of them had more than one sexual partner. We 427 do recommend HCWs to continue giving health talks about STIs to MSM during their health 428 facility visits and in any other forums to enlighten them about STIs. Also because other 429 studies have reported that knowledge alone is often insufficient to effect behaviour change. 430 There is need for active follow-up of high risk individuals to prevent STIs which can be done 431 by; 432

Recalling of MSM for follow-ups, who report engaging in high risk sexual behaviours via
phone calls, peer leaders or community health works and creating demand for STI prevention
cascade. Also bulky short messages reminder for offering STI testing to high risk MSM can
be facilitated by using prompts incorporated within electronic MSM records.

On the other hand because there is a higher likelihood of extra-genital STIs being missed out
by both the patient and clinicians; there is need for at least annual screening of extra-genital
sites (rectum and pharynx) among sexually active MSM.

Also because asymptomatic STIs can be missed out due to syndromic management of STIs;
we recommend quarterly STI screening for individuals who are at risk so that the
asymptomatic STIs can be detected and treated to curb the spread of STIs.

#### 443 Implications to Policy Makers

444 Clear policies and guidelines should be put in place to enable HCWs to know what to do once 445 an individual has self-identified himself as MSM. For instance, MSM can be linked to 446 programmes or health facilities that can follow them up for possible regular screening and 447 treatment of STIs.

Self-testing for STIs allows convenience, privacy and gives the patient an opportunity to avoid embarrassment while still accessing STI care. So they can implement the use of selfsampling kits for various STI testing as another study reported that HIV self –testing was accepted by similar populations (32).

#### 452 Conclusion

453 Participant's knowledge level regarding STIs was low. Even though other studies have 454 reported that individuals to reduce risk of STI infection would need not only knowledge 455 about STIs. Still we recommend HCWs to continue educating patients about STIs. This is 456 because patients who are enlightened have a higher probability of seeking health care 457 services compared to the once who are not. Hence stop further spread of the STIs to their458 sexual partners.

459

#### 460 Supporting Information

461 **Supplemental questionnaire:** indicating the questions assessing participant's knowledge

462 levels.

463

#### 464 Acknowledgements

This work was partially funded by IAVI with the generous support of USAID and other donors; a full list of IAVI donors is available at <u>www.iavi.org</u>. The contents of this manuscript are the responsibility of IAVI and co-authors and do not necessarily reflect the views of USAID or the US Government. We would like to acknowledge the participants and the research assistants who assisted in data collection. We also appreciate Kenneth Ekoru from the imperial university and Janet Muasya from the University of Nairobi for their input in the data analysis.

472

#### 473 Author Contributions

- 474 Conceptualization: Delvin Kwamboka Nyasani
- 475 Data curation: Delvin Kwamboka Nyasani and Laura Lunani Lusike
- 476 Formal analysis: Justus.O.Osero and Laura Lunani Lusike
- 477 **Funding acquisition:** Delvin Kwamboka Nyasani
- 478 **Project Administration:** Justus.O.Osero and Meshack Onyambu

479 **Resources:** 

- 480 **Supervision:** Justus.O.Osero and Meshack Onyambu
- 481 Validation: Matthew A. Price and Gaundensia Nzembi Mutua

- 482 Writing original draft: Delvin Kwamboka Nyasani
- 483 Writing, review & editing: Delvin Kwamboka Nyasani, Justus.O.Osero, Meshack

- 484 Onyambu, Geoffrey Oino Ombati, Matthew A. Price and Gaundensia Nzembi Mutua.

486	Appendix 1: Knowledge assessment score of study participants
	Knowledge Score

Knowledge Score					
Mean=12.25, Median=12.00					
No. of right answers	No. of	Score percentage	Cumulative score		
0	participants		percentage		
3	1	.2	.2		
4	5	1.2	1.5		
5	5	1.2	2.7		
6	16	3.9	6.7		
7	24	5.9	12.6		
8	30	7.4	20.0		
9	36	8.9	28.8		
10	40	9.9	38.7		
11	44	10.8	49.5		
12	33	8.1	57.6		
13	32	7.9	65.5		
14	46	11.3	76.8		
15	15	3.7	80.5		
16	15	3.7	84.2		
17	12	3.0	87.2		
18	14	3.4	90.6		
19	8	2.0	92.6		
20	5	1.2	93.8		
21	3	.7	94.6		
22	5	1.2	95.8		
23	5	1.2	97.0		
24	4	1.0	98.0		
25	8	2.0	100.0		
Total	406	100.0			

#### 493 **References**

- World Health Organization (WHO) 2021. World Health Organization (WHO), Sexually
   Transmitted Infections 2021. 2021.
- 496 2. World Health Organization (WHO 2021). Guidelines for the Management of Sexually
  497 Transmitted Infections. 2021.
- 498 3. Centres for Disease Control and Prevention. Sexually transmitted disease surveillance 2010.
   499 Sexually Transmitted Disease Surveillance 2011. 2011;
- Kidd S, Torrone E, Su J, Weinstock H. Reported Primary and Secondary Syphilis Cases in the
   United States: Implications for HIV Infection. Sex Transm Dis. 2018 Sep;45(9):S42–7.
- Newman L, Rowley J, Vander Hoorn S, Wijesooriya NS, Unemo M, Low N, et al. Global Estimates
   of the Prevalence and Incidence of Four Curable Sexually Transmitted Infections in 2012 Based
   on Systematic Review and Global Reporting. PloS One. 2015;10(12):e0143304.
- 505 6. WHO 2016. Global Health Sector Strategy on Sexually Transmitted Infections 2016-2021
   506 Towards Ending STIs. 2016.
- Smith DK, Grohskopf LA, Black RJ, Auerbach JD, Veronese F, Struble KA, et al. Antiretroviral
   postexposure prophylaxis after sexual, injection-drug use, or other nonoccupational exposure to
   HIV in the United States: recommendations from the U.S. Department of Health and Human
   Services. MMWR Recomm Rep Morb Mortal Wkly Rep Recomm Rep. 2005 Jan 21;54(RR-2):1–20.
- Surveillance Report. 2018.
   Centres for Disease Control & Prevention. HIV
- Public Health England. Public Health England.Sexually transmitted infections and screening for
   Chlamydia i England. England; 2017.
- 515 10. Purcell DW, Johnson CH, Lansky A, Prejean J, Stein R, Denning P, et al. Estimating the population
  516 size of men who have sex with men in the United States to obtain HIV and syphilis rates. Open
  517 AIDS J. 2012;6:98–107.
- 518 11. Scott-Sheldon LAJ CP, Chan PA. Increasing Sexually Transmitted Infections in the U.S.:A Call for
   519 Action for Research, Clinical, and Public Health Practice. Arch Sex Behav. 2019;
- 12. Hladik W, Barker J, Ssenkusu JM, Opio A, Tappero JW, Hakim A, et al. HIV Infection among Men
  Who Have Sex with Men in Kampala, Uganda–A Respondent Driven Sampling Survey. PLoS ONE
  [Internet]. 2012 May 31;7(5). Available from:
- 523 https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3364961/
- Muraguri N, Tun W, Okal J, Broz D, Raymond HF, Kellogg T, et al. HIV and STI Prevalence and Risk
   Factors Among Male Sex Workers and Other Men Who Have Sex With Men in Nairobi, Kenya.
   JAIDS J Acquir Immune Defic Syndr. 2015 Jan 1;68(1):91.
- 527 14. Ministry of Health & National AIDS Control Council 2016. Kenya AIDS Response Progress Report
   528 2016. 2016.

- 529 15. Seidu AA, Ahinkorah BO, Dadzie LK, Tetteh JK, Agbaglo E, Okyere J, et al. A multi-country cross 530 sectional study of self-reported sexually transmitted infections among sexually active men in
   531 sub-Saharan Africa. BMC Public Health. 2020 Dec 7;20(1):1884.
- 16. National Aids and STI Control Programme (NASCOP). Key Population Mapping and Size
   Estimation in Selected Counties in Kenya: Phase 1 Key Findings, Kenya. Phase 1. Kenya; 2019.
- Mutisya EM, Mutua G, Nyasani D, Nduta H, Kabuti RW, Muturi-Kioi V, et al. Willingness to
  participate in future HIV vaccine trials among men who have sex with men and female sex
  workers living in Nairobi, Kenya. PloS One. 2020;15(8):e0238028.
- 18. UNAIDS 2020. Global AIDS Update: Seizing the Moment; July 2020. UNAIDS. AIDSinfo website;
   accessed July 2020, available at: http://aidsinfo.unaids.org/. UNAIDS. Core Epidemiology Slides;
   July 2020. 2020.
- 540 19. Oguntibeju OO. Quality of life of people living with HIV and AIDS and antiretroviral therapy.
  541 HIVAIDS Auckl NZ. 2012;4:117–24.
- 542 20. Balán IC, Lopez-Rios J, Dolezal C, Rael CT, Lentz C. Low STI Knowledge, Risk Perception, and
  543 Concern about Infection among Men Who Have Sex with Men and Transgender Women at High
  544 Risk of Infection. Sex Health. 2019 Nov;16(6):580–6.
- 545 21. Mimiaga MJ, Goldhammer H, Belanoff C, Tetu AM, Mayer KH. Men who have sex with men:
  546 perceptions about sexual risk, HIV and sexually transmitted disease testing, and provider
  547 communication. Sex Transm Dis. 2007 Feb;34(2):113–9.
- 548 22. O'Donnell O (2018). Access to health care in developing countries:breaking down demand-side
  549 barriers.Utilization of Health and Medical Services:Factors influencinghealth care seeking
  550 behaviour and unmet needs in rural areas of Kenya. Theses DR.Masters.2007. [Kenya]; 2018.
- 551 23. Kimberly A. Workowski, Gail A. Bolan,. Sexually Transmitted Diseases Treatment Guidelines,
  552 2015 [Internet]. 2015 [cited 2022 Oct 13]. Available from:
  553 https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6403a1.htm
- 24. Carey C, O'Donnell K, Davoren M, Quinlan M, Igoe D, Barrett P. Factors associated with lower
   knowledge of HIV and STI transmission, testing and treatment among MSM in Ireland: findings
   from the MSM Internet Survey Ireland (MISI) 2015. Sex Transm Infect. 2021 Aug 1;97(5):351.
- 557 25. Mansor N, Ahmad N, Rahman HA. Determinants of knowledge on sexually transmitted infections
  558 among students in public higher education institutions in Melaka state, Malaysia. PLOS ONE.
  559 2020 Oct 29;15(10):e0240842.
- 26. Daneback K, Ross MW, Månsson SA. Bisexuality and Sexually Related Activities on the Internet. J
   Bisexuality. 2008 Sep 1;8(1–2):115–29.
- Feinstein BA, Dodge B. Meeting the sexual health needs of bisexual men in the age of
   biomedical HIV prevention: Gaps and priorities. Arch Sex Behav. 2020 Jan;49(1):217–32.
- 28. Wayal S, Reid D, Weatherburn P, Blomquist P, Fabiane S, Hughes G, et al. Association between
  knowledge, risk behaviours, and testing for sexually transmitted infections among men who
  have sex with men: findings from a large online survey in the United Kingdom. HIV Med. 2019
  Sep;20(8):523–33.

- 29. Parker RD, Rüütel K. Associations of high-risk behaviour and HIV status with HIV knowledge
  among persons in Tallinn, Estonia. Scand J Public Health. 2010 Nov 1;38(7):748–55.
- 30. Balán IC, Lopez-Rios J, Dolezal C, Rael CT, Lentz C. Low sexually transmissible infection
  knowledge, risk perception and concern about infection among men who have sex with men
  and transgender women at high risk of infection. Sex Health. 2019 Nov;16(6):580–6.
- 573 31. Holtgrave DR, Qualls NL, Curran JW, Valdiserri RO, Guinan ME, Parra WC. An overview of the
  574 effectiveness and efficiency of HIV prevention programs. Public Health Rep Wash DC 1974. 1995
  575 Apr;110(2):134–46.
- 576 32. Figueroa C, Johnson C, Verster A, Baggaley R. Attitudes and Acceptability on HIV Self-testing
  577 Among Key Populations: A Literature Review. AIDS Behav. 2015 Nov;19(11):1949–65.