- 1 *Effects of joint mobilization combined with acupuncture on pain,*
- 2 physical function, and depression in stroke patients with chronic
- 3 neuropathic pain: a randomized controlled trial
- 4
- 5 Ji-Eun Lee¹, Takayuki Akimoto², Ho-Seong Lee^{1,3*}
- ⁶ ¹Department of Exercise and Medical Science, Graduate School, Dankook University,
- 7 Cheonan, Republic of Korea
- ⁸ ²Laboratory of Muscle Biology, Faculty of Sport Sciences, Waseda University,
- 9 Tokorozawa, Japan
- ¹⁰ ³Institute of Medical-Sports, Dankook University, Cheonan, Republic of Korea

11

- 12 * Corresponding author
- 13 E-mail: hoseh28@dankook.ac.kr (HSL)

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

15

16 Abstract

17	Objective: To investigate the effectiveness of joint mobilization (JM) combined
18	with acupuncture (AC) for treatment pain, physical function and depression in
19	poststroke patients. Methods: A total of 69 poststroke patients were randomly
20	assigned to the JM+AC group (n = 23), the JM group (n = 23), and the control group
21	(n = 23). Patients in the JM+AC group and the JM group received JM for 30 minutes,
22	twice a week for 12 weeks, and the JM+AC group received AC for 30 minutes
23	separately once a week. The control group did not receive JM or AC. Pain (visual
24	analog scale, shoulder pain and disability index, Western Ontario and McMaster
25	universities osteoarthritis index), physical function (range of motion, 10-m walking
26	speed test, functional gait assessment, manual function test, activities of daily living
27	scale, instrumental activities of daily living scale), and depression (center for
28	epidemiologic studies depression scale, Beck depression inventory) were assessed for
29	each patient before and after the 12 weeks of intervention. Results: Pain and physical
30	function were improved significantly in the JM+AC group compared with the JM and
31	control groups. Physical function and depression were improved significantly in the
32	JM+AC and JM groups compared with the control group. Conclusion: The treatment
33	of JM combined with AC improved pain, depression, and physical function of
34	poststroke patients with chronic neuropathic pain in this study. This valuable finding

provides empirical evidence for the designing therapeutic interventions and
 identifying potential therapeutic targets.

37 **Keywords:** acupuncture, joint mobilization, neuropathic pain, poststroke

38

39 Introduction

Stroke is one of global healthcare problems that is common, serious, and 40 disabling. Motor recovery from stroke is frequently poor or insufficient, with only 41 one-third of poststroke patients regaining dexterity within the first 6 months [1]. Less 42 than 45% of stroke patients are likely to achieve complete functional recovery, and 43 the majority of patients has some residual impairment and an inability to perform 44 activities of daily living (ADL) [2-4]. In addition to motor control and sensory 45 46 deficits, stroke survivors also develop complications such as pain, spasticity, joint constraint, skin or vascular damage, and chronic neuropathic pain caused by nerve 47 damage, which are the main challenges in poststroke management [5,6]. 48

Shoulder pain experienced by more than 50% of poststroke patients, is a common sequelae [7], interfering with rehabilitation and reducing arm function and ADL [8]. Knee pain is the second most common pain in patients after stroke, and it has been reported to cause motor dysfunction and abnormal gait [9]. The recovery of gait function is a primary goal of rehabilitation, with the majority of poststroke

patients suffering from a reduction in gait velocity and independent gait is crucial in 54 carrying out ADL following stroke [10]. Additionally, sensorimotor dysfunction 55 causes limited joint range of motion (ROM) and muscle weakness in the lower 56 extremities of the affected side, leading to difficulties in performing functional 57 activities, such as sit to stand and gait [11,12]. Therefore, it is necessary to reduce 58 pain and increase the joint ROM to improve physical function in poststroke patients. 59 To tackle these problems and to help restore and improve the function of upper and 60 lower limbs, a wide range of treatments has been proposed to therapists, so far. 61

62 Mulligan first proposed mobilization-with-movement treatment as a joint mobilization (JM) technique [13]. JM is an orthopedic manipulation therapy used for 63 its mechanical effects to relieve pain, improve mobility, and treat contractures via the 64 third stage of the translatoric movement and the convex-concave rule [14]. According 65 to the convex-concave rule, gliding on a convex joint surface is performed in the 66 opposite direction to the bone's movement; gliding on a concave joint surface is 67 performed in the same direction as the bone's movement. When a concave joint 68 surface is distracted, the joint is pulled along the long axis of the bone; alternatively, 69 when a convex joint surface is distracted, the joint surface must be separated [14]. JM 70 has the mechanical effect of disrupting the contracture by direct movement of the joint 71 area, and the arthrokinetic reflex effect of suppressing or facilitating the related 72 muscles is achieved by stimulating the joint receptors [15]. However, there appears to 73

be a limit to how JM alone can reduce pain and restore physical function in poststrokepatients.

Acupuncture (AC) is commonly used to treat/manage chronic diseases in Korea 76 [16]. It is recognized to be effective in regulating physiological balance by controlling 77 muscles and acupuncture points or by stimulating the affected area [17]. And AC is 78 effective in reducing pain by stimulating the nerves in stroke patients and help to 79 restore motor function and improve the ADL score [18]. Studies suggest that AC is 80 also effective in improving pain and depression in poststroke patients [19]. 81 Particularly, AC combined with other therapies is more effective in poststroke 82 patients than AC alone [20]. Therefore, it is possible that AC combined with JM is 83 more effective to improve pain, depression, and physical function in poststroke 84 patients. Although there are many studies about effectiveness of JM for poststroke 85 patients, a few studies have examined the effects of JM combined with AC for these 86 patients. 87

In this work, we examined the changes in pain, physical function, and depression before and after the 12 weeks of intervention of JM combined with AC in poststroke patients. We hypothesized that combining JM with AC would improve physical function more compared with JM alone.

92

93 Methods

94 **1. Study design**

95	The study had a random sampling design. Participants were randomized
96	electronically using block sizes of 3 across groups. All investigators and data analysts
97	were blinded until the study and analysis were completed.

Written informed consent was obtained from all subjects and the signed informed consent documents were kept on file. The protocol was approved by the Dankook University ethics committee (IRB number: 2021-05-046) and it adhered to the tenets of the Declaration of Helsinki and its later amendments or comparable ethical standards. The individual in this manuscript has given written informed consent (as outlined in PLOS consent form) to publish these case details.

This clinical trial was registered with the Korean Clinical Research Information
 Service approved by the WHO. Registration number: KCT0007924. Date registered:
 22/11/2022, retrospectively registered. No dropouts or adverse events were reported.

107 2. Participants

This study included 69 adults who were diagnosed with stroke and were enrolled in a public health center and welfare center located in D city, South Korea. This group was further divided into the JM+AC, the JM, and the control groups (n = 23 each).

111	The inclusion criteria were: diagnosis of stroke ≥ 6 months ago; pain in the shoulders
112	and knees for more than 6 months; and a Korean Mini Mental State Examination
113	score of \geq 24. The exclusion criteria were: a risk of tumors or infections; a previous
114	history of orthopedic disease or ankle surgery; a previous history of orthopedic
115	disease or shoulder and knee joint surgery; and a length of ≥ 6.3 mm in the line
116	bisection test; taking medications for diseases other than stroke, such as antiplatelet
117	drugs and anticoagulants. The demographics and clinical characteristics of the study
118	patients were summarized at inclusion (Table 1). No significant differences in the
119	pretraining status of the control and experimental groups were noted.

Variables	JM+AC (n=23)	JM (n=23)	CON (n=23)	р
Age (years)	73.78 ± 6.52	74.13 ± 5.63	76.74 ± 6.62	0.23
Sex (male/female)	12 / 11	17 / 6	13 / 10	0.29
Affected side (right/left)	13 / 10	10 / 13	13 / 10	0.60
Onset time (month)	52.57 ± 24.39	61.35 ± 21.00	61.09 ± 22.20	0.33
MMSE-K (score)	26.17 ± 1.34	26.35 ± 1.44	25.87 ± 1.40	0.51

120 **Table 1. Participant characteristics.**

Values are means \pm standard deviation.

MMSE-K: Korean mini mental state examination.

121 **3. Interventions**

122 **3.1. Joint mobilization**

123 For 12 weeks, the JM+AC and JM groups received grades II and III JM for 30 min, twice weekly, to relieve pain in the shoulder and knee joints and to improve the 124 ROM [14]. Shoulder JM was performed on the glenohumeral joint and 125 scapulothoracic joint in the supine and prone positions. Lateral distraction, downward 126 glide, posterior glide, and anterior glide were performed on the glenohumeral joint. 127 Elevation, depression, protraction, retraction, upward rotation, and downward rotation 128 were performed on the scapulothoracic joint. Knee JM was performed on the 129 tibiofemoral and patellofemoral joints in the supine and prone positions. Lower 130 distraction, posterior glide, and anterior glide were performed on the tibiofemoral 131 joint. Downward, lateral, and medial glides were performed on the patellofemoral 132 joint. 133

134 **3.2 Acupuncture**

The JM+AC group received acupuncture and moxibustion treatments around the shoulder and the knee joints for 30 min, once a week for 12 weeks. Acupuncture treatment was performed using disposable sterile stainless-steel needles (0.20×30 mm, Dongbang, Korea) for 15 min on average (figure 1). The penetration depth varied according to the site and was 20 mm deep on average. For moxibustion treatment, mini moxibustion (Taegeuk, Korea) was performed for 15 min (fig 1).

141

Fig 1. Acupuncture and moxibustion treatment materials.

142 Disposable sterile needles (left picture). Disposable mini moxa rolls (right picture).

143 **4. Outcome Measures**

144 **4.1 Pain**

145	Pain severity was measured using a visual analog scale (VAS), the shoulder pain
146	and disability index (SPADI), and the Korean Western Ontario McMaster Universities
147	osteoarthritis index (KWOMAC) before and after 12 weeks of the experimental
148	period. The VAS used a 100-mm line on which a pain-free state was 0 and severe pain
149	was 100, and the reliability is 0.99 [21]. SPADI is an evaluation tool for assessing the
150	degree of shoulder pain and disability that comprises 13 evaluation items scored on a
151	scale of 0-10 [22]. A score above 100 (out of 130) indicates more shoulder pain and
152	disability. The reliability is 0.94 [23]. KWOMAC, a Korean version of WOMAC
153	[24], is an evaluation tool for assessing the degree of knee pain and function [25]. It
154	consists of 24 evaluation items scored on a scale of 0-4 points and a total score of 96
155	points. Higher scores indicate a worse knee condition. The reliability is 0.96 [25].

156 **4.2 Physical function**

The range of motion (ROM), 10-meter walk test (10-MWT), functional gait assessment (FGA), manual function test (MFT) for measuring physical function, ADL, and instrumental ADL (IADL) were evaluated before and after 12 weeks of the experimental period.

A goniometer (Saehan Corporation, Korea) was used to measure the ROM of the shoulder and knee joint. Active ROM for shoulder flexion (SF), shoulder extension (SE), shoulder abduction (SAB), shoulder adduction (SAD), and knee flexion (KF) were measured. The measurement was performed three times, and the average value was used. The reliability is 0.89 [26].

The 10-MWT requires the participant to walk independently over the middle 10 m portion on a 14-m path; the time is measured in seconds and the test is repeated three times. The reliability is 0.99 [27].

FGA was developed by Wrisley et al. [28] to evaluate gait and comprises 10 items scored on a scale of 0–3 points and a total score of 30 points. Specific items include walking on level surfaces, changing walking speed, turning the head to the side while walking, moving the head up and down while walking, walking with one foot as an axis, walking around obstacles, gait with a narrow base of support, walking with eyes closed, walking backward, and climbing stairs. The reliability is 0.93 [28].

MFT is a test tool developed to assess the overall condition of arm function in a stroke patient. It consists of eight items, and if subitem inspection is possible, it is measured as 1 point. The reliability is 0.95 [29].

ADL and IADL are tools developed by Won et al. [30] ADL comprising seven
items covering dressing, washing the face, bathing, eating, moving, using the toilet,

and controlling the bladder, with a total of 21 points. IADL comprises 30 items scored
on a scale of 1–3. Ten items cover grooming, chores, preparing meals, washing
clothes, going out a short distance, using transport, buying goods, managing money,
using a phone, and taking medicine. Higher scores indicate lower dependence in
evaluating the independence level in ADL. The reliability is 0.93 [30].

185 **4.3 Depression**

Depression was measured using the center for epidemiologic studies depression scale (CES-D) and the Beck depression inventory (BDI) before and after 12 weeks of the experimental period.

CES-D is a self-reported depression evaluation tool [31], comprising 20 evaluation items scored on a scale of 0–3 points. Scores higher than 21 (out of 60) indicate a higher frequency of depressive experiences. The reliability is 0.90 [31].

BDI is a tool for evaluating emotional, cognitive, motivational, and physiological depression, and the Korean version was used [32]. It comprises 21 evaluation items scored on a scale of 0–3 points and a total score of 63 points. The reliability is 0.84 [32].

196 **5. Statistical Analysis**

The data were analyzed statistically using SPSS 23.0 software (SPSS, Chicago,
IL, USA). General characteristics prior to the interventions were analyzed using 11

199	descriptive statistics. Two-way analysis of variance (2-way ANOVA) with repeated
200	measures was performed to verify the difference between groups and time periods.
201	Multiple comparisons were performed according to the post hoc test (Tukey HSD)
202	when there was a significant difference between groups and time periods. A paired t
203	test was used to compare variations in pain, physical function, and depression before
204	and after experimental period for each group. Statistical significance was reached
205	when $p < 0.05$.

206

207 **Results**

208 **1. Pain**

The changes in pain severity are shown in Table 2. The VAS was not statistically significant between groups, but SPADI decreased significantly in the JM+AC group compared to the JM group and the control group (p < 0.05), and KWOMAC decreased significantly in the JM+AC group compared to the control group (p < 0.05). The VAS, SPADI, and KWOMAC were decreased significantly in the JM+AC and JM groups after 12 weeks of intervention (p < 0.05).

Table 2. Pain before and after joint mobilization and joint mobilization 215

combined with acupuncture. 216

Variables	JM	+AC (n=	23)	J	M (n=23)	С	ON (n=2	3)	F	ES
Period	pre	post	t	pre	post	t	pre	post	t		
	75.65 ±	56.13 ±	5.18*	63.48±	56.30±	2.32*	62.83	67.17 ±	-1.50	0.97	0.28
VAS (mm)	18.55	5.80		3.45	4.35		±	18.52			
							14.99				
CDADI	68.90 ±	63.71 ±	6.11*	58.16 ±	56.79±	3.32*	58.56	58.86±	-1.31	4.78*	0.13
SPADI	0.12	8.40†‡		11.92	11.98		±	10.27			
(score)							10.32				
	73.40 ±	51.88 ±	5.93*	68.65 ±	59.57±	3.07*	68.61	71.43 ±	-1.70	3.32*	0.09
KWOMAC	5.22	12.46†		11.95	9.80		±	12.87			
(50010)							11.40				

Values are means \pm standard deviation.

VAS: visual analog scale, SPADI: shoulder pain and disability index, KWOMAC: Korean Western Ontario and McMaster universities osteoarthritis index.

*p < 0.05 vs. pre-exercise. p < 0.05 vs. CON. p < 0.05 vs. JM.

2. Physical function 217

The changes in physical functions are shown in Table 3. 218

Table 3. Physical function before and after joint mobilization and joint 219

mobilization combined with acupuncture. 220

Vari	ables	JM	+AC (n=	=23)	J	M (n=23	6)	С	ON (n=23	3)	F	ES
Per	riod	pre	post	t	pre	post	t	pre	post	t		
		119.91	125.26	-4.74*	128.17	129.17	-1.17	128.6	124.48	4.18	1.12	0.03
	SF (°)	± 14.00	± 13.26		± 14.46	± 13.21		5 ±	± 3.30			
								3.02				
	SE (0)	33.70 ±	38.96 ±	-5.79*	37.52 ±	37.83 ±	-0.36	34.48	32.39 ±	2.36	2.03	0.06
	SE (°)	8.13	6.64		8.63	1.40		± 1.76	1.40			
	GAD	109.13	114.70	-6.26*	117.44	118.52	-1.38	112.2	110.22	2.26	2.56	0.07
ROM	SAB	± 11.93	± 10.17		± 2.46	± 2.02		1 ±	± 2.64			
	(*)							2.41				
	SAD	40.83 ±	47.65 ±	-7.13*	39.91 ±	42.39 ±	-1.89	41.96	$40.00 \pm$	2.20	2.28	0.07
	(°)	6.34	6.77		1.64	1.47		± 1.10	0.89			
		117.31	127.17	-5.49*	119.91	125.65	-4.37*	112.8	112.39	0.44	7.27*	0.18
	KF	± 11.68	± 8.71†		± 2.02	± 1.70†		3 ±	± 2.65			
	(°)							2.61				
	10-M	25.83 ±	14.81 ±	7.44*	23.23 ±	18.13 ±	4.97*	24.55	24.04 ±	0.98	1.54	0.45
GAIT	WT	11.49	8.45		6.23	6.49		± 9.58	9.92			

(s)

medRxiv preprint doi: https://doi.org/10.1101/2023.02.06.23285552; this version posted February 8, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license . FGA $18.78 \pm 23.13 \pm -5.90^*$ $19.04 \pm 22.00 \pm -4.22^*$ 20.04 $19.74 \pm$ 0.79 0.75 0.22 (score 3.28 2.70 3.04 2.70 ± 3.83 4.17) **MFT** $20.78 \pm 22.91 \pm -6.16^*$ $20.43 \pm 21.52 \pm -4.64^*$ 20.00 $19.52 \pm$ 1.85 11.65 0.26 * (score 1.57 1.41 ± 1.71 1.68 1.73†‡ 1.62†) **ADL** $17.57 \pm 20.22 \pm -5.42^*$ $17.13 \pm 19.30 \pm -7.61^*$ 16.65 $16.57 \pm$ 0.33 0.30 13.93 * **ADL (score** 0.52 1.88^{+} 0.39 0.19† ± 0.36 0.40) IADL $24.35 \pm 26.96 \pm -1.13^* \ 23.13 \pm 23.57 \pm -0.93$ 22.96 $22.70 \pm$ 1.03 4.77* 0.13 (score 0.90 0.81 0.72 ± 0.80 0.70 0.39†‡)

Values are means \pm standard deviation.

ROM: range of motion, SF: shoulder flexion, SE: shoulder extension, SAB: shoulder abduction, SAD: shoulder adduction, KF: knee flexion, KE: knee extension, 10-MWT: 10-meter walking speed test, FGA: functional gait assessment, MFT: manual function test, ADL: activities of daily living scale, IADL: instrumental activities of daily living scale.

*p < 0.05 vs. pre-exercise. p < 0.05 vs. CON. p < 0.05 vs. JM.

221 **2.1 Range of motion**

222	SF, SE, SAB, and SAD were not statistically significant between groups, but KF
223	increased significantly in the JM+AC and JM groups compared to the control group
224	(p < 0.05). Within the JM+AC group, SF, SE, SAB, and SAD increased significantly
225	(p < 0.05), whereas KF increased significantly in the JM+AC and JM groups after 12
226	weeks of intervention ($p < 0.05$).

227 **2.2 Gait**

The 10-MWT and FGA were not statistically significant between groups, but 10-MWT and FGA improved significantly in the JM+AC and JM groups after 12 weeks of intervention (p < 0.05).

231

2.3 Activities of daily living

MFT increased significantly in the JM+AC group compared to the JM group and control groups (p < 0.05), as well as in the JM group compared to the control group (p < 0.05). ADL increased significantly in the JM+AC and JM groups compared to the control group (p < 0.05), and IADL increased significantly in the JM+AC group compared to the JM group and the control group (p < 0.05). Within the JM+AC and JM groups, MFT and ADL increased significantly (p < 0.05), whereas IADL increased significantly in the JM+AC group after 12 weeks of intervention (p < 0.05).

239 **3. Depression**

240	The changes in depression are shown in Table 4. CES-D and BDI decreased
241	significantly in the JM+AC and JM groups compared to the control group (p < 0.05),
242	and CES-D and BDI decreased significantly in the JM+AC group after 12 weeks of
243	intervention ($p < 0.05$).

244 Table 4. Depression before and after joint mobilization and joint mobilization

245 combined with acupuncture.

Variables	JM	+AC (n=	23)	J	M (n=23))	C	ON (n=2	3)	F	ES
Period	pre	post	t	pre	post	t	pre	post	t		
CES-D	26.83 ±	20.30 ±	5.24*	20.48 ±	20.43 ±	0.06	27.91	33.00 ±	-3.60	7.11*	0.18
(score)	8.63	8.87†		9.57	8.88		± 8.96	8.77			
	32.48 ±	22.35 ±	5.49*	30.61 ±	29.96 ±	1.19	33.70	39.43 ±	-3.33	5.97*	0.15
BDI (score)	11.04	7.97†		12.23	10.99		± 9.49	9.15			

Values are means \pm standard deviation.

CES-D: center for epidemiologic studies depression scale, BDI: Beck depression inventory.

*p < 0.05 vs. pre-exercise. †p < 0.05 vs. CON. ‡p < 0.05 vs. JM.

247 **Discussion**

The main finding of this study was that JM combined with AC treatment improved the subjective pain, depression, and physical function of poststroke patients more effectively than JM alone.

Shoulder and knee pain impedes rehabilitation and interferes with daily activities 251 in poststroke patients [7–9]. Kaltenborn et al. [14] indicated that JM was effective in 252 relieves shoulder and knee pain through direct movement of the joints. In AC 253 treatments, acupuncture treatment is reported to increase pain threshold by stimulating 254 the central nervous system, and moxibustion treatment has an analgesic effect through 255 thermal stimulation [18,33]. Additionally, previous studies reported that AC treatment 256 257 reduced SPADI and KOMAC [34,35]. Taken together, it was confirmed that JM combined with AC treatment had a more positive effect on pain in poststroke patients 258 than JM alone. 259

Kluding and Santos [36] stated that the JM technique improved movement without changing in joint ROM. Alternatively, Choi et al. [37] reported that AC treatment reduces pain and increases joint flexibility by regenerating the surrounding soft tissue. We found that there was a significant difference in ROM before and after the intervention in the JM+ AC group, but not in the JM group in this study. These results may imply that it is difficult to improve joint ROM with JM alone.

As mentioned in the introduction, gait is an important factor for independent 266 living and social participation. JM showed a positive effect on gait by improving 267 active ROM which is the ROM that can be achieved by patient himself/herself [38], 268 and AC treatment stimulated muscles and nerves to improve gait [39]. Additionally, 269 previous studies stated that acupuncture had a beneficial impact on the energy 270 production of joints during walking [40]. In this study, gait was significantly 271 improved after the intervention in the JM+ AC and JM groups, but there was no 272 difference between groups. Generally, gait necessitates movement of hip, knee, and 273 ankle joints. The lower extremity interventions in this study were limited to the knee 274 joint, which may not have been sufficient to influence group effect comparisons. In 275 the future, it would be prudent to intervene in all joints of lower extremity. 276

277 We employed several measurements to access physical function, which is important for daily living. MFT is an assessment of overall arm function [29], SPADI 278 is for shoulder pain and disability [22]. IADL is for evaluation of dressing, toilet 279 using, and movement [30]. It is thought that the reduction of SPADI restored arm 280 function in poststroke patients and affected MFT and IADL. Previous research has 281 shown that acupuncture has little effect on improving the lifestyle of stroke patients 282 [41], but it is believed that it had a positive effect on IADL in this study because joint 283 mobilization was combined with acupuncture. Arm function and daily life activities of 284 the patients in the JM+ AC group might be improved through the increase in muscle 285 activity induced by JM and the recovery of neurological damage by AC treatment. 286

Depression is associated with pain in poststroke patients [7], and depression in 287 these patients are associated with mortality, suicide risk, and quality of life [42–44]. 288 Thus, depression must be verified in rehabilitation process for functional recovery 289 after stroke, in general. The CES-D and BDI scores in the JM+ AC group were 290 significantly decreased after 12 weeks of the intervention, but not in the JM group. 291 This is most likely due to AC treatment to alleviate pain and improve ADL. A 292 previous study showed that JM improved depression by reducing stress factors 293 through improvement of unstable posture [45]. Furthermore, it has been reported that 294 acupuncture treatment was effective in suppressing changes in the autonomic nervous 295 system caused by mental stress [46]. It has also been reported that moxibustion 296 treatment relaxed and stabilized the body through its thermal action [18]. Thus, it is 297 considered that acupuncture and moxibustion treatment of AC reduced depression 298 through autonomic nerve inhibition and thermal stimulation. 299

300 Limitation

There are several limitations in our study. The major part of subjects in this study was older farmers living in rural areas, and it is not uncertain whether the results of this study can be applied to other elderly population living in urban areas. Additionally, it might be better if we assessed the effect of the treatment more frequently because the results of this study were obtained before and after the 12 weeks of intervention period.

307

308 Conclusion

309	Improvements in pain, depression, and physical function of poststroke patients in
310	the JM+ AC group were significantly greater than those in the JM and the control
311	groups, suggesting JM combined with AC was more effective in improving not only
312	pain but also the ability to physical function. JM combined with AC may also be
313	helpful for poststroke patients who have depression and decreased physical function
314	due to chronic neuropathic pain. Future studies should reveal the physiological
315	assessment of muscle strength and the effect of JM combined with AC on muscle tone
316	based on changes in physical function in stroke patients with chronic neuropathic
317	pain.

318

319 Author Contributions

320 Conceived and designed the experiments: JEL. Performed the experiments: JEL.

321 Analyzed the data: JEL, TA. Wrote the paper: JEL, TA. Revised paper: TA, HSL.

323 **Conflicts of Interest**

324 The authors declare that there are no conflicts of interest regarding the publication of

325 this paper.

326

327 Acknowledgments

328 We appreciate Kim Yeon-oh and Choi Jong-bin of the Dangjin City Public Health

329 Center for their assistance with this experiment.

330

331 Data Availability Statement

332 Access to data was prepared using the following data set and is available at the free,

333 public repository Figshare (https://doi.org/10.6084/m9.figshare.21789092).

334

335 Funding

336 This research did not receive any specific grant from funding agencies in the public,

337 commercial, or not-for-profit sectors.

339 **References**

340	1.	Dobkin BH. Rehabilitation after stroke. N Engl J Med. 2005;352(16):1677–84.
341	2.	Flynn RW V, MacWalter RSM, Doney ASF. The cost of cerebral ischaemia.
342		Neuropharmacology. 2008;55(3):250-6.
343	3.	Kelly-Hayes M, Beiser A, Kase CS, Scaramucci A, D'Agostino RB, Wolf PA.
344		The influence of gender and age on disability following ischemic stroke: the
345		Framingham study. J Stroke Cerebrovasc Dis. 2003;12(3):119–26.
346	4.	Marini C, Baldassarre M, Russo T, De Santis F, Sacco S, Ciancarelli I, et al.
347		Burden of first-ever ischemic stroke in the oldest old: evidence from a
348		population-based study. Neurology. 2004;62(1):77-81.
349	5.	De Ryck A, Brouns R, Fransen E, Geurden M, Van Gestel G, Wilssens I, et al. A
350		prospective study on the prevalence and risk factors of poststroke depression.
351		Cerebrovasc Dis Extra. 2013 Jan;3(1):1–13.
352	6.	Vér C, Hofgárt G, Menyhárt L, Kardos L, Csiba L. Ankle-foot continuous
353		passive motion device for mobilization of acute stroke patients. Open J Ther
354		Rehabil. 2015;3(02):23.
355	7.	Klit H, Finnerup NB, Overvad K, Andersen G, Jensen TS. Pain Following
356		Stroke: A Population-Based Follow-Up Study. PLoS One. 2011 Nov
357		15;6(11):e27607.

358	8.	Rand D. Proprioception deficits in chronic stroke—Upper extremity function
359		and daily living. PLoS One. 2018 Mar 30;13(3):e0195043.
360	9.	Caglar NS, Akin T, Aytekin E, Komut EA, Ustabasioglu F, Okur S, et al. Pain
361		syndromes in hemiplegic patients and their effects on rehabilitation results. J
362		Phys Ther Sci. 2016 Mar;28(3):731–7.
363	10.	Yoo C, Yong M hyun, Chung J, Yang Y. Effect of computerized cognitive
364		rehabilitation program on cognitive function and activities of living in stroke
365		patients. J Phys Ther Sci. 2015;27(8):2487–9.
366	11.	Fujita T, Iwata M, Fukuda M, Ikeda M. Relationship between lower extremity
367		muscle mass, leg extension strength and muscle power of hemiplegic stroke
368		patients. J Phys Ther Sci. 2011;23(2):277-82.
369	12.	Lin PY, Yang YR, Cheng SJ, Wang RY. The relation between ankle
370		impairments and gait velocity and symmetry in people with stroke. Arch Phys
371		Med Rehabil. 2006;87(4):562–8.
372	13.	Mulligan BR. Mobilisations with movement (MWMS) for the hip joint to restore
373		internal rotation and flexion. J Man Manip Ther. 1996;4(1):35-6.
374	14.	Kaltenborn FM, Evjenth O. Manual Mobilization of the Joints: The Kaltenborn
375		Method of Joint Examination and Treatment. Vol 1. spine 5th ed Oslo Norli.
376		2009;

377	15.	Ersoy U, Kocak UZ, Unuvar E, Unver B. The Acute Effect of Talocrural Joint
378		Mobilization on Dorsiflexor Muscle Strength in Healthy Individuals: A
379		Randomized Controlled Single-Blind Study. J Sport Rehabil. 2019;28(6):601–5.
380	16.	Kim JH, Kismali G, Gupta SC. Natural Products for the Prevention and
381		Treatment of Chronic Inflammatory Diseases: Integrating Traditional Medicine
382		into Modern Chronic Diseases Care. Evidence-Based Complement Altern Med.
383		2018;2018:9837863.
384	17.	Yang A, Wu HM, Tang J, Xu L, Yang M, Liu GJ. Acupuncture for stroke
385		rehabilitation. Cochrane Database Syst Rev. 2016;(8).
386	18.	Zuo CY, Lv P, Zhang CS, Lei RX, Zhou W, Wu QF, et al. Ipsi- and Contralateral
387		Moxibustion Generate Similar Analgesic Effect on Inflammatory Pain. Olajide
388		OA, editor. Evidence-Based Complement Altern Med. 2019;2019:1807287.
389	19.	Lee JA, Park SW, Hwang PW, Lim SM, Kook S, Choi KI, et al. Acupuncture for
390		shoulder pain after stroke: a systematic review. J Altern Complement Med. 2012
391		Sep;18(9):818–23.
392	20.	Heo KH, Hwang EH, Cho HW, Lee I, Hong JW, Shin YI, et al. An analysis of
393		the effectiveness of stroke between east-west integrative medicine hospital and
394		western medicine hospital by the data of brain rehabilitation registry. J Korean
395		Med Rehabil. 2013;23(3):117–24.

396	21.	Wagner DR, Tatsugawa K, Parker D, Young TA. Reliability and utility of a
397		visual analog scale for the assessment of acute mountain sickness. High Alt Med
398		Biol. 2007;8(1):27–31.
399	22.	Roach KE, Budiman-Mak E, Songsiridej N, Lertratanakul Y. Development of a
400		shoulder pain and disability index. Arthritis Rheum Off J Am Coll Rheumatol.
401		1991;4(4):143–9.
402	23.	Angst F, Goldhahn J, Pap G, Mannion AF, Roach KE, Siebertz D, et al.
403		Cross-cultural adaptation, reliability and validity of the German Shoulder Pain
404		and Disability Index (SPADI). Rheumatology. 2007;46(1):87-92.
405	24.	Bellamy N. WOMAC Osteoarthritis Index: User Guide IV. WOMAC.
406		Queensland, Aust. 2000;
407	25.	Bae SC, Lee HS, Yun HR, Kim TH, Yoo DH, Kim SY. Cross-cultural adaptation
408		and validation of Korean Western Ontario and McMaster Universities
409		(WOMAC) and Lequesne osteoarthritis indices for clinical research. Osteoarthr
410		Cartil. 2001;9(8):746–50.
411	26.	Riddle, D. L., Rothstein, J. M., Lamb RL. Goniometric reliability in a clinical
412		setting: shoulder measurements. Phys Ther. 67(5):668-73.
413	27.	Steffen TM, Hacker TA, Mollinger L. Age- and Gender-Related Test
414		Performance in Community-Dwelling Elderly People: Six-Minute Walk Test,

415		Berg Balance Scale, Timed Up & amp; Go Test, and Gait Speeds. Phys Ther.
416		2002 Feb 1;82(2):128–37.
417	28.	Wrisley DM, Marchetti GF, Kuharsky DK, Whitney SL. Reliability, internal
418		consistency, and validity of data obtained with the functional gait assessment.
419		Phys Ther. 2004;84(10):906–18.
420	29.	Miyamoto S, Kondo T, Suzukamo Y, Michimata A, Izumi SI. Reliability and
421		validity of the Manual Function Test in patients with stroke. Am J Phys Med
422		Rehabil. 2009;88(3):247–55.
423	30.	Chang-Won WON, Keum-Yeol Y, Yong-Gyun RHO, Soo-Young KIM,
424		Kyung-Hwan CHO, Ho-Cheol S, et al. The development of Korean activities of
425		daily living (K-ADL) and Korean instrumental activities of daily living
426		(K-IADL) scale. J Korean Geriatr Soc. 2002;107–20.
427	31.	Radloff LS. The CES-D scale: A self-report depression scale for research in the
428		general population. Appl Psychol Meas. 1977;1(3):385-401.
429	32.	Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for
430		measuring depression. Arch Gen Psychiatry. 1961;4(6):561-71.
431	33.	Lee JD, Chon JS, Jeong HK, Kim HJ, Yun M, Kim DY, et al. The
432		cerebrovascular response to traditional acupuncture after stroke. Neuroradiology.
433		2003;45(11):780–4.

434	34.	Dyson-Hudson TA	Shiflett SC.	Kirshblum SC.	Bowen JE.	Druin EL.
		_ / 2 0 0 0 0 0				

- 435 Acupuncture and Trager psychophysical integration in the treatment of
- 436 wheelchair user's shoulder pain in individuals with spinal cord injury. Arch Phys
- 437 Med Rehabil. 2001;82(8):1038–46.
- 438 35. Lo MY, Wu CH, Luh JJ, Wang TG, Fu LC, Lin JG, et al. The effect of
- 439 electroacupuncture merged with rehabilitation for frozen shoulder syndrome: A
- single-blind randomized sham-acupuncture controlled study. J Formos Med
- 441 Assoc. 2020;119(1, Part 1):81–8.
- 442 36. Kluding PM, Santos M. Effects of ankle joint mobilizations in adults poststroke:
 443 a pilot study. Arch Phys Med Rehabil. 2008 Mar;89(3):449–56.
- 444 37. Choi G cheol, Lee J won, Bae JE, Kim D jin, Hong J su, Kyung D hyun. The
- 445 Clinical Effect of Complex Korean Medical Admission Treatment in Patients
- 446 with Fractures of Distal Radius by Traffic Accident: 2 Cases Series Report. J
- 447 Korean Med Rehabil. 2021;31(1):187–93.
- 38. Ravichandran H, Janakiraman B. The Effects of Ankle Joint Mobilization on
 Dorsiflexion Range and Gait Parameters in Chronic Stroke Survivors: A
- 450 Systematic Review and Meta-analysis. J Stroke Med. 2021;4(1):15–24.
- 451 39. Kim MS, Moon BS. Effect of comprehensive Korean medicine with
- 452 rehabilitation in stroke patients: a retrospective study. J Physiol Pathol Korean
- 453 Med. 2016;30(5):355–9.

454	40.	Hauer K, Wendt I, Schwenk M, Rohr C, Oster P, Greten J. Stimulation of
455		acupoint ST-34 acutely improves gait performance in geriatric patients during
456		rehabilitation: a randomized controlled trial. Arch Phys Med Rehabil.
457		2011;92(1):7–14.
458	41.	Sibbritt D, Peng W, Lauche R, Ferguson C, Frawley J, Adams J. Efficacy of
459		acupuncture for lifestyle risk factors for stroke: A systematic review. PLoS One.
460		2018;13(10):e0206288.
461	42.	Kellermann M, Berecz R, Bereczki D. Does the severity of depressive symptoms
462		after stroke affect long-term survival? An 18-year follow-up. PLoS One.
463		2018;13(12):e0209157.
464	43.	Wilkins SS, Akhtar N, Salam A, Bourke P, Joseph S, Santos M, et al. Acute post
465		stroke depression at a Primary Stroke Center in the Middle East. PLoS One.
466		2018;13(12):e0208708.
467	44.	Kang SW, Lee SH, Lee YS, Kwon S, Bosch P, Lee YJ, et al. Association
468		between depression and quality of life in stroke patients: The Korea National
469		Health and Nutrition Examination Survey (KNHANES) IV-VII (2008-2018).
470		PLoS One. 2022;17(6):e0269010.
471	45.	Park D, Lee E, Lee K. Effects of Thoracic Flexibility Exercise Program on Pain,
472		Heart Rate Variability, and Depression in Patients with Chronic Low Back Pain.
473		J Korean Soc Integr Med. 2019;7(4):161–70.

- 474 46. Cho SY, Ha SY, Jang JY, Nam SS, Kim YS. Original Article: Effect of
- 475 Hwangryunhaedoktang Pharmacopuncture at CV17 (Jeonjung) for Mental
- 476 Stress on Short-term Analysis of Heart Rate Variability. J Korean Acupunct
- 477 Moxibustion Soc. 2009;26(5):50.

479

480 Supporting Information

- 481 S1 Fig. Acupuncture and moxibustion treatment materials. Disposable sterile
- 482 needles (left picture). Disposable mini moxa rolls (right picture).
- 483 S1 Table. Participant characteristics. MMSE-K: Korean mini mental state
 484 examination.
- 485 S2 Table. Pain before and after joint mobilization and joint mobilization
- 486 combined with acupuncture. VAS: visual analog scale, SPADI: shoulder pain and
- 487 disability index, KWOMAC: Korean Western Ontario and McMaster universities
- 488 osteoarthritis index.

489 S3 Table. Physical function before and after joint mobilization and joint

- 490 mobilization combined with acupuncture. ROM: range of motion, SF: shoulder
- 491 flexion, SE: shoulder extension, SAB: shoulder abduction, SAD: shoulder adduction,
- 492 KF: knee flexion, KE: knee extension, 10-MWT: 10-meter walking speed test, FGA:
- 493 functional gait assessment, MFT: manual function test, ADL: activities of daily living
- 494 scale, IADL: instrumental activities of daily living scale.

495 S4 Table. Depression before and after joint mobilization and joint mobilization

496 **combined with acupuncture.** CES-D: center for epidemiologic studies depression

497 scale, BDI: Beck depression inventory.

- 499 S1 Checklist. CONSORT Checklist. Checklist of information to include when
- 500 reporting a randomized trial.
- 501 S2 Checklist. CONSORT Flow Diagram. Flow diagram.



Figure