

1 **Dying to make a care policy: Community perception of causes of deaths using standardized**
2 **verbal autopsy method: Saudi type 2 diabetes mellitus register data**

3 Faleh Alyazidi ^{1,2*} , Deler Shakely ¹ , Max Petzold ¹ , Fawaz Alyazidi ³ , Laith Hussain-Alkhateeb ¹

4 ¹ School of Public Health and Community Medicine, Sahlgrenska Academy, University of Gothenburg,
5 Gothenburg, Sweden

6 ² Department of Public Health, College of Health Sciences at Al-Leith, Umm Al-Qura University, Al-Leith,
7 Kingdom of Saudi Arabia

8 ³ Infectious Diseases Control Department, Executive Directorate of Preventive Medicine, Makkah
9 Healthcare Cluster, Makkah, Kingdom of Saudi Arabia

10 * Corresponding author

11 Faleh.saud.alyazidi@gu.se

12

13

14

15

16

17

18

19

20 **Abstract:**

21 Diabetes mellitus (DM) is a serious global health issue which significantly impacts public health
22 and socioeconomic development. Exploring how the community perceives the causes of deaths
23 and their associated risk factors is crucial for public health. This study combines verbal autopsy
24 (VA) with the Type 2 Diabetes Mellitus (T2DM) register to explore community perceptions of
25 causes of death and associated influential factors in the western region of Saudi Arabia. 302 VA
26 interviews were conducted with relatives or caregivers of deceased who died between 2017
27 and 2021 based on T2DM medical register from Alnoor Specialist Hospital in Makkah city, Saudi
28 Arabia. Cause-specific mortality fractions (CSMFs) obtained from the VA using the InterVA-5
29 model were utilized to assess community perception. We used a multivariable logistic
30 regression model to determine factors influencing community perceptions of causes of death.
31 Lin's CCC with 95% CI was used to analyze the concordance for the CSMFs from verbal autopsy
32 causes of death (VACoD) as a presumed reference standard and family-reported causes of
33 death (FRCoD). The outcomes of this study demonstrate a generally broad spectrum of
34 community perceived mortalities, with some critical misconceptions based on the type of death
35 and the deceased's background and characteristics, with an overall CCC of 0.60 (95% CI: 0.20-
36 1.00; $p=0.03$). The study findings demonstrate that community perception is weak if the
37 deceased was male compared to female (aOR: 0.52; 95% CI: 0.26 –1.03) and if the deceased
38 was ≥ 80 years compared to 34–59 years (aOR: 0.48; 95% CI: 0.16-1.38), but it significantly
39 improves among married compared to single (aOR: 2.13; 95% CI: 1.02 –4.42). The Saudi
40 community perception of causes of death with reported type 2 diabetes was relatively plausible
41 but varied substantially based on the type of death, sex, age ≥ 80 years, and other vital events

42 like marital status. Higher or lower community perception is attributed to how people may
43 perceive risk factors associated with the causes of death, which can guide public health
44 planning and interventional programs. The study findings further emphasize the need to
45 employ robust and standardized VA methods within the routine medical services for a
46 systemized assessment of families' reported causes of death.

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62 **Introduction**

63 Diabetes mellitus (DM) remains a major global public health issue with a significant impact on
64 public health and socio-economic development. Diabetes is one of the top ten causes of death
65 worldwide. Together with cardiovascular diseases, respiratory diseases and cancer, these
66 conditions account for over 80% of all premature non-communicable diseases (NCDs) deaths
67 [1]. The International Diabetes Federation estimates that 536.6 million people were living with
68 diabetes in 2021, and this number is projected to increase by 46%, reaching 783.2 million in
69 2045 [2], with approximately 50% of all diabetic individuals are unaware of their conditions [3].
70 Several reports have attributed the prevalence of DM in the Middle East and North Africa
71 regions to the increased per capita income, urbanization, and pronounced lifestyle changes that
72 enabled physical inactivity and increased the obesity rate.

73
74 In the context of chronic illnesses and treatment management, community perception of health
75 and illness – which is the ability to perceive the world through sensory organs [4] – is critical for
76 public health administrative practices. A comprehensive and integrated approach to
77 understanding how communities perceive health and disease is crucial for developing relevant
78 and effective health intervention strategies and can empower individuals to seek and handle
79 health knowledge properly [5-7]. Individuals are more likely to take action to prevent illnesses if
80 they acknowledge their susceptibility to a particular form of illness or if they are likely to
81 experience serious consequences [8]. Lay concepts of health and illness provide essential
82 information to biomedical models [9, 10] and can significantly impact communities' health and
83 illness behaviors, which requires systematic means of assessment at population level [6, 10].

84 Saudi Arabia is associated with this global epidemic of NCDs [11, 12]. According to the World
85 Health Organization (WHO), Saudi Arabia has the second highest rate of diabetes in the Middle
86 East (7th highest in the world), with an estimated population of 7 million diabetics and more
87 than 3 million pre-diabetics [13], which is a serious public health issue. Previous Saudi reports
88 have shown that the success of national response strategies is heavily dependent on the
89 public's perceptions and attitudes toward the risk of an epidemic and the importance of
90 preventive measures [14]. Addressing the health literacy problems among a culturally diverged
91 Saudi population will be crucial by virtue of the undergoing health system re-engineering
92 towards the Saudi 2030 Vision. The Vision 2030 Health Sector Transformation Program was
93 initiated in 2021 with a five-year goal of reforming the health sector to become a
94 comprehensive, effective, and integrated health system that prioritizes the health of individual
95 and society, including citizens, residents, and visitors [15]. A growing body of evidence have
96 additionally demonstrated that inadequate health literacy is significant among the Saudi
97 population [16-18], whereby negative perceptions of diabetes risk factors can impede disease
98 management and prevention. Patients with poor health literacy skills are prevalent among the
99 Saudi population. A published cross-sectional survey conducted in Saudi Arabia to describe the
100 distribution of low health literacy in the Saudi population found that 46% of respondents were
101 classified as having low health literacy; this was associated with older age groups [17]. This
102 group demands actively engaged healthcare professionals to effectively simplify the care
103 management, which will only add more burden to the already overstretched health system
104 caused by COVID-19 and other emerging diseases.

105

106 Global statistics show that about half of annual deaths pass without formal medical certification
107 leaving crucial causes of death information outside any form of useful application for policy
108 making and public health program monitoring [19-23]. While this issue is broadly associated
109 with low- and middle-income countries (LMICs) due to insufficient investment in civil
110 registration and vital statistics (CRVS) systems [23], higher income countries are also prone to
111 misclassification of crucial international classification of disease (ICD) information [24]. Saudi
112 Arabia is among countries facing demographic transition, which can influence the doctor-
113 patient communications and hence the ultimate uptake of health information and their
114 classifications.

115
116 Verbal autopsy (VA) – standardized interviews conducted by trained fieldworkers or healthcare
117 staff with final caregivers or close relatives of the deceased on the medical signs, symptoms,
118 and circumstances surrounding deaths – has emerged in more recent time as a widely used
119 method for determining causes of death in settings where adequate reporting and reliable
120 classification of causes of death are distracted [25, 26]. While this set of VA information is
121 classically processed by physicians to determine a likely medical cause of death, computerized
122 tools such as *InterVA-5* have been validated in several settings, which demonstrate significant
123 advantages over the VA physician review process, mainly being cheap and consistent over time
124 and place [6, 27, 28]. VA is commonly conducted within registered and unregistered population
125 and can consistently elicit information through structured and narrative sections using the
126 standardized WHO VA questionnaires. At the end of the VA interview, the interviewer would
127 typically finish off with asking respondents about their own interpretation of the causes of

128 death of their deceased, such information that can form the basis of understanding community
129 perception of causes of death. Social autopsy (SA), a less structured method for investigating
130 death, includes questions on changeable social, cultural, and healthcare system aspects, which
131 shows great potential for routine public health applications [23, 29]. SA has often been used in
132 connection with VA in registered populations. Nevertheless, it tends to be time and cost
133 consuming if used as a sole tool, and unlike VA, it will not be amenable for large-scale
134 applications or automated interpretations, which led to its innovation as VA-SA integrated
135 process [29, 30].

136
137 Within the Saudi research and health policy interests, where VA is rarely practiced, no studies
138 have yet sought standardized tools such as the VA to explore the community perception of
139 causes of death, which can have significant implications in guiding public health planning and
140 novel interventional programs. This study will employ VA to explore the community perceptions
141 of causes of death based on T2DM register in the Western region of Saudi Arabia. To success
142 this study aim, VA causes of death (VACoD) as a presumed reference standard will be compared
143 to the family-reported causes of death (FRCoD), and the determinants of the community
144 perception will be further assessed across different causes of death groups and population
145 backgrounds and characteristics.

146

147 **Methodology**

148 **Study population and context**

149 The study conducted 302 Verbal autopsy interviews with relatives or caregivers of the deceased
150 who died between 2017 and 2021, according to T2DM medical register from the Alnoor
151 Specialist Hospital – a public hospital in Makkah city, Saudi Arabia, which is among three other
152 hospitals that constitute to major healthcare provider to a population size of 2.5 million. All
153 patients included in this study fulfilled the study criteria; (1) death occurred between 2017 and
154 2021 (small history interval to reduce recall bias during the VA process), (2) at least one clinical
155 diagnosis (ICD) of T2DM, and (3) informed consent to perform the verbal autopsy interview
156 given by the respondent.

157
158 Makkah is a large province and of a relatively dynamic population compared to other Saudi
159 provinces, mainly due to their close and constant interaction with people who come from all
160 over the world to perform religion pilgrimage (Hajj and Umrah). This frequent (short and long-
161 term) stay of a large population-scale creates a special demographic mix where migration of
162 culture, economy, diseases, and demography is highly prevalent in this region. This diverged
163 community has the potential to influence the community perceptions and beliefs about a
164 certain phenomenon, such as how they perceive illness or causes of death.

165 **Verbal autopsy interpretation**

166 For VA interpretation, several software programs are used for automated interpretation such as
167 '*Smart VA*', '*InSilicoVA*', and '*InterVA*' [31-33], with the *InterVA* being used more broadly [19,

168 33]. The 2016 WHO VA instrument has been updated to contain all the input variables needed
169 by all three software, making their use more consistent and standardized. [28, 34, 35]. In recent
170 years, a series of InterVA models have been created to interpret VA data using Bayesian
171 probabilistic modeling [36].
172
173 VA uses a Bayesian probability model to determine the most probable cause of death and their
174 corresponding likelihoods for each VA case [5, 26, 37]. The Bayesian concept operation permits
175 the combination of medical experts' views as prior and relevant available data, which can
176 collectively augment the model outputs [6, 38]. This methodological process used in the
177 InterVA model is explained in full elsewhere [36]. Using Bayes theorem, up to three probable
178 causes of death (likelihood) are typically assigned for each cause by the automated InterVA
179 tool, given the specific signs and symptoms reported [26, 33]. Each assigned cause has
180 corresponding likelihoods, and the sum likelihoods of all assigned causes has a maximum value
181 of 1.00 (or 100%). In cases where the likelihood proportions did not add up to 1.0 for a
182 particular case, any remaining margin of likelihood not accounted for by the likelihood of the
183 first, second, and third causes can therefore be included as a partial indeterminate component
184 in analyzing overall cause of death and CSMFs patterns. It has been suggested that this is a
185 more practical approach than aggregating the sum of small residual probabilities of unlikely
186 causes, which might lead to misleading conclusion in a large number of cases [36]. The Cause-
187 Specific Mortality Fraction (CSMF) is an output from the algorithm. CSMF corresponds to the
188 proportion that each cause of death category contributed to the total number of deaths [5, 6,
189 38].

190 **Data collection management**

191 Information on the deceased's background and characteristics were taken directly from the
192 medical register at Alnoor Specialist hospital. Death records contained data for capturing vital
193 events [deaths in and out-hospital, date of death, sex, age, ICD-10 code of causes of death, and
194 contact details of the next of kin]. Considering the heterogeneous and dynamic population in
195 this Province, other characteristics of the deceased such as nationality, marital status, and
196 education, were also considered. A total of pre-estimated sample of 302 relatives or close
197 caregivers of the deceased were randomly chosen from the medical register and consented to
198 conduct the VA interview. In addition, interviews were administered by the first author
199 together with a trained research assistant. The standard face-to-face VA interview process has
200 been challenged by the COVID-19 pandemic, which coincided with the strict physical distancing
201 preventive measures implemented in Saudi Arabia. Therefore, telephone interviews were
202 sought as a substitute for the standard face-to-face method. The interviews were conducted in
203 Arabic without audio-recording, but data were reviewed and safely stored at the university
204 electronic databases after removing all personal identifiers, and all participants were
205 anonymized. Closed-ended questions were collected using the Open Data Kit (ODK) on Android
206 smartphones. VA data were reported into a Microsoft Access database, and the participants'
207 study IDs were used to link the VA data with the background and characteristic information.

208 **Data management**

209 The deceased's age was categorized into four age groups (34-59, 60-69, 70-79, and ≥ 80 years
210 old), and the levels of education were grouped as *illiterate, basic education including (primary,*

211 *secondary school, and high school), and advanced education including university level. The*
212 *relationship between the VA informants and the deceased was classified into two groups (first-*
213 *degree relative and second-degree relative), whereby the deceased marital status was*
214 *dichotomized as single and married. The place of death was recorded and processed as binary*
215 *information (home, hospital/health facility). The causes of death were further categorized into*
216 *broader categories, described in S1 Table.*

217 **Data analysis**

218 Numerical and graphical descriptive statistics were generally sought for causes of death data in
219 overall and per groups. Each individual VA was processed using InterVA-5 model, which assigns
220 up to three likely causes of death per WHO VA cause categories [39], while FRCoDs were given
221 as a single or two causes of death. Crude and adjusted logistic regression for the binomial
222 agreement (agreement or not) between VACoD and FRCoD was analyzed against the
223 informant's background, characteristics, and socioeconomic factors for each cause of death
224 category. All potential confounders from the background and characteristic factors were
225 assessed in a forward-wise likelihood ratio test. Statistical analyses were performed using
226 STATA release 17.0 software using a significant level of 5% as a cut-off for the hypothesis test.

227
228 Generated causes of death likelihoods were utilized to calculate population-level CSMFs for
229 each cause of death group between VACoD and FRCoD, and their proportional absolute
230 differences and 95% confidence intervals were calculated. A concordance correlation
231 coefficient (CCC) for the CSMFs from VACoD and FRCoD was also calculated using Lin's CCC with
232 95% CI [40]. Based on Pearson's correlation, the CCC compares the viewpoints between the two

233 measures – just like other correlation tests, CCC has a range of -1 to 1, with 1 representing
234 perfect agreement [41]. The CCC combines precision and accuracy measures to determine how
235 far the observed data vary from the perfect concordance line (that is, the line at 45 degrees on
236 a square scatter plot). Lin's coefficient increases in value as a function of the data's reduced
237 major axis's closeness to the perfect concordance line (data accuracy) and the data's tightness
238 about its reduced major axis (the precision of the data) [40]. The CCC can essentially estimate
239 the agreement between VACOD and FRCOD from a population causes of death level.

240 **Ethical approval**

241 This study received the ethical approval to perform the VA interviews from the Institutional
242 Review Board (IRB) at Makkah Health Affairs, Saudi Arabia (IRB Number: H-02-K-076-0321-478).
243 The study was aligned with the Declaration of Helsinki. Informed consent was fully respected at
244 both individual and household levels, with the right to refuse or withdraw from interviews.
245 Additionally, data was only accessed by the principal researcher for the purposes of the study
246 to ensure the confidentiality and anonymity of the participants.

247 **Results**

248 A total of 302 deceased relatives or caregivers were contacted and consented to conduct the
249 VA interview, with three out of the respondents being female. Of these interviews, which were
250 completed successfully over 4 months, InterVA-5 assigned either single, two, or three likely
251 causes of death; a single cause of death to 299 individuals (99%), two likely causes to 26
252 individuals (8.60%), and three likely causes to 2 cases (0.66%). Three cases (0.99%) were

253 explicitly designated ‘indeterminate’ by InterVA-5, generally reflecting a lack of particular VA
254 data or contradictory evidence.

255
256 For the VA interviews where FRCoD was reported, 255 families (84.43%) gave a single cause of
257 death, and 124 families (41.05%) reported two causes. There were 36 families (11.92%) where
258 the given likely causes of death were considered ‘indeterminate’ since they did not match with
259 the International Classification of Diseases (ICD-10) codes.

260
261 The results from a multivariable logistic regression model for exploring factors of determining
262 the community perception are presented in Table 1. Based on the study findings, the odds ratio
263 of the agreement decreases if the deceased was male compared to female (aOR: 0.52; 95% CI:
264 0.26 –1.03) and it significantly increases if married compared to single (aOR: 2.13; 95% CI: 1.02
265 –4.42). Improved perception of causes of death was associated with deaths of people with
266 advanced education compared to illiterate (aOR:1.21; 95% CI: 0.45 – 3.20), and the death
267 occurring at hospital compared to deaths at home (aOR: 1.25; 95% CI: 0.56 – 2.80), despite
268 being not statistically significant. In addition, the odds of agreement tend to decrease if the
269 interview was conducted with a respondent of a second-degree relationship to the deceased
270 compared to a first-degree relationship (aOR: 0.91; 95% CI: 0.37 –2.20). Our results found that
271 the odd ratio of the agreement for age groups (60-69, 70–79, and >=80 years) increases if the
272 deceased aged 60-69 and 70-79 years compared to deaths aged 34-59 years, and the likelihood
273 tends to decrease if the deceased was >=80 years compared to the reference (aOR: 1.34; 95%
274 CI: 0.60 – 2.98, aOR:1.04; 95% CI: 0.45 – 2.41 and, aOR:0.48; 95% CI: 0.16 – 1.38, respectively).

275 **Table 1. Background and characteristics of 302 deaths in Makkah region, Saudi Arabia, and**
 276 **showing multivariable OR for agreement between FRCoD and VACoD.**

Background characteristics	Number of deaths <i>n</i> (%)	Crude OR (95% CI)	Adjusted OR (95% CI)
Age group			
34–59	68 (22.52)	Ref	Ref. ^a
60-69	88 (29.14)	1.37 (0.61 - 3.04)	1.34 (0.60-2.98)
70–79	84 (27.81)	1.09 (0.48 – 2.51)	1.04 (0.45-2.41)
>=80	62 (20.53)	0.50 (0.17 - 2.42)	0.48 (0.16-1.38)
Sex			
Female	126 (41.72)	Ref	Ref. ^b
Male	176 (58.28)	0.72 (0.40 - 1.31)	0.52 (0.26-1.03)
Respondent			
First degree relationship	260 (86.09)	Ref	Ref. ^c
Second degree relationship	42 (13.91)	0.90 (0.37 - 2.16)	0.91(0.37-2.20)
Education			
Illiterate	121 (40.07)	Ref	Ref. ^d
Basic education	137 (45.36)	1.09 (0. 51 - 2.32)	1.09 (0.52- 2.30)
Advance education	44 (14.57)	1.24 (0. 46 - 3.32)	1.21 (0.45- 3.20)
Year of death			
2018	28 (9.27)	Ref	Ref. ^c
2019	69 (22.85)	0.48 (0.14 - 1.54)	0.49 (0.15-1.62)

2020	161 (53.31)	0.90 (0.34 - 2.42)	0.90 (0.33-2.47)
2021	44 (14.57)	0.81 (0.24 – 2.66)	0.89 (0.26-2.99)
Marital status			
Single	116 (38.41)	Ref	Ref. ^c
Married	186 (61.59)	1.78 (0.93 – 3.41)	2.13 (1.02-4.42)*
Place of death			
Home	60 (19.87)	Ref	Ref. ^c
Hospital	242 (80.13)	1.29 (0.59 - 2.82)	1.25 (0.56-2.80)

277 **Abbreviation(s):** FRCoD: Family reported causes of death

278 VACoD: Verbal autopsy causes of death

279 a, adjusted for sex

280 b, adjusted for education, marital status and age groups

281 c, adjusted for age groups and sex

282 d, adjusted for age groups, sex and marital status

283

284 To evaluate the community perceptions of causes of death, CSMFs obtained from the VA using

285 the InterVA-5 model were utilized to assess FRCoD, along with the CSMF absolute differences

286 and their 95% confidence intervals. Table 2 shows that infectious diseases, stroke, and

287 circulatory-related deaths were found to be underestimated as probable causes of death

288 reported by families in relation to the VA reference standard. In contrast, CSMF of

289 indeterminate and renal-related deaths groups, were overestimated as probable causes of

290 death. In addition, accidental, NCD and diabetes related deaths groups were slightly

291 overestimated with a minimal absolute CSMF difference. Despite no ICD codes being assigned

292 by InterVA-5 for covid-19 (had no defined ICD code at the time) and mental illness-related

293 deaths categories, the CSMF attributed to covid-19 by families was as high as 13.18% of all
294 deaths.

295 **Table 2. Cause specific mortality fractions per disease categories derived from VACoD and**
296 **FRCoD for 302 deaths, Makkah region, Saudi Arabia**

Categories	VACoD	FRCoD	Absolute Difference 95%CI
Infectious	10.2	2.0	-8.2 (-12.0,-4.4)*
Cancer	5.2	4.4	-0.8 (-4.2,2.6)
Diabetes	14.3	15.8	1.5 (-4.2,7.2)
Circulatory	35.7	19.7	-16.0 (-23.0,-9.0)*
NCD	4.5	4.9	0.4 (-3.0,3.8)
Renal	2.8	9.2	6.4 (2.6,10.2)*
Accidental	2.5	1.0	-1.5 (-3.6,0.6)
Indeterminate	7.7	16.8	9.1 (3.9,14.3)*
Stroke	16.6	9.4	-7.2 (-12.5,-1.9)*
Covid-19	0.0	13.1	13.2 (9.4,17.0)*
Mental	0.0	3.1	3.1 (1.1,5.1)*

297 **Abbreviation(s):** FRCoD: Family reported causes of death

298 VACoD: Verbal autopsy causes of death

299 *Significant Absolute Difference

300

301

302 Based on findings from the CCC, there was an overall agreement between the two approaches
303 (VACoD and FRCoD) in terms of cause-specific mortality. The Concordance of FRCoD concerning
304 VACoD and fractions is shown in (Fig 1), with a CCC of 0.60 (95% CI: 0.20-1.00; p=003).

305

306


307

308

309

310

311



312

313 **Fig 1. Concordance correlation coefficient (log-logscale) between CSMF determined by VA and**
314 **respondent reports, in relation to the line of equivalence, for 302 deaths in Makkah city,**
315 **Saudi Arabia.**

316

317 Discussion

318 The year 2030 is expected to be an exceptional milestone for Saudi Arabia, with a
319 comprehensive agenda involving the assessment of essential national and global goals. The 17
320 Sustainable Development Goals (SDGs) including goal target 3.7 (*ensuring universal access to*
321 *health-care services including for family planning, information, and education as well as*
322 *vaccination*), and goal target 3.8 (*achieving Universal Health Coverage*), as well as the Saudi
323 National 2030 Vision goals will both be assessed and disseminated. Creating partnerships with

324 the private sector, academia, and civil society with an emphasis on strengthening civil
325 registration and vital statistic systems (CRVS) would be crucial elements for succeeding in these
326 national and global goals. However, gaps in registering vital data persist (a main task attributed
327 to the CRVS), and despite increasing calls for employing standardized VA concepts to
328 strengthen public health applications, including the CRVS, the VA approach is entirely novel in
329 Saudi Arabia and many other neighboring countries where gaps in causes of death information
330 exist. Exploring the perception of public health aspects, essentially how causes of death are
331 perceived within communities in Saudi Arabia is crucial while the country's health system is
332 now undergoing a significant re-engineering process towards national and global agenda.

333
334 This study findings show a generally broad spectrum of community perceived mortalities with
335 some crucial misconception based on the type of death and the background and characteristics
336 of the deceased. During the data collection period, the world has experienced a devastating
337 COVID-19 pandemic, which has ultimately re-profiled deaths and influenced the public and
338 medical opinions of causes of death in Saudi Arabia and elsewhere. While the concept of
339 community perception assessment is not new, understanding how local communities perceive
340 cause of deaths, and consequently recognizing their associated risk factors is now timely and
341 more crucial in view of the health and health system transitions due to COVID-19. Proposing
342 such a standardized assessment would also be of interest to the Saudi health system to
343 continuously monitor the health literacy of the people particularly within such dynamic Saudi
344 communities. Based on published reports, low health literacy – which is how people are able to
345 access, understand and utilize information and facilities to guide health-related decisions for

346 themselves and other members in their communities – contributes to a wide range of health
347 problems and is associated with poor health outcomes and ineffective chronic diseases
348 management and treatments, such as the case of diabetes [16, 42, 43].

349
350 The study utilized VA to explore the community perceptions, which is mostly amenable for
351 estimating CSMFs at population level and can adequately direct intended funds and
352 interventional public health programs [44-46]. This study findings showed relatively minimal
353 community misconception based on the derived CSMFs across diabetes, NCD, and accidental
354 causes of death categories, implying that communities were more aware of and able to
355 plausibly recognize these diseases as ultimate causes of death. Diabetes as a chronic illness
356 demands frequent presentation at the healthcare services for routine follow up and
357 treatments. High level of exposure to valid health information via professional medical
358 practitioners can positively contribute to the community health literacy level, which may
359 explain the higher level of perception for this cause of death category and likely other diseases
360 from the NCD category. It can be said that communities are influenced by what they remember
361 of signs and circumstances of death events whereas other factors such as exposure to local
362 beliefs or to professional health practices can influence the subject's memory [47]. For a
363 disease such as cancer for instance, mental representations of death can be exceptionally
364 influential and highlight how communities may perceive some inevitable illnesses [48]. This
365 study shows trivial CSMF differences due to cancer, where communities are arguably
366 attributing cancer illness to death in a plausible manner [48, 49]. Features of illness
367 representations involve views about a disease's etiology, time course, and curability [48].

368 For example, if individuals link cancer with death, they may avoid seeking health care services.

369 In fact, there is considerable evidence that a fatalistic view of cancer hinders adherence to

370 screening guidelines [50] and impeded engagement in cancer protective behaviours such as

371 physical activity, healthy diet and avoid smoking [51, 52]. Hence, addressing fatalistic beliefs

372 through communication about certain diseases like cancer can play an important role in cancer

373 prevention measures. Stroke as a cause of death, was however underestimated by

374 communities, a pattern found to be consistent with previous Saudi and non-Saudi studies [53-

375 55]. Difficulties in recalling or recognizing disease symptoms, rather than a mere lack of

376 knowledge, can possibly explain the observed pattern by communities in relation to stroke, but

377 may also reflect a need for a more effective stroke promotional campaign.

378 Public awareness of diseases that are associated with diabetes mellitus, like renal (kidney) and

379 circulatory diseases are important for public health interventional plans. Our results show that

380 respondents tend to overestimate renal diseases as causes of death, and this pattern is

381 consistent with prior cross-sectional research from Saudi Arabia [56, 57], which revealed a

382 dearth of knowledge about renal diseases among the Saudi population. While the prevalence of

383 renal diseases is relatively high in Saudi Arabia due to climatic, behavioral and genetic factors

384 [58], the Saudi health system has a comprehensive treatment and follow up management, both

385 in terms of dialysis services and kidney transplantation. This interaction between high disease

386 prevalence and adequate healthcare and the fact that most of the study population are

387 presumably diabetic (which is medically associated with renal diseases), seems to influence the

388 community's perception of causes of death related to renal diseases. Saudi communities have

389 underestimated circulatory diseases' cause of death as it was also observed in other published

390 reports revealing low perception of cardiovascular causes of death [59-61]. This
391 underestimation could be attributed to the disease complexity and general lack of information
392 and awareness. Although this may seem plausible from a medical point of view, failing to
393 recognize symptoms of common illnesses can have some public health implications since most
394 deaths have taken place at hospital settings where relatives or caregivers are presumably
395 informed about the illness of their deceased. This was also observed in other cross-sectional
396 studies linking low health literacy skills with unrealistic risk underestimation [62, 63]. Another
397 plausible explanation is that circulatory diseases are frequently asymptomatic with risk factors
398 slowly progressing over time leading to delayed recognition and perception. In line with a
399 previous study [64], which demonstrated a low awareness of major infectious diseases, the
400 present study revealed a low level of awareness in relation to infectious diseases, which is what
401 would be expected in such setting due to stigmatization, and discrimination. This pattern
402 supports the findings of previous studies in Saudi that reported a statistically significant positive
403 correlation between some infectious diseases and stigma [65-67]. Stigma and discrimination in
404 communities present a risk when reporting causes of death based on region, sex, and ethnicity,
405 especially deaths related to stigmatized diseases [68]. As such, stigma can hinder accurate
406 public awareness of some diseases and act as a barrier to people adopting health-promoting
407 behaviors, seeking health care, and adhering to treatment.

408 The study findings indicate that community perception tends to be low if the deceased was
409 ≥ 80 years compared to 34-59 years. Despite being insignificant, this may be due to older
410 people being more likely to develop chronic conditions, leading to disease recognition
411 complexity at this age. Consequently, relatives attribute death to “fate” and perceive it as a

412 natural part of the aging process rather than attributing it to a specific cause. We also found
413 that community perception improves significantly if the deceased were married compared to
414 single. Possibly, this is because deceased married individuals have a larger support network,
415 including their spouse, family, and friends. Despite the ethical implications, symptoms and
416 medical history of the deceased may have been shared with relatives, enabling a more accurate
417 determination of the cause of death. Nevertheless, it should be noted that the results may vary
418 according to the particular population being studied and the research methods used.

419
420 Although VA is commonly conducted in a registered population (a population being routinely
421 monitored over time and space such as the concept followed in health socio-demographic and
422 surveillance system), our study utilized medical records from the Saudi regional diabetes
423 register as a denominator and later applied the VA process. Thus, the study finding may be
424 overly pessimistic and have caused overlapping across the death categories. The method's
425 accuracy of cause of death determination is highly dependent on the type of death, the quality
426 of the interview, and the procedures used to assign causes of death [26, 69]. A variety of factors
427 related to the interviewer, the respondent's background, or both can impact the interview
428 quality and thereafter the VA results. The method has been shown to be reasonably practical
429 for identifying causes of death in infancy or due to specific conditions such as injuries or
430 maternal causes. On the other hand, medical causes of adult deaths share symptom complexes,
431 making it difficult to differentiate between various causes of death from such descriptions
432 when based solely on symptoms provided by relatives or occasionally by non-relative such as
433 employers or work colleagues mainly in deaths among expats. Subjects with type 2 diabetes

434 mellitus, for example, have a two- to four-fold increased risk of myocardial infarction and
435 sudden death compared to individuals without diabetes, whereby roughly half of all diabetes
436 deaths co-exist with some types of cardiovascular diseases [70]. This demonstrates the
437 disease's complexities and how the community may fail to recognize the true causes of death
438 without effective health promotion and educational programs.

439
440 Despite the use of an internationally recognized standard VA method, interpreting and
441 generalizing such findings in view of the small sample size can be challenging. The VA telephone
442 interviews were used instead of the traditional face-to-face method to comply with the
443 imposed COVID-19 pandemic's physical distancing measures at the time of the data collection.
444 Nevertheless, numerous studies have found that telephone interviews are as effective as face-
445 to-face interviews [71, 72]. According to one study, the results of VA interviews via phone calls
446 were consistent with previous literature, which found that the telephone interview method to
447 be feasible, accepted by caregivers and healthcare workers, and has a reliable level of data
448 quality [73]. Given the ethical nature of VA, it can arguably be said that conducting distant VA
449 interviews on a fairly small sample in an area where VA has never been practiced before can be
450 viewed as a strength in this study from an ethical perspective, which has the potential of
451 reducing unwanted emotional or cultural consequences.

452 **Conclusion**

453 Diabetes Mellitus is a leading cause of death globally. The Saudi community perception of
454 causes of death was relatively plausible but varied substantially based on the type of death, sex,

455 age \geq 80 years and other vital events like marital status. Higher or lower community perception
456 is a proxy of how people may perceive risk factors associated with the causes of death, which
457 can guide public health interventional programs. Unlike other inconsistent cross-sectional
458 studies, this study proposed a standardized and sustainable method for monitoring people's
459 health literacy and allows harmonized comparison over time and space in otherwise
460 unexplored population. While this study into VA respondents' perceptions of causes of death is
461 interesting for diabetes-related deaths, communities generated a considerably less complete
462 and consistent picture of other cause-specific mortality. This highlights the importance of
463 integrating rigorous and consistent VA methods within the routine medical services rather than
464 relying on individual opinions mainly for complex causes of death outcomes. Nevertheless,
465 translating cause-specific mortality data into effective policymaking remains challenging
466 without standardized assessment of non-medical causes such as social and health system
467 aspects contributing to death.

468 **Acknowledgments**

469 The authors gratefully acknowledge all the deceased's relatives who consented to involvement
470 in this study. We also appreciate the Saudi Ministry of Health, Makkah Health Affairs, for
471 approving this study.

472

473

474

475

476

References

- 477 1. Lin X, Xu Y, Pan X, Xu J, Ding Y, Sun X, et al. Global, regional, and national burden and
478 trend of diabetes in 195 countries and territories: an analysis from 1990 to 2025. *Scientific*
479 *reports*. 2020;10(1):1-11.
- 480 2. Sun H, Saeedi P, Karuranga S, Pinkepank M, Ogurtsova K, Duncan BB, et al. IDF Diabetes
481 Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and
482 projections for 2045. *Diabetes research and clinical practice*. 2022;183:109119.
- 483 3. Ogurtsova K, Guariguata L, Barengo NC, Ruiz PL-D, Sacre JW, Karuranga S, et al. IDF
484 diabetes Atlas: Global estimates of undiagnosed diabetes in adults for 2021. *Diabetes Research*
485 *and Clinical Practice*. 2022;183:109118.
- 486 4. Fish W. *Philosophy of perception: A contemporary introduction*: Routledge; 2010.
- 487 5. Fottrell E, Tollman S, Byass P, Golooba-Mutebi F, Kahn K. The epidemiology of
488 'bewitchment' as a lay-reported cause of death in rural South Africa. *J Epidemiol Community*
489 *Health*. 2012;66(8):704-9.
- 490 6. Hussain-Alkhateeb L, Fottrell E, Petzold M, Kahn K, Byass P. Local perceptions of causes
491 of death in rural South Africa: a comparison of perceived and verbal autopsy causes of death.
492 *Glob Health Action*. 2015;8:28302. Epub 2015/07/22. doi: 10.3402/gha.v8.28302. PubMed
493 PMID: 26193897; PubMed Central PMCID: PMC4507750.
- 494 7. Keikelame MJ, Swartz L. Lost opportunities to improve health literacy: observations in a
495 chronic illness clinic providing care for patients with epilepsy in Cape Town South Africa.
496 *Epilepsy & Behavior*. 2013;26(1):36-41.
- 497 8. Pujilestari CU, Ng N, Hakimi M, Eriksson M. "It is not possible for me to have diabetes"—
498 community perceptions on diabetes and Its risk Factors in Rural Purworejo District, Central
499 Java, Indonesia. *Global Journal of Health Science*. 2014;6(5):204.
- 500 9. Julia L. Lay experiences of health and illness: past research and future agendas.
501 *Sociology of health & illness*. 2003;25(3):23-40.
- 502 10. Piko BF, Bak J. Children's perceptions of health and illness: images and lay concepts in
503 preadolescence. *Health education research*. 2006;21(5):643-53.
- 504 11. Alhawaish AK. Economic costs of diabetes in Saudi Arabia. *Journal of family &*
505 *community medicine*. 2013;20(1):1.
- 506 12. Alotaibi A, Perry L, Gholizadeh L, Al-Ganmi A. Incidence and prevalence rates of diabetes
507 mellitus in Saudi Arabia: An overview. *Journal of epidemiology and global health*.
508 2017;7(4):211-8.
- 509 13. Alwin Robert A, Abdulaziz Al Dawish M, Braham R, Ali Musallam M, Abdullah Al Hayek A,
510 Hazza Al Kahtany N. Type 2 diabetes mellitus in Saudi Arabia: major challenges and possible
511 solutions. *Current diabetes reviews*. 2017;13(1):59-64.
- 512 14. Al-Hanawi MK, Angawi K, Alshareef N, Qattan AM, Helmy HZ, Abudawood Y, et al.
513 Knowledge, attitude and practice toward COVID-19 among the public in the Kingdom of Saudi
514 Arabia: a cross-sectional study. *Frontiers in public health*. 2020;8:217.
- 515 15. Alasiri AA, Mohammed V. Healthcare Transformation in Saudi Arabia: An Overview Since
516 the Launch of Vision 2030. *Health Serv Insights*. 2022;15:11786329221121214. Epub

- 517 2022/09/10. doi: 10.1177/11786329221121214. PubMed PMID: 36081830; PubMed Central
518 PMCID: PMCPMC9445529.
- 519 16. Abdel-Latif MM, Saad SY. Health literacy among Saudi population: a cross-sectional
520 study. *Health promotion international*. 2019;34(1):60-70.
- 521 17. Almubark R, Basyouni M, Alghanem A, Althumairi N, Alkhamis D, Alharbi LS, et al. Health
522 literacy in Saudi Arabia: Implications for public health and healthcare access. *Pharmacology
523 Research & Perspectives*. 2019;7(4):e00514.
- 524 18. AbuAlreesh A, Alburikan KA. Health literacy among patients with poor understanding of
525 prescription drug label instructions in Saudi Arabia. *Saudi Pharmaceutical Journal*.
526 2019;27(6):900-5.
- 527 19. Byass P, Hussain-Alkhateeb L, D'Ambruoso L, Clark S, Davies J, Fottrell E, et al. An
528 integrated approach to processing WHO-2016 verbal autopsy data: the InterVA-5 model. *BMC
529 Med*. 2019;17(1):102. Epub 2019/05/31. doi: 10.1186/s12916-019-1333-6. PubMed PMID:
530 31146736; PubMed Central PMCID: PMCPMC6543589.
- 531 20. D'Ambruoso L, Kahn K, Wagner RG, Twine R, Spies B, Van Der Merwe M, et al. Moving
532 from medical to health systems classifications of deaths: extending verbal autopsy to collect
533 information on the circumstances of mortality. *Global health research and policy*. 2016;1(1):1-
534 15.
- 535 21. D'Ambruoso L, Price J, Cowan E, Goosen G, Fottrell E, Herbst K, et al. Refining
536 circumstances of mortality categories (COMCAT): a verbal autopsy model connecting
537 circumstances of deaths with outcomes for public health decision-making. *Global health action*.
538 2021;14(sup1):2000091.
- 539 22. Hussain-Alkhateeb L, D'Ambruoso L, Tollman S, Kahn K, Van Der Merwe M, Twine R, et
540 al. Enhancing the value of mortality data for health systems: adding Circumstances Of Mortality
541 CATegories (COMCATs) to deaths investigated by verbal autopsy. *Glob Health Action*.
542 2019;12(1):1680068. Epub 2019/10/28. doi: 10.1080/16549716.2019.1680068. PubMed PMID:
543 31648624; PubMed Central PMCID: PMCPMC6818104.
- 544 23. Thomas L-M, D'Ambruoso L, Balabanova D. Verbal autopsy in health policy and systems:
545 a literature review. *BMJ global health*. 2018;3(2):e000639.
- 546 24. Elebro K, Röst M, Moussa K, Johnsdotter S, Essén B. Misclassified maternal deaths
547 among East African immigrants in Sweden. *Reproductive health matters*. 2007;15(30):153-62.
- 548 25. Baiden F, Bawah A, Biai S, Binka F, Boerma T, Byass P, et al. Setting international
549 standards for verbal autopsy. *SciELO Public Health*; 2007. p. 570-1.
- 550 26. Fottrell E, Byass P. Verbal autopsy: methods in transition. *Epidemiologic reviews*.
551 2010;32(1):38-55.
- 552 27. Bauni E, Ndila C, Mochamah G, Nyutu G, Matata L, Ondieki C, et al. Validating physician-
553 certified verbal autopsy and probabilistic modeling (InterVA) approaches to verbal autopsy
554 interpretation using hospital causes of adult deaths. *Population health metrics*. 2011;9(1):1-12.
- 555 28. Leitao J, Desai N, Aleksandrowicz L, Byass P, Miasnikof P, Tollman S, et al. Comparison of
556 physician-certified verbal autopsy with computer-coded verbal autopsy for cause of death
557 assignment in hospitalized patients in low-and middle-income countries: systematic review.
558 *BMC medicine*. 2014;12(1):1-9.

- 559 29. Kalter HD, Salgado R, Babilie M, Koffi AK, Black RE. Social autopsy for maternal and child
560 deaths: a comprehensive literature review to examine the concept and the development of the
561 method. *Population health metrics*. 2011;9(1):1-13.
- 562 30. Källander K, Kadobera D, Williams TN, Nielsen RT, Yevo L, Mutebi A, et al. Social
563 autopsy: INDEPTH Network experiences of utility, process, practices, and challenges in
564 investigating causes and contributors to mortality. *Population health metrics*. 2011;9(1):1-12.
- 565 31. McCormick TH, Li ZR, Calvert C, Crampin AC, Kahn K, Clark SJ. Probabilistic cause-of-
566 death assignment using verbal autopsies. *Journal of the American Statistical Association*.
567 2016;111(515):1036-49.
- 568 32. Hart JD, de André PA, de André CDS, Adair T, Barroso LP, Valongueiro S, et al. Validation
569 of SmartVA using conventional autopsy: a study of adult deaths in Brazil. *The Lancet Regional
570 Health-Americas*. 2022;5:100081.
- 571 33. Chandramohan D, Fottrell E, Leitao J, Nichols E, Clark SJ, Alsokhn C, et al. Estimating
572 causes of death where there is no medical certification: evolution and state of the art of verbal
573 autopsy. *Global Health Action*. 2021;14(sup1):1982486.
- 574 34. Aukes AM, Arion K, Bone JN, Li J, Vidler M, Bellad MB, et al. Causes and circumstances of
575 maternal death: a secondary analysis of the Community-Level Interventions for Pre-eclampsia
576 (CLIP) trials cohort. *Lancet Glob Health*. 2021;9(9):e1242-e51. Epub 2021/08/02. doi:
577 10.1016/s2214-109x(21)00263-1. PubMed PMID: 34332699; PubMed Central PMCID:
578 PMCPMC8370879.
- 579 35. Thomas LM, D'Ambruoso L, Balabanova D. Verbal autopsy in health policy and systems:
580 a literature review. *BMJ Glob Health*. 2018;3(2):e000639. Epub 2018/05/08. doi:
581 10.1136/bmjgh-2017-000639. PubMed PMID: 29736271; PubMed Central PMCID:
582 PMCPMC5935163.
- 583 36. Byass P, Chandramohan D, Clark SJ, D'ambrosio L, Fottrell E, Graham WJ, et al.
584 Strengthening standardised interpretation of verbal autopsy data: the new InterVA-4 tool.
585 *Global health action*. 2012;5(1):19281.
- 586 37. Fantahun M, Fottrell E, Berhane Y, Wall S, Högberg U, Byass P. Assessing a new
587 approach to verbal autopsy interpretation in a rural Ethiopian community: the InterVA model.
588 *Bulletin of the World Health Organization*. 2006;84(3):204-10.
- 589 38. Streatfield PK, Khan WA, Bhuiya A, Alam N, Sié A, Soura AB, et al. Cause-specific
590 mortality in Africa and Asia: evidence from INDEPTH health and demographic surveillance
591 system sites. *Global health action*. 2014;7(1):25362.
- 592 39. Leitao J, Chandramohan D, Byass P, Jakob R, Bundhamcharoen K, Choprapawon C, et al.
593 Revising the WHO verbal autopsy instrument to facilitate routine cause-of-death monitoring.
594 *Global health action*. 2013;6(1):21518.
- 595 40. Lawrence I, Lin K. A concordance correlation coefficient to evaluate reproducibility.
596 *Biometrics*. 1989:255-68.
- 597 41. Byass P, Herbst K, Fottrell E, Ali MM, Odhiambo F, Amek N, et al. Comparing verbal
598 autopsy cause of death findings as determined by physician coding and probabilistic modelling:
599 a public health analysis of 54 000 deaths in Africa and Asia. *Journal of global health*. 2015;5(1).
- 600 42. Berkman ND, Sheridan SL, Donahue KE, Halpern DJ, Crotty K. Low health literacy and
601 health outcomes: an updated systematic review. *Annals of internal medicine*. 2011;155(2):97-
602 107.

- 603 43. Kindig DA, Panzer AM, Nielsen-Bohlman L. Health literacy: a prescription to end
604 confusion. 2004.
- 605 44. Chandramohan D, Maude GH, Rodrigues LC, Hayes RJ. Verbal autopsies for adult deaths:
606 their development and validation in a multicentre study. *Tropical Medicine & International*
607 *Health*. 1998;3(6):436-46.
- 608 45. Quigley M, Schellenberg JA, Snow RW. Algorithms for verbal autopsies: a validation
609 study in Kenyan children. *Bulletin of the World Health Organization*. 1996;74(2):147.
- 610 46. Quigley MA, Chandramohan D, Rodrigues LC. Diagnostic accuracy of physician review,
611 expert algorithms and data-derived algorithms in adult verbal autopsies. *International journal*
612 *of epidemiology*. 1999;28(6):1081-7.
- 613 47. Kessels RP. Patients' memory for medical information. *Journal of the Royal Society of*
614 *Medicine*. 2003;96(5):219-22.
- 615 48. Moser RP, Arndt J, Han PK, Waters EA, Amsellem M, Hesse BW. Perceptions of cancer as
616 a death sentence: prevalence and consequences. *Journal of health psychology*.
617 2014;19(12):1518-24.
- 618 49. Robb KA, Simon AE, Miles A, Wardle J. Public perceptions of cancer: a qualitative study
619 of the balance of positive and negative beliefs. *BMJ open*. 2014;4(7):e005434.
- 620 50. Niederdeppe J, Levy AG. Fatalistic beliefs about cancer prevention and three prevention
621 behaviors. *Cancer Epidemiology Biomarkers & Prevention*. 2007;16(5):998-1003.
- 622 51. Clarke JN, Everest MM. Cancer in the mass print media: Fear, uncertainty and the
623 medical model. *Social science & medicine*. 2006;62(10):2591-600.
- 624 52. Miles A, Rainbow S, von Wagner C. Cancer Fatalism and Poor Self-Rated Health Mediate
625 the Association between Socioeconomic Status and Uptake of Colorectal Cancer Screening in
626 England Socioeconomic Status, Fatalism, Health, and CRC Screening. *Cancer epidemiology,*
627 *biomarkers & prevention*. 2011;20(10):2132-40.
- 628 53. Alaqeel A, AlAmmari A, AlSyefi N, Al-Hussain F, Mohammad Y. Stroke awareness in the
629 Saudi community living in Riyadh: prompt public health measures must be implemented.
630 *Journal of Stroke and Cerebrovascular Diseases*. 2014;23(3):500-4.
- 631 54. Basfar WM, Al-Sebyani AE, Aljawi GA, Milyani HA, Jan MM. Public knowledge of stroke
632 amongst a saudi population. *World Journal of Research and Review*. 2016;3(2):262912.
- 633 55. Krishnamurthi RV, Barker-Collo S, Barber PA, Tippet LJ, Dalrymple-Alford JC, Tunnage B,
634 et al. Community knowledge and awareness of stroke in New Zealand. *Journal of Stroke and*
635 *Cerebrovascular Diseases*. 2020;29(3):104589.
- 636 56. Al-Husayni F, Al-Zahrani A, Zwawy M, Alamri S, Aljedaani R, Almalki A. The awareness
637 and perception of chronic kidney disease in Jeddah, Saudi Arabia. *Saudi Journal of Kidney*
638 *Diseases and Transplantation*. 2021;32(2):488.
- 639 57. Alobaidi S. Knowledge of chronic kidney disease among the population of Saudi Arabia
640 evaluated using a validated questionnaire: a cross-sectional study. *Patient preference and*
641 *adherence*. 2021;15:1281.
- 642 58. Mousa D, Alharbi A, Helal I, Al-Homrany M, Alhujaili F, Alhweish A, et al. Prevalence and
643 Associated Factors of Chronic Kidney Disease among Relatives of Hemodialysis Patients in Saudi
644 Arabia. *Kidney international reports*. 2021;6(3):817.

- 645 59. Al-Baghli NA, Al-Ghamdi AJ, Al-Turki KA, El-Zubaier AG, Al-Mostafa BA, Al-Baghli FA, et
646 al. Awareness of cardiovascular disease in eastern Saudi Arabia. *Journal of Family and*
647 *Community Medicine*. 2010;17(1):15.
- 648 60. AL-Tamimi I, Bawazir A. Knowledge and Perceived Susceptibility of Cardiovascular
649 Diseases (CVDs) among Saudi Female Teachers. *Into Arch Public Health Community Med*.
650 2020;4(45):1-10.
- 651 61. Homko CJ, Santamore WP, Zamora L, Shirk G, Gaughan J, Cross R, et al. Cardiovascular
652 disease knowledge and risk perception among underserved individuals at increased risk of
653 cardiovascular disease. *Journal of Cardiovascular Nursing*. 2008;23(4):332-7.
- 654 62. Bekele DM, Goshu DY, Yalew AW, Higgins MK, Gary RA. Low Subjective Cardiovascular
655 Disease Risk Perceptions among Hypertensive Patients in Addis Ababa, Ethiopia. *Integrated*
656 *Blood Pressure Control*. 2022;15:81.
- 657 63. Webster R, Heeley E. Perceptions of risk: understanding cardiovascular disease. *Risk*
658 *Management and Healthcare Policy*. 2010;3:49.
- 659 64. Liu H, Li M, Jin M, Jing F, Wang H, Chen K. Public awareness of three major infectious
660 diseases in rural Zhejiang province, China: a cross-sectional study. *BMC infectious diseases*.
661 2013;13(1):1-9.
- 662 65. Alhazzani AA, Alqahtani AM, Abouelyazid A, Alqahtani AM, Alqahtani NA, Asiri KM, et al.
663 Public awareness, knowledge, and attitudes toward epilepsy in the Aseer region, Saudi Arabia—
664 a community-based cross-sectional study. *Epilepsy & Behavior*. 2016;63:63-6.
- 665 66. Almoayad F, Mahboub S, Amer LB, Alrabiah A, Alhashem A. Stigmatisation of COVID-19
666 in Riyadh, Saudi Arabia: a cross-sectional study. *Sultan Qaboos University Medical Journal*.
667 2021;21(4):525.
- 668 67. Raheel H. Stigma towards people living with HIV/AIDS (PLWAs) among adolescents of
669 Riyadh, Kingdom of Saudi Arabia. *J AIDS Clin Res*. 2016;7(612):2.
- 670 68. Hinga A, Marsh V, Nyaguara A, Wamukoya M, Molyneux S. The ethical implications of
671 verbal autopsy: responding to emotional and moral distress. *BMC medical ethics*. 2021;22(1):1-
672 16.
- 673 69. Yang G, Rao C, Ma J, Wang L, Wan X, Dubrovsky G, et al. Validation of verbal autopsy
674 procedures for adult deaths in China. *International journal of epidemiology*. 2006;35(3):741-8.
- 675 70. Vargas RB, Mangione CM, Asch S, Keesey J, Rosen M, Schonlau M, et al. Can a chronic
676 care model collaborative reduce heart disease risk in patients with diabetes? *Journal of general*
677 *internal medicine*. 2007;22(2):215-22.
- 678 71. Block ES, Erskine L. Interviewing by telephone: Specific considerations, opportunities,
679 and challenges. *International journal of qualitative methods*. 2012;11(4):428-45.
- 680 72. Holt A. Using the telephone for narrative interviewing: a research note. *Qualitative*
681 *research*. 2010;10(1):113-21.
- 682 73. binti Nasaruddin NH, Ganapathy SS, Awaluddin SM, Anuar MFM, binti Alias N, Mang CY,
683 et al. Conducting verbal autopsy by telephone interview during the pandemic to support
684 mortality surveillance: a feasibility study in Malaysia: Verbal autopsy telephone interview.
685 *Western Pacific Surveillance and Response*. 2022;13(2).
- 686

687

688 **Supporting information**

689 **S1 Table. Categorizing of causes of death reported by VA and families**

Category	Cause of death
Infectious diseases	Sepsis (non-obstetric) Acute resp infect incl pneumonia HIV/AIDS related death Pulmonary tuberculosis Other and unspecified infect diseases
Cancer	Digestive neoplasms Respiratory neoplasms Breast neoplasms Other and unspecified neoplasms Pancreatic cancer, Lung cancer Tumors, Head cancer
Diabetes mellitus	Diabetes mellitus Gangrene, Diabetic foot amputation, foot ulcer
Circulatory	Acute cardiac disease Other and unspecified cardiac disease Heart failure, heart attack cardiomyopathy or myocardial insufficiency Cardiac arrest, Heart attack, Hypertension, Hypotension, Heart disease
NCD	Chronic obstructive pulmonary disease Abdominal related Acute abdomen Other and unspecified NCD
Renal	Kidney disease, kidney failure, dialysis Renal failure
Accidental	Accidental fall Road traffic accident Accidental injuries Accidental expos to smoke fire & flame
Indeterminate	Any cause of death that has no ICD code
Stroke	Stroke
Covid-19	Covid-19
Mental illness	Dementia, Alzheimer




Fig 1. Concordance correlation coefficient (log-logscale) between CSMF determined by VA and respondent reports, in relation to the line of equivalence, for 302 deaths in Makkah city, Saudi Arabia.

Figure