

1 **The predictors of new-onset atrial fibrillation incidence after on-pump**
2 **coronary artery bypass graft surgery: A prospective study evaluating syntax**
3 **score**

4 **Running title:** Incidence of new-onset atrial fibrillation after coronary artery bypass graft
5 surgery could be predicted by syntax score.

6
7 Authors: Hoda Mombeini ¹, Alireza Ebrahimi ², Saeed Yazdankhah ³, Mohammad Ali Sheikhi ⁴,
8 Shahla Majidi ³, Majid Pakdin ^{2*}

- 9
10 1- Dept. Cardiology, Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran
11 2- Student Research Committee, Faculty of Medicine, Shiraz University of Medical
12 Sciences, Shiraz, Iran
13 3- Dept. Cardiology, Nikan Hospital, Tehran, Iran
14 4- Dept. Cardiovascular Surgery, Atherosclerosis Research Center, Jundishapur University
15 of Medical Sciences, Ahvaz, Iran

16
17 * Corresponding author: Majid Pakdin, M.D

18 Address: Student Research Committee, Shiraz Medical School, Karimkhan Zand St, Shiraz, Iran

19 Email: mp.pakdin@gmail.com

20
21
22 ***Competing interest***

23 The authors confirm that there is no conflict of interest to declare.

24 ***Funding***

25 The authors received no financial support for the investigation and authorship.

26
27
28
29

1 **Abstract**

2

3 **Background:** Atrial fibrillation (AF) is considered the most common supraventricular
4 arrhythmia in patients undergoing coronary artery bypass graft (CABG). The predictive value of
5 the SYNTAX score for post-CABG new-onset AF incidence has not been clearly evaluated. This
6 study aimed to assess this association in patients undergoing isolated on-pump CABG.

7 **Method:** This study was done in a single-center, randomized, and observational setting. A total
8 of 133 patients undergoing on-pump isolated CABG who were older than 18 years and had sinus
9 rhythm were enrolled. Demographic variables of patients were recorded, and the SYNTAX score
10 was measured for the participants. The multivariate logistic regression model was applied to
11 identify the predictors of post-CABG new-onset AF.

12 **Results:** The logistic regression model showed that SYNTAX score of more than 28.25 (p-
13 value= 0.001; OR= 14.25, 95% CI= 2.90_70.11), hypertension (p-value=0.02; OR = 6.59, 95%
14 CI = 1.23_34.57), and calcium channel blocker consumption (p-value=0.02; OR = 8.05, 95% CI
15 = 1.43_45.42) are predictors of new-onset AF after on-pump CABG.

16 **Conclusion:** This study demonstrated that patients with higher SYNTAX scores in coronary
17 angiography are more likely to develop new-onset AF after isolated on-pump CABG.

18

19 **Keywords:** Coronary Artery Bypass Grafting, SYNTAX Score, Postoperative Atrial Fibrillation

20

21

22

23

24

25

26

27

28

29

30

31

1 **1. Background**

2 Atrial fibrillation (AF) is the most common arrhythmia occurring after coronary artery bypass
3 graft (CABG) surgery, with an incidence of approximately 27-33% [1-3]. Several risk factors
4 were found to be related to postoperative AF occurrence, such as old age, diabetes mellitus,
5 hypertension, chronic obstructive pulmonary disease, paroxysmal AF history, left ventricular
6 dysfunction, cardiopulmonary bypass time, and cross-clamp time [2]. Patients are more
7 vulnerable to developing AF arrhythmias within the first 4-5 postoperative days, particularly
8 during the first 72 hours after the surgery [4]. Furthermore, post-CABG AF is accompanied by
9 higher complications such as thromboembolism, renal failure, heart failure, mortality, and
10 extended hospital stay, leading to lower quality of life for patients and increased treatment
11 costs [5, 6]. The survival probability was found to be decreased in patients who had suffered
12 from new-onset post-CABG AF [7].

13 The synergy between percutaneous coronary intervention with taxus and cardiac surgery
14 (SYNTAX) score is based on angiographic imaging that assesses the complexity of coronary
15 lesions and the severity of coronary artery disease (CAD) [8]. This scoring system was first
16 introduced to compare two therapeutic methods of percutaneous coronary intervention with taxus
17 stents versus CABG [6]. SYNTAX is also widely applied to predict the long-term consequences
18 of CAD and decide on appropriate therapeutic interventions [8, 9]. Previous studies mentioned
19 that patients undergoing CABG with higher SYNTAX scores are at greater mortality risk [10].

20 Although numerous investigations showed that SYNTAX score could be used as a predictive
21 index for several post-CABG complications, such as myocardial infarction and stent thrombosis,
22 little has been done regarding using this score as a predictor of new-onset post-CABG AF. The
23 present study aimed to determine the association of the score with AF incidence after on-pump
24 CABG.

25 **2. Methods**

26 **2.1. Study Design and Population**

27 A total of 133 patients undergoing on-pump isolated CABG surgery at the cardiac surgery ward
28 in Golestan hospital, Ahvaz, Iran, were randomly enrolled in this observational study and were
29 subsequently evaluated for post-CABG AF. Inclusion criteria were defined as individuals older
30 than 18 years old with cardiac sinus rhythm who had CABG. The exclusion criteria included a
31 history of thyroid disease, paroxysmal AF arrhythmia, previous cardiac surgery, percutaneous

1 coronary intervention (PCI) during the past six months, use of amiodarone or digoxin, moderate
2 to severe mitral regurgitation, having implantable cardioverter defibrillator or permanent
3 pacemaker, or left atrial size more than 4 cm.

4 Patients were observed by continuous electrocardiogram (EKG) monitoring during their stay in
5 the intensive care unit (ICU). After detecting AF during constant monitoring, a 12-lead-EKG
6 was taken to confirm arrhythmia. After being discharged from ICU, patients were also followed
7 in the ward by radial pulse examination at least four times daily in terms of rate and regularity,
8 which was done by expert staff. Should an irregular pulse be observed or patients become
9 symptomatic, a 12-lead-ECG was taken to assess arrhythmia. In this study, AF was described as
10 those that lasted for a minimum of 5 minutes. Ultimately, patients were classified into two
11 groups: patients with AF arrhythmia and those without AF. The patients' SYNTAX scores
12 compared with each other.

13 **2.2. SYNTAX Score Measurement**

14 To calculate the SYNTAX score, we scored each coronary lesion with >50% stenosis in vessels
15 >1.5 mm diameter. Each lesion could involve more than one diseased segment. Other
16 deterministic factors were coronary dominance, total occlusions, trifurcations, bifurcations,
17 aorto-ostial lesions, severe tortuosity, severe calcifications, thrombus, and diffuse diseases.

18 All angiographic variables were calculated by an expert team consisting of a general cardiologist,
19 an interventional cardiologist, and a cardiac surgeon based on the latest version of SYNTAX
20 score. This version has been designed to improve long-term outcomes prediction following
21 coronary revascularization by subsuming seven clinical variables, including age, peripheral
22 vascular disease, chronic obstructive pulmonary disease, unprotected left main CAD, left
23 ventricular ejection fraction, creatinine clearance, and sex, into the original SYNTAX score [11].

24 **2.3. Statistical Analysis**

25 The results were analyzed using Statistical Package for Social Sciences (SPSS), version 22. The
26 relationship between post-CABG AF and different factors was analyzed using Chi-squared test,
27 two-tailed *t*-test, Fisher's exact test, or Mann-Whitney U test as appropriate. Afterward, the
28 multivariate logistic regression model was performed to identify the predictors of post-CABG
29 AF. The predictive power of SYNTAX score for post-CABG AF was tested by receiver

1 operating characteristic (ROC) curve analysis. P value < 0.05 was considered statistically
2 significant.

3 **2.4. Ethical Considerations**

4 The study protocol followed national and international guidelines. Informed consent was
5 obtained from all participants, their anonymity was guaranteed, and the study protocol was
6 approved by the Ethical Committee of Ahvaz Jundishapur University of Medical Sciences (Reg.
7 No. IR.AJUMS.REC.1394.91).

8 **3. Results**

9 One hundred thirty-three patients with a mean age of 59 ± 8.5 years participated in our study.
10 Most participants were male, and the prevalence rates of diabetes mellitus, hypertension,
11 dyslipidemia, and cigarette smoking were 51.1%, 75.2%, 69.2%, and 25.6%, respectively. The
12 left ventricular ejection fraction (LVEF) ranged from 20% to 60% (Mean \pm SD: $47.7 \pm 8.8\%$).
13 The mean number of involved vessels was 2.6 ± 0.52 , and the SYNTAX score ranged from 7 to
14 55.5 (mean: 28.66 ± 9.95). The left internal thoracic artery was used in 131 (98.5%) patients,
15 while 23 (17.3%) and 80 (60.2%) patients received one and two saphenous vein grafts,
16 respectively. Three and four saphenous vein grafts were used for 25 (18.8%) and 4 (3%),
17 respectively. One remaining patient received no vein graft.

18 New-onset post-CABG AF incidence in our study was 21.8%. Preoperative variables and
19 medications, as well as perioperative variables, are demonstrated in Tables 1, 2, and 3. During
20 the investigation, there were three cases of mortality which all had developed AF arrhythmia
21 during hospitalization. The cause of two deaths was postoperative acute renal failure. These two
22 cases developed AF on days 2 and 3 of hospitalization. The mortality of the third case, who
23 developed AF on day 7, was due to pneumonia and respiratory failure after prolonged admission
24 in the ICU.

25 The ROC curve analysis showed good accuracy for the SYNTAX score in predicting post-
26 CABG AF, with the area under the curve being 0.71 [95% CI: (0.69, 0.81) $p < 0.001$]. The
27 sensitivity and specificity were 89% and 56%, respectively, for SYNTAX scores greater than
28 28.25 (Figure 1).

29 Multivariate analysis of the risk of new-onset AF incidence was also performed. Ten factors,
30 including aspirin and calcium channel blockers, hypertension, left circumflex (LCX) artery

1 involvement, SYNTAX score, creatinine level, LVEF, platelet count, cardiopulmonary bypass
2 time, and the number of anastomoses were considered in our analysis (Table 4).

3 The logistic regression model showed that SYNTAX score of more than 28.25 (p-value= 0.001;
4 OR= 14.25, 95% CI= 2.90_70.11), hypertension (p-value=0.02; OR = 6.59, 95% CI =
5 1.23_34.57), and calcium channel blocker consumption (p-value=0.018; OR = 8.05, 95% CI =
6 1.43_45.42) were the predictors of AF after on-pump CABG.

7 **4. Discussion**

8 AF is considered the most common cardiac arrhythmia and accounts for mechanical heart
9 dysfunction and subsequent complications [12]. The prevalence of AF in the general population
10 has been measured to be 1-2% [12]. Among patients who had CABG, the incidence of AF is
11 much higher, about 25-40% [13], and it has been reported that AF is the leading cause of post-
12 CABG mortality [14]. A range of investigations has examined different prognostic AF indexes in
13 patients undergoing CABG. Consistent with previous findings, our study showed a clinically
14 significant correlation between post-CABG AF and hypertension, use of calcium channel
15 blockers, and high SYNTAX score.

16 Different scoring systems, including HATCH score, EusoSCORE, and SYNTAX score, have
17 been used to assess the relationship of the obtained score with post-CAB AF incidence in
18 previous studies [12, 13, 15-17]. In the current study, the latter has been used for this purpose.
19 Priorly, SYNTAX score was classified as a reliable predictor of major cardiovascular event rate
20 after revascularization [18-20].

21 Although the association of SYNTAX score and new-onset post-CABG AF has not been
22 evaluated extensively, few studies have shown the relationship between higher SYNTAX score
23 and new-onset post-CABG AF incidence in patients who underwent on-pump CABG [12, 13,
24 21]. The present study showed that a higher SYNTAX score could be used as a prognostic factor
25 of new-onset post-CABG AF incidence in patients who underwent on-pump CABG. The cutoff
26 level of SYNTAX score to predict post-CABG AF was 28.25 in our study, which is higher than
27 the determined cutoff level in previous studies [12, 13, 21]. Contrarily, Fukui *et al.* reported that
28 both high SYNTAX score and high Euroscore II were independent predictors of early major
29 complications after CABG; however, according to their investigation, new-onset post-CABG AF
30 was not increased in the group with both high SYNTAX score and high Euroscore II than other
31 groups [22]. Compared to our study, they used a combination of Euroscore II and SYNTAX
Mombeini, H. et al.

1 score, and their study was not designed exclusively to evaluate the correlation between
2 SYNTAX score and new-onset post-CABG AF; as a result, the patients with preoperative AF
3 arrhythmia had not been excluded from the study. Furthermore, the majority of patients in their
4 study had undergone off-pump CABG, and as previously documented, those cases undergoing
5 off-pump CABG are at lower risk of developing AF [23].

6 Several studies have identified hypertension as a risk factor for post-CABG AF [24, 25].
7 Consistent with the current body of evidence, our data suggest hypertension as an independent
8 risk factor for post-CABG AF (OR = 6.5). This association may be due to hypertension-related
9 structural changes. A number of changes in the hypertrophied heart, such as fibrosis, may act as
10 a substrate for reentry arrhythmias [26, 27]. Subsequently, AF could be maintained by these
11 focal reentry firings [28].

12 According to previous studies, non-dihydropyridine calcium channel blockers (CCBs) could
13 significantly reduce the incidence of post-cardiac surgery supraventricular arrhythmias [29].
14 Additionally, prophylactic diltiazem could reduce new-onset post-CABG AF [30]. In contrast to
15 these findings, a comprehensive evaluation of different pharmacotherapies for the prevention of
16 post-CABG AF reported that CCB therapy could not decrease the odds of new-onset post-CABG
17 AF [31]. In our study, CCB consumers were found to have higher rates of postoperative AF than
18 non-consumers. The following aspects could explain this finding: (1) In our analysis, all patients
19 were on dihydropyridine treatment, not nondihydropyridines. (2) With the paucity of information
20 on postoperative adherence to medications, it may be possible that this result is due to low
21 postoperative continuation of calcium channel blockers and hypertension crisis following
22 calcium channel blocker withdrawal, as the majority of our patients used them for hypertension.

23 There were several limitations in the present study. First, our study population could not be
24 considered representative of all post-CABG patients owing to the fact that our cases were
25 obtained only from one center and are subject to sampling bias. Second, we stated each patient's
26 cardiac rhythm status based on the preoperative ECG findings relying on the medical charts to
27 determine whether any prior episodes of AF arrhythmia had been noted. As a result, we might
28 have missed potential asymptomatic paroxysmal AF arrhythmia in individuals who showed
29 normal sinus rhythm on the preoperative ECG. Finally, continuous Holter ECG monitoring was
30 not continued after ICU discharge; however, radial pulses and symptoms were assessed in the

1 ward at least four times daily, so there is little likelihood of sustained episodes of AF being
2 missed.

3 **5. Conclusion**

4 The results of this study reveal that SYNTAX score ≥ 28.25 , hypertension, and use of calcium
5 channel blockers are predictors of post-CABG AF. Therefore, it seems sensible to calculate
6 preoperative SYNTAX scores and consider using anti-arrhythmic agents for those with high
7 SYNTAX scores, which may ultimately decrease post-CABG AF occurrence.

8 **6. Acknowledgement**

9 The authors appreciate the nurses and physicians at the cardiac surgery ward of Golestan
10 Hospital, Ahvaz, who helped to conduct this study.

11 **7. References**

- 12 1. Mathew JP, Parks R, Savino JS, Friedman AS, Koch C, Mangano DT, et al. Atrial
13 fibrillation following coronary artery bypass graft surgery: predictors, outcomes, and resource
14 utilization. *Jama*. 1996;276(4):300-6.
- 15 2. Peretto G, Durante A, Limite LR, Cianflone D. Postoperative arrhythmias after cardiac
16 surgery: incidence, risk factors, and therapeutic management. *Cardiology research and practice*.
17 2014;2014.
- 18 3. Mathew JP, Fontes ML, Tudor IC, Ramsay J, Duke P, Mazer CD, et al. A multicenter
19 risk index for atrial fibrillation after cardiac surgery. *Jama*. 2004;291(14):1720-9.
- 20 4. Podrid PJ. Prevention of postoperative atrial fibrillation: what is the best approach. *J Am*
21 *Coll Cardiol*. 1999;34(2):340-2.
- 22 5. Creswell LL, Damiano Jr RJ. Postoperative atrial fibrillation: an old problem crying for
23 new solutions. *The Journal of thoracic and cardiovascular surgery*. 2001;121(4):638-41.
- 24 6. Gungor H, Eryilmaz U, Akgullu C, Zencir C, Kurtoglu T, Selvi M, et al. Preoperative
25 poor coronary collateral circulation can predict the development of atrial fibrillation after
26 coronary artery bypass graft surgery. *Coronary artery disease*. 2013;24(7):572-6.
- 27 7. Villareal RP, Hariharan R, Liu BC, Kar B, Lee V-V, Elayda M, et al. Postoperative atrial
28 fibrillation and mortality after coronary artery bypass surgery. *Journal of the American College*
29 *of Cardiology*. 2004;43(5):742-8.
- 30 8. Bundhun PK, Sookharee Y, Bholee A, Huang F. Application of the SYNTAX score in
31 interventional cardiology: a systematic review and meta-analysis. *Medicine*. 2017;96(28).

- 1 9. Farooq V, Head SJ, Kappetein AP, Serruys PW. Widening clinical applications of the
2 SYNTAX Score. *Heart*. 2014;100(4):276-87.
- 3 10. Yang H, Zhang L, Xu CH. Use of the SYNTAX Score II to predict mortality in
4 interventional cardiology: A systematic review and meta-analysis. *Medicine*. 2019;98(2).
- 5 11. Farooq V, Van Klaveren D, Steyerberg EW, Meliga E, Vergouwe Y, Chieffo A, et al.
6 Anatomical and clinical characteristics to guide decision making between coronary artery bypass
7 surgery and percutaneous coronary intervention for individual patients: development and
8 validation of SYNTAX score II. *The Lancet*. 2013;381(9867):639-50.
- 9 12. Engin M, Aydın C. Investigation of the effect of HATCH score and coronary artery
10 disease complexity on atrial fibrillation after on-pump coronary artery bypass graft surgery.
11 *Medical Principles and Practice*. 2021;30(1):45-51.
- 12 13. Geçmen Ç, Babür Güler G, Erdoğan E, Hatipoğlu S, Güler E, Yılmaz F, et al. SYNTAX
13 score predicts postoperative atrial fibrillation in patients undergoing on-pump isolated coronary
14 artery bypass grafting surgery. *The Anatolian Journal of Cardiology*. 2016.
- 15 14. El-Chami MF, Kilgo P, Thourani V, Lattouf OM, Delurgio DB, Guyton RA, et al. New-
16 onset atrial fibrillation predicts long-term mortality after coronary artery bypass graft. *Journal of*
17 *the American College of Cardiology*. 2010;55(13):1370-6.
- 18 15. Bundhun PK, Sookharee Y, Bholee A, Huang F. Application of the SYNTAX score in
19 interventional cardiology: A systematic review and meta-analysis. *Medicine (Baltimore)*.
20 2017;96(28):e7410.
- 21 16. Emren V, Aldemir M, Duygu H, Kocabaş U, Tecer E, Cerit L, et al. Usefulness of
22 HATCH score as a predictor of atrial fibrillation after coronary artery bypass graft. *Kardiologia*
23 *Polska (Polish Heart Journal)*. 2016;74(8):749-53.
- 24 17. Selvi M, Gungor H, Zencir C, Gulasti S, Eryilmaz U, Akgullu C, et al. A new predictor
25 of atrial fibrillation after coronary artery bypass graft surgery: HATCH score. *Journal of*
26 *Investigative Medicine*. 2018;66(3):648-52.
- 27 18. Sianos G, Morel M-A, Kappetein AP, Morice M-C, Colombo A, Dawkins K, et al. The
28 SYNTAX Score: an angiographic tool grading the complexity of coronary artery disease.
29 *EuroIntervention*. 2005;1(2):219-27.
- 30 19. Valgimigli M, Serruys PW, Tsuchida K, Vaina S, Morel M-A, van den Brand MJ, et al.
31 Cyphering the complexity of coronary artery disease using the syntax score to predict clinical
Mombeini, H. et al.

- 1 outcome in patients with three-vessel lumen obstruction undergoing percutaneous coronary
2 intervention. *The American journal of cardiology*. 2007;99(8):1072-81.
- 3 20. Serruys PW, Morice M-C, Kappetein AP, Colombo A, Holmes DR, Mack MJ, et al.
4 Percutaneous coronary intervention versus coronary-artery bypass grafting for severe coronary
5 artery disease. *New England journal of medicine*. 2009;360(10):961-72.
- 6 21. Cerit L, Duygu H, Gulsen K, Kemal HS, Ozcem B, Balcioglu O, et al. Is SYNTAX score
7 predictive of atrial fibrillation after on-pump coronary artery bypass graft surgery? *Korean*
8 *circulation journal*. 2016;46(6):798-803.
- 9 22. Fukui T, Uchimuro T, Takanashi S. EuroSCORE II with SYNTAX score to assess risks
10 of coronary artery bypass grafting outcomes. *European Journal of Cardio-Thoracic Surgery*.
11 2015;47(1):66-71.
- 12 23. Møller CH, Penninga L, Wetterslev J, Steinbrüchel DA, Gluud C. Off-pump versus
13 on-pump coronary artery bypass grafting for ischaemic heart disease. *Cochrane Database of*
14 *Systematic Reviews*. 2012(3).
- 15 24. Aranki SF, Shaw DP, Adams DH, Rizzo RJ, Couper GS, VanderVliet M, et al. Predictors
16 of atrial fibrillation after coronary artery surgery. Current trends and impact on hospital
17 resources. *Circulation*. 1996;94(3):390-7.
- 18 25. Sobczyk D, Sadowski J, Sniezek-Maciejewska M. [Causes of atrial fibrillation early after
19 coronary artery bypass grafting]. *Przegl Lek*. 2005;62(3):141-7.
- 20 26. McLenachan JM, Dargie HJ. Ventricular arrhythmias in hypertensive left ventricular
21 hypertrophy. Relationship to coronary artery disease, left ventricular dysfunction, and
22 myocardial fibrosis. *Am J Hypertens*. 1990;3(10):735-40.
- 23 27. Toyoshima H, Park YD, Ishikawa Y, Nagata S, Hirata Y, Sakakibara H, et al. Effect of
24 ventricular hypertrophy on conduction velocity of activation front in the ventricular myocardium.
25 *Am J Cardiol*. 1982;49(8):1938-45.
- 26 28. Iwasaki Y-k, Nishida K, Kato T, Nattel S. Atrial fibrillation pathophysiology:
27 implications for management. *Circulation*. 2011;124(20):2264-74.
- 28 29. Wijeyesundera DN, Beattie WS, Rao V, Karski J. Calcium antagonists reduce
29 cardiovascular complications after cardiac surgery: a meta-analysis. *Journal of the American*
30 *College of Cardiology*. 2003;41(9):1496-505.

1 30. Dobrilovic N, Vadlamani L, Buchert B, Wright C. Diltiazem prophylaxis reduces
 2 incidence of atrial fibrillation after coronary artery bypass grafting. *Journal of Cardiovascular*
 3 *Surgery*. 2005;46(5):457.

4 31. Piccini JP, Zhao Y, Steinberg BA, He X, Mathew JP, Fullerton DA, et al. Comparative
 5 effectiveness of pharmacotherapies for prevention of atrial fibrillation following coronary artery
 6 bypass surgery. *The American Journal of Cardiology*. 2013;112(7):954-60

7 8. Tables

8

Table 1. Pre-operative variables and laboratory data.

| | Group Non-AF | Group AF | P- value |
|---|--|---|--------------------|
| | N(%) or Mean \pmSD (n=104) | N(%) or Mean \pmSD (n=29) | |
| Age | 58.7 \pm 8.5 | 60.6 \pm 8.7 | 0.48 ¹ |
| Gender (% male) | 70 (67.3%) | 19 (65.5%) | 0.86 ² |
| Diabetes Mellitus | 52 (50.0%) | 16 (55.2%) | 0.62 ² |
| Hypertension | 73 (70.2%) | 27 (93.1%) | 0.01 ² |
| Dyslipidemia | 71 (68.3%) | 21 (72.4%) | 0.67 ² |
| Body mass index | 26.7 \pm 3.4 | 28 \pm 5.3 | 0.24 ¹ |
| Cigarette Smoking | 29 (27.9%) | 5 (17.2%) | 0.25 ² |
| Stroke history | 10 (9.6%) | 4 (13.8%) | 0.51 ² |
| Myocardial infarction history | 36 (34.6%) | 12 (41.4%) | 0.51 ² |
| Left main disease | 12 (11.5%) | 2 (6.9%) | 0.73 ² |
| Left anterior descending disease | 104 (100%) | 29 (100%) | >0.99 ² |
| Left circumflex disease | 85 (81.7%) | 27 (93.1%) | 0.16 ² |
| Ramus intermedius artery | 4 (3.8%) | 1 (3.4%) | 0.92 ² |
| Right coronary artery disease | 87 (83.7%) | 25 (86.2%) | 0.74 ² |
| Number of involved vessels | | | |
| One | 4 (3.8%) | 0 (0.0%) | 0.53 ¹ |
| Two | 28 (26.9%) | 6 (20.7%) | |

| | | | |
|---|---------------|----------------|--------------------|
| Three | 72 (69.2%) | 23 (79.3%) | |
| Left ventricular ejection fraction | 48.12± 6.76 | 45.17± 9.11 | 0.08 ¹ |
| SYNTAX Score | 27.25± 10.27 | 33.76± 6.66 | 0.012 ¹ |
| Creatinine | 1.01± 0.24 | 1.21± 0.61 | 0.12 ¹ |
| Glomerular filtration rate | 82.93± 24.93 | 76.44± 29.43 | 0.24 ¹ |
| Hemoglobin | 12.3± 1.53 | 12.3± 1.60 | 0.94 ¹ |
| Platelet | 262.38± 77.35 | 299.00± 114.35 | 0.12 ¹ |

¹ Two-tailed T-test

² χ^2

1

Table 2. Preoperative medications.

| | Group non-AF | Group AF | P- value ¹ |
|--|-----------------|----------------|-----------------------|
| | N(%) (n=104) | N(%) (n=29) | |
| Beta-blocker | 90 (86.5%) | 23 (79.3%) | 0.34 |
| Aspirin | 96 (92.3%) | 24 (82.8%) | 0.13 |
| Statin | 87 (83.7%) | 24 (82.8%) | 0.91 |
| Renin-angiotensin system inhibitors | 57 (54.8%) | 17 (58.6%) | 0.72 |
| Calcium channel blocker | 8 (7.7%) | 5 (17.2%) | 0.13 |

¹ χ^2

2

Table 3. Peri-operative variables.

| | Group non-AF | Group AF | P- value |
|---|---------------------------------|--------------------------------|---------------------|
| | N (%) or mean ±SD (n=104) | N (%) or mean ±SD (n=29) | |
| Cardio-pulmonary bypass time | 74.5±21.93 | 84.00±32.60 | 0.14 ¹ |
| Aortic cross-clamp time | 42.80±11.80 | 43.83±10.15 | 0.51 ¹ |
| Use of left internal thoracic artery | 102 (98.1%) | 29 (100%) | > 0.99 ² |
| Number of anastomoses | 2.97±0.74 | 3.31±0.54 | 0.008 ¹ |

¹ Two-tailed t-test

χ^2

1
2
3

Table 4. Multivariate analysis of the risk of new-onset AF incidence.


| | P-value ¹ | Odds ratio | 95% CI |
|---|----------------------|------------|---------------|
| Aspirin | 0.210 | 0.336 | 0.62- 1.835 |
| Calcium channel blocker | 0.018 | 8.045 | 1.425- 45.416 |
| Hypertension | 0.026 | 6.587 | 1.225- 34.568 |
| Left circumflex artery disease | 0.677 | 1.472 | 0.238- 9.106 |
| Syntax score (> 28.25) | 0.001 | 14.253 | 2.898- 70.111 |
| Creatinine | 0.136 | 4.689 | 0.615- 35.737 |
| Left ventricular ejection fraction | 0.165 | 0.956 | 0.897- 1.019 |
| Platelet | 0.066 | 1.005 | 1.000- 1.011 |
| Cardiopulmonary bypass time | 0.330 | 0.988 | 0.963- 1.013 |
| Number of anastomoses | 0.090 | 2.471 | 0.869- 7.027 |

¹ Fischer's exact test

4
5
6
7
8
9

9. Figures

Figure 1. Sensitivity and specificity of SYNTAX Score at Post CABG atrial fibrillation diagnosis.



1