

1 A pilot investigation of enteric pathogens and stunting and
2 malnutrition using a combination of hospital surveillance and
3 a birth cohort of children in Dili, Timor-Leste.

4
5 Short title: Enteric pathogens and malnutrition in children from
6 Timor-Leste.

7
8 Danielle M. Cribb^{1*}, Nevio Sarmiento^{2,3}, Almerio Moniz¹, Nicholas S. S. Fancourt², Kathryn Glass¹,
9 Anthony D. K. Draper^{1,2,4}, Joshua R. Francis², Milena M. Lay dos Santos³, Endang Soares da Silva³,
10 Benjamin G. Polkinghorne¹, Virginia de Lourdes da Conceição^{2,3}, Feliciano da Conceição⁵, Paulino da
11 Silva², Joanita Jong⁵, Martyn D. Kirk¹, Samantha Colquhoun¹.

12

- 13 1. National Centre for Epidemiology and Population Health, Australian National University,
14 Canberra, Australia
15 2. Global and Tropical Health Division, Menzies School of Health Research, Charles Darwin
16 University, Darwin, Australia
17 3. Timor-Leste Ministry of Health, Dili, Timor-Leste
18 4. Northern Territory Centre for Disease Control, Northern Territory Government Department
19 of Health, Darwin, Australia
20 5. Ministry of Agriculture and Fisheries, Dili, Timor-Leste

21

22 * Corresponding author

23 Email: Danielle.Cribb@anu.edu.au (DC)

24

25

26 ABSTRACT

27 Background

28 Enteric pathogens contribute to child malnutrition in low-to-middle-income countries. In Timor-Leste,
29 there has been limited study of this relationship.

30 Methodology/Principal findings

31 We investigated enteric disease, stunting, and malnutrition in children in Dili, Timor-Leste (July 2019
32 – October 2020). Sixty infants received up to four home visits. We collected faecal samples,
33 demographics, anthropometrics, food and water sources, feeding practices, and animal husbandry
34 details. For 160 children admitted to Hospital Nacional Guido Valadares with clinical diagnosis of
35 severe diarrhoea or severe acute malnutrition (SAM), we collected faecal samples, diagnostics, and
36 anthropometrics. Faeces were tested by polymerase chain reaction (PCR) using the BioFire®
37 FilmArray® Gastrointestinal Panel. Descriptive analyses and regression analyses were conducted using
38 R.

39 We detected enteric pathogens in 68.8% (95% confidence interval [CI] 60.4–76.2%) of infants, 88.6%
40 of children with SAM (95% CI 81.7–93.3%), and 93.8% of children with severe diarrhoea (95% CI 67.7–
41 99.7%). Diarrhoeagenic *Escherichia coli* and *Campylobacter* spp. were most frequently detected. A
42 median weight-for-height Z-score (WHZ) of -1.1 (interquartile range [IQR] -2.0 to -0.1) for infants
43 across all home visits, -3.9 (IQR -4.7 to -3.0) for children with SAM and -2.6 (IQR -4.2 to -1.1) for children
44 with severe diarrhoea was observed.

45 The most common admission diagnosis was SAM (88.1%, 95% CI 81.7–92.5%). Hospitalised children
46 had increased odds of *Shigella*/enteroinvasive *E. coli* (OR 11.4, 95% CI 2.8–47.3) and *Giardia lamblia*
47 detection (OR 31.7, 95% CI 3.6–280) compared with infants. For home visits, bottle feeding was
48 associated with increased odds of pathogen detection (OR 8.2, 95% CI 1.1–59.7).

49 **Conclusions/Significance**

50 We detected high prevalence of enteric pathogens and signs of malnutrition in children in Dili. Our
51 pilot is proof of concept for a study to fully explore the risk factors and associations between enteric
52 pathogens and malnutrition in Timor-Leste.

53 Keywords: enteric pathogens, Timor-Leste, children, malnutrition, stunting, *Campylobacter*

54

55 **Author summary**

56 Enteric pathogens are a contributing factor to child malnutrition in low-to-middle-income countries.
57 This study investigated the relationship between enteric disease, malnutrition, and stunting in
58 children in Dili, Timor-Leste. Sixty infants received up to four home visits and 160 children were
59 admitted to the hospital with a clinical diagnosis of severe diarrhoea or severe acute malnutrition.
60 Faecal samples were collected and tested for enteric pathogens and risk factors were explored. The
61 study found that enteric pathogens were detected in most infants and children with malnutrition and
62 diarrhea. The most commonly detected pathogens were diarrheagenic *Escherichia coli* and
63 *Campylobacter* spp. Most children exhibited signs of malnutrition from infancy, and hospitalised
64 children had a higher likelihood of having certain enteric pathogens. Feeding practices, water use, and
65 animal cohabitation may pose risks for enteric pathogen infection. The results suggest that enteric
66 pathogens are a significant problem in Dili, Timor-Leste, and further research is needed to explore the
67 complexity of risk factors.

68 Introduction

69 Childhood undernutrition is a major challenge for many countries and is associated with high levels of
70 morbidity and mortality (1). In 2020, 22.0% of children aged under five years worldwide suffered from
71 stunting (low height for age) and 6.7% suffered from wasting (low weight for height) (2). Children in
72 low- and middle-income countries (LMICs) are disproportionately impacted by stunting (38.8%) and
73 wasting (12.9%) (3). Undernutrition can lead to cognitive, physical, and metabolic developmental
74 issues, potentially reducing physical and intellectual ability and economic productivity in adulthood
75 (1).

76 Socioeconomic and environmental factors contribute to child undernutrition, including poor maternal
77 health and education, early termination of breastfeeding, low household wealth, and unimproved
78 sanitation (3, 4). There is growing evidence for an association between infection with enteric bacteria,
79 viruses, and parasites and child stunting and wasting, particularly in LMICs (5, 6). Enteric bacterial
80 pathogens contracted early in life may damage the lining of the small and large intestines, reducing
81 the absorption of nutrients from food and promoting chronic inflammation, leading to undernutrition
82 and a compromised immune system (5, 6). Such enteric pathogens potentially involved in stunting and
83 malnutrition include enterotoxigenic *Escherichia coli* (ETEC), *Shigella*, and *Campylobacter* species (7-
84 9).

85 *Campylobacter* is the most common cause of foodborne bacterial gastroenteritis globally, causing an
86 estimated 96 million cases and >21,000 deaths annually (10). In high-income settings, infections are
87 often sporadic in nature and largely attributed to consuming undercooked chicken (11). In low-income
88 settings, *Campylobacter* is frequently endemic and while still largely attributed to poultry, infection
89 determinants are from more varied environmental and zoonotic sources (11, 12). Recent studies have
90 further implicated *Campylobacter* in child stunting and malnutrition. A Peruvian cohort exhibited poor
91 weight gain with infection and marginal poor linear growth with symptomatic infection (7, 9).
92 Similarly, 84.9% of the multisite Malnutrition and Enteric Disease Study (MAL-ED) cohort presented

93 with a *Campylobacter*-positive faecal sample by one year of age, with those diagnosed with multiple
94 *Campylobacter* infections being more likely to exhibit poor linear growth (12).

95 Timor-Leste is one of the lowest-income countries in Asia and has a population of 1.34 million circa
96 2022 (13, 14). The 2016 Demographic and Health Survey (DHS) estimated that the proportion of
97 children with stunting was 46.5%, wasting 24.2% and underweight (low weight for age) 40.4% (15).
98 Diarrhoeal disease is prevalent in Timor-Leste with 10.7% of children represented in the DHS
99 presenting with diarrhoea within the two weeks preceding the survey, peaking at 17.8% in children
100 aged 12-23 months (16).

101 We conducted a pilot investigation to examine the feasibility of a birth cohort and hospital-based
102 surveillance study in Timor-Leste. We aimed to i) estimate the proportion of children from either a
103 birth cohort or hospital admissions with severe acute malnutrition (SAM) and/or severe diarrhoea
104 who tested positive for enteric pathogens on stool, ii) estimate the proportion of children from a birth
105 cohort who were malnourished or stunted at any time during the first 12 months of life, iii) quantify
106 malnourishment in children admitted to hospital with SAM and/or severe diarrhoea, iv) describe the
107 social, environmental, food safety, and food security issues associated with enteric pathogen
108 infections in Dili, Timor-Leste, and v) support capacity building in research and laboratory skills. This
109 investigation formed the basis of a larger study to understand the impact of enteric pathogens on
110 stunting and malnutrition.

111

112 **Methods**

113 This study was carried out at the national referral hospital in Dili, Timor-Leste, Hospital Nacional Guido
114 Valadares (HNGV). HNGV has a catchment population of 500,000 people with approximately 4,500
115 births per year.

116 **Birth cohort**

117 **Study setting and recruitment**

118 We recruited newborns into the longitudinal birth cohort between July and September 2019.
119 Newborn infants were eligible for inclusion if they were born at ≥ 38 weeks gestation at HNGV and if
120 they were reported to live within a 10-kilometre radius of HNGV. The local research team approached
121 parents in the maternity unit at HNGV after delivery and prior to discharge. Parents provided informed
122 consent to enrol their infants. Home visits occurred when infants were approximately one, four, seven,
123 and 12 months of age.

124 **Equipment and questionnaire**

125 The local study team were trained to use a calibrated portable baby scale and a standardised mat for
126 measuring weight (Seca 334 Baby Scales) and height (Seca 210 Mobile Measuring Mat for Babies and
127 Toddlers). All infants were weighed and measured using a standardised procedure. In addition, the
128 team interviewed parents using a standardised questionnaire at each visit to investigate feeding
129 practices, household size, water, sanitation, and hygiene (WASH) practices and accessibility, and
130 household animal ownership and containment (16). All data were collected on an android tablet using
131 Research Electronic Data Capture (REDCap) (17, 18), synchronised online, and stored on a secure
132 server hosted at the Australian National University (ANU).

133 **Specimen collection and testing**

134 Faecal samples were collected in sample pots from all infants at the time of the field visit or in the
135 following days and transported to the lab in an insulated United Nations Children’s Fund container.
136 Parents and/or guardians were provided with instructions in their native language, Tetum to ensure
137 samples were correctly collected and stored prior to transport. Samples were transported to the
138 National Health Laboratory (NHL) for polymerase chain reaction (PCR) testing and culture for
139 *Campylobacter*, *Shigella* or *Salmonella* isolates. Animal faecal samples were collected from the
140 environment proximal to homes where a positive *Campylobacter*, *Shigella* or *Salmonella* infant faecal
141 sample was detected. Samples were transported to the Timor-Leste Diagnostics Laboratory for Animal
142 Health (DLAH) for preparation and isolate culture on appropriate selective media.

143 **Hospital-based surveillance**

144 **Study setting and recruitment**

145 We recruited children admitted to HNGV with SAM, diarrhoea with severe dehydration (DWSD), or
146 dysentery between October 2019 to July 2020 to investigate enteric pathogens in children with these
147 conditions in Dili. Children were eligible for inclusion if they were aged less than five years at time of
148 hospital admission, if they resided within Dili, and if they met the adapted World Health Organization
149 (WHO) criteria for diagnosis of SAM, DWSD and/or severe dysentery (19).

150 **Specimen and information collection**

151 Participants were identified for eligibility by paediatricians during the daily paediatric ward round.
152 Following informed consent from parents or guardians, ward nurses collected faecal samples from
153 eligible children and the study team recorded vital statistics and demographic characteristics from the
154 medical notes and transported the samples to the NHL for PCR testing.

155 **Pathogen detection**

156 The NHL used a BioFire® FilmArray® Gastrointestinal Panel PCR assay to test all faecal samples from
157 birth cohort and hospitalised children to detect 22 enteric pathogens and parasites (20). Samples

158 positive by PCR for selected enteric bacteria (*Campylobacter*, *Salmonella* and *Shigella*/EIEC) were
159 reflex cultured on appropriate selective media.

160 **Data analysis**

161 All data were cleaned and analysed using R statistical software (21). We converted anthropometric
162 measures to weight-for-height (WHZ), height-for-age (HAZ), and weight-for-age (WAZ) Z scores and
163 mapped these against the WHO child growth standards (22, 23) using R package “zscorer” (24). We
164 performed Fisher’s exact and Chi-square tests to analyse differences between groups and visits. We
165 calculated odds ratios (OR) using generalised estimating equations (GEE) to account for repeated
166 observations with the R package “geepack”, where appropriate (25). GEE were used to analyse risk
167 factors in the birth cohort against the outcomes of i) pathogen detection and ii) moderate wasting,
168 defined as a WHZ of -2 or below (26). Additionally, GEE were used to analyse associations between
169 hospitalised SAM cases and non-hospitalised children for pathogen detections, demographics and
170 faecal consistency. GEE models used an exchangeable correlation structure and binomial distribution
171 with logistic link. We used the “arsenal” package to calculate generalised linear models, where
172 appropriate (27). A p value of $p < 0.05$ was considered significant for Fisher’s and Chi-square tests, and
173 an OR with confidence intervals (CI) that did not cross over one was considered significant for GEE.

174 **Ethics**

175 The Timor-Leste National Institute of Health-Research Ethics & Technical Committee (reference
176 462/MS-INS/GDE/IV/2019) for the Timor-Leste Instituto Nacional de Saúde (INS), and the Australian
177 National University Ethics Committee (reference 2019/221) provided ethical approval. The field study
178 team were trained in Good Clinical Practice for research by the project lead (28). All parents or legal
179 guardians provided informed consent prior to study enrolment.

180 Results

181 Birth cohort

182 We recruited 32 male and 28 female infants into the birth cohort. Only 53.3% (32/60) were retained
183 until the final home visit. At birth, the median WHZ was 0.9 (interquartile range [IQR] -0.5 – 1.6), with
184 no scores calculated for four newborns with lengths below 45cm (S1 Table). The median WHZ over all
185 four home visits decreased to -1.1 (IQR -2.0 to -0.1). Of the 32 infants who were weighed and
186 measured at all four visits, 31.3% (10/32) exhibited poor weight gain (WHZ of ≤ -3.0) on at least one
187 visit. While the birth cohort tracked the WHO 50th percentile for height-for-age, weight-for-age and
188 weight-for-height tracked between the 15th and 25th percentiles (**Error! Reference source not found.**).

189 **Fig 1. (a) Height-for-age World Health Organization standards compared with Timor-Leste birth cohort,**
190 **2019–2020; (b) Weight-for-age WHO standards compared with Timor-Leste birth cohort, 2019–2020; (c)**
191 **Weight-for-height WHO standards compared with Timor-Leste birth cohort, 2019–2020.** WHO percentiles are
192 annotated on figure. 50th percentile for WHO standards is presented in bold. Grey shading represents the 95%
193 confidence interval for the Timor-Leste birth cohort.

194 We collected 141 infant faecal specimens for PCR testing from 161 home visits (87.6% sample
195 collection rate), with at least one faecal sample collected from 45/60 (75.0%) of participants. We
196 detected at least one enteric pathogen in 68.8% (97/141; 95%CI 60.4%–76.2%) of samples, and of
197 these 63.9% (62/97; 95% CI 53.5%–73.2%) were positive for multiple pathogens.

198 We observed significant differences in pathogen occurrence between home visits, with the number of
199 pathogens detected within the cohort and the number of multiple pathogen detections in individual infants
200 generally increasing with age ($p < 0.05$;

201

202

203 **Table 1).** Twenty-two study participants had samples PCR tested at each visit, with each having at least
204 one positive result for an enteric pathogen and 40.9% (9/22) positive at every visit. There was a weak
205 relationship between WHZ and the number of pathogens detected per sample at each visit (**Fig 2**).

206

207

208

209 **Table 1: Number and proportion of pathogens detected by multiplex PCR at each birth cohort home visit and**
210 **stool presentation of samples, Dili, Timor-Leste, 2019–2020.**

Pathogen	Visit 1 n (%)	Visit 2 n (%)	Visit 3 n (%)	Visit 4 n (%)	Total n (%)
Bacteria					
<i>Campylobacter</i> spp.*	3 (6.7)	1 (2.9)	3 (8.3)	8 (30.8)	15 (10.6)
<i>Clostridioides difficile</i>	1 (2.2)	4 (11.8)	4 (11.1)	4 (15.4)	13 (9.2)
<i>Salmonella</i>	0 (0)	0 (0)	4 (11.1)	0 (0)	4 (2.8)
Shiga toxin-producing <i>Escherichia coli</i> (STEC)	0 (0)	0 (0)	0 (0)	2 (7.7)	2 (1.4)
Enteroaggregative <i>E. coli</i>	11 (24.4)	21 (61.8)	30 (83.3)	15 (57.7)	77 (54.6)
Enteropathogenic <i>E. coli</i>	3 (6.7)	10 (29.4)	21 (58.3)	15 (57.7)	49 (34.8)
Enterotoxigenic <i>E. coli</i> lt/st	1 (2.2)	4 (11.8)	12 (33.3)	5 (19.2)	22 (15.6)
<i>Shigella</i> /Enteroinvasive <i>E. coli</i>	1 (2.2)	0 (0)	1 (2.8)	0 (0)	2 (1.4)
<i>Plesiomonas shigelloides</i>	0 (0)	0 (0)	1 (2.8)	0 (0)	1 (0.7)
<i>Vibrio</i> spp. †	1 (2.2)	0 (0)	4 (11.1)	0 (0)	5 (3.5)
Viruses					
Adenovirus F40/41	0 (0)	0 (0)	2 (5.6)	4 (15.4)	6 (4.3)
Norovirus GI/GII	0 (0)	3 (8.8)	7 (19.4)	2 (7.7)	12 (8.5)
Rotavirus A	0 (0)	3 (8.8)	2 (5.6)	4 (15.4)	9 (6.4)
Sapovirus	0 (0)	0 (0)	4 (11.1)	0 (0)	4 (2.8)
Parasites					
<i>Cryptosporidium</i> spp.	0 (0)	0 (0)	0 (0)	3 (11.5)	3 (2.1)
<i>Giardia lamblia</i>	0 (0)	0 (0)	1 (2.8)	0 (0)	1 (0.7)
Total samples	45	34	36	26	141
* <i>Campylobacter</i> spp. includes <i>C. jejuni</i> , <i>C. coli</i> , and <i>C. upsaliensis</i> . † <i>Vibrio</i> spp. includes <i>V. parahaemolyticus</i> , <i>V. vulnificus</i> , and <i>V. cholerae</i> . N.B. percentages and counts do not equal total samples due to multiple pathogen detections in some samples.					

211

212 **Fig 2. Weight-for-height Z-score by number of pathogens detected for each birth cohort visit, Dili, Timor-**
213 **Leste, 2019–2020.**

214 Diarrhoeagenic *E. coli* (DEC) strains (Shiga toxin-producing *E. coli* [STEC], EPEC, EAEC, and ETEC) were
215 the most common pathogens detected. Six infants were PCR positive for DEC at all four visits, with
216 EAEC detected in 96.0% (23/24) of their faecal samples, indicating possible chronic carriage. Fifteen
217 samples were PCR positive for *Campylobacter* spp., four samples of *Salmonella*, and two samples of
218 *Shigella*/enteroinvasive *E. coli* (EIEC) across the entire cohort. During the first home visits, three
219 samples collected from animal faeces were successfully cultured and *Campylobacter* spp. isolated by
220 the DLAH team. No isolates were cultured from human clinical samples.

221 Full descriptive results and ORs for these analyses are available in S2 Table and S3 Table. Compared to
222 children aged <3 months, we found that those aged 3–<6 months (OR 5.4, 95% CI 2.1–14.3), 6–<9
223 months (OR 18.1, 95% CI 4.8–68.8), and ≥12-months (OR 12.1, 95% CI 3.1–47.0) had higher odds of
224 pathogen detection, i.e., the odds of pathogen detection increased with age. We controlled for age in
225 analysis of all additional risk factors. We found that exclusive bottle feeding was associated with
226 increased odds of enteric pathogen detection (OR 8.2, 95% CI 1.1–59.7) compared with breastfeeding.
227 No other factors were found to be statistically significant.

228 For the outcome of wasting, controlled for age, we found that households with five or fewer members
229 were associated with lower odds of wasting (OR 0.2, 95% CI 0.1–1.0) compared to those with 6–10
230 household members. No other factors were significant.

231 **Hospital-based surveillance**

232 We recruited 160 children in hospital-based surveillance. One child was excluded as they were aged
233 above five years. We included 159 children in descriptive analyses including those that were missing
234 anthropometric data including gender (n=2), middle-upper arm circumference (n=5), and height (n=1).
235 Females represented 52.9% (83/157) of participants with known gender. We received PCR results
236 from 148 participants as two patients were discharged before specimen collection and nine were
237 missing laboratory PCR data.

238 The mean age at admission was 17.7 months (range 1.0–55.7 months, standard deviation (SD) ±11.4).
239 We diagnosed 140 (88.1%, 95% CI 81.7–92.5%) children with SAM and 19 (11.9%, 95% CI 7.5–18.3%)
240 with severe diarrhoeal disease (13 with DWSD, and six with dysentery). The median WHZ for those
241 diagnosed with SAM (-3.9, IQR -4.7 to -3.0) was lower than those with severe diarrhoea (-2.6, IQR -4.2
242 to -1.1). Six WHZ were not calculated for SAM participants due to unknown gender (n=2), unknown
243 height (n=1) or a height below 45cm (n=3).

244 Of the 148 participants with complete multiplex PCR results, 88.6% (117/132; 95% CI 81.7–93.3%) of
245 SAM participants were positive for at least one enteric pathogen, with 73.5% (97/132; 95% CI 65.0–
246 80.6%) positive for multiple pathogens. Of those diagnosed with severe diarrhoea, 93.8% (15/16; 95%
247 CI 67.7–99.7%) were positive for at least one enteric pathogen and all but one were positive for
248 multiple pathogens (14/16; 87.5%, 95% CI 60.4–97.8%). We detected EAEC and/or EPEC in 81.8%
249 (9/11) of loose faecal samples. The age at admission for participants varied between the type of
250 pathogen detected, with viral pathogens detected in admissions with a mean age of 13.8 months (SD
251 ± 8.1), bacteria at 17.8 months ($SD \pm 11.2$), and parasites at 19.3 months ($SD \pm 11.7$). There were no
252 statistically significant differences for age at admission by pathogen group between SAM and severe
253 diarrhoea cases ($p > 0.05$) (S4 Table).

254 The mean number of pathogens detected per individual was 2.8 (range 0–7). The most detected
255 pathogens were DEC species followed by *Campylobacter* spp., *Shigella*/EIEC, and *Giardia lamblia* (

256 Table 2). Two *Campylobacter* spp. and two *Salmonella* isolates were cultured.

257

258 **Table 2: Hospital-based surveillance data of diagnosis-specific pathogen detections and total pathogen**
 259 **detections from PCR testing, Dili, Timor-Leste, 2019-2020.**

Pathogen	Pathogen detections in SAM cases n (%)	Pathogen detections in severe diarrhoea [‡] cases n (%)	All pathogen detections n (%)
Bacteria			
<i>Campylobacter</i> spp.*	33 (25.0)	3 (18.8)	36 (24.3)
<i>Clostridioides difficile</i>	6 (4.5)	0 (0.0)	6 (4.1)
<i>Plesiomonas shigelloides</i>	1 (0.8)	0 (0.0)	1 (0.7)
<i>Salmonella</i>	4 (3.0)	1 (6.3)	5 (3.4)
Shiga toxin-producing <i>Escherichia coli</i> (STEC)	4 (3.0)	0 (0.0)	4 (2.7)
Enteroaggregative <i>E. coli</i>	84 (63.6)	10 (62.5)	94 (63.5)
Enteropathogenic <i>E. coli</i>	70 (53.0)	10 (62.5)	80 (54.1)
Enterotoxigenic <i>E. coli</i> lt/st	41 (31.1)	4 (25.0)	45 (30.4)
<i>E. coli</i> O157	2 (1.5)	0 (0.0)	2 (1.4)
<i>Shigella</i> /Enteroinvasive <i>E. coli</i>	23 (17.4)	7 (43.8)	30 (20.3)
<i>Vibrio</i> spp. [†]	5 (3.8)	3 (18.8)	8 (5.4)
Viruses			
Adenovirus F40/41	3 (2.3)	1 (6.3)	4 (2.7)
Astrovirus	6 (4.5)	1 (6.3)	7 (4.7)
Norovirus GI/GII	12 (9.1)	1 (6.3)	13 (8.8)
Rotavirus A	6 (4.5)	1 (6.3)	7 (4.7)
Sapovirus	13 (9.8)	1 (6.3)	14 (9.5)
Parasites			
<i>Cryptosporidium</i> spp.	17 (12.9)	1 (6.3)	18 (12.2)
<i>Giardia lamblia</i>	31 (23.5)	2 (12.5)	33 (22.3)
<i>Cyclospora cayetanensis</i>	12 (9.1)	0 (0.0)	12 (8.1)
<i>Entamoeba histolytica</i>	0 (0.0)	1 (6.3)	1 (0.7)
Total samples	132	16	148
* <i>Campylobacter</i> spp. includes <i>C. jejuni</i> , <i>C. coli</i> , and <i>C. upsaliensis</i> . † <i>Vibrio</i> spp. includes <i>V. parahaemolyticus</i> , <i>V. vulnificus</i> , and <i>V. cholerae</i> . ‡Severe diarrhoea includes diarrhoea with severe dehydration and dysentery diagnoses. PCR = polymerase chain reaction test. SAM = severe acute malnutrition. N.B. percentages and counts do not equal total samples due to multiple pathogen detections in some samples.			

260

261 Comparing hospitalised and non-hospitalised children

262 Hospitalised children had increased odds of both *Shigella*/EIEC (OR 11.4, 95% CI 2.8–47.3) and *Giardia*
 263 *lamblia* infection (OR 31.7, 95% CI 3.6–280) compared with the non-hospitalised birth cohort (S5
 264 Table).

265 Discussion

266 Infants and young children from a birth cohort and hospital-based surveillance study exhibited high
267 enteric pathogen loads and signs of wasting and undernutrition from the first months of life. Ninety
268 percent of children admitted to hospital for treatment of DWSD, dysentery, or SAM were positive for
269 at least one enteric pathogen and 70% of infants visited at home also returned a positive PCR test for
270 at least one enteric pathogen over the 12-month study. A high proportion of samples across both
271 study arms had multiple pathogens recovered. Pathogen detection generally increased with age
272 within the birth cohort, likely influenced by exposure to new nutrition sources and environmental
273 factors. While the mechanisms of child malnutrition remain complex, our pilot study has
274 demonstrated the ability to conduct longitudinal research in Timor-Leste to explore and understand
275 the relationship between enteric pathogens and malnutrition.

276 We detected a variety of enteric pathogens in children regardless of the presence of wasting and
277 diarrhoeal disease. Across symptomatic and asymptomatic faecal samples, *Campylobacter* spp., DEC
278 strains EAEC, EPEC, and ETEC, *Shigella*/EIEC, norovirus, *Giardia*, and *Cryptosporidium* were most
279 prevalent. This is consistent with results from the MAL-ED study (29) and the Global Enteric
280 Multicenter Study (GEMS)(30). Diarrhoeal faecal samples in our study frequently had DEC strains
281 present, suggesting these may be driving disease presentation in combination with non-pathogenic
282 factors. Pathogenic *E. coli* is commonly identified as a cause of both acute and persistent paediatric
283 diarrhoea in low- and middle-income countries (31-34), particularly in malnourished children (35, 36).
284 Further, GEMS found that diarrhoea from typical EPEC and heat stable ETEC infections were associated
285 with malnutrition and an increased risk of death in infants (34). There is evidence of chronic carriage
286 or reinfection with EAEC in the birth cohort. While persistent EAEC infections are known to be
287 implicated in long-term diarrhoeal disease in children, this was not evident in our study (37-39).
288 Children diagnosed with DWSD or dysentery in hospital-based surveillance had a high prevalence of
289 DEC, *Campylobacter* spp. and/or *Shigella*, pathogens previously implicated with these conditions (40).

290 Despite high prevalence of certain pathogens, we did not detect any significant relationships between
291 individual pathogens and wasting or diarrhoeal disease presentation. This may indicate that these
292 outcomes are due to multiple pathogen interactions or additional nutritional or environmental
293 factors. Pathogen virulence likely also plays an important part in the distribution of pathogens
294 between malnourished and/or diarrhoeal children versus those that are relatively healthy (41).

295 Most pathogens were recovered from faecal samples with limited or no evidence of diarrhoeal
296 disease. While diarrhoeal disease is implicated in child malnutrition, subclinical infections with enteric
297 pathogens may be more strongly linked to poor growth outcomes (12). The MAL-ED study detected
298 high prevalence of *Campylobacter* from an early age and a negative association between high
299 pathogen load and growth, particularly in asymptomatic cases (12). Carriage of enteric pathogens like
300 EAEC and *Campylobacter* can lead to enteric and systemic inflammation and malabsorption of
301 nutrients resulting in malnutrition and faltered growth, particularly in prolonged cases or repeat
302 infections (42-44). Detection of a wide range of pathogens in the absence of diarrhoeal symptoms
303 warrants further investigation to understand the relationship between enteric pathogens and
304 malnutrition (29).

305 Within Timor-Leste and neighbouring Southeast Asian countries, there is limited evidence on the
306 prevalence of enteric pathogens both in clinical and environmental settings. A multisite study of
307 children aged under five years conducted in Indonesia between 2009 and 2012 detected a high
308 prevalence of rotavirus (37.4%), adenovirus (34.9%) and ETEC (10.0%) (45). Another Indonesian case-
309 control study of diarrhoeal disease conducted between 1997 and 1999 detected very low levels of
310 enteric pathogens in paediatric patients using culture-based methods (i.e., less than 8.0% prevalence
311 for total pathogens)(46), however, it is worth noting that culture-based methods can be lower in
312 sensitivity than PCR-based methods. Our present study aimed to explore environmental and zoonotic
313 pathogen sources in addition to risk factors in the home. Flooding is common in the region, particularly
314 Timor-Leste. This combined with ubiquitous pit latrines and improper garbage disposal provides ample

315 opportunities for transmission of enteric pathogens in the environment. This is supported by a
316 Nigerian study that tested groundwater sources for enteric pathogens. High levels of bacteria
317 including *E. coli*, *Salmonella*, *Shigella*, and *Campylobacter* spp. were found in boreholes that likely
318 weren't adequately maintained and had contamination from nearby septic tanks, sludge systems, and
319 waste dumpsites (47). An Egyptian study also isolated these bacteria from ground water in rural areas
320 (48). While efforts to improve personal practice with water, sanitation, and hygiene measures, food
321 sources and preparation, and animal husbandry are important, exploring the endemicity of enteric
322 pathogens in the environment should also be explored.

323 There are several limitations of our study. This study was conducted to examine acceptability for
324 undertaking a larger birth cohort study. It was under-powered to examine some important
325 relationships. Timor-Leste is prone to flash-flooding making it difficult to undertake home visits and,
326 on a few occasions, HNGV and NHL experienced flash-flooding. We had reasonably high attrition of
327 our birth cohort, due to highly mobile families especially during the COVID-19 pandemic when the
328 study was interrupted between April and July 2020, leading to substantial loss to follow up. Limited
329 animal faecal samples were able to be collected due an outbreak in African Swine fever in the first six
330 months of the study. In addition, multiplex PCR testing is very sensitive and detects presence of
331 pathogen nucleic acid but not necessarily viable pathogen. The Biofire GI PCR panel has shown high
332 sensitivity and specificity in detection of diarrhoeal pathogens (49). We intended to culture all human
333 clinical *Salmonella*, *Campylobacter* and *Shigella* samples at NHL, however due to a lack of available
334 resources this was not completed in most cases. In the absence of culture results to compare to the
335 positive PCR results for *Campylobacter*, *Shigella* and *Salmonella* we were unable to determine the
336 validity of the PCR results in this setting. However, similar studies have shown that multiplex PCR
337 testing can provide an accurate description of enteric pathogen prevalence in low- and middle-income
338 countries (50, 51).

339 In this study, we demonstrated the possibility of undertaking longitudinal field research in Timor-Leste
340 and produced useful and practical information that will be used to assist in guiding the strategy and
341 approach of future research. Nutrition awareness, choices and availability, food and water safety and
342 exposure to animal and environmental pathogens affect lives of Timor-Leste families and contribute
343 to the multiple malnutrition risk factors. We recommend conducting a larger-scale study to investigate
344 infant and child dietary practices, food safety, WASH, gender, cultural and environmental hygiene
345 conditions in relation to One Health and food safety and security, with a focus on the risk of enteric
346 pathogens. This project should comprehensively investigate malnutrition and stunting risk factors,
347 including zoonotic transmission of enteric pathogens to provide an evidence base for targeted
348 interventions in urban and rural settings in Timor-Leste, utilising a One Health collaborative approach
349 to enhance collaboration and capacity.

350 **Acknowledgements**

351 The authors thank all persons and organisations that provided support for this study. These include:
352 Professor Néilson Martins and the National Institute of Health-Research Ethics & Technical Committee;
353 Dr Dongbao Yu and Dr Rajesh Pandav at the World Health Organization; Dr Merita Monteiro at
354 Departamentu Kontrolu Moras Hada'et (CDC); Maria Angela Varela Niha at Departamentu Vijilansia
355 Epidemiolojia, Ministério da Saúde; Dr Jeremy Beckett at Maluk Timor; local research team members
356 Francisca Soares and Antoninho Gusmão; Menzies staff including Dr Jennifer Yan, Salvador Amaral, Dr
357 Ian Marr, Lucsendar Alves, Dr Jo Wapling, Karen Champlin, and Dr Shawn Ting; Ismael Barreto and all
358 staff from the National Health Laboratory; all staff from the Diagnostics Laboratory for Animal Health;
359 all staff at HNGV including Dr Flavio Brandão de Araujo, Dr Ingrid Bucens, Dr Amelia Ferreria Pinto and
360 the HNGV Paediatrics team; Ms Melanie McVean and the St John of God Midwifery team; the HNGV
361 Obstetrics and Midwifery team, and; Dr Anna Okello, Dr Francette Geraghty-Dusan, and Bethany Lees
362 from the Australian Centre for International Agricultural Research.

363

364 References

- 365 1. Kinyoki DK, Osgood-Zimmerman AE, Pickering BV, Schaeffer LE, Marczak LB, Lazzar-Atwood A,
366 et al. Mapping child growth failure across low- and middle-income countries. *Nature*.
367 2020;577(7789):231-4. doi: 10.1038/s41586-019-1878-8
- 368 2. United Nations. The Sustainable Development Goals Report 2022. 2022.
- 369 3. Li Z, Kim R, Vollmer S, Subramanian SV. Factors Associated With Child Stunting, Wasting, and
370 Underweight in 35 Low- and Middle-Income Countries. *JAMA Netw Open*. 2020;3(4):e203386. doi:
371 10.1001/jamanetworkopen.2020.3386
- 372 4. Mizumoto K, Takahashi T, Kinoshita Y, Higuchi M, Bachroen C, Da Silva V. A qualitative study
373 of risk factors related to child malnutrition in Aileu District, Timor-Leste. *Asia Pac J Public Health*.
374 2015;27(2):NP1398-408. doi: 10.1177/1010539513486175
- 375 5. MAL-ED Network Investigators. The MAL-ED Study: A Multinational and Multidisciplinary
376 Approach to Understand the Relationship Between Enteric Pathogens, Malnutrition, Gut Physiology,
377 Physical Growth, Cognitive Development, and Immune Responses in Infants and Children Up to 2 Years
378 of Age in Resource-Poor Environments. *Clin Infect Dis*. 2014;59(suppl_4):S193-S206. doi:
379 10.1093/cid/ciu653
- 380 6. Petri WA, Jr., Miller M, Binder HJ, Levine MM, Dillingham R, Guerrant RL. Enteric infections,
381 diarrhea, and their impact on function and development. *J Clin Invest*. 2008;118(4):1277-90. doi:
382 10.1172/JCI34005
- 383 7. Lee G, Olortegui MP, Yori PP, Black RE, Caulfield L, Chavez CB, et al. Effects of *Shigella*-,
384 *Campylobacter*-and ETEC-associated diarrhea on childhood growth. *Pediatr Infect Dis J*.
385 2014;33(10):1004-9. doi: 10.1097/INF.0000000000000351
- 386 8. Black RE, Brown KH, Becker S. Effects of diarrhea associated with specific enteropathogens on
387 the growth of children in rural Bangladesh. *Pediatrics*. 1984;73(6):799-805. doi:
- 388 9. Lee G, Pan W, Peñataro Yori P, Paredes Olortegui M, Tilley D, Gregory M, et al. Symptomatic
389 and Asymptomatic *Campylobacter* Infections Associated with Reduced Growth in Peruvian Children.
390 *PLoS Negl Trop Dis*. 2013;7(1):e2036. doi: 10.1371/journal.pntd.0002036
- 391 10. World Health Organization. WHO estimates of the global burden of foodborne diseases:
392 foodborne disease burden epidemiology reference group 2007-2015. Geneva, Switzerland; 2015.
- 393 11. Kaakoush NO, Castaño-Rodríguez N, Mitchell HM, Man SM. Global Epidemiology of
394 *Campylobacter* Infection. *Clin Microbiol Rev*. 2015;28(3):687-720. doi: 10.1128/CMR.00006-15

- 395 12. Amour C, Gratz J, Mduma E, Svensen E, Rogawski ET, McGrath M, et al. Epidemiology and
396 Impact of *Campylobacter* Infection in Children in 8 Low-Resource Settings: Results From the MAL-ED
397 Study. Clin Infect Dis. 2016;63(9):1171-9. doi: 10.1093/cid/ciw542
- 398 13. Provo A, Atwood S, Sullivan EB, Mbuya N. Malnutrition in Timor-Leste: A review of the burden,
399 drivers, and potential response World Bank; 2017.
- 400 14. UNFPA Timor-Leste. Timor-Leste Population and Housing Census 2022: Preliminary Results.
401 Government of Timor-Leste; 2022.
- 402 15. Hall A, Hyder Z, Karyadi E. Malnutrition in young children and their mothers in Timor-Leste.
403 Washington, D.C; 2020.
- 404 16. General Directorate of Statistics. Timor-Leste Demographic and Health Survey 2016. Dili,
405 Timor-Leste: GDS and ICF; 2018.
- 406 17. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture
407 (REDCap)—A metadata-driven methodology and workflow process for providing translational
408 research informatics support. J Biomed Inform. 2009;42(2):377-81. doi: 10.1016/j.jbi.2008.08.010
- 409 18. Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, et al. The REDCap consortium:
410 Building an international community of software platform partners. J Biomed Inform.
411 2019;95:103208. doi: 10.1016/j.jbi.2019.103208
- 412 19. World Health Organization. Management of severe malnutrition: a manual for physicians and
413 other senior health workers. Geneva; 1999.
- 414 20. Biomérieux. The BioFire® FilmArray® Gastrointestinal (GI) Panel 2022 [cited 2022 1
415 September]. Available from: <https://www.biofire.com/products/the-filmarray-panels/filmarraygi/>.
- 416 21. R Core Team. R: A language and environment for statistical computing Vienna, Austria 2022
417 [cited 2022 1 September]. Available from: <https://www.R-project.org/>.
- 418 22. World Health Organization. WHO child growth standards : length/height-for-age, weight-for-
419 age, weight-for-length, weight -for-height and body mass index-for-age : methods and development.
420 2006 [cited 2022 1 July]. Available from: <https://www.who.int/publications/i/item/924154693X>.
- 421 23. World Health Organization. WHO child growth standards : head circumference-for-age, arm
422 circumference-for-age, triceps skinfold-for-age and subscapular skinfold-for-age : methods and
423 development. 2007 [cited 2022 1 July]. Available from:
424 <https://www.who.int/publications/i/item/9789241547185>.
- 425 24. Myatt MaG, E. zscorer: Child Anthropometry z-Score Calculator 2019 [cited 2022 1
426 September]. Available from: <https://CRAN.R-project.org/package=zscorer>.
- 427 25. Højsgaard S, Halekoh U, Yan J. The R Package geepack for Generalized Estimating Equations. J
428 Stat Soft. 2005;15(2):1-11. doi: 10.18637/jss.v015.i02

- 429 26. Croft T, Marshall, A, Allen, C et al. Guide to DHS Statistics: DHS-7 (version 2): The Demographic
430 and Health Surveys Program. Rockville, Maryland, USA: ICF; 2020.
- 431 27. Heinzen E, Sinnwell, J, Atkinson, E, Gunderson, T, Dougherty, G, Votruba, P, Lennon, R,
432 Hanson, A, Goergen, K, Lundt, E, Broderick, B, McCullough, M. arsenal: An Arsenal of 'R' Functions for
433 Large-Scale Statistical Summaries 2021 [cited 2022 1 September]. Available from: [https://CRAN.R-](https://CRAN.R-project.org/package=arsenal)
434 [project.org/package=arsenal](https://CRAN.R-project.org/package=arsenal).
- 435 28. Global Health Training Centre. Research Ethics Online Training 2022 [cited 2022 1 December].
436 Available from: <https://globalhealthtrainingcentre.tghn.org/elearning/research-ethics/>.
- 437 29. Platts-Mills JA, Babji S, Bodhidatta L, Gratz J, Haque R, Havt A, et al. Pathogen-specific burdens
438 of community diarrhoea in developing countries: a multisite birth cohort study (MAL-ED). *Lancet Glob*
439 *Health*. 2015;3(9):e564-75. doi: 10.1016/s2214-109x(15)00151-5
- 440 30. Kotloff KL, Nasrin D, Blackwelder WC, Wu Y, Farag T, Panchalingham S, et al. The incidence,
441 aetiology, and adverse clinical consequences of less severe diarrhoeal episodes among infants and
442 children residing in low-income and middle-income countries: a 12-month case-control study as a
443 follow-on to the Global Enteric Multicenter Study (GEMS). *Lancet Glob Health*. 2019;7(5):e568-e84.
444 doi: 10.1016/S2214-109X(19)30076-2
- 445 31. Nataro JP, Kaper JB. Diarrheagenic *Escherichia coli*. *Clin Microbiol Rev*. 1998;11(1):142-201.
446 doi: 10.1128/CMR.11.1.142
- 447 32. Ellis SJ, Crossman LC, McGrath CJ, Chattaway MA, Holken JM, Brett B, et al. Identification and
448 characterisation of enteroaggregative *Escherichia coli* subtypes associated with human disease. *Sci*
449 *Rep*. 2020;10(1):7475. doi: 10.1038/s41598-020-64424-3
- 450 33. Gomes TA, Elias WP, Scaletsky IC, Guth BE, Rodrigues JF, Piazza RM, et al. Diarrheagenic
451 *Escherichia coli*. *Braz J Microbiol*. 2016;47(Suppl 1):3-30. doi: 10.1016/j.bjm.2016.10.015
- 452 34. Kotloff KL, Nataro JP, Blackwelder WC, Nasrin D, Farag TH, Panchalingam S, et al. Burden and
453 aetiology of diarrhoeal disease in infants and young children in developing countries (the Global
454 Enteric Multicenter Study, GEMS): a prospective, case-control study. *Lancet*. 2013;382(9888):209-22.
455 doi: 10.1016/S0140-6736(13)60844-2
- 456 35. Berkley JA, Lowe BS, Mwangi I, Williams T, Bauni E, Mwarumba S, et al. Bacteremia among
457 children admitted to a rural hospital in Kenya. *N Engl J Med*. 2005;352(1):39-47. doi:
458 10.1056/NEJMoa040275
- 459 36. Chisti MJ, Tebruegge M, La Vincente S, Graham SM, Duke T. Pneumonia in severely
460 malnourished children in developing countries - mortality risk, aetiology and validity of WHO clinical
461 signs: a systematic review. *Trop Med Int Health*. 2009;14(10):1173-89. doi: 10.1111/j.1365-
462 3156.2009.02364.x

- 463 37. Jensen BH, Olsen KEP, Struve C, Krogfelt KA, Petersen AM. Epidemiology and Clinical
464 Manifestations of Enteroaggregative *Escherichia coli*. Clin Microbiol Rev. 2014;27(3):614-30. doi:
465 10.1128/CMR.00112-13
- 466 38. Roche JK, Cabel A, Sevilleja J, Nataro J, Guerrant RL. Enteroaggregative *Escherichia coli* (EAEC)
467 impairs growth while malnutrition worsens EAEC infection: a novel murine model of the infection
468 malnutrition cycle. J Infect Dis. 2010;202(4):506-14. doi: 10.1086/654894
- 469 39. Huang DB, Mohanty A, DuPont HL, Okhuysen PC, Chiang T. A review of an emerging enteric
470 pathogen: enteroaggregative *Escherichia coli*. J Med Microbiol. 2006;55(10):1303-11. doi:
471 10.1099/jmm.0.46674-0
- 472 40. François R, Yori PP, Rouhani S, Sigvas Salas M, Paredes Olortegui M, Rengifo Trigoso D, et al.
473 The other Campylobacters: Not innocent bystanders in endemic diarrhea and dysentery in children in
474 low-income settings. PLoS Negl Trop Dis. 2018;12(2):e0006200. doi: 10.1371/journal.pntd.0006200
- 475 41. Tickell KD, Sharmin R, Deichsel EL, Lamberti LM, Walson JL, Faruque ASG, et al. The effect of
476 acute malnutrition on enteric pathogens, moderate-to-severe diarrhoea, and associated mortality in
477 the Global Enteric Multicenter Study cohort: a post-hoc analysis. Lancet Glob Health. 2020;8(2):e215-
478 e24. doi: 10.1016/s2214-109x(19)30498-x
- 479 42. Okhuysen PC, Dupont HL. Enteroaggregative *Escherichia coli* (EAEC): a cause of acute and
480 persistent diarrhea of worldwide importance. J Infect Dis. 2010;202(4):503-5. doi: 10.1086/654895
- 481 43. MAL-ED Network Investigators. Relationship between growth and illness, enteropathogens
482 and dietary intakes in the first 2 years of life: findings from the MAL-ED birth cohort study. BMJ Glob
483 Health. 2017;2(4):e000370. doi: 10.1136/bmjgh-2017-000370
- 484 44. Kosek MN, Ahmed T, Bhutta Z, Caulfield L, Guerrant R, Houghton E, et al. Causal Pathways from
485 Enteropathogens to Environmental Enteropathy: Findings from the MAL-ED Birth Cohort Study.
486 EBioMedicine. 2017;18:109-17. doi: 10.1016/j.ebiom.2017.02.024
- 487 45. Puspandari N, Amalia, N, Hartoyo, Y, Nursufiah, S, Sunarno, S, Sariadji, K. Enteric pathogen
488 among children under five years old with diarrheal diseases in Indonesia. IOP Conf Ser Earth Environ
489 Sci. 2021;913. doi: 10.1088/1755-1315/913/1/012098
- 490 46. Oyofe BA, Subekti D, Tjaniadi P, Machpud N, Komalarini S, Setiawan B, et al. Enteropathogens
491 associated with acute diarrhea in community and hospital patients in Jakarta, Indonesia. FEMS
492 Immunol Med Microbiol. 2002;34(2):139-46. doi: 10.1111/j.1574-695X.2002.tb00615.x
- 493 47. Olalemi AO, Ige OM, James GA, Obasoro FI, Okoko FO, Ogunleye CO. Detection of enteric
494 bacteria in two groundwater sources and associated microbial health risks. J Water Health.
495 2021;19(2):322-35. doi: 10.2166/wh.2021.212

- 496 48. Elfadaly HA, Hassanain NA, Hassanain MA, Barakat AM, Shaapan RM. Evaluation of primitive
497 ground water supplies as a risk factor for the development of major waterborne zoonosis in Egyptian
498 children living in rural areas. *J Infect Public Health*. 2018;11(2):203-8. doi: 10.1016/j.jiph.2017.07.025
- 499 49. Jo SJ, Kang HM, Kim JO, Cho H, Heo W, Yoo IY, et al. Evaluation of the BioFire Gastrointestinal
500 Panel to Detect Diarrheal Pathogens in Pediatric Patients. *Diagnostics (Basel)*. 2021;12(1). doi:
501 10.3390/diagnostics12010034
- 502 50. Liu J, Platts-Mills JA, Juma J, Kabir F, Nkeze J, Okoi C, et al. Use of quantitative molecular
503 diagnostic methods to identify causes of diarrhoea in children: a reanalysis of the GEMS case-control
504 study. *Lancet*. 2016;388(10051):1291-301. doi: 10.1016/S0140-6736(16)31529-X
- 505 51. Gaensbauer JT, Lamb M, Calvimontes DM, Asturias EJ, Kamidani S, Contreras-Roldan IL, et al.
506 Identification of Enteropathogens by Multiplex PCR among Rural and Urban Guatemalan Children with
507 Acute Diarrhea. *Am J Trop Med Hyg*. 2019;101(3):534-40. doi: 10.4269/ajtmh.18-0962
- 508
- 509

510 **Supporting Information**

511 **S1 Table. Weight-for-height, height-for-age, and weight-for-age z-score for birth cohort infants at each home**
512 **visit in Dili, Timor-Leste, 2019-2020.**

513 **S2 Table. Adjusted univariate odds ratios using a generalised estimating equations model for risk factors**
514 **associated with pathogen detection for infants in a birth cohort, Dili, Timor-Leste, 2019-2020.**

515 **S3 Table. Adjusted univariate odds ratios using a generalised estimating equations model for risk factors**
516 **associated with moderate wasting for infants in a birth cohort, Dili, Timor-Leste, 2019-2020.**

517 **S4 Table. Mean age in months at admission by diagnosis and type of enteric pathogen detected for hospital-**
518 **based surveillance cases, Dili, Timor-Leste, 2019–2020.**

519 **S5 Table. Adjusted univariate odds ratios using a generalised estimating equations model for pathogens and**
520 **host factors associated with hospitalisation for infants in a birth cohort, Dili, Timor-Leste, 2019-2020.**




Fig1c




Fig2




Fig1a




Fig1b