Multilevel determinants of Covid-19 vaccine hesitancy and undervaccination among

2 marginalized populations in the United States: A scoping review

- 4 Peter A. Newman^{1*}, Thabani Nyoni¹, Kate Allan¹, Sophia Fantus², Duy Dinh³, Suchon Tepjan⁴,
- 5 Luke Reid¹, Adrian Guta⁵

1

3

6

12

- 7 ¹ Factor-Inwentash Faculty of Social Work, University of Toronto, Toronto, Ontario, Canada
- 8 ² School of Social Work, University of Texas at Arlington, Texas, USA
- 9 1,3 Faculty of Health Sciences, Queen's University, Kingston, Ontario, Canada
- 10 ^{1,4} VOICES-Thailand Foundation, Chiang Mai, Thailand
- 11 ⁵ School of Social Work, University of Windsor, Windsor, Ontario, Canada
- 13 *Corresponding author
- 14 E-mail: p.newman@utoronto.ca

Abstract

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

Background: Amid persistent disparities in Covid-19 vaccination, we conducted a scoping review to identify multilevel determinants of Covid-19 vaccine hesitancy (VH) and undervaccination among marginalized populations in the U.S. **Methods:** We utilized the scoping review methodology developed by the Joanna Briggs Institute and report all findings according to PRISMA-ScR guidelines. We developed a search string and explored 7 databases to identify peer-reviewed articles published from January 1, 2020–October 31, 2021, the initial period of U.S. Covid-19 vaccine availability. We combine frequency analysis and narrative synthesis to describe factors influencing Covid-19 vaccination among marginalized populations. **Results:** The search captured 2,496 non-duplicated records, which were scoped to 50 peerreviewed articles: 11 (22%) focused on African American/Black people, 9 (18%) people with disabilities, 4 (8%) justice-involved people, and 2 (4%) each on Latinx, people living with HIV/AIDS, people who use drugs, and LGBTO+ people. Forty-four articles identified structural factors, 36 social/community, 27 individual, and 40 vaccine-specific factors. Structural factors comprised medical mistrust (of healthcare systems, government public health) and access barriers due to unemployment, unstable housing, lack of transportation, no/low paid sick days, low internet/digital technology access, and lack of culturally and linguistically appropriate information. Social/community factors including trust in a personal healthcare provider (HCP), altruism, family influence, and social proofing mitigated VH. At the individual level, low perceived Covid-19 threat and negative vaccine attitudes were associated with VH. **Discussion:** This review indicates the importance of identifying and disaggregating structural factors underlying Covid-19 undervaccination among marginalized populations, both crosscutting and population-specific—including multiple logistical and economic barriers in access,
and systemic mistrust of healthcare systems and government public health—from individual and
social/community factors, including trust in personal HCPs/clinics as reliable sources of vaccine
information, altruistic motivations, and family influence, to effectively address individual
decisional conflict underlying VH as well as broader determinants of undervaccination.

Keywords

Vaccination hesitancy, Covid-19 vaccines, medical mistrust, social determinants of health,

46

access to care, ethnic and racial minorities

Introduction

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

Vaccination remains an essential component of public health strategies to control the Covid-19 pandemic. Vaccination has reduced Covid-19-related morbidity and mortality in the United States (U.S.) [1], and globally—with an estimated 63% (14.4–19.8 million) reduction in deaths in the first year of Covid-19 vaccination alone [2]. Vaccine booster doses [3], and bivalent vaccine boosters compared with past monovalent vaccination [4], have further reduced Covid-19-associated hospitalization and death. Yet, despite the tremendous successes of Covid-19 vaccination, pervasive challenges ascribed to vaccine hesitancy (VH), and broader Covid-19 undervaccination, threaten the effectiveness of vaccines in controlling Covid-19 and other vaccine-preventable diseases. The World Health Organization (WHO) Strategic Advisory Group of Experts (SAGE) on Immunization defines vaccine hesitancy (VH) as "delay in acceptance or refusal of vaccination despite availability of vaccination services" [5,p.4161]. In addition to individual-level correlates of VH, SAGE [6] along with other research initiatives [7,8] have drawn attention to the critical importance of vaccine availability and access—amid systematic variations in vaccine coverage across countries and subpopulations—and issues that may pertain to specific vaccines. Collectively, these challenges support the need to identify and examine multilevel factors that contribute to Covid-19 VH and broader Covid-19 undervaccination (e.g., due to inaccessibility. unavailability) [9]. Disparities in Covid-19 morbidity and mortality, as well as lower rates of Covid-19 vaccination, are well documented among marginalized populations in the U.S. [10-13]. Marginalization is defined by systemic processes driven by ideologies such as racism, sexism, heterosexism, cissexism, ableism, and xenophobia, that exclude certain populations or relegate

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

them to the periphery of political and socioeconomic resources [14,15]. The structural and social inequities produced by marginalization create adverse social determinants of health (SDOH) such as residential segregation and housing insecurity, employment precarity, income insecurity, lack of access to affordable healthcare, barriers to physical mobility, and stigmatization—all of which increase vulnerability to poor health outcomes [14]. Adverse SDOH exacerbate vulnerability to Covid-19 through the disproportionate prevalence of underlying conditions associated with worse Covid-19 outcomes, lower levels of health insurance and healthcare access, and by impeding the ability to practice public health measures to reduce viral transmission [16-19]. Although heightened vulnerability indicates an increased need for vaccination, these same SDOH may contribute to *lower* levels of Covid-19 vaccination [20,21]. In fact, across 45 U.S. states, 0.1%–24.9% lower vaccination was documented among Blacks vs. Whites, 0.0%–37.6% lower among Hispanics vs. Whites, as of May 2021—with state-level rate differences significantly associated with a metric of structural racism [21]. In a national study, eligible healthcare facilities and community pharmacies in counties with higher Black composition were significantly less likely to provide Covid-19 vaccination [20]. Although some of the racial/ethnic disparities in Covid-19 vaccination documented earlier in the pandemic have reportedly narrowed over time, attributed in part to the success of targeted interventions [11], disparities persist in lower administration of booster doses among Black, Hispanic/Latinx, and American Indian populations versus Whites [22]. Consistent with the "inverse equity hypothesis," with each innovation in Covid-19 vaccines, uptake is predicted to be greater among wealthier and more socially connected segments of the population, thereby increasing rather than reducing inequity [23,24]. Nevertheless, population-level disparities in Covid-19 vaccination are often conceptualized as contingent on individual-level factors (e.g.,

low perceived threat, vaccine attitudes) to the relative underemphasis of structural and systemic factors [20,21,25,26]. Moreover, trust/mistrust in a personal HCP as a source of reliable information [27] is often conflated with systemic medical mistrust: distrust of healthcare systems, government public health, medical treatments, and/or the pharmaceutical industry arising from present-day experiences of discrimination and/or historical experiences of discrimination and injustice in the healthcare system [28-30].

Building on the SAGE "Vaccine Hesitancy Determinants Matrix" and recommendations to approach VH as a population- and context-specific phenomenon [6,31], along with critiques of the lack of clarity in the definition and application of VH [32-34], we conducted a scoping review to examine multilevel factors associated with Covid-19 VH and broader Covid-19 undervaccination among marginalized populations in the U.S.

Methods

A scoping review was conducted given our aim of mapping the evidence landscape [35] on factors associated with Covid-19 VH and undervaccination among marginalized populations. A scoping review protocol, including our search strategy and review process, has been previously published [36]. We utilized the scoping review methodology described by Arksey and O'Malley [37], further developed by the Joanna Briggs Institute [38], with all results reported in accordance with PRISMA Extension for Scoping Reviews (PRISMA-ScR) guidelines [39].

Research question

This scoping review was guided by the question, "What are the determinants of Covid-19 vaccine hesitancy' reported among marginalized populations in the U.S. and Canada?", with

sub-questions, "what are the focal populations in studies of Covid-19 VH and undervaccination among marginalized populations?" and "what are the contexts and determinants of Covid-19 VH and undervaccination by structural, social and community, individual, and vaccine-specific factors?" [36].

Information sources and search strategy

We identified the following list of databases to search in consultation with a specialist research librarian: Medline, Embase, Cochrane CENTRAL, Cochrane Covid-19 Study Register, PsychINFO, Sociological Abstracts, and the International Bibliography of the Social Sciences (IBSS). We then developed a search string with keywords and synonyms related to vaccine hesitancy (i.e., "vaccine hesitancy", "vaccine refusal", "vaccine confidence", "vaccine distrust", "vaccine mistrust", "vaccine barriers") and Covid-19 (i.e., "corona virus" or "coronavirus" or "COVID" or "nCoV" along with adjacent terms "19" or "2019, or "SARS-CoV-2" or "SARS Coronavirus 2", etc.). S1 Table shows an example search string for Medline and Embase. We adapted the search string to accommodate differences in syntax and parameters among the selected databases.

Study selection criteria

We developed inclusion and exclusion criteria prior to conducting the search for peer-reviewed articles published between January 1, 2020 and October 31, 2021: 1) populations in the U.S. or Canada; 2) adults ≥18 years; 3) focused on (>50%) marginalized populations, 4) focused on Covid-19 vaccine hesitancy, refusal, confidence, mistrust, or barriers. Articles were excluded if they were 1) published prior to 2020; 2) not written in English; 3) focused on healthcare

workers, staff, or students/interns; or 4) focused on children (<18 years) or parents' vaccination of children. We selected search dates corresponding to the declaration of Covid-19 as a public health emergency through the initial authorization of Covid-19 vaccines for the adult population in the U.S. and Canada in December 2020, until U.S. FDA and Health Canada recommendations for the first booster vaccines, beginning in late September 2021 to early November 2021 [40,41]. We identified marginalized populations informed by a review of populations particularly vulnerable to discrimination and exclusion in healthcare [42] or due to broader adverse SDOH [14]. The populations include those marginalized in relation to their race or ethnicity, sexual orientation or gender identity, physical or mental disability, persons living with HIV, immigrants/refugees, persons who experience homelessness, justice-involved people, and people who use drugs, along with their intersections (e.g., Latinx immigrants).

Study selection process

Search results of peer-reviewed articles were uploaded into Covidence software. Multiple research team discussions ensured consistent application of inclusion/exclusion criteria. Initially groups of two reviewers (among LR, TN, DD, KA, SF or ST) independently screened abstracts and titles until we attained inter-rater reliability of 90% [43]; this was achieved after two rounds of abstract review and team discussions to achieve consensus. Subsequently, in accordance with rapid review methods, one reviewer screened each abstract/title to determine inclusion or exclusion [44]. One reviewer (TN, DD, KA, SF or ST) then screened each full-text for inclusion. Full-text articles designated for exclusion were then reviewed by a single arbitrator (PN), with discrepancies resolved by consensus among the research team [44,45].

Data extraction and synthesis

We abstracted data on publication characteristics (i.e., author(s), study sites, sample size, populations, methods, and domains of VH factors identified (see Table 1). We then reviewed the abstracted data using quantitative (i.e., frequency) analysis to describe study locales, sample sizes, focal populations, methods, and VH factor domains, and qualitative (i.e., narrative) analysis to synthesize the evidence on VH and undervaccination among each subpopulation.

All authors categorized publications by domains of factors identified, using a four-component framework informed by the "Vaccine Hesitancy Determinants Matrix" [6,31] and recommendations to disaggregate multilevel phenomena [8,33,34]. Domains comprised structural (e.g., access to vaccination site, medical mistrust); social/community (e.g., family support, trust in personal HCP or clinic, social proofing [i.e., based on actions of others in one's community], community benefits); individual (e.g., perceived Covid-19 threat, perceived transmission risk, general vaccine beliefs); and Covid-19 vaccine-specific factors (e.g., vaccine safety, side effects, efficacy). One change to data extraction (but not inclusion) was made from the initial protocol in that we differentiated between medical mistrust at the "system-level" and trust in a personal HCP/clinic.

Results

The search captured 4,399 peer-reviewed articles in total, with 2,496 abstracts screened after removal of duplicates. We identified 363 full-text articles to be screened for eligibility, with a total of 50 peer-reviewed articles included [12,26,46-93]. The PRISMA flow diagram shows the process of identifying relevant peer-reviewed journal articles (see Fig 1).

Description of studies

Among the 50 articles included, 86% were quantitative studies, 10% qualitative, and 4% mixed methods (see Table 1). Sample sizes ranged from n=24 to n=164,283 (N=499,237; median=437).

All studies were conducted in the U.S., including 24 (48%) national studies, 8 (16%) in the West, 7 (14%) Northeast, 5 (10%) Midwest, 5 (10%) Southeast, and 1 (2%) in the Southwest region.

No studies were identified from Canada that met inclusion criteria within the timeframe of the review.

196

197

Fisher	1668	National	Online	Racial/ethnic min.	X				(+)		
Garcia	1515	National	Online: US Hemodialysis Facilities	Racial/ethnic min.	Х				(+/-)	(-)	(-)
Garcia	46	W	Telephone: OR	Latinx		х		(-)	(+)		(-)
Geana	25	MW	Telephone: Midwest USA	JI		Х		(-)	(-)	(-)	(-)
Ghaffari-Rafi	359	W	Telephone: HI	PWD	Х			(-)	(-)	(+)	(-)
Guidry	788	National	Online	Racial/ethnic min.	х			(-)			(-)
Hagan	164,283	National	Data obtained from Electronic Medical Records in Federal Bureau of Prisons	JI	х			(+/-)	(+)	(-)	(+)
Iadarola	825	NE	Online: NY	PWD	Х			(-)	(-)	(-)	(-)
Jimenez	111	NE	Online: NJ	AA/Black & Latinx	х			(-)	(-)		(-)
Johnson	248	SE	Clinic: Baton Rouge, LA	AA/Black	X			(-)	(-)		(-)
Jones	94	SE	Telephone/Online: Miami, FL	PLHA	х			(-)	(+)	(+/-)	(-)
Kricorian	1950	National	Online	AA/Black & Latinx	х			(-)	(+)		(-)
Kuhn	90	W	Telephone: Los Angeles, CA	PEH	Х			(-)	(-)	(-)	(-)
Luo	6715	National	Data obtained from the Medicare Current Beneficiary Survey	PWD	X			(-)		(+)	(-)
Masson	258	W	California residential substance use treatment programs	PWUD	X			(-)		(+)	(-)
Momplaisir	24	NE	Online: Philadelphia, PA	AA/Black		х		(-)	(+/-)	(-)	(-)
Moore	257	SE	Community Events: The Central Savannah River Area	AA/Black	X			(-)		(-)	
Myers	439	National	Online	PWD	х			(-)	(+)	(+)	(-)
Okoro	183	MW	Online: MN and WI	AA/Black			X	(-)	(+/-)	(-)	(-)
Razzaghi	135,968	National	Data obtained from the Vaccine Safety Datalink (VSD)	Racial/ethnic min.	Х			(-)			(-)

Rodriguez	2301	National	Emergency Departments in Hospitals	Racial/ethnic min.	х		(-)	(-)		(-)
Rodriguez	175	SE	Online: Miami, FL	PLHA	X		(-)	(+)	(-)	(-)
Rungkitwattanakul	90	National	Hemodialysis centres	AA/Black	х		(-)	(-)	(-)	(-)
Ryerson	56,749	National	Telephone	PWD	х		(-)	(+)	(+)	
Salmon	2525	National	Online	Racial/ethnic min.	х		(-)	(+/-)	(+)	(-)
Stephenson	290	National	Data obtained from the Love and Sex in the Time of COVID-19 study	LGBTQ+	х			(+)	(+)	
Stern	5110	National	Correctional and detention facilities	JI	Х		(-)	(-)	(-)	(-)
Stoler	1040	National	Online	AA/Black	х		(-)	(+)		
Teixeira da Silva	1350	National	Online	LGBTQ+	x		(-)	(+/-)		(-)
Tsapepas	664	NE	Telephone: New York City, NY	Racial/ethnic min.		х	(-)			(-)
Uhr	701	National	Online	PWD	X		(+/-)	(+)	(-)	(-)
Wagner	1138	MW	Data obtained from the Detroit Metro Area Community Study	AA/Black	х		(-)			(-)
Wang	293	NE	Online: DE	AA/Black	X		(-)			(-)
Xiang	401	W	Online: Mainly OR, WA, and CA	PWD	Х			(+/-)	(+)	(-)
Yang	387	National	Online	PWUD	X		(-)	(+/-)	(-)	(+)
Zhang	435	National	Online: Refugee communities	IMM	Х		(-)	(+)		(-)

AA, African Americans; IMM, immigrants and refugees; JI, justice-involved people; Latinx, Latinx/Hispanic people; LGBTQ+, Lesbian, gay, bisexual, transgender, queer and other sexual/gender minority people; PEH, people experiencing homelessness; PWD, people with disabilities; PLHA, people living with HIV/AIDS; PWUD, people who use drugs; Racial/Ethnic min., racial/ethnic minority.

Table 2 shows the number of focal articles by subpopulation, with 11 (22%) focused on African American/Black people, 9 (18%) on people with disabilities, 4 (8%) on justice-involved people, and 2 (4%) each on Latinx, people living with HIV/AIDS (PLHA), people who use drugs (PWUD), and lesbian, gay, bisexual, transgender, and queer (LGBTQ+) people. We allocated findings from articles broadly focused on racial/ethnic minorities to each of the subpopulations (i.e., African American/Black, Latinx, Asian, American Indian) for which disaggregated data were provided (i.e., one article could be a "source" for several populations). Four additional articles addressing VH pertaining to two subpopulations (e.g., African American/Black PLHA) were each allocated as two sources. Table 2 shows the total number of sources (n=73) addressing population-specific factors associated with VH across the 50 articles. Among these, 41.1% of sources (n=30) identified VH factors among African American/Black populations, 17.8% (n=13) Latinx populations, 12.3% (n=9) people with disabilities, and 5.5% (n=4) each Asian, justice-involved, and PLHA.

Table 2. Focal articles and additional sources by subpopulation

215

216217

Subpopulation	Focal articles on each sub- population	Additional sources addressing each subpopulation	Number (%) of total sources by sub- population (n=73)		
African American/Black	11	19	30 (41)		
Latinx	2	11	13 (18)		
People with disabilities	9	0	9 (12)		
Justice involved	4	0	4 (5)		
People living with HIV/AIDS	2	2	4 (5)		
Asian people	0	4	4 (5)		
People who use drugs	2	1	3 (4)		
Immigrants/refugees	1	1	2 (3)		
LGBTQ+ people	2	0	2 (3)		
People experiencing homelessness	1	0	1 (1)		
American Indians/Native	0	1	1(1)		
Subtotal	34	39	73		

LGBTQ+, Lesbian, gay, bisexual, transgender, queer and other sexual/gender minority people

Fig 2 shows the number of articles addressing factors associated with VH in each of four domains: structural, social/community, individual, and Covid-19-vaccine-specific. Table 3 shows the domains of factors associated with VH in each subpopulation. Next, we describe findings based on a narrative synthesis of factors associated with VH and undervaccination for each focal population.

Fig 2. Count of articles identifying factors in each of the four domains of Covid-19 vaccine hesitancy and underutilization

Table 3. Population by domains of factors identified in Covid-19 vaccine hesitancy and

undervaccination

Subpopulation	Domains							
	Structural	Social/Community	Individual	Vaccine-specific				
African American/Black	X	X	X	X				
American Indian/Native American	X			X				
Asian	X	X		X				
Latinx	X	X		X				
LGBTQ+	X	X	X	X				
Immigrant/Refugee	X	X		X				
Justice-involved people	X	X	X	X				
People experiencing homelessness	X	X	X	X				
People living with HIV/AIDS	X	X	X	X				
People with disabilities	X	X	X	X				
People who use drugs	X		X	X				

LGBTQ+, Lesbian, gay, bisexual, transgender, queer and other sexual/gender minority people

Black and African American people

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

Eleven articles focused exclusively on Covid-19 vaccination among Black/African Americans [26,56,59,69,74,75,77,81,85,89,90], and 19 additional articles provided disaggregated data—a total of 30 sources [46-48,49,51-53,55,60,65-68,70,73,82,84,86,92]. At the structural level, 27 sources addressed the negative impacts of medical mistrust and distrust of government on Covid-19 vaccination among Black and African Americans [26,46-49,51-53,55,59,65-70,73-75,77,81,82,84-86,89,92]. Several studies described Covid-19 VH, such as due to fears of being experimented on with Covid-19 vaccines without consent, "based on...knowledge of historic mistreatment and unethical research practices that adversely impacted black patients and participants" [74, p.1786], including the legacy of the Tuskegee Study of untreated syphilis [53,67,68,77]. Eleven sources addressed barriers to present-day healthcare and Covid-19 vaccines for Black/African Americans, including unemployment, housing insecurity, limited interactions with HCP, and difficulties travelling to vaccination sites [48,52,53,56,68-70,75,77,89,90]. At the social/community level, nine sources indicated the role of trust in HCPs in Covid-19 vaccination intentions [26,46,51,60,69,70,74,82,85]. Characteristic of many studies, "Blacks and Hispanics were more likely than Whites to state that it was 'extremely' important that a medical professional of their same race/ethnicity endorse the vaccine before they took it" [26,p.8]. Seven studies addressed the influence of families and communities on Covid-19 vaccination [48,52,59,70,74,77,81], and further that "governmental agencies are less wellreceived as information sources compared to friends or family" [59, p.1103]. In one study, family influence promoted VH by casting doubt on public health information, rather than motivating vaccination [81].

At the individual level, four studies addressed negative impacts of religious beliefs, including preference for 'natural' medicines over vaccines, and general vaccine aversion on Covid-19 vaccination [67,74,77,81]. Covid-19 vaccine-specific concerns, identified in 14 studies, included vaccine safety, adverse effects, and uncertain efficacy [26,51-53,59,67-70,73,74,77,81,90], and the importance of evidence of Covid-19 vaccine safety and efficacy specific to Black/African Americans: "any data that's connected with some of the conditions that are prevalent in our community" [53, p.8].

Latinx/Hispanic people

Two studies focused on Covid-19 vaccination among Latinx/Hispanic people [50,61], with an additional 11 sources providing data specific to this population [26,46,48,49,53,60,67,68,70,73,82]. At the structural level, five studies identified negative impacts of medical mistrust [48,49,61,67,82], including fear of being used as "guinea pigs (conejillos de indias)" and of being sterilized [61]. Limited access to information in one's preferred language, complicated online vaccination sign-up procedures, lack of insurance, and lack of transportation to vaccination sites posed barriers to Covid-19 vaccination [48,53,61,68,70]. Two studies underscored the importance of tailored vaccine education and accessible information [68]: "[a family member] doesn't speak English...so, a lot of the information that he may have heard about the vaccine and Covid probably came off of Univision and Telemundo" [48, p.6].

At the social/community level, five studies described the important role of personal HCPs in Covid-19 vaccination among Latinx communities [26,46,60,70,82], including endorsement

from an HCP of one's own ethnicity [26]. Two studies addressed altruism and wanting to protect

one's community as a motivator of Covid-19 vaccination [53,61]. Families and close communities were described as playing an important role in Covid-19 vaccination knowledge and decision-making [48,61,70]: "79% of the youth had discussed health with their families for an average of 50 minutes [per week]" [61, p.753]. No individual-level factors were described, but Covid-19 vaccine-specific concerns arose around safety, adverse effects, and efficacy [26,61,67,68,70,73].

Asian people

No studies focused exclusively on Covid-19 vaccination among people of Asian descent in the U.S.; four sources provided disaggregated data on Asian people [46,53,64,67]. At the structural level, barriers included difficulties in accessing reliable translations and messengers of Covid-19 vaccine information [53]. Beyond linguistic translation, a Filipinx participant explained, "In terms of translators...maybe those who may not be of the same culture or the way that they're explaining the medical terminology may be intimidating" [53, p.6]. Medical mistrust was identified as negatively impacting vaccination intention [53,67], with trust in personal HCPs as sources of vaccine information positively associated with Covid-19 vaccination intention among Asian people [46]. Covid-19 vaccine-specific concerns arose around safety, adverse effects, and efficacy [53,64,67], including the need for population-specific evidence: "There has been a lot of concerns in my family on how the vaccine works for people with heart disease, which really affects...a lot of the Filipino community, and also those with respiratory diseases" [53, p.6].

American Indians

No studies among those reviewed focused on Covid-19 vaccination among American Indians or Alaskan Natives; one source provided data on American Indians [53]. Structural factors arose regarding distrust of government, desire for autonomy ("to be respected)", challenges in access to digital technology and the internet, lack of quality translations and tailored information, lack of employment benefits, insurance, and transportation to get vaccinated [53]. In addition to distrust ascribed to the politicization of vaccination, a participant decried structural-level challenges for elders: "Some elders...they're houseless...there's no cell phone, they ride public transportation. How is it going to get distributed [to them]...the most vulnerable... it feels like our culture always gets the shitty end of the stick" [53, p.6]. Covid-19 vaccine-specific concerns arose around side effects, efficacy, and multiple doses—associated with challenges for Native families due to the need for several trips to vaccination sites.

LGBTQ+ people

Two sources focused on Covid-19 vaccination among LGBTQ+ people [83,86]. At the structural level, medical mistrust based on historical discrimination against sexual and gender minorities was negatively associated with Covid-19 vaccination intentions [86]. Loss of employment and housing insecurity were described as factors that underscored the threat of the pandemic among LGBTQ+ people, thus motivating vaccination. At the social and community level, altruistic motivations were positively associated, while anticipated stigma was negatively associated with vaccination intentions [83,86]. Individual-level factors impacting vaccination were described in perceived threat of Covid-19: "negative experiences, along with the reported high levels of perception of the seriousness of Covid-19 infection, are likely driving the high

levels of willingness to be vaccinated" [83,p.9]. Covid-19 vaccine-specific concerns included adverse effects and safety [86].

People with disabilities

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

Nine articles focused on challenges in Covid-19 vaccination for people with disabilities (PWD) [12,57,58,64,67,72,76,88,91]. At the structural level, five studies addressed the impact of trust in government: three reported a positive association with Covid-19 vaccination [57,58,88], while two described governmental distrust contributing to VH [67,72]. Strict public health measures and widespread reliance on digital technology during the pandemic posed structural barriers in access to Covid-19 vaccination for PWD [12,72,76,88]. While PWD were described as more likely than people without a disability to endorse Covid-19 vaccination, they were more likely to report difficulties booking online vaccination appointments and locating and travelling to vaccination sites [12]. At the social and community level, having a trusted HCP or clinic exerted a positive influence on Covid-19 vaccination [12,57,76,88,91]. For example, among people living with multiple sclerosis (MS), MS-specialist HCPs and the National MS Society were deemed trusted sources: "well-positioned to provide accurate education about Covid-19 disease and vaccine considerations...specific to the MS population as well as their individual patients" [57,p.3]. At the individual level, perceived threat of Covid-19 was a positive correlate of Covid-19 vaccination among PWD, including individuals with chronic diseases and multiple comorbidities [12,57,58,64,72,76,91]. Covid-19 vaccine-specific concerns included adverse effects, speed of vaccine development, vaccine safety, and efficacy [58,64,67,72,76,88,91]. In addition to "safety and efficacy...the top Covid-19 vaccination concerns", apprehension arose around "the effect of

the vaccine on MS symptoms and implications for [immunosuppressive] DMT [disease-modifying therapy] use" [88, p.1078].

Justice-involved people

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

Four articles focused on Covid-19 vaccination among justice-involved (JI) individuals (i.e., currently or formerly convicted or incarcerated) [54,63,66,84]. At the structural level, medical mistrust and distrust of government exerted a negative impact on Covid-19 vaccination among individuals in detention facilities [54,63,84] "based on their interactions with law enforcement or the justice system or their experiences with institutional racism" [84, p.476]. A study with incarcerated women described "a direct correlation between trust in the pharmaceutical industry and willingness to receive the vaccine": "I've watched too many people die because pharmaceuticals... So I don't know. Pharmaceutical companies just see a lot of big money in there" [63, p.895]. One study described mitigation of structural-level barriers to Covid -19 vaccination through tailored vaccine education and "early access to vaccine through a dedicated federal allocation" for incarcerated individuals and facility staff [66,p.5889]. At the social and community level, family and community influence was associated with Covid-19 VH [54.63.66] among JI individuals in the absence of tailored public health messaging: "The probable low health and informational literacy of these women leave them vulnerable to influence from family and friends, many sharing and promoting disinformation and conspiracy theories frequently shared by homegrown anti-vaccine organizations" [63, p.895]. Social proofing exerted a positive influence: "some people may have been encouraged to accept vaccination because they could see their peers being vaccinated" [54,66, p.5889].

At the individual level, all four studies reported low perceived risk of contracting Covid-

19 as a correlate of VH among JI people [54,63,66,84]. A study in correctional and detention facilities in 4 states reported that of participants (n=1823/5110) who would refuse Covid-19 vaccination, 18.9% (n=344/1823) "did not perceive themselves to be at risk for Covid-19 or thought vaccination was unnecessary" [84, p.475]. Covid-19 vaccine-specific concerns arose around vaccine safety and efficacy [54,63,84], which was associated with "lack of access to reliable information" [54, p.375].

People living with HIV/AIDS

Two articles focused on factors impacting Covid-19 vaccination intentions among PLHA [70,80], with two additional sources providing PLHA-specific data [51,83]. At the structural level, medical mistrust and distrust of government among PLHA were associated with lower Covid-19 vaccination intentions [51,70,80]. A study with HIV-positive and predominantly sexual minority Black Americans (n=101) indicated that "The most prevalent general mistrust beliefs (endorsed by about half or more than half) concerned withholding information or a lack of honesty by the government," which was correlated with VH [51, p.203]. A study with HIV-positive Black and Latinx Americans described unemployment being exacerbated by the pandemic, with negative impacts on access to healthcare and Covid-19 vaccination [70]. At the social and community level, the positive influence of personal HCPs as trusted sources of information on Covid-19 vaccination was reported [51,70,80]: "Lower hesitancy scores were present in participants who reported that they would get vaccinated if their provider recommended it" [80, p.98].

At the individual level, perceived threat was associated with Covid-19 vaccination intentions: "...HIV-positive men may have more concerns about the seriousness and perceived

impact of Covid-19 infection on their health" [80, 83, p.9]. Covid-19 vaccine-specific concerns arose around vaccine safety, adverse effects, and efficacy [51,70,80], with some PLHA concerned that their health condition may make Covid-19 vaccination unsafe [80].

People who use drugs

Two studies focused on Covid-19 vaccination among people who use drugs (PWUD) [73,92], with one additional source providing disaggregated data on PWUD [64]. At the structural level, barriers to accessing healthcare and Covid-19 vaccines, as well as medical mistrust and distrust in government were associated with lower Covid-19 vaccination intentions [64,73]. One study noted vulnerability to misinformation among PWUD amid medical mistrust: "As a marginalized population, patients with illicit drug use are often detached from and mistrust the healthcare system—by extension, these patients may be more reliant on illegitimate information sources" [64, p.803]. At the individual level, mask wearing was associated with vaccine trust, both described as a function of accurate Covid-19 information [73]. Covid-19 vaccine-specific factors arose around safety and efficacy concerns that may negatively impact uptake [73].

Immigrants and refugees

One study focused on Covid-19 vaccination among immigrants and refugees [93], with one additional source providing disaggregated data [61]. At the structural level, barriers to healthcare and Covid-19 vaccine access were described as negatively impacting vaccination [61,93]. A study with Latinx immigrant families identified "fear of encountering law enforcement/immigration as barriers to access health care" [61, p.750]. Limited access to

information in one's preferred language, lack of reliable information from trusted sources, constraints in access to the internet and digital technology, and lack of transportation and employment benefits (i.e., paid sick days) were identified as barriers to Covid-19 vaccination [93]. At the social and community level, family influence was demonstrated in a national online survey of refugees (n=435): among those who reported intention to receive a Covid-19 vaccine, nearly two-thirds (65.0%; n=199/306) indicated that it was motivated in part to protect their family members [93]. Covid-19 vaccine-specific concerns about vaccine safety, adverse effects, and efficacy were associated with VH, with two-thirds (71.3%; n=92/129) of refugees who reported no intention or being unsure of accepting Covid-19 vaccination indicating concerns about adverse effects [93].

People experiencing homelessness

One article focused on challenges to Covid-19 vaccination for people experiencing homelessness [71]. At the structural level, Covid-19 VH was described as a function of "a more general distrust of systems (e.g., health and homeless services) that are not designated to meet their needs" [71, p.6]. Trusting the information received from friends, family, and social media rather than "official news sources" was associated with greater Covid-19 VH [71]. At the individual level, general vaccine aversion and low Covid -19 risk perception, and vaccine-specific concerns about safety, adverse effects, and efficacy were associated with VH among people experiencing homelessness [71].

Discussion

This scoping review of Covid-19 vaccine hesitancy and broader underutilization of

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

Covid-19 vaccination provides evidence of multilevel determinants of delays or refusal of Covid-19 vaccination across marginalized populations in the U.S. Overall, a preponderance of structural-level factors were identified that impact Covid-19 vaccination across marginalized populations. Logistical and economic constraints including lack of transportation to distant locales, lack of paid sick days, lack of availability of vaccination at familiar/local community venues, and barriers in access due to being unemployed, unstably housed, uninsured/underinsured, and the digital divide (i.e., access to digital technology and the internet, and technical and financial ability to utilize it [94]) were compounded by lack of culturally and linguistically appropriate information and access to trusted HCPs. Collectively, these findings suggest that approaches to undervaccination among marginalized populations that focus predominantly on individual-level, psychological factors may fail to capture the breadth and impact of determinants of Covid-19 vaccination. Importantly, we identified evidence for differential impacts of medical mistrust, as a structural factor [29], and trust in a personal HCP or clinic, as a social/community factor. The influence of medical mistrust reported with respect to government public health, healthcare systems, and the pharmaceutical industry, emerged across multiple marginalized populations including African American, Latinx, LGBTQ+ populations, and their demographic intersections—attributed in part to historical discrimination and past unethical medical research and interventions, as well as present-day discrimination in healthcare and government institutions. Nevertheless, having a trusted HCP, particularly with shared race or ethnicity [26,51,69,70,74,82,85], or a familiar source of primary healthcare, mitigated Covid-19 VH despite broader mistrust of healthcare systems and government public health. JI populations, PEH, and PWUD expressed medical mistrust due to generally negative experiences with

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

government and healthcare institutions. However, as in the case of JI populations, proactive interventions to promote tailored education along with policy decisions to prioritize Covid-19 vaccination in prisons served to mitigate VH and promote vaccination [66]. Personal HCPs and usual care venues were widely reported to assuage concerns and reduce VH, indicating a crucial role for primary healthcare [95]. In this light, the absence of information from trusted sources tailored to address healthcare needs experienced as most relevant to one's own community—reported among Black and Pilipino populations, PLHA and PWD—and the lack of availability of Covid-19 and vaccine information in one's primary language (reported among Latinx, Asian, and immigrant/refugee populations) emerged as substantial missed opportunities to promote Covid-19 vaccination. Access to Covid-19 vaccination from a known HCP or clinic may be particularly important for marginalized communities, who are more predisposed towards medical mistrust due to negative interactions with the healthcare system and other societal institutions, and more vulnerable to discrimination in healthcare [96,97]. Overall, the conflation of structural factors—documented in multifaceted barriers in access to Covid-19 vaccination and systemic mistrust of healthcare systems—with individual and social/community factors, such as perceived threat of Covid-19, perceived risk of transmission, general vaccine attitudes, endorsement from one's personal HCP, and family influence, each of which may impact on "decisional conflict" more specifically aligned with VH [32], may fail to foster solutions that effectively address the identified problems [34,95,98]. Moreover, the attribution of structural-level barriers to individual volition and deficits may exacerbate alienation and mistrust among populations already marginalized due to systemic discrimination, stigma, and adverse SDOH [96].

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

Interestingly, some of the same ostensible factors showed negative or positive associations with VH depending on the population and context. For example, family influence exerted a positive impact on Covid-19 vaccination among Latinx communities and immigrant/refugee populations through familial sharing of health information and desire to protect one's family. However, among other populations, the context of medical mistrust and the virtual vacuum created by the absence of access to culturally tailored public health information rendered family influence a liability, resulting in promulgation of conspiracy theories and mistrust of Covid-19 vaccines among some African Americans [81], PEH [71], and JI persons [63]. These examples underscore the importance of approaching VH as a population-specific and context-dependent phenomenon [6,7]—in addition to disentangling the assessment of systemic factors from individual-level and social/community concerns [99]. Consequently, the same intervention applied to what might be understood as the same factor may exert opposite effects on Covid-19 VH across different populations. Finally, facilitators of Covid-19 vaccination emerged despite constraints. For example, findings in common across PLHA and PWD—among whom one might reasonably expect heightened concerns about vaccination due to underlying comorbidities and potential interactions with routine medications—included the benefits of regular contact with a personal HCP or clinic; this promoted trust in the vaccine information received and confidence in the HCP's ability to address population-specific concerns. Furthermore, intersectional strengths reported regarding Latinx immigrant and refugee populations included motivations to protect one's community and cultural norms around familial sharing of health information, which served to mitigate VH. Beyond a deficit model, community assets and sociocultural facilitators may be leveraged [100] to support Covid-19 vaccination.

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

Findings from this scoping review should be understood in the context of limitations. First, as a scoping review, our objective was to map and synthesize evidence on Covid-19 VH and undervaccination among marginalized populations; we cannot draw causal conclusions about the associations identified, nor was our aim to quantify Covid-19 vaccination intentions or uptake, more apropos of meta-analysis. However, our scoping review methods reveal nuanced information through synthesis of quantitative and qualitative findings on population-specific barriers to Covid-19 vaccination and suggest important directions for future research. Second, additional populations to those included might be characterized by marginalization, although we based our focal populations on previous evidence and a priori criteria [42,43]. Third, while our initial search comprised studies from the U.S. and Canada, no studies were identified from Canada that met inclusion criteria during the specified time frame, possibly due to the later availability of Covid-19 vaccination to the general adult population [41]. Studies published after the time frame of our review might include additional populations, data, and study locales, although we purposely focused on the initial introduction of Covid-19 vaccines before subsequent booster doses and bivalent vaccines. Lastly, factors associated with VH and underutilization of monovalent and bivalent boosters may differ from those revealed for initial Covid-19 vaccination, although vaccine inequities remain [23]. In conclusion, we identified multilevel factors that influence Covid-19 VH and broader underutilization of Covid-19 vaccination, several of which may apply to other vaccinations. This scoping review suggests that distinguishing among individual and social/community factors that may fuel decisional ambivalence characteristic of Covid-19 VH—"a state of indecision and uncertainly about vaccination before a decision is made to act (or not act)" [35, p.58]—and factors attributable to systemic and structural barriers in access to vaccination and historicallyjustified mistrust of healthcare institutions, government public health, and the pharmaceutical industry, may support evidence informed strategies to promote Covid-19 vaccination among marginalized communities. Nevertheless, distilling and applying lessons from Covid-19 vaccination to effectively and equitably address VH and broader challenges in vaccine access may be contingent on resourcing and scaling up community partnerships and local health infrastructures [23,25,96,100], national population-specific public health initiatives to promote vaccination [101,102], and structural reforms beyond those traditionally associated with public health.

Acknowledgements

We would like to thank Nalini Singh, University of Toronto Research Librarian, for assistance in designing the literature search strategy and Dares Wotansirigul for uploading sources into Covidence software.

Funding

This work was supported by the Canadian Institutes of Health Research [GA3-177731].

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Peter A. Newman is a member of the Editorial Board of PLOS ONE.

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

References 1. Steele MK, Couture A, Reed C, Iuliano D, Whitaker M, Fast H, et al. Estimated number of COVID-19 infections, hospitalizations, and deaths prevented among vaccinated persons in the US, December 2020 to September 2021. JAMA Netw Open. 2022;5(7):e2220385. https://doi.org/10.1001/jamanetworkopen.2022.20385 Watson OJ, Barnsley G, Toor J, Hogan AB, Winskill P, Ghani AC. Global impact of the 2. first year of COVID-19 vaccination: a mathematical modelling study. Lancet Infect Dis. 2022:22(9):1293-302. https://doi.org/10.1016/S1473-3099(22)00320-6 3. Arbel R, Hammerman A, Sergienko R, Friger M, Peretz A, Netzer D, Yaron S. BNT162b2 vaccine booster and mortality due to Covid-19. N Engl J Med. 2021;385(26):2413-20. https://www.nejm.org/doi/10.1056/NEJMoa2115624 Surie D, DeCuir J, Zhu Y, Gaglani M, Ginde AA, Douin DJ, et al. Early estimates of 4. bivalent mRNA vaccine effectiveness in preventing COVID-19–associated hospitalization among immunocompetent adults aged ≥65 Years — IVY Network, 18 States, September 8-November 30, 2022. MMWR Morb Mortal Wkly Rep. 2022;71(5152):1625-30. http://dx.doi.org/10.15585/mmwr.mm715152e2 MacDonald NE; SAGE Working Group on Vaccine Hesitancy. Vaccine hesitancy: 5. definition, scope and determinants. Vaccine. 2015;33(34):4161-4. https://doi.org/10.1016/j.vaccine.2015.04.036 SAGE Working Group. Report of the SAGE Working Group on vaccine hesitancy. 6. Geneva: WHO; 2014 Nov 12. Available from: https://www.assetscienceinsociety.eu/sites/default/files/sage working group revised report vaccine hesitan cy.pdf [accessed 4 January 2023].

- 573 7. Larson HJ, Clarke RM, Jarrett C, Eckersberger E, Levine Z, Schulz WS, et al. Measuring
- trust in vaccination: a systematic review. Hum Vaccin Immunother. 2018;14(7):1599-609.
- 575 https://doi.org/10.1080/21645515.2018.1459252
- 576 8. Thomson A, Robinson K, Vallée-Tourangeau G. The 5As: a practical taxonomy for the
- determinants of vaccine uptake. Vaccine. 2016;34(8):1018-24.
- 578 https://doi.org/10.1016/j.vaccine.2015.11.065
- 579 9. Gostin LO, Hodge JG Jr, Bloom BR, El-Mohandes A, Fielding J, Hotez P, et al. The public
- health crisis of underimmunisation: a global plan of action. Lancet Infect Dis.
- 581 2020;20(1):e11-6. https://doi:10.1016/S1473-3099(19)30558-4
- 582 10. Baack BN, Abad N, Yankey D, Kahn KE, Razzaghi H, Brookmeyer K, et al. COVID-19
- vaccination coverage and intent among adults aged 18-39 years United States, March-
- 584 May 2021. MMWR Morb Mortal Wkly Rep. 202170(25):928-33.
- 585 http://dx.doi.org/10.15585/mmwr.mm7025e2
- 586 11. Kriss JL, Hung MC, Srivastav A, Black CL, Lindley MC, Lee JT, et al. COVID-19
- vaccination coverage, by race and ethnicity National Immunization Survey Adult COVID
- Module, United States, December 2020-November 2021. MMWR Morb Mortal Wkly Rep.
- 589 2022;71(23):757-63. http://dx.doi.org/10.15585/mmwr.mm7123a2
- 590 12. Ryerson AB, Rice CE, Hung MC, Patel SA, Weeks JD, Kriss JL, et al. Disparities in
- 591 COVID-19 vaccination status, intent, and perceived access for noninstitutionalized adults,
- by disability status National Immunization Survey Adult COVID Module, United States,
- 593 May 30-June 26, 2021. MMWR Morb Mortal Wkly Rep. 2021;70(39):1365-71.
- 594 https://doi.org/10.15585/mmwr.mm7039a2

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

Shariff SZ, Richard L, Hwang SW, Kwong JC, Forchuk C, Dosani N, et al. COVID-19 vaccine coverage and factors associated with vaccine uptake among 23,247 adults with a recent history of homelessness in Ontario, Canada: a population-based cohort study. Lancet Public Health. 2022;7(4):e366-77. https://doi.org/10.1016/s2468-2667(22)00037-8 Baah FO, Teitelman AM, Riegel B. Marginalization: conceptualizing patient vulnerabilities in the framework of social determinants of health—an integrative review. Nurs Inq. 2019;26(1):e12268. https://doi.org/10.1111/nin.12268 Sharma M. Developing an integrated curriculum on the health of marginalized populations: 15. successes, challenges, and next steps. J Health Care Poor Underserved. 2014;25(2):663-9. https://doi.org/10.1353/hpu.2014.0102 Lau DT, Sosa P. Disparate impact of the COVID-19 pandemic and health equity data gaps. 16. Am J Public Health. 2022;112(10):1404-6. https://doi.org/10.2105/AJPH.2022.307052 World Health Organization. COVID-19 and the social determinants of health and health 17. equity: evidence brief. Geneva: WHO, Dec 6, 2021. Available from: https://www.who.int/publications/i/item/9789240038387 [accessed 5 January 2023]. Mackey K, Ayers CK, Kondo KK, Saha S, Advani SM, Young S, et al. Racial and ethnic disparities in COVID-19-related infections, hospitalizations, and deaths: a systematic review. Ann Intern Med. 2021;174(3):362-73. https://doi.org/10.7326/M20-6306 Mude W, Oguoma VM, Nyanhanda T, Mwanri L, Njue C. Racial disparities in COVID-19 pandemic cases, hospitalisations, and deaths: a systematic review and meta-analysis. J Glob Health. 2021;11:05015. https://doi.org/10.7189/jogh.11.05015 Hernandez I, Dickson S, Tang S, Gabriel N, Berenbrok LA, Guo J. Disparities in distribution of COVID-19 vaccines across US counties: a geographic information system-

- based cross-sectional study. PLoS Med. 2022;19(7):e1004069.
- https://doi.org/10.1371/journal.pmed.1004069
- 620 21. Siegel M, Critchfield-Jain I, Boykin M, Owens A, Muratore R, Nunn T, et al. Racial/ethnic
- disparities in state-level COVID-19 vaccination rates and their association with structural
- racism. J Racial Ethn Health Disparities. 2022;9(6):2361-74.
- 623 https://doi.org/10.1007/s40615-021-01173-7
- 624 22. Centers for Disease Control and Prevention. COVID data tracker. CDC, 2022. Available
- from: https://covid.cdc.gov/covid-data-tracker/#vaccination-demographics-trends [accessed]
- 626 5 January 2023].
- 627 23. Underhill K, Johnson OCA. Vaccination equity by design. Yale Law J. 2021;131:53-88.
- 628 24. Victora CG, Joseph G, Silva ICM, Maia FS, Vaughan JP, et al. The inverse equity
- hypothesis: analyses of institutional deliveries in 286 national surveys. Am J Public Health.
- 630 2018;108(4):464-71. https://doi.org/10.2105/AJPH.2017.304277
- 631 25. Khubchandani J, Macias Y. COVID-19 vaccination hesitancy in Hispanics and African-
- Americans: a review and recommendations for practice. Brain Behav Immun Health.
- 633 2021;15:100277. https://doi.org/10.1016/j.bbih.2021.100277
- 634 26. Kricorian K, Turner K. COVID-19 vaccine acceptance and beliefs among Black and
- Hispanic Americans. PLoS One. 2021;16(8):e0256122.
- https://doi.org/10.1371/journal.pone.0256122
- 637 27. Mohottige D, Boulware LE. Trust in American medicine: a call to action for health care
- 638 professionals. Hastings Cent Rep. 2020;50(1):27-9. https://doi.org/10.1002/hast.1081

- 639 28. Benkert R, Cuevas A, Thompson HS, Dove-Meadows E, Knuckles D. Ubiquitous yet
- unclear: a systematic review of medical mistrust. Behav Med. 2019;45(2):86-101.
- https://doi.org/10.1080/08964289.2019.1588220
- 642 29. Jaiswal J, Halkitis PN. Towards a more inclusive and dynamic understanding of medical
- mistrust informed by science. Behav Med. 2019;45(2):79-85.
- 644 https://doi.org/10.1080/08964289.2019.1619511
- 645 30. Williams DR, Mohammed SA. Racism and health I: pathways and scientific evidence. Am
- Behav Sci. 2013;57(8):1152-73. https://doi.org/10.1177/0002764213487340
- 31. Schuster M, Eskola J, Duclos P; SAGE Working Group on Vaccine Hesitancy. Review of
- vaccine hesitancy: Rationale, remit and methods. Vaccine. 2015;33(34):4157-60.
- https://doi.org/10.1016/j.vaccine.2015.04.035
- 650 32. Larson HJ, Gakidou E, Murray CJL. The vaccine-hesitant moment. N Engl J Med.
- 651 2022;387(1):58-65. https://doi.org/10.1056/nejmra2106441
- 652 33. Dubé E, Laberge C, Guay M, Bramadat P, Roy R, Bettinger J. Vaccine hesitancy: an
- overview. Hum Vaccin Immunother. 2013;9(8):1763-73. https://doi.org/10.4161/hv.24657
- 654 34. Bedford H, Attwell K, Danchin M, Marshall H, Corben P, Leask J. Vaccine hesitancy,
- refusal and access barriers: the need for clarity in terminology. Vaccine. 2018;36(44):6556-
- 8. https://doi.org/10.1016/j.vaccine.2017.08.004
- 657 35. Colquhoun HL, Levac D, O'Brien KK, Straus S, Tricco AC, Perrier L, et al. Scoping
- reviews: time for clarity in definition, methods, and reporting. J Clin Epidemiol.
- 659 2014;67(12):1291-4. https://doi.org/10.1016/j.jclinepi.2014.03.013

- 660 36. Newman PA, Reid L, Tepjan S, Fantus S, Allan K, Nyoni T, et al. COVID-19 vaccine
- hesitancy among marginalized populations in the U.S. and Canada: protocol for a scoping
- review. PLoS One. 2022;17(3):e0266120. https://doi.org/10.1371/journal.pone.0266120
- 663 37. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. Int J Soc
- Res Methodol. 2005;8(1):19-32. https://doi.org/10.1080/1364557032000119616
- 38. Joanna Briggs Institute. Joanna Briggs institute reviewers' manual: 2015
- edition/supplement. Adelaide: The Joanna Briggs Institute; 2015.
- 39. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA
- Extension for Scoping Reviews (PRISMA-ScR): checklist and explanation. Ann Intern
- Med. 2018;169(7):467-73. https://doi.org/10.7326/m18-0850
- 40. U.S. Department of Health & Human Services. Coronavirus: COVID-19 vaccines.
- Washington: HHS, 2022 Dec 12. Available from: https://www.hhs.gov/coronavirus/covid-
- 672 <u>19-vaccines/index.html</u> [accessed 5 January 2023].
- 41. Health Canada. Health Canada authorizes first COVID-19 vaccine. Ottawa: Government of
- Canada; 2020 Dec 9. Available from: https://www.canada.ca/en/health-
- 675 <u>canada/news/2020/12/health-canada-authorizes-first-covid-19-vaccine0.html</u> [accessed 5
- 676 January 2023].
- 677 42. Palmer Kelly E, McGee J, Obeng-Gyasi S, Herbert C, Azap R, Abbas A, et al.
- Marginalized patient identities and the patient-physician relationship in the cancer care
- context: a systematic scoping review. Support Care Cancer. 2021;29(12):7195-207.
- 680 https://doi.org/10.1007/s00520-021-06382-8
- 43. Landis JR, Koch GG. The measurement of observer agreement for categorical data.
- Biometrics. 1977;33(1):159-74. https://pubmed.ncbi.nlm.nih.gov/843571/

- 683 44. Tricco AC, Antony J, Zarin W, Strifler L, Ghassemi M, Ivory J, et al. A scoping review of
- rapid review methods. BMC Med. 2015;13:224. https://doi.org/10.1186/s12916-015-0465-
- 685
- 686 45. Garritty C, Gartlehner G, Nussbaumer-Streit B, King VJ, Hamel C, Kamel C, et al.
- Cochrane Rapid Reviews Methods Group offers evidence-informed guidance to conduct
- rapid reviews. J Clin Epidemiol. 2021;130:13-22.
- https://doi.org/10.1016/j.jclinepi.2020.10.007
- 690 46. Allen JD, Abuelezam NN, Rose R, Fontenot HB. Factors associated with the intention to
- obtain a COVID-19 vaccine among a racially/ethnically diverse sample of women in the
- USA. Transl Behav Med. 2021;11(3):785-92. https://doi.org/10.1093/tbm/ibab014
- 693 47. Arvanitis M, Opsasnick L, O'Conor R, Curtis LM, Vuyyuru C, Yoshino Benavente J, et al.
- Factors associated with COVID-19 vaccine trust and hesitancy among adults with chronic
- conditions. Prev Med Rep. 2021;24:101484. https://doi.org/10.1016/j.pmedr.2021.101484
- 696 48. Balasuriya L, Santilli A, Morone J, Ainooson J, Roy B, Njoku A, et al. COVID-19 vaccine
- acceptance and access among Black and Latinx communities. JAMA Netw Open.
- 698 2021;4(10):e2128575. https://doi.org/10.1001/jamanetworkopen.2021.28575
- 699 49. Battarbee AN, Stockwell MS, Varner M, Newes-Adeyi G, Daugherty M, Gyamfi-
- Bannerman C, et al. Attitudes toward COVID-19 illness and COVID-19 vaccination among
- pregnant women: a cross-sectional multicenter study during August-December 2020. Am J
- Perinatol. 2022;39(01):075-083. https://doi.org/10.1055/s-0041-1735878
- 703 50. Bauer C, Zhang K, Lee M, Jones M, Rodriguez A, de la Cerda I, et al. Real-time geospatial
- analysis identifies gaps in COVID-19 vaccination in a minority population. Sci Rep.
- 705 2021;11(1):18117. https://doi.org/10.1038/s41598-021-97416-y

- 706 51. Bogart LM, Ojikutu BO, Tyagi K, Klein DJ, Mutchler MG, Dong L, et al. COVID-19
- related medical mistrust, health impacts, and potential vaccine hesitancy among Black
- Americans living with HIV. J Acquir Immune Defic Syndr. 2021;86(2):200-7.
- 709 https://doi.org/10.1097/gai.000000000002570
- 710 52. Bogart LM, Dong L, Gandhi P, Klein DJ, Smith TL, Ryan S, et al. COVID-19 vaccine
- 711 intentions and mistrust in a national sample of Black Americans. J Natl Med Assoc.
- 712 2022;113(6):599-611. https://doi.org/10.1016/j.jnma.2021.05.011
- 713 53. Carson SL, Casillas A, Castellon-Lopez Y, Mansfield LN, Morris D, Barron J, et al.
- 714 COVID-19 vaccine decision-making factors in racial and ethnic minority communities in
- Los Angeles, California. JAMA Netw Open. 2021;4(9):e2127582.
- 716 https://doi.org/10.1001/jamanetworkopen.2021.27582
- 717 54. Chin ET, Leidner D, Ryckman T, Liu YE, Prince L, Alarid-Escudero F, et al. Covid-19
- vaccine acceptance in California state prisons. N Engl J Med. 2021;385(4):374-6.
- 719 https://doi.org/10.1056/NEJMc2105282
- 720 55. Clark LR, Sawkin MT, Melton BL, Meyer EG. The COVID-19 vaccine conundrum: an
- assessment of vaccine hesitancy amongst patients at a federally qualified health center. J
- Pharm Pract. 2022:8971900221116193. https://doi.org/10.1177/08971900221116193
- 56. Crozier J, Christensen N, Li P, Stanley G, Clark DS, Selleck C. Rural, underserved, and
- minority populations' perceptions of COVID-19 information, testing, and vaccination:
- report from a southern state. Popul Health Manag. 2022;25(3):413-22.
- 726 https://doi.org/10.1089/pop.2021.0216

- 57. Ehde DM, Roberts MK, Herring TE, Alschuler KN. Willingness to obtain COVID-19
- vaccination in adults with multiple sclerosis in the United States. Mult Scler Relat Disord.
- 729 2021;49:102788. https://doi.org/10.1016/j.msard.2021.102788
- 730 58. Ehde DM, Roberts MK, Humbert AT, Herring TE, Alschuler KN. COVID-19 vaccine
- hesitancy in adults with multiple sclerosis in the United States: a follow up survey during
- the initial vaccine rollout in 2021. Mult Scler Relat Disord. 2021;54:103163.
- 733 https://doi.org/10.1016/j.msard.2021.103163
- 734 59. Fernández-Penny FE, Jolkovsky EL, Shofer FS, Hemmert KC, Valiuddin H, Uspal JE, et
- al. COVID-19 vaccine hesitancy among patients in two urban emergency departments.
- 736 Acad Emerg Med. 2021;28(10):1100-7. https://doi.org/10.1111/acem.14376
- 737 60. Fisher KA, Nguyen N, Crawford S, Fouayzi H, Singh S, Mazor KM. Preferences for
- COVID-19 vaccination information and location: associations with vaccine hesitancy, race
- and ethnicity. Vaccine. 2021;39(45):6591-4. https://doi.org/10.1016/j.vaccine.2021.09.058
- 740 61. Garcia J, Vargas N, de la Torre C, Magana Alvarez M, Clark JL. Engaging Latino families
- about COVID-19 vaccines: a qualitative study conducted in Oregon, USA. Health Educ
- 742 Behav. 2021;48(6):747-57. https://doi.org/10.1177/10901981211045937
- 62. Garcia P, Montez-Rath ME, Moore H, Flotte J, Fults C, Block MS, et al. SARS-CoV-2
- vaccine acceptability in patients on hemodialysis: a nationwide survey. J Am Soc Nephrol.
- 745 2021;32(7):1575–81. https://doi.org/10.1681/asn.2021010104
- 63. Geana MV, Anderson S, Ramaswamy M. COVID-19 vaccine hesitancy among women
- leaving jails: a qualitative study. Public Health Nurs. 2021;38(5):892-6.
- 748 https://doi.org/10.1111/phn.12922

- 749 64. Ghaffari-Rafi A, Teehera KB, Higashihara TJ, Morden FTC, Goo C, Pang M, et al.
- Variables associated with coronavirus disease 2019 vaccine hesitancy amongst patients
- with neurological disorders. Infect Dis Rep. 2021;13(3):763-810.
- 752 https://doi.org/10.3390/idr13030072
- 753 65. Guidry JPD, Laestadius LI, Vraga EK, Miller CA, Perrin PB, Burton CW, et al.
- Willingness to get the COVID-19 vaccine with and without emergency use authorization.
- 755 Am J Infect Control. 2021 Feb;49(2):137-42. https://doi.org/10.1016/j.ajic.2020.11.018
- 756 66. Hagan LM, Dusseau C, Crockett M, Rodriguez T, Long MJ. COVID-19 vaccination in the
- 757 Federal Bureau of Prisons, December 2020-April 2021. Vaccine. 2021;39(40):5883-90.
- 758 https://doi.org/10.1016/j.vaccine.2021.08.045
- 759 67. Iadarola S, Siegel JF, Gao Q, McGrath K, Bonuck KA. COVID-19 vaccine perceptions in
- New York State's intellectual and developmental disabilities community. Disabil Health J.
- 761 2022;15(1):101178. https://doi.org/10.1016/j.dhjo.2021.101178
- 762 68. Jimenez ME, Rivera-Núñez Z, Crabtree BF, Hill D, Pellerano MB, Devance D, et al. Black
- and Latinx community perspectives on COVID-19 mitigation behaviors, testing, and
- 764 vaccines. JAMA Netw Open. 2021;4(7):e2117074.
- 765 https://doi.org/10.1001/jamanetworkopen.2021.17074
- 766 69. Johnson KD, Akingbola O, Anderson J, Hart J, Chapple A, Woods C, Yeary K, McLean A.
- Combatting a "Twin-demic": a quantitative assessment of COVID-19 and influenza
- vaccine hesitancy in primary care patients. Health Promot Perspect. 2021;11(2):179-85.
- 769 https://doi.org/10.34172/hpp.2021.22
- 770 70. Jones DL, Salazar AS, Rodriguez VJ, Balise RR, Starita CU, Morgan K, et al. Severe acute
- respiratory syndrome coronavirus 2: vaccine hesitancy among underrepresented racial and

- ethnic groups with HIV in Miami, Florida. Open Forum Infect Dis. 2021;8(6):ofab154.
- 773 https://doi.org/10.1093/ofid/ofab154
- 774 71. Kuhn R, Henwood B, Lawton A, Kleva M, Murali K, King C, et al. COVID-19 vaccine
- access and attitudes among people experiencing homelessness from pilot mobile phone
- 376 survey in Los Angeles, CA. PLoS One. 2021;16(7):e0255246.
- 777 https://doi.org/10.1371/journal.pone.0255246
- 778 72. Luo H, Qu H, Basu R, Rafferty AP, Patil SP, Cummings DM. Willingness to get a COVID-
- 19 vaccine and reasons for hesitancy among Medicare beneficiaries: results from a national
- survey. J Public Health Manag Pract. 2022;28(1):70-6.
- 781 https://doi.org/10.1097/phh.00000000001394
- 782 73. Masson CL, McCuistian C, Straus E, Elahi S, Chen M, Gruber VA, et al. COVID-19
- vaccine trust among clients in a sample of California residential substance use treatment
- 784 programs. Drug Alcohol Depend. 2021;225:108812.
- 785 https://doi.org/10.1016/j.drugalcdep.2021.108812
- 786 74. Momplaisir F, Haynes N, Nkwihoreze H, Nelson M, Werner RM, Jemmott J.
- 787 Understanding drivers of coronavirus disease 2019 vaccine hesitancy among Blacks. Clin
- 788 Infect Dis. 2021;73(10):1784-9. https://doi.org/10.1093/cid/ciab102
- 789 75. Moore JX, Gilbert KL, Lively KL, Laurent C, Chawla R, Li C, et al. Correlates of COVID-
- 790 19 vaccine hesitancy among a community sample of African Americans living in the
- southern United States. Vaccines. 2021;9(8):879. https://doi.org/10.3390/vaccines9080879
- 792 76. Myers A, Ipsen C, Lissau A. COVID-19 vaccination hesitancy among Americans with
- disabilities aged 18-65: an exploratory analysis. Disabil Health J. 2022;15(1):101223.
- 794 https://doi.org/10.1016/j.dhjo.2021.101223

795 77. Okoro O, Kennedy J, Simmons G Jr, Vosen EC, Allen K, Singer D, et al. Exploring the 796 scope and dimensions of vaccine hesitancy and resistance to enhance COVID-19 797 vaccination in Black communities. J Racial Ethn Health Disparities. 2022;9(6):2117-30. 798 https://doi.org/10.1007/s40615-021-01150-0 799 78. Razzaghi H, Meghani M, Pingali C, Crane B, Naleway A, Weintraub E, et al. COVID-19 800 vaccination coverage among pregnant women during pregnancy - eight integrated health 801 care organizations, United States, December 14, 2020-May 8, 2021. MMWR Morb Mortal 802 Wkly Rep. 2021;70(24):895-9. https://doi.org/10.15585/mmwr.mm7024e2 803 Rodriguez RM, Torres JR, Chang AM, Haggins AN, Eucker SA, O'Laughlin KN, et al. The 804 rapid evaluation of COVID-19 vaccination in emergency departments for underserved 805 patients study. Ann Emerg Med. 2021;78(4):502-10. 806 https://doi.org/10.1016/j.annemergmed.2021.05.026 807 Rodriguez VJ, Alcaide ML, Salazar AS, Montgomerie EK, Maddalon MJ, Jones DL. 80. 808 Psychometric properties of a vaccine hesitancy scale adapted for COVID-19 vaccination 809 among people with HIV. AIDS Behav. 2022;26(1):96-101. https://doi.org/10.1007/s10461-810 021-03350-5 811 81. Rungkitwattanakul D, Yabusaki A, Singh D, Lawson P, Nwaogwugwu U, Iheagwara OS, 812 et al. COVID-19 vaccine hesitancy among African American hemodialysis patients: a 813 single-center experience. Hemodial Int. 2021;25(3):410-2. 814 https://doi.org/10.1111/hdi.12922 815 Salmon DA, Dudley MZ, Brewer J, Kan L, Gerber JE, Budigan H, et al. COVID-19 82. 816 vaccination attitudes, values and intentions among United States adults prior to emergency

817

818

819

820

821

822

823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

838


839


use authorization. Vaccine. 2021;39(19):2698-711. https://doi.org/10.1016/j.vaccine.2021.03.034 Stephenson R, Sullivan SP, Pitter RA, Hunter AS, Chavanduka TM, COVID-19 pandemic optimism and vaccine willingness among an online sample of US gay, bisexual, and other men who have sex with men. Vaccines. 2021;9(7):745. https://doi.org/10.3390/vaccines9070745 Stern MF, Piasecki AM, Strick LB, Rajeshwar P, Tyagi E, Dolovich S, et al. Willingness to 84. receive a COVID-19 vaccination among incarcerated or detained persons in correctional and detention facilities - four states, September-December 2020. MMWR Morb Mortal Wkly Rep. 2021;70(13):473-7. https://doi.org/10.15585/mmwr.mm7013a3 85. Stoler J, Enders AM, Klofstad CA, Uscinski JE. The limits of medical trust in mitigating COVID-19 vaccine hesitancy among Black Americans. J Gen Intern Med. 2021;36(11):3629-31. https://doi.org/10.1007/s11606-021-06743-3 Teixeira da Silva D, Biello K, Lin WY, Valente PK, Mayer KH, Hightow-Weidman L, et al. COVID-19 vaccine acceptance among an online sample of sexual and gender minority men and transgender women. Vaccines. 2021;9(3):204. https://doi.org/10.3390/vaccines9030204 87. Tsapepas D, Husain SA, King KL, Burgos Y, Cohen DJ, Mohan S. Perspectives on COVID-19 vaccination among kidney and pancreas transplant recipients living in New York City. Am J Health Syst Pharm. 2021;78(22):2040-5. https://doi.org/10.1093/ajhp/zxab272 Uhr L, Mateen FJ. COVID-19 vaccine hesitancy in multiple sclerosis: a cross-sectional 88. survey. Mult Scler. 2022;28(7):1072-80. https://doi.org/10.1177/13524585211030647

- 840 89. Wagner AL, Sheinfeld Gorin S, Boulton ML, Glover BA, Morenoff JD. Effect of vaccine
- effectiveness and safety on COVID-19 vaccine acceptance in Detroit, Michigan, July 2020.
- 842 Hum Vaccin Immunother. 2021;17(9):2940-5.
- https://doi.org/10.1080/21645515.2021.1917233
- 844 90. Wang SX, Bell-Rogers N; FNP-c, Dillard D, Harrington MA. COVID-19 vaccine hesitancy
- in Delaware's underserved communities. Dela J Public Health. 2021;7(4):168-75. doi:
- 846 10.32481/djph.2021.09.022
- 91. Xiang XM, Hollen C, Yang Q, Brumbach BH, Spain RI, Wooliscroft L. COVID-19
- vaccination willingness among people with multiple sclerosis. Mult Scler J Exp Transl
- 849 Clin. 2021;7(2):20552173211017159. https://doi.org/10.1177/20552173211017159
- 92. Yang Y, Dobalian A, Ward KD. COVID-19 vaccine hesitancy and its determinants among
- adults with a history of tobacco or marijuana use. J Community Health. 2021;46(6):1090-8.
- https://doi.org/10.1007/s10900-021-00993-2.
- 853 93. Zhang M, Gurung A, Anglewicz P, Subedi P, Payton C, Ali A, et al. Acceptance of
- COVID-19 vaccine among refugees in the United States. Public Health Rep.
- 855 2021;136(6):774-81. https://doi.org/10.1177/00333549211045838
- 856 94. Ramsetty A, Adams C. Impact of the digital divide in the age of COVID-19. J Am Med
- 857 Informatics Assoc. 2020;22(7):1147-8. https://doi.org/10.1093/jamia/ocaa078
- 858 95. Ratzan S, Schneider EC, Hatch H, Cacchione J. Missing the point how primary care can
- overcome Covid-19 vaccine "hesitancy". N Engl J Med. 2021;384(25):e100.
- https://doi.org/10.1056/nejmp2106137

861 96. Dada D, Djiometio JN, McFadden SM, Demeke J, Vlahov D, Wilton L, et al. Strategies 862 that promote equity in COVID-19 vaccine uptake for Black communities: a review. J 863 Urban Health. 2022;99(1):15-27. https://doi.org/10.1007/s11524-021-00594-3 864 97. Rodriguez RM, Nichol G, Eucker SA, Chang AM, O'Laughlin KN, Pauley A, et al. Effect 865 of COVID-19 vaccine messaging platforms in emergency departments on vaccine 866 acceptance and uptake: a cluster randomized clinical trial. JAMA Intern Med. 2022. 867 https://doi.org/10.1001/jamainternmed.2022.5909 868 Peretti-Watel P, Larson HJ, Ward JK, Schulz WS, Verger P. Vaccine hesitancy: clarifying 98. 869 a theoretical framework for an ambiguous notion. PLoS Curr. 870 2015;7:ecurrents.outbreaks.6844c80ff9f5b273f34c91f71b7fc289. 871 https://doi.org/10.1371/currents.outbreaks.6844c80ff9f5b273f34c91f71b7fc289 872 99. Sturgis P, Brunton-Smith I, Jackson J. Trust in science, social consensus and vaccine 873 confidence. Nat Hum Behav. 2021;5(11):1528-34. https://doi.org/10.1038/s41562-021-874 01115-7 875 100. Rubincam C, Lacombe-Duncan A, Newman PA. Taking culture seriously in biomedical 876 HIV prevention trials: a meta-synthesis of qualitative studies. Expert Rev Vaccines. 877 2015;15(3):331-47. https://doi.org/10.1586/14760584.2016.1118349 878 101. Demeke J, Ramos SR, McFadden SM, Dada D, Nguemo Djiometio J, Vlahov D, et al. 879 Strategies that promote equity in COVID-19 vaccine uptake for Latinx communities: a 880 review. J Racial Ethn Health Disparities. 2022:1-9. https://doi.org/10.1007/s40615-022-881 <u>01320-8</u> 882 102. U.S. Department of Health and Human Services. Campaign approach to reaching Black 883 audiences. Washington: HHS, 2022 Sep 19. Available from:

884 https://wecandothis.hhs.gov/resource/campaign-approach-to-reaching-black-audiences 885 [accessed 5 January 2023]. 886 Centers for Disease Control and Prevention. Guidance for vaccinating older adults and 887 people with disabilities: ensuring equitable COVID-19 vaccine access. Atlanta: CDC, 2022 888 Jan 20. Available from: https://www.cdc.gov/vaccines/covid-19/clinical-889 considerations/older-adults-and-disability/access.html [accessed 5 January 2023]. 890 **Supporting information** 891 892 S1 Table. Sample search string





medRxiv preprint doi: https://doi.org/10.1101/2023.02.23.23286342; this version posted February 23, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.