Coping with drug resistant tuberculosis alongside COVID-19 and other stressors in Zimbabwe: a qualitative study.

4 Collins Timire ^{1,2,3}

1

2

- 5 Katharina Kranzer ^{1,3,4}
- 6 Debora Pedrazzoli ⁵
- 7 Fungai Kavenga²
- 8 Samuel Kasozi ^{2,6}
- 9 Fredrick Mbiba³
- 10 Virginia Bond ^{7,8}
- 11 Clinical Research Department, Faculty of Infectious & Tropical Diseases, London School of
- 12 Hygiene & Tropical Medicine (LSHTM), London, UK
- ² AIDS & TB Department, Ministry of Health and Child Care, Harare, Zimbabwe
- ³ The Health Research Unit; Biomedical Research & Training Institute, Harare, Zimbabwe
- ⁴ Division of Infectious Diseases & Tropical Medicine, Medical Centre of the University of
- 16 Munich, Munich, Germany.
- ⁵ Department of Infectious Disease Epidemiology, Faculty of Global and Public Health, London
- 18 School of Hygiene & Tropical Medicine, UK
- 19 ⁶ The Global Fund To Fight AIDS, TB and Malaria, Geneva, Switzerland
- ⁷ Department of Global Health and Development, Faculty of Public Health and Policy, LSHTM,
- 21 London, UK
- 22 ⁸ Zambart, Lusaka, Zambia
- 23 Corresponding author details
- 24 Collins TIMIRE

29

- 25 Collins.Timire@lshtm.ac.uk +4477 1972 1145 or +263777687328
- Paper contents: Abstract=299 words; Main text=5371 words; References=56
- 27 **Keywords:** Multi-drug resistant tuberculosis; Zimbabwe; COVID-19; financial catastrophe.
- 28 **Short title:** Coping with drug resistant tuberculosis and COVID-19 in Zimbabwe

30	ORCID numbers	
31	Collins Timire	https://orcid.org/0000-0003-0997-9799
32	Katharina Kranzer	https://orcid.org/0000-0001-5691-7270
33	Debora Pedrazzoli	https://orcid.org/0000-0002-6434-4457
34	Fungai Kavenga	
35	Samuel Kasozi	https://orcid.org/0000-0001-9467-3692
36	Fredrick Mbiba	https://orcid.org/0000-0002-1666-2979
37	Virginia Bond	http://orcid.org/0000-0002-6815-4239
38		
39		

Abstract 40 **Background** 41 42 Households in low-resource settings are more vulnerable to events which adversely affect their livelihoods, including shocks such as the death of a family member, inflation, droughts and more 43 recently COVID-19. Drug Resistant Tuberculosis (DR-TB) is also another shock that inflicts 44 physical, psychological and socioeconomic burden on individuals and households. We describe 45 experiences and coping strategies among people affected by DR-TB and their households in 46 Zimbabwe during the COVID-19 pandemic, 2020 to 2021. 47 Methods 48 We conducted 16 in-depth interviews with adults who had just completed or were completing 49 50 treatment. Interview themes included health seeking behaviour, impact of DR-TB on livelihoods 51 and coping strategies adopted during treatment. We analysed data using thematic analyses. 52 **Results** 53 Health seeking from providers outside the public sector, extra-pulmonary TB and health system factors resulted in delayed DR-TB diagnosis and treatment and increased financial drain on 54 households. DR-TB reduced productive capacity and narrowed job opportunities leading to 55 income loss that continued even after completion of treatment. Household livelihood was further 56 adversely affected by lockdowns due to COVID-19, outbreaks of bird flu and cattle disease. 57 58 Stockouts of DR-TB medicines, common during COVID-19, exacerbated loss of productive time 59 and transport costs as medication had to be accessed from other clinics that were further away. Reversible coping strategies included: reducing number of meals; relocating in search of 60 61 caregivers and/or family support; spending savings; negotiating with school authorities to keep children in school. Some households had to adopt irreversible coping strategies such as selling 62 63 productive assets and withdrawing children from school. Conclusion 64 DR-TB combined with COVID-19 and other stressors pushed households into deeper poverty, 65 and vulnerability. Multi-sectoral approaches that combine health systems, psychosocial and 66 67 economic interventions are crucial to mitigate diagnostic delays and suffering, and meaningfully support people with DR-TB and their households to compensate the loss of livelihoods during 68 and post DR-TB treatment. 69

70 **Background** Tuberculosis (TB) is a global public health challenge, and until the COVID-19 pandemic, was 71 72 the leading cause of death from a single infectious disease (ranking above HIV).(1) An estimated 73 10.6 million people fell ill with TB in 2021 and 1.6 million (15%) of them died.(1) Treatment 74 success (cure and treatment completion) for people with drug resistant TB (DR-TB) is worse, (averaging around 60% globally and around 54% in Zimbabwe)(2-4), compared to 86% for 75 76 people with drug-susceptible TB (DS-TB).(1,5) People with DR-TB have higher risk of loss to follow up, death and post-TB disabilities including poor lung function, auditory complications 77 and, depending on site of infection, bone deformities.(6–8) 78 79 Successful treatment outcomes among people with DR-TB depend on completing appropriate 80 and efficacious course of medicines. Until very recently, treatment of DR-TB was much longer 81 and more toxic than that of drug susceptible TB (DS-TB), lasting up to 18-24 months. As a 82 result, people affected by DR-TB and their households often have to overcome even greater health system, psychosocial and socioeconomic barriers over an extended time before 83 completing treatment.(9–12) Surveys of cost incurred by TB patients and their households have 84 shown that a higher proportion of people affected by DR-TB incur catastrophic costs than those 85 86 affected by DS-TB (82% vs 47%).(13–15) While the greater proportion of TB related costs are indirect costs such as income loss due to reduced productivity and time spent health seeking, (16) 87 88 people and their households also incur out-of-pocket costs which may be substantial to poor people who are less likely to have medical insurance, savings or stable earnings.(17) Health 89 system approaches that have been implemented globally to reduce TB related costs include: i) 90 early diagnosis of TB through systematic screening and universal access to drug susceptibility 91 92 testing (18), ii) decentralisation of TB treatment to reduce transport costs, iii) provision of free TB medication and diagnostics and iv) provision of psycho-social and economic support. 93 94 Notwithstanding these interventions, costs of hospitalisation, radiology services (chest X-rays), ancillary medicines, consultations and blood tests are often not considered, (19) and individuals 95 96 and their primary caregivers also incur substantial income loss (or indirect costs) related to taking time off work to seek and receive TB care.(20,21) Such indirect costs, together with costs 97 98 related to food and transport to/from the health facility, constitute the major cost drivers.(19–21) In addition to economic losses, households also experience worsening social relations and 99 stigma.(24) Further shocks and stressors may affect households either concurrently or 100

102

103

104

105

106

107

108109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126 127

128

129

130

sequentially.(25) Such shocks may be spatial or covariate (e.g., COVID-19, drought, floods) when they affect communities, nations, regions or the whole world or may be temporal based on their occurrence (annual, seasonal).(26) Often shocks have synergistic relationships that determine household coping strategies.(25) Poor and distressed households — usually without medical insurance or savings — may adopt reversible coping strategies such as delaying health seeking for chronic diseases, relocating in search of food or caregivers and spending savings, and faced with acute infections or global shocks, may be forced to mobilise and expend their resources rapidly.(27) Under such circumstances, they may be forced to adopt irreversible coping strategies such as taking loans at exorbitant interests, withdrawing children from school and selling productive assets, thereby deepening their vulnerability to current and future shocks.(27,28) COVID-19 is a global stressor and has been shown to be a stress multiplier.(29) Poor households that experience a convergence of DR-TB, COVID-19 and other stressors may have both a limited choice of coping strategies and limited coping capacity.(30) Zimbabwe, a resource limited country in sub Saharan Africa wracked with economic hardship since 2000, had an incidence of TB and DR-TB of 190 per 100,000 population and 4.9 per 100,000 population, respectively, in 2021, and TB/HIV co-infection was 50%.(5,31,32) Around 90% of the population is unemployed and earn a living through informal jobs (subsistence farming, informal businesses as vendors and cross border trading), and 72% live in poverty.(33) From March 2020-August 2020, the Zimbabwean government imposed a stringent lockdown with the aim to control the spread of COVID-19. During this time, businesses were shut and public transport almost completely ceased. Poor people, especially those in urban areas, faced an increased risk of contracting COVID-19 owing to overcrowded living conditions (34) and bore the brunt of lockdowns since they mostly live from daily earnings and are informally employed with no social security. In the absence of social support, COVID-19 thus negatively affected household income and food security. It is well established that poverty fuels both COVID-19 and TB and in turn, the two diseases deepen impoverishment and irreversible coping strategies. (27,30) Previous studies have looked at the impact of DR-TB, but not in the context of global shocks such as COVID-19.(9,35,36) COVID-19 affected whole populations and impacted the health systems, livelihood and transportation. To study the joint impact of DR-TB, COVID-19 and other stressors, we conducted 16 in-depth interviews with people affected by DR-TB in

- 231 Zimbabwe in 2020-2021 to trace their experiences and coping strategies during the pandemic
- and their period of treatment.

Methods 133 *Management of DR-TB in Zimbabwe* 134 135 In Zimbabwe, DR-TB care is provided at primary health care level. Prior to 2017, treatment duration was 18-24 months with daily injections for the first six months. From 2017, shorter (9-136 137 12 months) injection-based regimens were introduced followed by the introduction of a short all oral 9 months regimen in 2020.(37) Xpert MTB/Rif (Cepheid, Sunnyvale, CA, USA) is the 138 139 primary diagnostic test for people presenting with signs and symptoms of TB enabling not just diagnosis of TB, but also detection of rifampicin resistance.(37) People on DR-TB treatment are 140 eligible for USD 25 conditional cash transfers and an additional USD 20 worth of grocery 141 vouchers (from 2021 onwards) every month for the duration of treatment. However, delays and 142 inconsistencies in cash disbursements and food hampers have been reported especially during the 143 COVID-19 pandemic, with only two-thirds (64%) of those registered effectively receiving 144 cash.(38) 145 Study design, population and recruitment 146 147 This was a qualitative phenomenological study to gather lived experiences of people with DR-148 TB. Men and women aged ≥18 years who were within two months of completing DR-TB treatment or had completed treatment within two months were identified from TB registers 149 150 within health facilities in Harare (urban) and Matebeleland South (rural) provinces in Zimbabwe. 151 The study team met or called the potential participant and appraised them about the study. Those 152 who expressed willingness to participate in the study were asked to identify a convenient place and time so that the study team could obtain written informed consent and conduct in-depth 153 154 interviews. 155 Data collection 156 In-depth interviews were conducted from October 2020 to March 2021 by CT (first author) and FM (social scientist), with CT conducting 12 interviews and FM four interviews. The interview 157 158 guide, translated into two local languages (Shona and Ndebele), included the following themes: 159 health seeking patterns, experiences with DR-TB treatment and DR-TB disease, and coping strategies adopted during treatment. The interviews lasted 35-45 minutes and were conducted in 160 Shona or Ndebele. All interviews were audio-recorded and were conducted in private locations 161 162 suggested by participants. Participants received USD3 for refreshments, and bus fares were also

reimbursed. After each interview, the two researchers (CT and FM) had debriefing meetings to discuss the findings.

Ethical approval

Ethical approval was obtained from the London School of Hygiene and Tropical Medicine Research Ethics Committee (Ref 22579), the Biomedical Research and Training Institute Institutional Review Board (AP160/2020) and the Medical Research Council of Zimbabwe (MRCZ/A/2645). Permission to access DR-TB registers was obtained from the Permanent Secretary for Health, Zimbabwe. All participants gave written informed consent to take part in the study and for the use of an audio-recorder. Names used in quotes and tables in this paper are pseudonyms.

Data analysis

All audio-recordings were transcribed into local language before translating the transcripts into English. Data were analyzed using thematic analysis. First, the transcripts were read several times aiming to get familiar with the data. Second, data were coded manually and using NVivo version 12 software (QSR International). The codes were both data driven and concept driven,

(39,40) and were grouped into themes to form thematic maps.(41) Divergent views were noted.

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199 200

201

202

203

204

205

206

207

208209

Results Sixteen people (eight women) were enrolled. Twelve were co-infected with HIV. Their characteristics are summarized in **Table 1**. We first describe the experiences of people with DR-TB starting from delays in treatment initiation (due to underdiagnoses, misdiagnosis of TB or health seeking from alternative providers), physical impact of DR-TB disease and treatment itself, and changes in livelihoods as a result of DR-TB episodes. Next we explore the coping strategies adopted in response to DR-TB alongside other shocks, including COVID-19. Delays in diagnosis and starting DR-TB treatment Ten of the 16 participants experienced delays of up to four months before they received a diagnosis and started DR-TB treatment. Reasons for delays included misdiagnosis, HIV coinfection and health seeking from alternative health providers such as traditional healers. Health system diagnostic delays Reasons for delays in diagnosing DR-TB were multifactorial including the use of tests with low sensitivity (i.e., smear microscopy), lack of healthcare worker awareness and limited health system resources. The use of smear microscopy instead of Xpert MTB/Rif resulted in underdiagnosis of TB and DR-TB. Smear microscopy has a lower sensitivity than Xpert MTB/Rif, especially in people living with HIV, and additionally cannot detect rifampicin resistance. That's when I went back to the same clinic and to the same nurses who had counselled me and I told them that I had not gained weight and they said, 'Maybe you have TB. Let's test you for TB'. When the results came out...they told me that I was TB negative. I was shocked to hear that I did not have TB. They asked me if I was taking my ARVs correctly. Ruth_F Though Ruth was finally diagnosed with TB on the basis of smear microscopy, she was initially started on first line TB treatment. Similarly, Godfrey who had had TB in the past and according to the National Guidelines, should have been prioritised for Xpert MTB/Rif testing, was diagnosed using smear microscopy. Hence, he was started on first line treatment. While admitted in hospital, he benefitted from a doctor visiting the facility who requested an Xpert MTB/Rif test. Godfrey felt that he had wasted time on the wrong treatment.

He [the doctor] said, 'I am not talking about the one [smear test] that is done 210 211 at this hospital. You have to send a sputum sample to the reference laboratory'. They said they hadn't. He then said, 'This hospital does not have 212 213 capacity to do some conclusive tests. What you are doing here are mere jokes'. 214 Godfrey_M ... I had to forget about the previous two months of admission in hospital. It 215 216 was a joke. [Laughs]. Godfrey_M Even where Xpert MTB/Rif was available as a primary diagnostic test, Spiwe was wrongly 217 started on first line TB treatment because healthcare workers misinterpreted the Xpert MTB/Rif 218 219 result. I took it [first line TB treatment] from November, December to January. Then 220 when I had gone to collect my re-supply in February they told me that I was 221 222 taking a wrong treatment and was supposed to be on DR-TB treatment. 223 Spiwe F 224 Diagnosing DR-TB in people who have extrapulmonary TB requires invasive sampling. The equipment, consumables and expertise is not readily available in primary health care or district 225 226 hospitals. Four participants with extra-pulmonary TB experienced delays in diagnosis and initiating appropriate treatment. All four were hospitalised before definite diagnoses were made 227 and required costly laboratory tests and medical procedures that were outsourced to private 228 229 facilities, as reflected in the experience of Talent. 230 ... the injection [needle] they used was thick and long. They inserted it into my 231 stomach and extracted some fluid from there. They took it to a private 232 laboratory and that is how they discovered that I had TB in my stomach. Talent_F 233 234 Five participants experienced delays in turnaround time for follow-up laboratory results, especially during the COVID-19 pandemic. Changes from intensive to continuation phase 235 regimens were only done once follow-up results were received. Longer intensive phase meant 236 237 that participants were exposed to more drugs and side effects. 238so it [long processing times] delays you again from getting your results 239 even though they always come late because of COVID. Felistus_F 240 Poor health seeking behaviour 241 Diagnosis of TB was also delayed among participants who first sought help from traditional 242 healers or private clinics. Financial resources for such participants were depleted by the time

244245

246

247

248

249 250

251

252

253

254

255

256257

258

259

260

261

262

263

264

265

266

267

268 269

270

271

272

273

274275

276

they were referred to public health facilities. Caleb's experience shows how some traditional healers would refer people to hospitals. We were also advised by traditional healers to consult hospitals and other traditional healers. They mostly advised us to go to hospitals. That's when we started going to X [a private clinic]. Caleb_M We had spent all the money we had by the time I was referred to the doctors at X [TB referral centre] who confirmed that it was TB. Moses M By contrast, those who had had a previous TB episode accessed care earlier at public health facilities, since they were quick to recognise 'possible TB' symptoms. Such participants incurred little or no costs and were more likely to be physically fit when they started treatment. I experienced chest pains and night sweats for less than a month. When I started having these symptoms, I said to myself, 'Maybe the TB that previously infected me was not fully treated. Let me go to the clinic and get tested'. Then I went for testing and the results came out MDR-TB positive. Felistus F Health service factors undermining quality of TB care Once people were started on DR-TB treatment, they reflected on how they endured long waiting times at clinics and lack of privacy during administration of injections. There is need for some form of privacy. I can't pull down my trousers and get an injection while other nurses are watching from one of the rooms. Peter_M Three participants were frustrated by lack of service continuity especially in urban clinics. Participants would meet new nurses at each clinic visit. These became more common during COVID-19 when some facilities were closed for fumigation and nurses redeployed to other facilities or services such as COVID-19 duties. We knew each other very well with the nurse who was there and was supposed to administer my injections the week that followed. But that nurse did not come to work because she had been transferred and there was a new nurse. I had to start explaining to her all over again. Peter_M Five participants reported that clinics experienced stock-outs of DR-TB drugs during the COVID-19 pandemic. They incurred additional costs because they were asked to travel to neighbouring clinics in search for medicines. Sometimes I would board two commuter omnibuses to get to the clinic only to be told that the drugs were out of stock. They would then refer me to other clinics to check if the tablets were in stock. Moses_M

278

279

280

281

282

283

284

285

286

287

288

289

290 291

292

293

294

295

296

297

298299

300

301

302

303

304

305

306

307

308

309

Participants living in urban settings were more likely to afford transportation costs compared to those in rural areas. A woman living in a rural village failed to access clofazimine, which meant that she had to go intermittently (during drug stockouts) without clofazimine. She was very frail although she was taking the medicines (except clofazimine) she managed to access religiously. ...I'm not getting all the drugs. There is one which was always in short supply; it is called clofazimine. So I wasn't getting all the drugs for my treatment. I started getting them later. They gave me a full course last month. All along I was getting on an alternate basis. Spiwe_F Talent who was on second-line antiretroviral therapy also experienced stock-outs of abacavir from public clinics resulting in a viral load rebound. She incurred substantial costs because she had to purchase abacavir from private pharmacies. In the second line I had a challenge. I was taking Lopinavir and Abacavir. Abacavir is not available in all places. It's hard to find. So I had a challenge in that abacavir was always out of stock. Being out of stock meant I had to buy it myself. So I would buy monthly supplies at USD7. So this caused my viral load to go down...to increase tremendously [correcting herself]. Talent_F Physical and socioeconomic impact of DR-TB The impact of DR-TB on households was exacerbated by treatment delays, income instabilities and additional non-TB shocks. People who delayed DR-TB treatment were too frail to walk and work. They were often hospitalised and had to hire cars to visit health facilities because they could not walk from their house to bus stops and from bus stops to health facilities. I wasn't able to walk coz of weak joints and shortness of breath. So I had to hire a vehicle every day from July-December at R40 [USD2.50] per day. Tecla F Because of the side effects of DR-TB treatment, one man likened the medicines to 'bulldozers'. Adherence to DR-TB medicines was difficult because of those side effects. The side effects also affected people's productivity and ability to work. A participant explained that he had to make a tough choice between adhering to medicines and returning to work. Those ones are like a bulldozer. They hit like a big hammer...[laughs]...That treatment hit me hard and made me very very weak and the injection caused me to sleep the whole day. Godfrey_M Some of us we are in informal trades. Sometimes you would want to be consistent on [adhere to] your medication but it will force you to stop taking

311

312

313

314

315

316

317 318

319

320

321

322

323

324

325

326

327

328 329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

your medication and go for work because you can't get on the street [informal job] when you have taken your medication. Peter M Households experienced huge losses of income during DR-TB treatment. Losses were related to reductions in productivity due to sickness and productive time lost while frequenting healthcare facilities and when household members could not attend to their work or daily chores because they were looking after a sick person. Loss of income was exacerbated by the informal nature of employment which does not reward and allow for absenteeism. ...my wife would support me to walk and would carry me on her back. So there was nothing else we could do for our living...My wife stopped doing piece jobs because she was looking after me. Moses_M ...everything changed as soon as I was diagnosed with TB. You can't get paid when you are not going to...to work. If you don't go to work everything will also be on hold. Peter M. The physical, psychological and socioeconomic burden associated with DR-TB caused changes in traditional gender roles and strained marital relationships. This in turn led, in some cases, to abandonment by spouses and/or depression. Three women were abandoned by their husbands and Caleb, a man, who had a supportive wife and was bedridden for a long time contemplated suicide.after 8 months into my treatment that's when my husband ran away from us and I was left alone. Life got tough for me and children. Up to now he hasn't come back. I am still alone. Agnella_F ... I said to my wife, 'Umn, I have tried a lot to get healed.... I have cried a lot. It's better for me that I go. May God give me rest; I am in pain!' Caleb_M In addition to the dire socioeconomic situation caused by DR-TB, COVID-19 lockdowns led to closure of businesses and forced households to spend their savings and to sell assets. Lockdowns closed a household business for Caleb and shattered job prospects for Godfrey who had recovered from DR-TB and was about to start a new job as a driver. It took a long time before my wife re-opened her kitchen [canteen]. During lockdown, especially the first lockdown, people were not allowed to move....that's when our father sold his car. Life was tough. Caleb_M Actually we depend on menial jobs. So as of now I... I was thinking of... what's the name of those companies that are advertised in..? Those earth moving [machines]. So I was ready to go there unfortunately the COVID pandemic forced everything to a standstill. Godfrey_M.

345

346

347

348 349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369 370

371

372

373

374

375

376

Loss of income was aggravated when participants were abandoned by their breadwinner spouses. Even post DR-TB treatment job opportunities were narrowed with limited prospects to recuperate to their pre-treatment income levels. Ruth who used to earn her livelihood as a hair stylist and was abandoned by her husband felt anxious to start working again. It affected my marriage and my work. I was supposed to be busy plaiting hair right now. If I decide to plait hair, one of the clients may have dandruffs, and with this TB, I may get sick again. So it really affected me. Ruth_29F Rural households experienced additional shocks such as drought and pestilences. Pestilences, namely bird flu and tickborne diseases (theileriosis), killed poultry and cattle and amplified the negative effects of DR-TB on household livelihood. That time our chickens got affected by bird flu. We used to rely on chickens for our relish...if you get sick you don't want to eat substandard foods. Moses_M Children were psychologically affected by seeing their parents suffer and in pain. Also they feared their parents might die. Talent explained that education of her child was affected in two ways when the child refused to go to school and when the family could not afford school fees due to depleted financial resources. 'What if I come back from school and find out that my mother is dead' E-eh right now she was supposed to be in Form 2 but she is in Grade 7 because she refused to go to school. The other ones dropped out of school because as their mother I was not fit enough to run around looking for jobs and assistance. Talent F Coping strategies adopted by people and their households Coping strategies were categorised into either reversible or irreversible. All households adopted short term reversible coping strategies such as reducing number of meals and spending savings. Six people relocated to their families in search of support and/or caregivers. It's like family support. Whenever I took my medication I'd get hungry and I was supposed to cook. I was supposed to... You know? I couldn't do all of this. So I realized that it was better for me to go back home for maternal care and support. Peter_M Support was often provided by immediate families. ...my father supported me a lot because that is when some of his pension came out. All of it was spent on my treatment. Loveness_F Talent disclosed her DR-TB status to neighbours who assisted her with food and reminders to go to clinic to pick up her medications.

My neighbours would bring me food. In some cases, they would remind me that it was 377 almost time to get to the clinic. I would thank them and tell them that I had fallen asleep. 378 Then I would go [to the clinic]. Talent F 379 380 Households that either did not have savings or had spent their savings at the start of treatment 381 initially sold non-productive and then productive assets to raise money for food and transport. We sold chickens to get money to buy sugar, cooking oil or rice. That's how we survived 382 till we reached a point when she sold our spanners. She'd look around the house to find 383 out if there was something else to sell so that we could get money for bus fare. Moses_M 384 Caleb's livelihood as a private DJ was severely affected following the distress sale of his sound 385 system. He deeply regretted the sale. He coped by making claims on family members to avert 386 387 further sales of assets. ... I ended up selling assets that I wasn't prepared to dispose. That PA (sound 388 system) system! I am now thinking of replacing the speakers that I sold. 389 Caleb M 390 *Uum.* We realised that we had burdened people. So my wife ended up phoning my mother 391 in-law. My mother-in-law would ask my father-in-law in Harare to send us some money 392 393 for groceries. That helped us a lot. Caleb_M 394 Some households managed to register their children on a government educational social 395 assistance programme. Those who failed to register their children tried to keep their children in 396 school by negotiating with school authorities. In terms of school fees BEAM [Basic Education Assistance Module] came to my rescue. I 397 used to worry and to depend on my mother to pay school fees for my child...Tecla_F 398 We wanted to register them on BEAM but it seems one needs to have connections in 399 order to get registered. Each time they were sent away from school for not paying fees, 400 my wife would immediately go to the school to plead for their readmission. Moses M 401

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431432

Discussion Our study revealed huge socioeconomic, physical and psychological impacts of DR-TB among people and their household members, including children. DR-TB pushes households into a downward spiral of poverty and debt traps in a context such as Zimbabwe that has little safety nets and is wracked with economic challenges. Stock-outs of DR-TB drugs were frequently experienced during COVID-19. COVID-19 lockdowns affected people's livelihoods as people depleted their savings. The convergence of COVID-19, economic challenges and other shocks such as pestilences amplified the impact of DR-TB on households, accelerating irreversible coping strategies. Diagnostic delays were a common theme in this study. Those diagnosed more rapidly had less severe disease at the time of treatment initiation and incurred less costs. Delays in starting TB treatment have been described in several settings.(42,43) People delay care-seeking because they perceive the quality of services in public clinics as poor or believe that their symptoms are due to other causes. (44) In Zimbabwe, time between onset of symptoms and start of DR-TB treatment was reported to be up to four months as people often sought care from private pharmacies and clinics first before they were diagnosed and referred to public clinics for treatment.(42) Our study shows that health system related factors contribute significantly to these delays. (43) Delayed diagnosis increases the likelihood of physical debility, (45) community transmission of DR-TB, hospitalisation and income loss due to reduced productive capacity by sick people or when households members assume carer roles.(46) As expected, longer diagnostic journeys stretched household resources for a long time such that financial resources were depleted at commencement of DR-TB treatment.(47) DR-TB may break households resilience and may strain social relations with family, neighbours or spouses leading to abandonment, also documented in our study.(45) The End TB strategy aims to reduce TB morbidity and mortality by introducing rapid diagnostics to reduce the time to effective treatment.(18) The full benefits of such technology may not be realised unless delays and structural barriers within the health system are addressed.(42,44) TB has long-term impacts on income and employment. DR-TB affected households are economically vulnerable even post TB treatment.(6) DR-TB narrows job opportunities and increases likelihood of post TB disabilities such as chronic lung disease. The disabilities result in households continuing to incur health care related costs. (7,8,48,49) In the same way, chronic

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

disabilities as a result of TB or other diseases, recurring TB or additional shocks e.g. COVID-19 and pestilences may break household resilience to shocks faster, rendering households vulnerable to current and future TB.(35) The socioeconomic situation of DR-TB affected households, already strained due to their disease and related costs, was worsened by COVID-19. Studies have shown that households respond by utilising all resources available to them, (27,28,50) and as observed in this study, may be forced to adopt irreversible coping strategies which had extensive impact on their livelihoods. As observed elsewhere, the impact of DR-TB, combined with COVID-19 was greatest on households without regular income and were food insecure, as they depleted their assets and exhausted short term coping strategies such as borrowing as they were no longer creditworthy.(16,50,51) In the context of distress sales, households may sell more assets to raise the amount of cash they need as assets are usually sold below their market values.(35,50) Though successful treatment outcomes were attained by all participants in this study, this was at the expense of their livelihoods. Effective TB treatment requires uninterrupted supplies of medicines. Participants reported drug stock-outs during the first few months of the COVID-19 pandemic. This forced them to seek medicines in neighbouring clinics or go without certain medicines. Stockouts of drugs and diagnostics during COVID-19 occurred globally and were enhanced by transport and travel interruptions. (52) The National TB programme decided to dispense three-month supplies of medicines to reduce people's contact with health facilities. (53) In the context of global supply chain disruptions and increased drug dispensing, stock-outs were therefore inevitable. This study was conducted between the first (March-July 2020) and second (December-February 2021) COVID-19 lockdowns, reducing the likelihood of recall bias. Most participants included in the study were initiated on DR-TB treatment during the first lockdown which allowed us to understand the challenges at the start of the pandemic. By including participants from rural and urban settings we were able to identify experiences and coping strategies which were setting specific. Participants who were interviewed were either at the end of treatment or had completed treatment providing insights into their diagnostic and treatment journeys. However, by interviewing participants at the end of successful treatment we introduced survival bias. Therefore, the challenges and coping strategies described here may not be representative of those

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

doing less well on treatment. Studies have reported that DR-TB inflicts huge physical, socioeconomic and psychological impact on household members. (12) However, we only managed to infer the impact of DR-TB on household members based on what the participants affected by DR-TB reported since we did not interview any household members, spouses or children. Our study has several implications for policy and practice. Firstly, since TB diagnosis is the entry point into treatment and care, delays in diagnosis should be addressed by fostering collaborations between private and public healthcare sectors and alternative health providers such as traditional healers (42,54), investing resources into earlier diagnosis and raising community awareness about signs and symptoms of TB and benefits of early health seeking from public facilities. Secondly, the impact of DR-TB on physical, mental, socioeconomic wellbeing of people and households affected by DR-TB calls for multisectoral approaches even beyond the treatment period, focusing on medical and social services for post TB care and measures to mitigate re-infection with DR-TB.(36) There are various social support systems provided by government agencies and nongovernmental organisations in Zimbabwe including the USD25 cash disbursement every month for the duration of treatment for people on DR-TB treatment. However, evidence from this study and the patient cost survey conducted in Zimbabwe shows that this amount is unlikely to mitigate the costs and income loss experienced by DR-TB households. (14) Importantly, the coverage of the cash scheme is suboptimal and delays and inconsistencies in disbursements are frequent.(38) Additional safety nets in the form of educational assistance; payment of medical bills and extra food and money during and possibly post DR-TB treatment are therefore required. The Ministry of Public Service Labour and Social Welfare manages the Basic Education Assistance Module (for educational social assistance); the Harmonised Social Cash Transfers (for vulnerable households) and the Assisted Treatment Medical Order facility (for formal workers who experience work related disabilities). However, the Assisted Treatment Medical Order does not cover chronic conditions e.g. HIV and TB.(55) Several non-governmental organisations are providing various forms of assistance to vulnerable households within the communities in which we collected data, but their reach is chequered, and coordination between programs is limited and sometimes non-existent. Effective stakeholder mapping and interministerial collaborations aimed at providing social protection to people and households with

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

DR-TB should be included in the National Strategic Plan. People with DR-TB will need to be linked to available social protection. Given that many participants and households included in this study had experienced an episode of TB before and were vulnerable to future DR-TB, social protection should take a broader focus and should be availed before people and households are affected by DR-TB. Such TB sensitive approaches, (56) that focus on food insecure or vulnerable households may have greater impact on TB epidemiology and control in Zimbabwe and will likely prevent future TB/DR-TB episodes. Conclusion DR-TB pushes households into poverty, and increases their vulnerability to shocks. Other stressors such as COVID-19, divorce and pestilences enhanced their vulnerability resulting in irreversible coping strategies. The impact of DR-TB extends beyond the duration of DR-TB treatment. Multisectoral approaches combing health system, psychosocial and economic interventions are crucial to reduce diagnostic delays and mitigate physical and mental consequences and loss of livelihoods due to DR-TB and other shocks. Acknowledgements We would like to thank the participants who took part in this study. **Author contributions** Conceptualised the study: CT, KK, VB. Reviewed the draft manuscript: CT, KK, DP, VB Data collection: CT, FM Data analysis: CT, VB, KK Critical review of manuscript: CT, KK, FM, VB, FK, SK, DP. All authors reviewed and approved the final manuscript. Data availability statement The transcripts used in this study have been shared as Supporting information (Transcripts_Coping with DR-TB and COVID-19). **Conflicts of interest** None declared **Funding statement**

Collins TIMIRE was funded by the Fogarty International Centre of the National Institutes of Health (NIH; Bethesda, Maryland, MD, USA) under Award Number *D43 TW009539* (https://www.nih.gov/about-nih). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- 1. World Health Organisation. Global TB Report 2022. Geneva, Switzerland; 2022.
- 2. World Health Organisation. Global Tuberculosis Report 2017 [Internet]. Geneva, Switzerland; 2017. Available from: WHO/HTM/TB/2017.23
- 3. World Health Organisation. Global Tuberculosis Report 2019 [Internet]. Geneva, Switzerland; 2019. Available from: WHO/CDS/TB/2019.15
- 4. World Health Organization. Global Tuberculosis Report 2016 Tuberculosis Country Profiles. Geneva, Switzerland; 2016.
- 5. World Health Organisation. WHO (2022) Global TB Report. Tuberculosis profile:

 Zimbabwe [Internet]. 2022. Available from:

 https://worldhealthorg.shinyapps.io/tb_profiles/?_inputs_&entity_type=%22country%22&lan=%22EN%22&iso2=%22KG%22
- 6. Meghji J, Gregorius S, Madan J, Chitimbe F, Thomson R, Rylance J, et al. The long term effect of pulmonary tuberculosis on income and employment in a low income, urban setting. Thorax. 2021;76:387–395.
- 7. Byrne AL, Marais BJ, Mitnick CD, Garden FL, Lecca L, Contreras C, et al. Chronic airflow obstruction after successful treatment of multidrug-resistant tuberculosis. ERJ Open Res [Internet]. 2017;3:1–11. Available from: http://dx.doi.org/10.1183/23120541.00026-2017
- 8. Jabeen K. Pulmonary infections after tuberculosis. Int J Mycobacteriology. 2016;Suppl 1(S73).
- 9. Isaakidis P, Rangan S, Pradhan A, Ladomirska J, Reid T, Kielmann K. "I cry every day": Experiences of patients co-infected with HIV and multidrug-resistant tuberculosis. Trop Med Int Heal. 2013;18(9):1128–33.
- 10. Chan E D, V L, Strand M J, Chan J F, Huynh M N, Goble M. Treatment and outcome analysis of 205 patients with multidrug-resistant tuberculosis. Am J Respir Crit Care Med. 2004;169:1103–9.
- 11. Ahuja S D, Ashkin, Aventano, Benerjee, Bauer, Bayona. Multidrug resistant pulmonary tuberculosis treatment regimens and patient outcomes: an Individual patient data meta-analysis of 9,153 Patients. PLoS Med. 2012;9(8):e1001300.
- 12. Yellappa V, Lefèvre P, Battaglioli T, Narayanan D, Van Der Stuyft P. Coping with

- tuberculosis and directly observed treatment: a qualitative study among patients from South India. BMC Health Serv Res [Internet]. 2016;16:1–11. Available from: http://dx.doi.org/10.1186/s12913-016-1545-9
- 13. Fuady A, Houweling TAJ, Mansyur M, Richardus JH. Catastrophic total costs in tuberculosis- affected households and their determinants since Indonesia 's implementation of universal health coverage. Infect Dis Poverty. 2018;7(3):1–14.
- 14. Timire C, Ngwenya M, Chirenda J, Metcalfe JZ, Kranzer K, Pedrazzoli D, et al. Catastrophic costs among tuberculosis affected households in Zimbabwe: a national health facility-based survey. Trop Med Int Heal [Internet]. 2021;26(10):1248–55. Available from: https://doi.org/10.1111/tmi.13647
- 15. World Health Organisation. National surveys of costs faced by TB patients and their households [Internet]. [cited 2023 Jan 30]. Available from: https://www.who.int/teams/global-tuberculosis-programme/tb-reports/global-tuberculosis-report-2022/uhc-tb-determinants/6-2-national-surveys-of-costs-faced-by-tb-patients-and-their-households
- 16. Ramma L, Cox H, Wilkinson L, Foster N, Cunnama L, Vassall A, et al. Patients' costs associated with seeking and accessing treatment for drug-resistant tuberculosis in South Africa. Int J Tuberc Lung Dis [Internet]. 2015 Dec;19(12):1513–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26614194
- 17. University Research Co. LLC. Providing Comprehensive Patient-Centred Care: A conceptual framework for social support for TB patients [Internet]. Bethesda, USA; 2014. Available from: www.urc-chs.com/sites/default/files/Providing Comprehensive Care.pdf
- 18. STOP TB Partnership. The paradigm shift 2016-2020: Global plan to End TB [Internet]. Geneva, Switzerland; 2015. Available from: http://www.stoptb.org/global/plan/plan2/
- 19. Pedrazzoli D, Lalli M, Boccia D, Houben R, Kranzer K. Can tuberculosis patients in resource-constrained settings afford chest radiography? Eur Respir J. 2017;49:1601877.
- 20. Ghazy RM, Saeh HM El, Abdulaziz S, Hammouda EA, Elzorkany AM, Khidr H. OPEN A systematic review and meta □ analysis of the catastrophic costs incurred by tuberculosis patients. Sci Rep [Internet]. 2022;12(558):1–16. Available from: https://doi.org/10.1038/s41598-021-04345-x
- 21. World Health Organization. Tuberculosis patient cost surveys: a handbook. Geneva,

- Switzerland; 2018.
- Ministry of Health Kenya. The First Kenya Tuberculosis Patient Cost Survey, 2017.
 Nairobi, Kenya; 2017.
- 23. Pedrazzoli D, Siroka A, Boccia D, Bonsu F, Nartey K, Houben R, et al. How affordable is TB care? Findings from a nationwide TB patient cost survey in Ghana. Trop Med Int Heal. 2018;23(8):870–8.
- 24. Hatherall B, Newell JN, Emmel NB, Khan SC, Amir M. "Who Will Marry a Diseased Girl?" Marriage, Gender, and Tuberculosis Stigma in Asia. Qual Health Res [Internet]. 2019;29(8):1109–19. Available from: https://doi.org/10.1177/1049732318812427
- 25. Olsson L, Maggie O, Tschakert P, Agrawal A, Eriksen, H S, Ma S, et al. Livelihoods and poverty. In: Climate Change 2014: Impacts, Adaptation, and Vulnerability Part A: Global and Sectoral Aspects Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge, United Kingdom: Cambridge University Press; 2014.
- 26. PEP-CBMS Network Coordinating Team. Definition and types of shocks and coping strategies to be monitored [Internet]. Technical Workshop on "Monitoring Household Coping Strategies during Complex Crises." 2011 [cited 2020 Mar 8]. Available from: http://www.pep-net.org/sites/pep-net.org/files/typo3doc/pdf/CBMS_country_proj_profiles/Philippines/poverty_maps/Copin g/Session2_Shocks_Coping_To_Monitor.pdf
- 27. Diwakar V. From pandemics to poverty: the implications of coronavirus for the furthest behind [Internet]. 2018 [cited 2022 May 5]. Available from: https://odi.org/en/insights/from-pandemics-to-poverty-the-implications-of-coronavirusfor-the-furthest-behind/
- 28. Russell S. Ability to pay for healthcare: concepts and evidence. Health Policy Plan. 2014;11(3):219–37.
- 29. Shah SMA, Mohammad D, Qureshi MFH, Abbas MZ, Aleem S. Prevalence, Psychological Responses and Associated Correlates of Depression, Anxiety and Stress in a Global Population, During the Coronavirus Disease (COVID-19) Pandemic. Community Ment Health J [Internet]. 2021;57(1):101–10. Available from: http://www.ncbi.nlm.nih.gov/pubmed/33108569

- 30. Diwakar V. From pandemics to poverty: hotspots of vulnerability in times of crisis. London, UK; 2020.
- 31. World Health Organisation. Global TB Report 2020: Country profiles [Internet]. Geneva, Switzerland; 2020. Available from: https://www.who.int/docs/default-source/hq-tuberculosis/global-tuberculosis-report-2020/country-profile-2020-final-web-min.pdf?sfvrsn=b4137a1c_0
- 32. World Health Organisation. Global Tuberculosis Report 2022. Geneva, Switzerland; 2022.
- 33. Ministry of Health and Child Care. Zimbabwe National Health Financing Policy: resourcing pathway to universal health coverage. Harare, Zimbabwe; 2015.
- 34. Chirisa I, Mutambisi T, Chivenge M, Mabaso E, Matamanda AR, Ncube R. The urban penalty of COVID-19 lockdowns across the globe: manifestations and lessons for Anglophone sub-Saharan Africa. GeoJournal [Internet]. 2022;87(2):815–28. Available from: http://www.ncbi.nlm.nih.gov/pubmed/32868960
- 35. Vanleeuw L, Zembe-Mkabile W, Atkins S. "I'm suffering for food": Food insecurity and access to social protection for TB patients and their households in Cape Town, South Africa. PLoS ONEoS one [Internet]. 2022;17(4):1–16. Available from: http://www.ncbi.nlm.nih.gov/pubmed/35472210
- 36. Loveday M, Hlangu S, Larkan L-M, Cox H, Daniels J, Mohr-Holland E, et al. "This is not my body": Therapeutic experiences and post-treatment health of people with rifampicin-resistant tuberculosis. PLoS One [Internet]. 2021;16(10):e0251482. Available from: http://www.ncbi.nlm.nih.gov/pubmed/34662887
- 37. Ministry of Health and Child Care. Zimbabwe tuberculosis management guidelines. Harare, Zimbabwe; 2016.
- 38. Timire C, Sandy C, Ferrand RA, Mubau R, Shiri P, Mbiriyawanda O, et al. Coverage and effectiveness of conditional cash transfer for people with drug resistant tuberculosis in Zimbabwe: a mixed methods study. PLoS Glob Public Heal. 2022;2(12):e0001027.
- 39. Bazeley P. Analysing Qualitative Data□: More Than 'Identifying Themes .' Malaysian J Qual Res. 2016;2(2):6–22.
- 40. Gibbs GR. Analysing Qualitative Data. London, UK: Sage Publications; 2007.
- 41. Braun V, Clarke V. Using thematic analysis in psychology. Qual Res Psychol [Internet].

- 2006;3(2):77–101. Available from: http://dx.doi.org/10.1191/1478088706qp063oa
- 42. Tadokera R, Huo S, Theron G, Timire C, Manyau-Makumbirofa S, Metcalfe JZ. Health care seeking patterns of rifampicin-resistant tuberculosis patients in Harare, Zimbabwe: A prospective cohort study. PLoS One [Internet]. 2021;16(7):e0254204. Available from: http://www.ncbi.nlm.nih.gov/pubmed/34270593
- 43. Shewade HD, Shringarpure KS, Parmar M, Patel N, Kuriya S, Shihora S, et al. Delay and attrition before treatment initiation among MDR-TB patients in five districts of Gujarat, India. Public Heal Action [Internet]. 2018;8(2):59–65. Available from: http://dx.doi.org/10.5588/pha.18.0003
- 44. Naidoo P, van Niekerk M, du Toit E, Beyers N, Leon N. Pathways to multidrug-resistant tuberculosis diagnosis and treatment initiation: a qualitative comparison of patients' experiences in the era of rapid molecular diagnostic tests. BMC Health Serv Res [Internet]. 2015 Oct 28;15:488. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26511931
- 45. Bond V, Chileshe M, Sullivan C, Magazi B. The Converging Impact of Tuberculosis, HIV/AIDS, and Food Insecurity in Zambia and South Africa. Renewal Working Paper. London, UK; 2009.
- 46. Department for International Development (DFID). Sustainable livelihoods guidance sheets [Internet]. 1999. Available from: www.livelihoodscentre.org
- 47. Tanimura T, Jaramillo E, Weil D, Raviglione M, Lönnroth K. Financial burden for tuberculosis patients in low- and middle-income countries: a systematic review. Eur Respir J. 2014;43(6):1763–75.
- 48. Chin AT, Rylance J, Makumbirofa S, Meffert S, Vu T, Clayton J, et al. Chronic lung disease in adult recurrent tuberculosis survivors in Zimbabwe: a cohort study. Int J Tuberc Lung Dis. 2019;23(2):203–11.
- 49. Tadyanemhandu C, Chirenda J, Garvey C, Metcalfe J. Treatment success, but living with the consequences of post-tuberculosis sequelae. Int J Tuberc Lung Dis. 2020;24:657–658.
- 50. Sweeney S, Mukora R, Candfield S, Guinness L, Grant DA, Vassall A. Measuring income for catastrophic cost estimates: Limitations and policy implications of current approaches. Soc Sci Med [Internet]. 2018;215:7–15. Available from: https://doi.org/10.1016/j.socscimed.2018.08.041

- 51. Foster N, Vassall A, Cleary S, Cunnama L, Churchyard G, Sinanovic E. The economic burden of TB diagnosis and treatment in South Africa. Soc Sci Med [Internet]. 2015

 Apr;130:42–50. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25681713
- 52. European Medicines Agency. Availability of medicines during COVID-19 pandemic [Internet]. [cited 2023 Jan 8]. Available from: https://www.ema.europa.eu/en/human-regulatory/overview/public-health-threats/coronavirus-disease-covid-19/availability-medicines-during-covid-19-pandemic
- 53. Sandy C, Takarinda KC, Timire C, Mutunzi H, Dube M, Dlodlo RA, et al. Preparing national tuberculosis control programmes for COVID-19. Int J Tuberc Lung Dis [Internet]. 2020;24(6):634–6. Available from: http://www.ncbi.nlm.nih.gov/pubmed/32553013
- 54. Shringarpure KS, Isaakidis P, Sagili KD, Baxi RK, Das M, Daftary A. "When Treatment Is More Challenging than the Disease": A Qualitative Study of MDR-TB Patient Retention. PLoS One [Internet]. 2016;11(3):e0150849. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26959366
- 55. Ministry of Health and Child Care. Contact investigation guidelines. Harare, Zimbabwe; 2019.
- 56. Boccia D, Pedrazzoli D, Wingfield T, Jaramillo E, Lönnroth K, Lewis J, et al. Towards cash transfer interventions for tuberculosis prevention, care and control □: key operational challenges and research priorities. BMC Infect Dis [Internet]. 2016;16(307):1–12. Available from: http://dx.doi.org/10.1186/s12879-016-1529-8

Legends of Tables

Table 1: Characteristics of DR-TB patients who were enrolled in the study, Zimbabwe, 2020-2021

Table 2: Additional quotes and themes

medRxiv preprint doi: https://doi.org/10.1101/2023.02.24.23286187; this version posted February 27, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Table 1: Characteristics of DR-TB patients who were enrolled in the study, Zimbabwe, 2020-2021

Identifier	HIV status	Regimen	Marital status	History of TB	Employed	Site of TB	Residency	Received DR-TB allowance	Stage of interview
Agnella_F‡	Positive	18-mo	Single mother of 2 children	No	No	PTB	Urban	Yes§	Post-treatment
Ruth_F	Positive	9-11mo	Single mother	Yes	No	PTB	Urban	Yes§	Post-treatment
Felistus_F	Negative	18-mo	Single, never married	Yes	No	PTB	Urban	Yes§	During treatment
Moses_M	Negative	9-11mo	Married	No	No	PTB	Urban	Yes§	During treatment
Peter_M	Positive	18-mo	Single, never married	No	No	PTB	Urban	No	Post-treatment
Caleb_M	Positive	9-11mo	Married	No	No	PTB	Rural	Yes §	During treatment
Talent_F	Positive	18-mo	Single mother	No	No	EPTB	Urban	No	Post-treatment
Lydia_F	Positive	18-mo	Single, never married	No	Yes	PTB	Rural	Yes within 3 months	During treatment
Godfrey_M	Positive	18-mo	Single, never married	Yes	No	PTB	Rural	No	Post-treatment
Sipho_M	Positive	18-mo	Married	Yes	No	PTB	Rural	Yes§	During treatment
Spiwe_F	Negative	9-11mo	Married	No	No	PTB	Rural	No	During treatment
Loveness_F	Negative	18-mo	Single mother (1child)	No	No	EPTB	Urban	Yes within 3 months	Post-treatment
Tecla_F	Positive	18-mo	Single mother (1child)	No	No	EPTB	Urban	Yes within 3 months	Post-treatment
Nyenge_M	Positive	9-11mo	Single	Yes	No	PTB	Rural	Yes within 3 months	During treatment
Wembe_M	Positive	9-11mo	Married	New Patient	No	ЕРТВ	Rural	No	Post-treatment
Fanuel_M	Positive	9-11mo	Married with 2 children	New patient	No	PTB	Rural	Yes within 3 months	During treatment

‡=Pseudonym and age; PTB=Pulmonary TB; EPTB=extra-pulmonary TB; § First cash disbursement came after at least 8 months

Table 2: Additional quotes and themes

Themes	Subtheme	Quotes supporting themes
Health seeking	Alternative providers	We were also advised by the traditional healers that we are supposed to go and consult hospitals and other traditional healers who helpthey mostly advised us to go to hospitals. That's when we started to go to Y [a private clinic]. Caleb_M
		we consulted another herbalist and he informed us that there was need for two white goats with which he performed the ritualsHe charged us R2500 (USD 170) but my father negotiated with him. He gave him R1000 (USD 70) and promised to bring the balance of R1000. Caleb_M
		On the first days when I was able to walk but was still weak, we visited a certain traditional healer. Caleb_M
	Recognition of TB symptoms	On that day I didn't even drink alcohol. I tried but I couldn't. I couldn't even smoke. What is happening here? Could it be that thing again? It's that thing again! I was certain about that. I decided to wait for some change. If I didn't see any change I would go back to the hospital. I realized there was no change. Then I went back [to the hospital]. Godfrey_M
Health system constraints	Misdiagnosis and diagnostic delays	On that day they called and advised that my treatment be changed. That's how I was diagnosed with MDR-TB. So they spent a few days hesitating to tell me that I was here to stay. This was no longer time for jokes. I had to forget about the previous two months of admission. It was a joke [laughs]. I was going to be admitted for six months because of DR-TB. Godfrey_M
		When I had completed treatment, they told me that I was supposed to be injected. Ruth_F
	Long waiting times	So they isolated us. After they have finished whatever they would be doing, that's when they would come to the back to attend to us. After that we would go back home. Ruth_F
		Those with common TB were served first but initially the sister-in-charge had told us that we were to be served first. But some of the times we had to wait for others to finish firstto wait for about 45 minutes to an hour then they would come and administer our injections. Then we would leave. Peter_M
	Lack of privacy	they would inject us while we were outside where we could be seen by people who stay close to us. There are some houses which do not have dura-walls, some are fencedpeople would watch as we received our injections. Loveness_F
	Delays in laboratory results	I am confused because if I give them a sputum sample so that my treatment is changed and I do not get the results it means I stay on the same treatment. Spiwe F

	Drug stock- outs	they would ask us to go to X [clinic] and other poly clinics like Y. They would say, "We have run out of a certain drug." Then we would rush to either Z or to Y to look for it. This happened several timesAgnella_F.
Psychosocial, physical and economic impacts	Separation from partner	It affected me very much because it affected my marriage. The moment I got unwell, my husband could not accept it. He lacked the will to help me. I just don't know why. Maybe he thought I was going to die and he had nothing to do with me. Because the moment I got sick he ran away and went to South Africa. He never came back. Ruth_F
	Hospitalisation and huge medical bills Side effects of	Yes on the first instance I was admitted for 10 days and we paid the required fee. On the second occasion money was needed for a blood transfusion that's when I was admitted for 3 days whilst we were waiting for the blood. So when the blood was delivered a transfusion was conducted and we paid for the blood. Then on the third occasion I was admitted for 5 days because my kidneys were no longer functioning properly and we paid for the admission and the medicines. Tecla_F I managed to finish my medication even though I have some side
	TB treatment	effects, I experienced lots of side effects on my legsthey are numb such that I can't walk, for me to walk long distances. Agnella_F.
		Like now, I have problem with my legs. The toes shiver on their own. You see that? I don't know what's causing the problem. I also have numbness from here [knees], and also in my joints and hands. I am supposed to consult a doctor or clinic to get help but I don't have the money. I need medical attention and food. Loveness_F
	Narrowed job opportunities	I was once employed by company X where they collect garbage bins. The work environment has a lot of dust. The moment I got there, it didn't take me a long time before I got sick. I think I only worked for just a week and started coughing again I later on quit the job. So this sickness affected me a lot. I can no longer get employed to do every other job except only those that pose less risk to my health. Ruth_F I used to work as a DJ but ever since I was diagnosed with TB last
		year, I have never worked. In addition, I was advised that I must not mix and mingle with people in dusty environments before I finish my treatment since I might meet a TB infected person and may get infected. Caleb_M
Coping strategies	Selling assets	I sold them for 700 each and got R1,400 for both speakers. I bought them for R3,500 each meaning R7, 000 for the two. I was looking forward to sell them for R2, 000 each and would get R4, 000 for the 2 speakers. So, I sold them for R700 each because things were tough for us. Caleb_M

	I sold 2 speakers. The old man [my father] sold a car. He sold his [Toyota] Wish. That time money was hard to come by. It took a long time before my wife got a kitchen [canteen] business. Caleb_M
	We sold two goats. We sold them under pressure because I wanted to go to X Mission hospital. The goats had a combined value of around R1200 but we sold them at R1000 because I wanted to get money to go to hospital. Spiwe_F
	I sold kitchen chairs because I wanted to buy food. At that time I really wanted food, so I sold my kitchen chairs. I sold them for USD25. The market price around that time was USD 120 but I didn't have a choice. Talent_F
	I sold my TV set for USD50 so that I could get money for food. I sold property. At that time it was supposed to fetch at least USD90. Agnella_F.
Relocation in search of a caregiver, family support and to reduce	I went to stay with my mother, the reason being I was looking for someone to assist me, someone who because I couldn't do anything for myself. I was bathed, sometimes I defecated on myself so I was monitored there. I wanted a caregiver to be around me every time. Talent_F
cost of transport.	There was a time I decided to relocate from I realized that I had run out of money for transport costs from X to Y. By then I was able get up and walk slowly on my own. Then I finished my course of tablets at the clinic that is in the suburb where I currently live. Talent_F
Support from social networks.	My neighbours would bring me food. In some cases, they would remind me that it was almost time to get to the clinic. I would thank them and tell them that I had fallen asleep. Then I would go [to the clinic]. Talent_F
Spending savings and defaulting debt payments.	She explained about my illness to the rotating savings and credit group and fortunately they all know me and they agreed that she be excused in making contributions since she was faced with a problem. They said she had to solve the problem at her home and she would pay back the money when she gets back on her feet. So she was given that lump sum of R11, 000. Up to now she hasn't paid it back. Caleb_M
Reduced spending on children's education.	My daughter failed to sit for her exams that year. She sat for her exams the year that followed. She passed but we could not afford fees for her to continue with education because life was very tough. Even now, life is still tough. Agnella_F