#### Factors associated with the utilisation of mosquito nets amongst rural adults: A cross-sectional

### study in East Nusa Tenggara Province Indonesia

Authors: Robertus Dole Guntur<sup>1</sup>, Maria Lobo<sup>1</sup>, Fakir M Amirul Islam<sup>2</sup>

#### **Affiliations:**

<sup>1</sup> Mathematics Study Program, Faculty of Science and Engineering, Nusa Cendana

University, Kupang NTT, Indonesia

<sup>2</sup> Department of Health Science and Biostatistics, School of Health Sciences, Swinburne University

of Technology, Melbourne, Australia

### Address for Correspondence:

Robertus Dole Guntur, Ph.D.

Mathematics Study Program

Faculty of Science and Engineering, Nusa Cendana University

Jln. Adisucipto Penfui Kupang NTT, Indonesia Post Code 85001

Email: robertus\_guntur@staf.undana.ac.id

T:+6281325525489

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

#### Abstract

**Backgrounds**: The use of mosquito nets has shown a significant impact on the reduction to malaria burden; however, the utilisation of this kind prevention measures greatly depend on the community behaviour which is limited to be studied in Indonesia. This study explored the factors associated with the use of mosquito nets in the rural of East Nusa Tenggara Province (ENTP) Indonesia.

**Methods**: The community-based cross-sectional study to 1503 rural adults was carried out from October to December 2019. Data on the ownership and the utilisation of mosquito nets were collected using a standardized questionnaire. Logistic regression method was applied to determine the factors associated with the use of mosquito nets.

**Results**. The ownership (utilisation) of any mosquito nets, long-lasting insecticide treated nets (LLINs), and non-LLINs was 95.8% (82.3%), 71.6% (54.8%) and 42.7% (27.6%), respectively. The likelihood of the using of mosquito nets was significantly higher for adults in low malaria endemic settings (MES) than in high MES (adjusted odds ratio [AOR]: 10.1;95% CI, 6.17 to 16.5 for any mosquito nets, AOR: 11.1;95% CI, 7.32 to 16.7 for non-LLINs). The use of LLINs was significantly higher for adults living in Hills areas (AOR: 2.11;95%CI, 1.37 to 3.25).

**Conclusion**. The coverage of the ownership nets was not followed by the utilisation of the nets in this province. To progress to malaria elimination in ENTP, the coverage and the utilisation of any mosquito nets should be high. Public health promotion to improve the awareness of the using of these nets should be prioritized, particularly for those living in high MES.

Keywords: Mosquito nets; rural adults; malaria elimination

### **INTRODUCTIONS**

Malaria is most prevalence in low-and middle-income countries <sup>1</sup>. The World Health Organization (WHO) estimated that the number of malaria cases in 2020 was 241 million <sup>2</sup>. However, the global trend of malaria deaths decreased about 30% in the last two decades, from 896,000 in 2000 to 627,000 in 2020 <sup>2</sup>. Based on the WHO report in South East Asia Region, the number of malaria deaths was projected about 8,879 in 2020, of which 83% were from India and 16% were from Indonesia <sup>2</sup>.

Indonesia which is administratively composed of 514 districts in total has a national commitment to achieve malaria elimination by 2030<sup>3</sup>. Sixty-two percent of the total number of districts in the country were classified as malaria elimination zones, while the rest of the districts were classified as low, moderate, and high malaria endemic settings (MES)<sup>4</sup>. However, this progress shows a significant disparity between Western and Eastern part of the country. Most of districts in the Western part of the country were classified as malaria free zone and low MES. Whilst, most of districts with moderate MES and all districts with high MES were in the Eastern part of the country that includes the East Nusa Tenggara Province (ENTP)<sup>4</sup>.

ENTP was the third-highest contributor of the malaria burden to the country in 2020 with the total of malaria cases in this province at the end of that year was 15,304 cases <sup>4</sup>. Under the collaboration with the National Malaria Control Program of Indonesian government, local authority of ENTP has applied various efforts to reduce the burden of malaria in this region. This includes the introduction to apply long lasting insecticide treated net (LLINs) which has been conducted since 2005 <sup>5</sup>.

The use of LLINs has shown a positive impact on the reduction to malaria burden since 2000 and is considered as the best measure to prevent malaria compared with other malaria prevention approaches <sup>6</sup>. Therefore, improving the coverage and the utilisation of LLINs is highly recommended by WHO to control malaria vector in malaria endemic countries and the coverage of universal access to LLINs should obtain at least 80% to accelerate malaria elimination effort <sup>7</sup>. However, in the case the distribution

of LLINs was limited, the use of non-LLINS could also be applied as it has also shown a positive impact to reduce the malaria burden <sup>8,9</sup>

Indonesia applied LLINs as the main vector control intervention in high MES through routine distribution in antenatal care and immunisation activities particularly for pregnant women and children <sup>3</sup>. The coverage of long-lasting insecticide treated net in high MES has improved significantly from less than 50% in 2007 to 75% in 2017 <sup>5</sup>. In spite of the fact that the distribution of LLINs increased significantly from 56,337 nets in 2015 to 4,376,636 in 2017 i.e., almost eight times higher, the literature on the coverage and the utilisation of this treated net on the population level was limited in Indonesia <sup>10</sup> and the use of this malaria prevention measures greatly depend on human behaviour <sup>11</sup>. All these yet to be thoroughly investigated in Indonesia.

Some limited study on mosquito nets has been conducted in ENTP [12-16]. A study on village level to explore the behaviour of local community in using mosquito nets indicated that there was an association between family attitude and the use of mosquito nets <sup>12</sup>. Previous findings related to the current study showed that the knowledge of treated mosquito nets was low [12]-13], and the utilisation of mosquito nets was also low <sup>15</sup>, as well as the awareness to sleep under mosquito nets varied between ethnicities <sup>14,16</sup>. However, factors associated with the use of mosquito nets have yet to be investigated in this province. Understanding the contributing factors related to the utilisation of mosquito nets will help the local authority designs an evidence-based approach to boost malaria elimination in this province. This research aimed to fill this gap by exploring the prominent factors associated with the use of mosquito nets malaria elimination in this province. This research aimed to fill this gap by exploring the prominent factors associated with the use of mosquito nets to support the national commitment to achieve malaria elimination by 2030.

#### **METHODS**

#### **Study settings**

The current community-based cross-sectional study was conducted in three districts in ENTP from October to December 2019. The eligible participants were 15-85 years of age. Of three districts, firstly, East Sumba district has an estimated number of populations of 258,490 people with 39% had no formal

education; secondly, Belu district has an estimated number of populations of 223,176 people with 32.7% had no formal education; and finally, East Manggarai district has an estimated number of population of 287,210 people with 18.4% had no education <sup>17</sup>

#### Study population and sample size calculation

All adults in three selected districts were the target population for this study. In total, the number of respondents enrolled for this study was 1503 adults. This number was obtained after considering of malaria prevalence in ENTP, intra-class correlation of malaria prevention study in Indonesia, design effect and participation rate of participants. The comprehensive calculation of sample size was previously presented elsewhere <sup>18</sup>.

#### **Data collection procedure**

A data collection tool was adapted from validated questionnaire <sup>19,20</sup> with some modifications. Originally the questionnaire was prepared in English. Then, the local language expert and the main author of this study translated the questionnaire into local language (Indonesian). They then combined the translation version of the questionnaire. The combined version of the questionnaire was used as a final tool to collect data. Nine enumerators who had background in nursing school and had working experience in the local public health centres were recruited and trained about the intention of the study, how to approach participants and get their written consent, how to complete the questionnaire. Data collection was supervised strictly by the main author of this study.

#### **Dependent Variables**

The dependent variables of the study were whether participants sleeping under any mosquito net, LLINs, or non-LLINs the night before the survey. These outcomes were obtained based on the response of participants on two questions (did you sleep under mosquito net last night? and what type of mosquito did you use last night?)

## **Independent Variables**

The independent variables of the study were gender, age group, education level, occupation, family size, socio-economic status (SES), the nearest health facilities, the distance to the nearest health facilities, household income in relation to the provincial minimum wages, household income in relation to Indonesia poverty line and geographical condition.

#### **Data Analysis**

Socio-demographic factors, including age, gender and education of participants were reported by descriptive statistics. The percentage of participants having the possession and utilisation of mosquito nets was computed with its 95% confidence interval (CI). To investigate the potential factors affecting the utilisation of mosquito net, a univariate and multivariate binary logistic regression analysis was applied. Adjusted odds ratio with 95% CI and p value less than 5% was used to affirm the significance of each variable.

### **Ethics Statement**

This research was approved by Human Ethics Committee of the Swinburne University of Technology (reference number 20191428-1490) and the Indonesian Ministry of Health (reference letter: 164 LB.02.01/2/KE.418/2019). Informed consent was submitted by all participants when they were enrolled.

#### RESULTS

The distribution of participants having the ownership of mosquito nets based on the socio-demographic and environmental background is presented in Table 1. People with ownership of mosquito nets were younger, had better education level, Entrepreneurs or government or non-government employees, better SES, from larger family size, and from high or low malaria endemic settings.

The ownership and the use of mosquito nets in rural adults of ENTP is showed in Table 2. Overall, the percentage of participants with the possession of mosquito nets was high at 95.8% with a 95% confidence interval (CI): 94.8 - 96.8. The proportion of participants having LLINs was high, accounting

for 71.6% with a 95% CI: 68.9 - 74.3, whilst the ownership of non-LLINs was only 42.7% with a 95% CI: 38.8 - 46.5. The use of any mosquito nets by rural adults was high, accounting for 82.3% with a 95% CI :80.2- 84.5, however only about half of participants (54.8% with a 95% CI: 51.4 - 58.2) utilised LLINs. This was worsened for non-LLINs. Merely about the quarter of participants (27.6% with a 95% CI: 23.2 - 31.9) applied non-LLINs for preventing malaria.

Factors associated with the use of any mosquito nets in rural of ENTP was presented in Table 3. After controlling all confounding variables in the multivariable analysis, the following results were obtained. Having at least diploma education level (adjusted odds ratio (AOR) = 5.68, 95% CI: 2.6 - 12.41), senior high school education level (AOR = 5.41, 95% CI: 3.07 - 9.53), junior high school education level (AOR = 2.87, 95% CI: 1.78 - 4.63), primary education level (AOR = 2.52, 95% CI: 1.79 - 3.55), living in low endemic settings (AOR = 10.1, 95% CI: 6.17 - 16.5), living in moderate endemic settings (AOR = 2.09, 95% CI: 1.50 - 2.90), having small family members (AOR = 1.73, 95% CI: 1.30 - 2.30), living closed to rice field (AOR = 2.36, 95% CI: 1.30 - 4.30), living closed to hills (AOR = 2.10, 95% CI: 1.32 - 3.36) were considerably associated with the use of any mosquito nets.

Factors associated with the use of non-LLINs in rural of ENTP was depicted in Table 4. After controlling all confounding variables in multivariable analysis, the following results were obtained. Having at least diploma education (adjusted odds ratio (AOR) = 2.37, 95% CI: 1.14 - 4.93), senior high school education level (AOR = 2.58, 95% CI: 1.57 - 4.23), junior high school education level (AOR = 2.69, 95% CI: 1.69 - 4.30), primary education level (AOR = 2.26, 95% CI: 1.52 - 3.35), living in low endemic settings (AOR = 11.1, 95% CI: 7.32 - 16.7), living in moderate endemic settings (AOR = 3.52, 95% CI:2.27 - 5.46), living closed to public health centre (AOR = 2.25, 95% CI: 1.58 - 3.20), living closed to subsidiary public health centre (AOR = 1.79, 95% CI: 1.22 - 2.63), living closed to village health post (AOR = 2.06, 95% CI:1.39 - 3.05), living less than 1 km from the nearest health service (AOR = 1.51, 95% CI: 1.05 - 2.18), living between 1 to 2 km from the nearest health service (AOR = 2.43, 95% CI: 1.67 - 3.54), living between 2 to 3 km from the nearest health service (AOR = 2.16, 95% CI: 1.42 - 3.30), were significantly associated with the use of non-LLINs.

Factors associated with the use of LLINs in rural of ENTP was portrayed in Table 5. After controlling all confounding variables in multivariable analysis, the following results were obtained. Living in low endemic settings (AOR = 0.60, 95% CI:0.45 -0.80) and living closed to hills (AOR = 2.11, 95% CI:1.37 - 3.25) were substantially associated with the use of LLINs.

### DISCUSSION

To our knowledge this is the first population-based study to explore the ownership and the utilisation of mosquito nets in rural adult of East Nusa Tenggara Province Indonesia. This study shows that the ownership of any mosquito nets was high in this area, however the utilisation of this nets particularly long-lasting insecticide treated net (LLINs) was low. This study also demonstrated that the high coverage of LLINs was not followed by the high use of LLINs. The prominent factors associated with a higher likelihood to utilise mosquito nets were low malaria endemic settings, higher education level, and better socio-economic status.

The difference in coverage of LLINs and the use of LLINs was consistent with previous findings in some other countries <sup>21–24</sup>. Our study shows that 72% percent of rural population have the possession of LLINs. This coverage was just below the coverage that recommended by WHO. Improving the coverage of these treated nets is critical to boost malaria elimination in this setting. This study also further demonstrated that the use of LLINs by rural population was low. The improvement of awareness of rural community to use LLINs is critical considering that the treated nets have been applied by the Indonesian government as the main intervention to control malaria in this country <sup>25</sup>. Failure to educate community on the benefit of the treated nets led to misuse of them including for food storage as indicated in other research <sup>26</sup>.

The results of this study showed that the use of mosquito nets was significantly higher in low MES compared to high MES. This result is consistent with findings in other settings <sup>23,27,28</sup>. The high proportion of adults' people in low MES sleeping under mosquito nets might be contributed to high

awareness of malaria for those communities. Previous studies showed that adults in low MES in this region has high malaria awareness compared to those in high MES <sup>13</sup>. The finding of this study is unexpected by the authors since the national government of Indonesia has allocated more resources in distributing mosquito net for high MES in the Eastern part of Indonesia including in this area since 2005 until now <sup>5,25</sup>. This signalled that the distribution of mosquito nets in this area might not be followed by the adequate health promotion activities to encourage local community taking the benefit of the mosquito net in preventing malaria.

The present study demonstrated that the use of any mosquito nets was in line with the increase of education level of participants. This is consistent with other studies <sup>29-32</sup>, indicating that there was a positive association between education level and the use of mosquito nets for preventing malaria. This study revealed that the likelihood of the use of mosquito net for participants having at least diploma education level was almost six times higher than those without education level. Participants having senior high school education level were five times higher to use mosquito net compared with those having no education level. The possible explanation for this could be that people with higher education level have a high level of malaria knowledge <sup>16,33</sup>, and they have many opportunities to be exposed to multiple sources of information <sup>34</sup>, so that they could understand the benefit of sleeping under mosquito net for preventing malaria. This research has demonstrated that more than half of rural adults in this province have low level of education (no formal school or only primary school) and other studies indicated that children dropping out of school was high in this area <sup>35,36</sup>. These situations need to be addressed by local authority to encourage rural population using mosquito nets to support malaria elimination effort in this area.

One of the important aspects to assess the benefit of the health intervention programs is the socioeconomic status of the community. The intervention programs should reach the poor as many times as they do to the relatively rich people. In this research, we found that the ownership of mosquito nets rose with the increasing of socio-economic status of participants. In contrast to the possession of mosquito nets, this study demonstrated that the poorest was more likely to use mosquito nets compared to the

richest group. This finding is consistent with other finding in other settings <sup>37,38</sup>. The high use of mosquito nets in the poor group might be connected with their perceived vulnerability as perceived susceptibility to malaria has been revealed to be higher in poor families. Moreover, people with high SES often have easy access to other methods for preventing malaria and might therefore not use of mosquito nets. One study indicated that the richest group living in houses with have screening in door and window often believed that they were adequately shielded from mosquito bites and therefore they did not use mosquito nets <sup>39</sup>.

While this study provides some significant findings to guide malaria elimination programs in this province, it also has some weaknesses. First, the ownership and the utilisation of mosquito nets were assessed based on the information provided by one adult in a household. Each participant was asked how many mosquito nets they have and whether all family members slept under mosquito nets in the night prior survey. The response to these questions might prone to recall bias, which is a common type of bias in a cross-sectional study design. Secondly, this study was conducted during the rain season in this area. Similarly, the other provinces in Indonesia, this province has two seasons, rain and dry season. The behaviour of rural population might show a different pattern in the use of mosquito nets, as indicated in other studies <sup>40</sup>. Therefore, the comprehensive pattern in the use mosquito nets throughout the year could not be revealed by this study.

In conclusion, this study demonstrated that the high coverage of mosquito nets particularly LLINs is not followed by the high use of this treated net. The important factors affecting the utilisation of mosquito nets were malaria endemic settings and education level of participants. To boost malaria elimination in ENTP, the coverage of any mosquito nets should be high. Health education promotion to improve the awareness rural community of the using of these nets should be prioritized for those living in high MES and low education level.

## Acknowledgments

We would like to thank the Australia Awards Scholarship for supporting this research and School of Health Sciences, Swinburne University of Technology for providing funding for primary data collection for this study. The funders had no role in the designing of the study, data collection, analysis, or interpretation of data, or writing the paper. We further would like to express our gratitude to the Health Ministry of Indonesia, the governor of ENTP, head of East Sumba, Belu, and East Manggarai district, nine head of sub-districts, and forty-nine village leaders for allowing this research to be conducted in their regions.

## **Conflict of Interest**

The authors have no conflicts of interest associated with the material presented in this paper.

#### **Author Contributions**

Conceptualization: RDG. Data curation: RDG, FMAI. Formal analysis: RDG, FMAI. Funding acquisition: RDG, FMAI. Methodology: RDG. Project administration: RDG, ML. Writing - original draft: RDG. Writing - review & editing: RDG, ML, FMAI

## **ORCID iDs**

Robertus Dole Guntur: https://orcid.org/0000-0002-1597-3644 Maria Lobo : https://orcid.org/0000-0002-2046-449X Fakir M. Amirul Islam: <u>https://orcid.org/0000-0003-3897-3302</u>

# REFERENCES

 Gutman JR, Lucchi NW, Cantey PT, et al. Malaria and Parasitic Neglected Tropical Diseases: Potential Syndemics with COVID-19? The American Journal of Tropical Medicine and Hygiene
 Epub ahead of print June 1, 2020. DOI: 10.4269/ajtmh.20-0516.

- World Health Organization. World Malaria Report 2021. Available from: https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2021. 2021.
- Asia Pacific Malaria Elimination Network. Indonesia National Strategic Plan 2020-2024.
   December 2020 . Available from: https://www.apmen.org/resources/indonesia-national-strategicplan-2020-2024. 2021.
- Ministry of Health Republic of Indonesia. Indonesia Health Profile 2020. Jakarta . Available from: https://pusdatin.kemkes.go.id/folder/view/01/structure-publikasi-data-pusat-data-daninformasi.html (Indonesian). 2021.
- Sitohang V, Sariwati E, Fajariyani SB, et al. Malaria elimination in Indonesia: halfway there. The Lancet Global Health 2018;6:e604–e606.
- 6. Wangdi K, Furuya-Kanamori L, Clark J, et al. Comparative effectiveness of malaria prevention measures: a systematic review and network meta-analysis. Parasites Vectors 2018;11:210.
- World Health Organization. Achieving and maintaining universal coverage with longlasting insecticidal nets for malaria control. Available from: https://apps.who.int/iris/bitstream/handle/10665/259478/WHO-HTM-GMP- 2017.20eng.pdf;jsessionid=90E881F5A4D639FDD106233F1960F66F?sequence=1. 2017.
- Okumu F. The fabric of life: what if mosquito nets were durable and widely available but insecticide-free? Malar J 2020;19:260.
- 9. Van Remoortel H, De Buck E, Singhal M, et al. Effectiveness of insecticide-treated and untreated nets to prevent malaria in India. Trop Med Int Health 2015;20:972–982.
- World Health Organization. World malaria report 2018. Geneva: WHO. World Health Organization, 2018.

- Monroe A, Olapeju B, Moore S, et al. Improving malaria control by understanding human behaviour. Bull World Health Organ 2021;99:837–839.
- Paschalia YP, Doondori AK, Irfan I, et al. Behavior of the Use of Mosquito Net as a Prevention of Malaria in Ondorea Village, Nanga Panda Sub-district. Open Access Maced J Med Sci 2021;9:620–623.
- Guntur RD, Kingsley J, Islam FMA. Malaria awareness of adults in high, moderate and low transmission settings: A cross-sectional study in rural East Nusa Tenggara Province, Indonesia. PLoS ONE 2021;16:e0259950.
- Guntur RD. The Variation of Behavioural Aspects of Malaria in East Nusa Tenggara Province Indonesia: Implication for Malaria Elimination in the Province [dissertation]. Swinburne University of Technology . Available from: http://hdl.handle.net/1959.3/470529. 2022.
- Ipa M, Widawati M, Laksono AD, et al. Variation of preventive practices and its association with malaria infection in eastern Indonesia: Findings from community-based survey. PLoS ONE 2020;15:e0232909.
- Guntur RD, Kingsley J, Islam FMA. Ethnic Variation and Its Association With Malaria Awareness: A Cross-sectional Study in East Nusa Tenggara Province, Indonesia. J Prev Med Public Health 2022;55:68–79.
- ENTP TCB of S. Nusa Tenggara Timur Province in Figures 2020. Available from: https://ntt.bps.go.id/publication/2020/04/27/080aa7bde0454f07e7848b49/provinsi-nusatenggara-timur-dalam-angka-2020.html. 2020.
- Guntur RD, Kingsley J, Islam FMA. Epidemiology of Malaria in East Nusa Tenggara Province in Indonesia: Protocol for a Cross-sectional Study. JMIR Res Protoc 2021;10:e23545.

- Manana PN, Kuonza L, Musekiwa A, et al. Knowledge, attitudes and practices on malaria transmission in Mamfene, KwaZulu-Natal Province, South Africa 2015. BMC Public Health 2017;18:41.
- Hlongwana KW, Mabaso ML, Kunene S, et al. Community knowledge, attitudes and practices (KAP) on malaria in Swaziland: A country earmarked for malaria elimination. Malar J 2009;8:29.
- Tapera O. Determinants of long-lasting insecticidal net ownership and utilization in malaria transmission regions: evidence from Zimbabwe Demographic and Health Surveys. Malar J 2019;18:278.
- 22. Asumah M, Akugri F, Akanlu P, et al. Utilization of insecticides treated mosquito bed nets among pregnant women in Kassena-Nankana East municipality in the upper east region of Ghana. Public Health Toxicol 2021;1:1–11.
- Aung PL, Win KM, Show KL. Utilization of insecticide-treated bed nets among pregnant women in Myanmar–analysis of the 2015–2016 Demographic and Health Survey. PLoS ONE 2022;17:e0265262.
- Maung TM, Oo T, Wai KT, et al. Assessment of household ownership of bed nets in areas with and without artemisinin resistance containment measures in Myanmar. Infect Dis Poverty 2018;7:19.
- Indonesia M of H. Indonesia national strategic plan 2020–2024. Available from: https://www.apmen.org/resources/indonesia-national-strategic-plan-2020-2024. 2020.
- Dhiman S, Veer V. Culminating anti-malaria efforts at long lasting insecticidal net? J Infect Public Health 2014;7:457–464.

- Aung PL, Win KM, Pumpaibool T. Malaria Preventive Practices among People Residing in Different Malaria-Endemic Settings in a Township of Myanmar: A Mixed-Methods Study. TropicalMed 2022;7:353.
- Kanyangarara M, Hamapumbu H, Mamini E, et al. Malaria knowledge and bed net use in three transmission settings in southern Africa. Malar J 2018;17:41.
- 29. Muhumuza E, Namuhani N, Balugaba BE, et al. Factors associated with use of malaria control interventions by pregnant women in Buwunga subcounty, Bugiri District. Malar J 2016;15:342.
- Haileselassie W, Habtemichael M, Adam R, et al. Regional and socio-economic disparity in use of insecticide-treated nets to prevent malaria among pregnant women in Kenya. International Health 2022;ihac024.
- 31. Kuse KA, Chikako TU, Bacha RH, et al. Multilevel Modelling of Individual, Community and Regional Level Factors Associated with Insecticide-Treated Net Usage among Pregnant Women in Ethiopia. Healthcare 2022;10:1418.
- 32. Baume CA, Franca-Koh AC. Predictors of mosquito net use in Ghana. Malar J 2011;10:265.
- Sixpence A, Nkoka O, Chirwa GC, et al. Levels of knowledge regarding malaria causes, symptoms, and prevention measures among Malawian women of reproductive age. Malar J 2020;19:225.
- Suka M, Odajima T, Okamoto M, et al. Relationship between health literacy, health information access, health behavior, and health status in Japanese people. Patient Education and Counseling 2015;98:660–668.
- Guntur RD, Lobo M. Statistical Modelling for Dropped Out School Children (DOSC) in East Nusa Tenggara Province Indonesia. J Phys: Conf Ser 2017;812:012073.

- Guntur RD, Lobo M. An Analysis of Factors that Affect Out of School Junior High Aged Children Using Logistic Regression Method. Asia-Pacific Collaborative Education Journal 2015;11:23–36.
- 37. Kanmiki EW, Awoonor-Williams JK, Phillips JF, et al. Socio-economic and demographic disparities in ownership and use of insecticide-treated bed nets for preventing malaria among rural reproductive-aged women in northern Ghana. PLoS ONE 2019;14:e0211365.
- Singh M, Brown G, Rogerson SJ. Ownership and use of insecticide-treated nets during pregnancy in sub-Saharan Africa: a review. Malar J 2013;12:268.
- Goesch JN, Schwarz NG, Decker M-L, et al. Socio-economic status is inversely related to bed net use in Gabon. Malar J 2008;7:60.
- 40. Ricotta E, Oppong S, Yukich JO, et al. Determinants of bed net use conditional on access in population surveys in Ghana. Malar J 2019;18:63.

## Tables

## Table 1. The ownership of mosquito nets by Socio-demographic and Environmental

Characteristics of Respondents in East Nusa Tenggara Province (N = 1,495)

Characteristics	Total	Ownership of mosqu	p-value	
Characteristics	Total_	No	Yes	
Overall	1,495	63(4.21)	1432(95.8)	
Gender				
Male	727	27(3.71)	700(96.3)	0.349
Female	768	36(4.69)	732(95.3)	
Age group				
< 30	205	6(2.93)	199(97.1)	0.005

30–39	418	12(2.87)	406(97.1)	
40–49	371	12(3.23)	359(96.8)	
50–59	295	24(8.14)	271(91.9)	
$\geq 60$	206	9(4.37)	197(95.6)	
Level of education				
No education	279	23(8.24)	256(91.8)	< 0.001
Primary school	678	31(4.57)	647(95.4)	
Junior high school	229	6(2.62)	223(97.4)	
Senior high school	210	3(1.43)	207(98.6)	
Diploma or above	99	0(0.00)	99(100.0)	
Main occupation				
Farmer	831	26(3.13)	805(96.9)	< 0.001
Housewife	403	32(7.94)	371(92.1)	
Entrepreneurs	48	0(0.00)	48(100.0)	
Other	62	4(6.45)	58(93.5)	
Gov. or non-gov. workers	151	1(0.66)	150(99.3)	
Socio-economic status				
Low	449	27(6.01)	422(94)	0.007
Average	860	35(4.07)	825(95.9)	
High	186	1(0.54)	185(99.5)	
Family size				
< = 4	803	49(6.10)	754(93.9)	< 0.001
> 4	692	14(2.02)	678(98.0)	
†HH Income in relation to PM	W			
< PMW	1,342	63(4.69)	1279(95.3)	0.006
≥ PMW	153	0(0.00)	153(100.0)	

## **‡HH Income in relation to IPL**

< IPL	1,395	63(4.52)	1332(95.5)	0.03
> = IPL	100	0(0.00)	100(100.0)	
The nearest health service				
Village maternity posts	386	23(5.96)	363(94.0)	0.026
Village health post	302	10(3.31)	292(96.7)	
Subsidiary public health centres	338	19(5.62)	319(94.4)	
Public health centres	469	11(2.35)	458(97.7)	
Distance to the nearest health se	ervices			
< 1 km	578	28(4.84)	550(95.2)	0.816
1–2 km	400	15(3.75)	385(96.3)	
2–3 km	204	8(3.92)	196(96.1)	
$\geq$ 3 km	313	12(3.83)	301(96.2)	
Geographical conditions				
Coastal area	203	8 (3.94)	195 (96.1)	0.04
Rice Field	198	5 (2.53)	193 (97.5)	
Hills	986	40 (4.06)	946 (95.9)	
Forest	108	10 (9.26)	98 (90.7)	
Malaria endemic settings				
High	495	11(2.22)	484(97.8)	< 0.001
Moderate	500	48(9.60)	452(90.4)	
Low	500	4(0.80)	496(99.2)	

†The Provincial Minimum Wages (PMW) in 2019 is IDR 1,795,000 monthly; ‡The Indonesia Poverty Line (IPL) in 2019 is defined all households having an average income less than IDR 1,990,170 monthly.

## Table 2. The ownership and the use of mosquito nets of the respondents in East Nusa Tenggara

**Province (N = 1,495)** 

#	Questions	n	% (95%CI)
1	The ownership of mosquito nets		
	Yes	1432	95.8(94.8, 96.8)
	No	63	4.20(0.00, 9.20)
2	Types of mosquito nets		
	Only LLINs	794	53.1(49.6, 56.6)
	Only Non LLINs	362	24.2(19.8, 28.6)
	LLINs and Non-LLINs	276	18.5(13.9, 23.1)
3	Respondent sleeping under bed net last night		
	Yes	1231	82.3(80.2, 84.4)
	No	264	17.6(13.0, 22.2)
4	Types of bed net used last night		
	LLINs	819	54.8(51.4, 58.2)
	Non-LLINs	412	27.6(23.3, 31.9)
	Respondents have owned any mosquito nets	1432	95.8(94.7, 96.8)
	Respondents have owned non-LLINs	638	42.7(38.8, 46.5)
	Respondents have owned LLINs	1070	71.6(68.9, 74.3)
	Respondents sleeping under Non LLINs last		
	night	412	27.6(23.2, 31.9)
	Respondents sleeping under LLINs last night	819	54.8(51.4, 58.2)
	Respondent sleeping under Any bed net last		
	night	1231	82.3(80.2, 84.5)

## Table 3. Factors associated with the use of any mosquito nets in East t Nusa Tenggara Province

(N = 1,495)

	Using of any mosquito nets				
Characteristics	No at risk	n (%)	$\mathbf{COR}^{\dagger}$	AOR <sup>‡</sup>	
Overall	1,495	1231 (82.3)			
Gender					
Male	727	606 (83.4)	1.15 (0.88, 1.50)	1.19 (0.89, 1.59)	
Female	768	625 (81.4)	1.00	1.00	
Age group					
< 30	205	171 (83.4)	1.37 (0.83, 2.24)	0.84 (0.48, 1.45)	
30–39	418	359 (85.9)	1.65 (1.07, 2.55)	1.32 (0.82, 2.12)	
40–49	371	298 (80.3)	1.11 (0.73, 1.69)	0.97 (0.61, 1.52)	
50–59	295	241 (81.7)	1.21 (0.78, 1.89)	1.05 (0.66, 1.68)	
$\geq 60$	206	162 (78.6)	1.00	1.00	
Level of education					
No education	279	189 (67.7)	1.00	1.00	
Primary school	678	570 (84.1)	2.51 (1.82, 3.48)	2.52 (1.79, 3.55)	
Junior high school	229	195 (85.2)	2.73 (1.76, 4.25)	2.87 (1.78, 4.63)	
Senior high school	210	188 (89.5)	4.07 (2.45, 6.76)	5.41 (3.07, 9.53)	
Diploma or above	99	89 (89.9)	4.24 (2.10, 8.54)	5.68 (2.6, 12.41)	
Socio-economic status					
Low	449	373 (83.1)	1.52 (1.00, 2.31)	2.18 (1.34, 3.54)	
Average	860	716 (83.3)	1.54 (1.05, 2.26)	2.14 (1.39, 3.28)	
High	186	142 (76.3)	1.00	1.00	
Family size					
< = 4	803	691 (86.1)	1.74 (1.33, 2.27)	1.73 (1.30, 2.30)	

 medRxiv preprint doi: https://doi.org/10.1101/2023.02.26.23286476; this version posted February 27, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

 It is made available under a CC-BY-NC-ND 4.0 International license .

 > 4
 692
 540 (78.0)
 1.00
 1.00

 Geographical conditions

Coastal area	203	146 (71.9)	1.18 (0.71, 1.96)	0.84 (0.48, 1.45)
Rice Field	198	171 (86.4)	2.91 (1.64, 5.17)	2.36 (1.30, 4.30)
Hills	986	840 (85.2)	2.64 (1.70, 4.12)	2.10 (1.32, 3.36)
Forest	108	74 (68.5)	1.00	1.00
Malaria endemic settings				
High	495	334 (67.5)	1.00	1.00
Moderate	500	418 (83.6)	2.46 (1.82, 3.33)	2.09 (1.50, 2.90)

 $^{\dagger}COR = crude odds ratio, ^{\ddagger}AOR = adjusted odds ratio; Main occupation, household income, the nearest health facilities, and the distance to the nearest health facilities were not significant in the final model.$ 

## Table 4. Factors associated with the use of non-LLINs in East t Nusa Tenggara Province (N =

### 1495)

	Using of non - LLINs			
Characteristics	No at risk	n (%)	COR <sup>†</sup>	AOR <sup>‡</sup>
Overall	1,495	412 (27.6)		
Gender				
Male	727	209 (28.7)	1.12 (0.90, 1.41)	0.96 (0.70, 1.31)
Female	768	203 (26.4)	1.00	1.00
Age group				
< 30	205	57 (27.8)	1.34 (0.86, 2.10)	1.02 (0.61, 1.68)

30–39	418	133 (31.8)	1.62 (1.10, 2.39)	1.17 (0.77, 1.79)
40–49	371	90 (24.3)	1.11 (0.74, 1.67)	0.91 (0.59, 1.41)
50–59	295	86 (29.2)	1.43 (0.95, 2.16)	1.18 (0.76, 1.83)
≥ 60	206	46 (22.3)	1.00	1.00
Level of education				
No education	279	39 (14.0)	1.00	1.00
Primary school	678	188 (27.7)	2.36 (1.62, 3.45)	2.26 (1.52, 3.35)
Junior high school	229	73 (31.9)	2.88 (1.86, 4.46)	2.69 (1.69, 4.30)
Senior high school	210	70 (33.3)	3.08 (1.97, 4.80)	2.58 (1.57, 4.23)
Diploma or above	99	42 (42.4)	4.53 (2.69, 7.65)	2.37 (1.14, 4.93)
Main occupation				
Farmer	831	214 (25.8)	1.00	1.00
Housewife	403	105 (26.1)	1.02 (0.77, 1.33)	0.96 (0.67, 1.38)
Entrepreneurs	48	10 (20.8)	0.76 (0.37, 1.55)	0.78 (0.37, 1.65)
Other	62	14 (22.6)	0.84 (0.46, 1.56)	0.93 (0.48, 1.81)
Gov. or non-gov. workers	151	69 (45.7)	2.43 (1.70, 3.46)	1.81 (1.06, 3.11)
The nearest health service				
Village maternity posts	386	66 (17.1)	1.00	1.00
Village health post	302	86 (28.5)	1.93 (1.34, 2.78)	2.06 (1.39, 3.05)
Subsidiary public health centres	338	88 (26.0)	1.71 (1.19, 2.45)	1.79 (1.22, 2.63)
Public health centres	469	172 (36.7)	2.81 (2.03, 3.89)	2.25 (1.58, 3.20)
Distance to the nearest health se	ervices			
< 1 km	578	140 (24.2)	1.35 (0.96, 1.89)	1.51 (1.05, 2.18)
1–2 km	400	140 (35.0)	2.27 (1.60, 3.22)	2.43 (1.67, 3.54)
2–3 km	204	72 (35.3)	2.30 (1.54, 3.44)	2.16 (1.42, 3.30)
$\geq$ 3 km	313	60 (19.2)	1.00	1.00

**Geographical conditions** 

medRxiv preprint doi: https://doi.org/10.1101/2023.02.26.23286476; this version posted February 27, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license . Coastal area 203 50 (24.6) 1.28 (0.73, 2.25) 0.56 (0.30, 1.04) **Rice Field** 198 82 (41.4) 2.76 (1.60, 4.78) 0.54 (0.28, 1.04) Hills 986 258 (26.2) 1.39 (0.85, 2.26) 0.37 (0.21, 0.65) 108 1.00 1.00 Forest 22 (20.4)

Malaria endemic settings				
High	495	44 (8.89)	1.00	1.00
Moderate	500	120 (24.0)	3.24 (2.23, 4.69)	3.52 (2.27, 5.46)
Low	500	248 (49.6)	10.1 (7.07, 14.4)	11.1 (7.32, 16.7)

<sup>†</sup>COR = crude odds ratio, <sup>‡</sup>AOR = adjusted odds ratio; Main occupation, Social economic status,

Household income, the nearest health facilities, and the distance to the nearest health facilities were not significant in the final model.

	Using of LLINs				
Characteristics	No at risk	n (%)	COR <sup>†</sup>	AOR <sup>‡</sup>	
Overall	1,495	819 (54.8)			
Gender					
Male	727	397 (54.6)	0.99 (0.81, 1.21)	1.1 (0.89, 1.37)	
Female	768	422 (54.9)	1.00	1.00	
Age group					
< 30	205	114 (55.6)	0.97 (0.66, 1.44)	1.00 (0.66, 1.50)	
30–39	418	226 (54.1)	0.91 (0.65, 1.28)	1.01 (0.71, 1.44)	
40–49	371	208 (56.1)	0.99 (0.70, 1.40)	1.02 (0.72, 1.45)	
50–59	295	155 (52.5)	0.86 (0.60, 1.23)	0.90 (0.62, 1.30)	
$\geq$ 60	206	116 (56.3)	1.00	1.00	
Geographical conditions					
Coastal area	203	96 (47.3)	0.97 (0.61, 1.54)	1.14 (0.70, 1.86)	
Rice Field	198	89 (44.9)	0.88 (0.55, 1.41)	1.51 (0.91, 2.52)	
Hills	986	582 (59)	1.55 (1.04, 2.31)	2.11 (1.37, 3.25)	
Forest	108	52 (48.1)	1.00	1.00	
Malaria endemic settings					
High	495	290 (58.6)	1.00	1.00	
Moderate	500	298 (59.6)	1.04 (0.81, 1.34)	0.98 (0.74, 1.31)	
Low	500	231 (46.2)	0.61 (0.47, 0.78)	0.60 (0.45, 0.80)	

## Table 5. Factors associated with the use of LLINs in East t Nusa Tenggara Province (N = 1495)

<sup>†</sup>COR = crude odds ratio, <sup>‡</sup>AOR = adjusted odds ratio; Education level, Main occupation, Social economic status, Household income, family size, the nearest health facilities, and the distance to the nearest health facilities were not significant in the final model.



