

1 **Characterization of Long COVID among U.S. Medicare Beneficiaries using Claims Data**

2

3 Yun Lu¹, PhD; Arnstein Lindaas², MA; Hector S Izurieta¹, MD, PhD; Myrna Cozen², MPH; Mikhail Menis¹,
4 PharmD; Xiangyu Shi², MS; Whitney Steele¹, PhD; Michael Wernecke², BA; Yoganand Chillarige² MPA;
5 Jeffrey A Kelman³, MD; and Richard A Forshee¹, PhD

6

7 Affiliations: (1) Center for Biologics Evaluation and Research, Food and Drug Administration, Silver
8 Spring, MD, USA; (2) Acumen LLC, Burlingame, CA, USA; (3) Centers for Medicare & Medicaid
9 Services, Washington, DC, USA

10

11 Address correspondence to: Yun Lu, PhD, FDA/CBER, 10903 New Hampshire Avenue, Silver Spring,
12 MD 20993; email: Yun.Lu@fda.hhs.gov; phone: 240-402-8704

13

14 Running title: Long COVID in Medicare Beneficiaries

15 **ABSTRACT**

16 This retrospective study utilized healthcare claims data to investigate the incidence, patient demographics,
17 and concurrent diagnoses associated with long COVID in the U.S. Medicare population. Nearly 194,000
18 (0.6%) beneficiaries had post-COVID condition diagnoses, with higher rates among nursing home
19 residents. Of those medically attended for COVID-19, 3-5% were diagnosed with post-COVID
20 conditions. We observed minimal demographic differences between those with and without long COVID.
21 When comparing diagnoses concurrent with long COVID and COVID-19, certain codes (G72 and J84)
22 for myopathies and interstitial pulmonary diseases were disproportionately present with long COVID.

23 **Keywords:** long COVID, post-COVID conditions, Medicare beneficiaries, claims data, real-world data

24 **BACKGROUND**

25 Evidence has emerged suggesting that coronavirus disease 2019 (COVID-19) can have significant long-
26 term sequelae [1-7]. Many individuals with a history of COVID-19 report symptoms weeks or months
27 after their initial infection; a growing body of literature substantiates these reports and points to a broad
28 range of symptoms and conditions that may emerge [3-6]. These persistent symptoms are commonly
29 referred to as long COVID or post-COVID conditions [8]. While a standardized case definition of long
30 COVID is still evolving, the broad range of sequelae experienced by patients subsequent to initial
31 COVID-19 infection highlights the need to elucidate the natural history of this phenomenon across
32 different populations.

33 This study aims to descriptively summarize the occurrence of long COVID among Medicare
34 beneficiaries, who represent the vast majority of the older population in the U.S.

35 **METHODS**

36 We utilized Medicare enrollment data and Fee-for-Service claims submitted between April 1, 2020 and
37 May 21, 2022. For claims to be included, both the submission and service dates must fall in the date
38 range. Medical conditions were assessed using *International Classification of Diseases, Tenth Revision,*
39 *Clinical Modification (ICD-10-CM)* diagnosis codes. Medically attended COVID-19 cases had at least
40 one claim with code U07.1 and post-COVID condition cases had either code B94.8 or U09.9 (eTable 1).
41 Code B94.8 is a general code for sequelae of infectious diseases, used for post-COVID conditions prior to
42 the introduction of the U09.9 code on October 1, 2021. This code was rarely used prior to the COVID-19
43 pandemic, which suggests that it was predominantly used to diagnose post-COVID conditions during the
44 pandemic (eFigure 1). However, to reduce the impact of misclassification, we restricted the use of B94.8
45 to periods (August 1, 2020 to February 28, 2022) where $\geq 70\%$ of cases had a prior COVID-19 diagnosis.
46 We also excluded <10 U09.9 claims with service start dates much earlier (on or prior to June 30, 2021)
47 than the diagnosis code effective date.

48 We first assessed the rate of post-COVID condition diagnoses amongst all Medicare Fee-for-Service
49 beneficiaries with continuous enrollment since April 1, 2020, separately for community-dwelling, nursing
50 home, and end-stage renal disease (ESRD) populations. Next, we defined a long COVID case as a post-
51 COVID condition diagnosis made ≥ 28 days after an initial COVID-19 claim, consistent with the
52 definition for long COVID used by the Centers for Disease Control and Prevention [8]. We compared the
53 demographic characteristics and concurrent diagnoses of this long COVID population with those who
54 survived for ≥ 28 days following a COVID-19 diagnosis and did not develop post-COVID conditions.
55 Among Medicare beneficiaries age 65 years or older, we also analyzed concurrent diagnoses using three-
56 character *ICD-10-CM* codes to understand the broad categories of conditions associated with long
57 COVID. We used concurrent diagnoses that appeared on the same claim without any cleaning for pre-
58 existing conditions. Finally, we compared the prevalence of concurrent diagnoses between long COVID
59 cases and COVID-19 cases without post-COVID conditions.

60 **RESULTS**

61 From April 1, 2020, to May 21, 2022, 193,691 (0.6%) of 31,847,927 studied Medicare beneficiaries had
62 been diagnosed with post-COVID conditions, as identified by the diagnosis codes on claims (eTable 2).
63 This proportion was higher among nursing home residents (1.0%) than among community-dwelling
64 beneficiaries (0.6%). Most beneficiaries diagnosed with a post-COVID condition had a previous COVID-
65 19 claim. Specifically, 81.0% of community-dwelling, 95.3% of nursing home, and 93.2% of ESRD
66 beneficiaries with a post-COVID condition diagnosis had a COVID-19 diagnosis on a prior claim.
67 Among beneficiaries with a COVID-19 claim, 142,255 (4.5%) community-dwelling, 13,384 (2.5%)
68 nursing home, and 3,774 (4.7%) ESRD Medicare beneficiaries were subsequently diagnosed with post-
69 COVID conditions. Around 35% of beneficiaries received their first post-COVID condition diagnosis
70 within 28 days of their initial COVID-19 claim (eTable 3). Among beneficiaries with COVID-19 and
71 post-COVID condition diagnoses, 105,826 (74.4%) community-dwelling and 10,380 (77.6%) nursing

72 home residents met our long COVID definition with at least one post-COVID condition diagnosis ≥ 28
73 days following their first COVID-19 claim. Using this case definition, the long COVID rate for
74 community-dwelling beneficiaries (3.4%) was higher than nursing home residents (2.0%) among
75 beneficiaries with a prior COVID-19 claim.

76 There were no significant sex, age, or race differences between long COVID cases and COVID-19 cases
77 without post-COVID conditions. Among long COVID cases in the community-dwelling population,
78 24.7% were 80 years or older, 55.8% were female, 86.1% were white, and 18.2% were dually eligible for
79 Medicare and Medicaid. Examining COVID-19 cases without post-COVID conditions, 22.8% were 80
80 years or older, 55.2% were female, 84.0% were white, and 19.1% were dually eligible. This same trend
81 was seen in the nursing home population, with no substantial differences between long COVID and
82 COVID-19 cases without post-COVID conditions; the exception being dual eligibility. For nursing home
83 beneficiaries, long COVID cases had a lower proportion of dual eligibility than COVID-19 cases without
84 post-COVID conditions (60.3% vs. 70.3%) (Table 1).

85 When comparing the rate ratio of diagnoses concurrent with long COVID and COVID-19, the diagnoses
86 disproportionately present among those with long COVID included “other and unspecified myopathies”,
87 “other interstitial pulmonary diseases”, “pressure ulcers”, “other pulmonary heart diseases”, and various
88 encounter reason codes typically associated with procedures (Table 2 and eTable 4). Certain codes (G72
89 and J84) for myopathies and interstitial pulmonary diseases were disproportionately present in both
90 community-dwelling and nursing home residents.

91 **DISCUSSION**

92 In this large study of the Medicare population, we found that, regardless of prior COVID-19 diagnoses,
93 almost 194,000 (0.6%) studied Medicare beneficiaries had been diagnosed with post-COVID conditions,
94 identified by diagnoses on health care claims. The rate was higher among nursing home residents (1.0%)

95 compared to community-dwelling beneficiaries (0.6%), which highlights the substantial burden of
96 COVID-19 on nursing homes.

97 Among beneficiaries with a prior COVID-19 claim, long COVID rates were lower for nursing home
98 residents than community-dwelling beneficiaries. This finding could be due to the combined effect of
99 multiple factors, including a higher likelihood to detect mild/asymptomatic COVID-19 cases in nursing
100 homes (as residents are in daily contact with health care personnel) and a lower ascertainment of post-
101 COVID conditions among individuals with multiple existing comorbidities. Our finding that community-
102 dwelling beneficiaries with post-COVID conditions were around four times more likely to lack a prior
103 COVID-19 code, with approximately 19% lacking a code, supports other studies suggesting that post-
104 COVID conditions could occur following mild cases of COVID-19 [6, 9-12].

105 We found lower rates of post-COVID conditions than those in prior studies of older individuals or other
106 populations [1, 13-15]. These lower rates could be due to cases of long COVID for which health care
107 providers did not provide a post-COVID condition code in the diagnoses in claims; this would be more
108 likely to occur during the earlier period of the pandemic before long COVID was widely recognized and
109 before a specific post-COVID condition diagnosis code was available. The discrepancy in long COVID
110 occurrence across studies may also reflect the ongoing evolution in both the definition of long COVID
111 and its recognition among health care providers.

112 For beneficiaries with a COVID-19 claim, we did not find major differences by age, sex or race between
113 those with long COVID compared to those without post-COVID condition diagnoses. This suggests that
114 on average susceptibility to long COVID may not be affected by major population demographics in this
115 population. Given that the post-COVID condition code can represent many conditions, future studies are
116 needed to identify and evaluate specific long COVID outcomes and related risk factors. Our finding that
117 dual eligibility status is only associated with long COVID among nursing home residents also requires
118 further investigation.

119 Among both community-dwelling and nursing-home-residing older individuals, diagnoses found to be
120 disproportionately associated with long COVID, as compared with COVID-19 cases without post-
121 COVID conditions, included certain codes (G72 and J84) for myopathies and interstitial pulmonary
122 diseases. This finding is consistent with a recent study of the Medicare Advantage population that
123 identified a substantial increase in respiratory failure (including interstitial pulmonary diseases) following
124 COVID-19 [13]. Our findings deserve further investigation in inferential studies.

125 While this study provides insights into the characterization of long COVID, there are limitations. This is a
126 descriptive study, not designed to differentiate between underlying health conditions associated with long
127 COVID and symptoms of long COVID. The study also relies on administrative health care data.

128 This large study of the Medicare population captured long COVID cases using claims data, finding only
129 minimal demographic differences between those with and without long COVID. Our finding that the
130 overall percentage of individuals with post-COVID conditions was higher among nursing home residents
131 highlights the substantial burden of COVID-19 on nursing homes. Our finding that certain codes for
132 myopathies and interstitial pulmonary diseases were differentially associated with long COVID deserves
133 investigation in inferential studies.

FUNDING

This work was supported by the Food and Drug Administration (FDA) as part of the SafeRx Project, a joint initiative of the Centers for Medicare & Medicaid Services and FDA.

ACKNOWLEDGEMENT

We thank Derick Ambarsoomzadeh, BA, Acumen LLC for contribution to writing and review of the report; Pablo Freyria Duenas, MA, Xi Li, MS, and Yue Zhang, MS, Acumen LLC for assistance with statistical programming; and Chad Buskirk, Centers for Medicare & Medicaid Services for his excellent program management.

Author's contributions: Yun Lu, Hector S. Izurieta, Mikhail Menis, Whitney Steele, Jeffrey A. Kelman, and Richard A. Forshee contributed to the study conception and design, data interpretation, manuscript writing and review. Arnstein Lindaas, Myrna Cozen, Xiangyu Shi, Michael Wernecke, and Yoganand Chillarige contributed to study design; data collection, analysis, and interpretation; and manuscript writing and review.

POTENTIAL CONFLICTS OF INTEREST

None of the authors reported potential conflicts of interest.

CORRESPONDING AUTHOR CONTACT INFORMATION

Address correspondence to: Yun Lu, PhD. CBER/FDA, 10903 New Hampshire Avenue, Silver Spring, MD 20993; email: Yun.Lu@fda.hhs.gov

ALTERNATE CORRESPONDING AUTHOR CONTACT INFORMATION

Arnstein Lindaas, MA, Acumen LLC, 500 Airport Boulevard Suite 100, Burlingame, CA 94010; email: alindaas@acumenllc.com

REFERENCES

1. Bull-Otterson L, Baca S, Saydah S, et al. Post-COVID Conditions Among Adult COVID-19 Survivors Aged 18–64 and ≥ 65 Years — United States, March 2020–November 2021. *MMWR Morb Mortal Wkly Rep* **2022**; 71: 713–7.
2. Carfi A, Bernabei R, Landi F. Persistent Symptoms in Patients After Acute COVID-19. *JAMA* **2020**; 324(6): 603-5.
3. Groff D, Sun A, Ssentongo AE, et al. Short-term and Long-term Rates of Postacute Sequelae of SARS-CoV-2 Infection: A Systematic Review. *JAMA Netw Open* **2021**; 4(10): e2128568.
4. Nalbandian A, Sehgal K, Gupta A, et al. Post-acute COVID-19 syndrome. *Nature Medicine* **2021**; 27(4): 601-15.
5. Logue JK, Franko NM, McCulloch DJ, et al. Sequelae in Adults at 6 Months After COVID-19 Infection. *JAMA Netw Open* **2021**; 4(2): e210830.
6. Subramanian A, Nirantharakumar K, Hughes S, et al. Symptoms and risk factors for long COVID in non-hospitalized adults. *Nature Medicine* **2022**; 28(8): 1706-14.
7. Al-Aly Z, Xie Y, Bowe B. High-dimensional characterization of post-acute sequelae of COVID-19. *Nature* **2021**; 594(7862): 259-64.
8. CDC. Long COVID or Post-COVID Conditions. Available at: <https://www.cdc.gov/coronavirus/2019-ncov/long-term-effects/index.html>. Accessed September 21, 2022.
9. Havervall S, Rosell A, Phillipson M, et al. Symptoms and Functional Impairment Assessed 8 Months After Mild COVID-19 Among Health Care Workers. *JAMA* **2021**; 325(19): 2015-6.
10. Boscolo-Rizzo P, Guida F, Polesel J, et al. Sequelae in adults at 12 months after mild-to-moderate coronavirus disease 2019 (COVID-19). **2021**; 11(12): 1685-8.

11. van Kessel SAM, Olde Hartman TC, Lucassen PLBJ, van Jaarsveld CHM. Post-acute and long-COVID-19 symptoms in patients with mild diseases: a systematic review. *Family Practice* **2021**; 39(1): 159-67.
12. Augustin M, Schommers P, Stecher M, et al. Post-COVID syndrome in non-hospitalised patients with COVID-19: a longitudinal prospective cohort study. *The Lancet Regional Health - Europe* **2021**; 6: 100122.
13. Cohen K, Ren S, Heath K, et al. Risk of persistent and new clinical sequelae among adults aged 65 years and older during the post-acute phase of SARS-CoV-2 infection: retrospective cohort study. *BMJ (Clinical research ed)* **2022**; 376: e068414.
14. Ioannou GN, Baraff A, Fox A, et al. Rates and Factors Associated With Documentation of Diagnostic Codes for Long COVID in the National Veterans Affairs Health Care System. *JAMA Network Open* **2022**; 5(7): e2224359-e.
15. CDC. Nearly One in Five American Adults Who Have Had COVID-19 Still Have "Long COVID". Available at: https://www.cdc.gov/nchs/pressroom/nchs_press_releases/2022/20220622.htm#:~:text=Overall%2C%20in%2013%20adults,to%20their%20COVID%2D19%20infection. Accessed September 21, 2022.

**Table 1. Demographic Summary for COVID-19 cases with and without Long COVID –
Community-Dwelling and Nursing Home Medicare Beneficiaries of All Ages**

Demographic Characteristics	Community Dwelling				Nursing Home			
	COVID-19 Only ^a		Long COVID ^b		COVID-19 Only		Long COVID	
	No.	%	No.	%	No.	%	No.	%
Total Beneficiaries	2,986,027		105,826		511,836		10,380	
Age								
Age below 65	434,163	14.5	13,901	13.1	57,344	11.2	1,116	10.8
Age 65-69	625,248	20.9	20,631	19.5	50,929	10.0	1,153	11.1
Age 70-74	730,601	24.5	25,712	24.3	66,284	13.0	1,455	14.0
Age 75-79	514,356	17.2	19,438	18.4	73,783	14.4	1,619	15.6
Age 80-84	341,572	11.4	13,198	12.5	81,373	15.9	1,756	16.9
Age 85-89	203,282	6.8	8,074	7.6	83,097	16.2	1,642	15.8
Age 90-94	102,387	3.4	3,749	3.5	65,478	12.8	1,125	10.8
Age 95+	34,418	1.2	1,123	1.1	33,548	6.6	514	5.0
Sex								
Female	1,648,628	55.2	59,067	55.8	323,069	63.1	6,372	61.4
Male	1,337,399	44.8	46,759	44.2	188,767	36.9	4,008	38.6
Race/Ethnicity								
Asian	40,836	1.4	1,011	1.0	7,693	1.5	146	1.4
Black	233,343	7.8	7,540	7.1	67,609	13.2	1,240	11.9
Hispanic	78,022	2.6	2,352	2.2	10,153	2.0	172	1.7
American Indian/Alaska Native	24,718	0.8	1,032	1.0	3,023	0.6	65	0.6
White	2,509,157	84.0	91,096	86.1	415,043	81.1	8,597	82.8
Other	36,631	1.2	1,116	1.1	5,181	1.0	86	0.8
Unknown	63,320	2.1	1,679	1.6	3,134	0.6	74	0.7

Demographic Characteristics	Community Dwelling				Nursing Home			
	COVID-19 Only ^a		Long COVID ^b		COVID-19 Only		Long COVID	
	No.	%	No.	%	No.	%	No.	%
Medicare Dual Eligibility								
Dual	571,417	19.1	19,311	18.2	359,708	70.3	6,254	60.3
Non-Dual	2,414,610	80.9	86,515	81.8	152,128	29.7	4,126	39.7
Reason for Entering Medicare								
Aged in without ESRD	2,252,609	75.4	78,203	73.9	345,883	67.6	6,931	66.8
Aged in with ESRD	503	0.0	36	0.0	542	0.1	14	0.1
Disabled without ESRD	712,844	23.9	26,565	25.1	160,521	31.4	3,302	31.8
Disabled with ESRD	9,988	0.3	488	0.5	2,969	0.6	73	0.7
ESRD only	10,082	0.3	534	0.5	1,921	0.4	60	0.6
Missing	1	0.0	-	-	-	-	-	-

Abbreviations: ESRD, end-stage renal disease.

^a COVID-19 only refers to individuals with a COVID-19 diagnosis without a subsequent post-COVID condition diagnosis.

^b Long COVID refers to individuals with a post-COVID condition diagnosis ≥ 28 days after their first COVID-19 diagnosis.

Table 2. Concurrent Three Character ICD-10-CM Diagnosis Summary – COVID-19 Cases with and without long COVID among Community-Dwelling and Nursing Home Beneficiaries Aged ≥65 Years

Diagnosis Code ^a	Diagnosis Code Description	COVID-19	Long	Rate Ratio ^d
		Only ^b	COVID ^c	
		Claims, %	Claims, %	
Community-Dwelling Beneficiaries		n = 13,663,828	n = 261,417	
G72	Other and unspecified myopathies	0.3	5.3	16.3
Z43	Encounter for attention to artificial openings	0.1	1.6	13.4
J84	Other interstitial pulmonary diseases	0.5	7.0	13.1
Z93	Artificial opening status	0.3	1.5	5.0
L89	Pressure ulcer	0.8	3.8	4.8
R43	Disturbances of smell and taste	0.2	0.9	4.1
M25	Other joint disorder, not elsewhere classified	0.7	3.0	4.1
R26	Abnormalities of gait and mobility	2.1	8.6	4.0
M62	Other disorders of muscle	2.5	10.2	4.0
G62	Other and unspecified polyneuropathies	0.5	2.0	4.0
R13	Aphagia and dysphagia	1.3	5.1	3.8
I27	Other pulmonary heart diseases	0.6	2.4	3.8
Z12	Encounter for screening for malignant neoplasms	0.2	0.9	3.7
Z86	Personal history of certain other diseases	3.1	11.5	3.7
Z48	Encounter for other postprocedural aftercare	0.2	0.7	3.7
Nursing Home Beneficiaries		n = 4,378,517	n = 26,411	
G72	Other and unspecified myopathies	0.2	3.5	18.5
J84	Other interstitial pulmonary diseases	0.4	2.5	6.9
Z43	Encounter for attention to artificial openings	0.4	2.4	5.7

Diagnosis Code ^a	Diagnosis Code Description	COVID-19	Long	Rate Ratio ^d
		Only ^b	COVID ^c	
		Claims, %	Claims, %	
Z86	Personal history of certain other diseases	4.2	18.1	4.3
Z87	Personal history of other diseases and conditions	2.8	9.4	3.3
Z68	Body mass index [BMI]	1.6	5.1	3.2
Z73	Problems related to life management difficulty	0.3	1.0	3.1
I27	Other pulmonary heart diseases	0.7	2.2	3.1
J15	Bacterial pneumonia, not elsewhere classified	0.7	2.2	3.0
I13	Hypertensive heart and chronic kidney disease	1.7	5.0	3.0
Z99	Dependence on enabling machines and devices, not elsewhere classified	3.1	8.6	2.8
J43	Emphysema	0.6	1.5	2.7
I11	Hypertensive heart disease	2.7	7.1	2.6
M48	Other spondylopathies	1.0	2.6	2.6
Z79	Long term (current) drug therapy	5.7	14.6	2.6

Abbreviations: *ICD-10-CM, International Classification of Diseases, Tenth Revision, Clinical Modification*

^a Diagnoses were summarized using the first three characters of *ICD-10-CM* diagnosis codes.

^b COVID-19 only refers to claims for individuals with a COVID-19 diagnosis without a subsequent post-COVID condition diagnosis.

^c Long COVID refers to claims for individuals with a post-COVID condition diagnosis ≥ 28 days after their first COVID-19 diagnosis.

^d The rate ratio was ordered in descending order among the top 100 most frequent diagnoses on Long COVID and COVID-19 only claims.