

RELATIONSHIP BETWEEN THE LEVEL OF SPIRITUALITY AND BLOOD PRESSURE CONTROL AMONG ADULT HYPERTENSIVE PATIENTS IN RURAL SOUTHWESTERN NIGERIA

Authors: Adetunji Omonijo^{1*}, Paul Olowoyo², Azeez Oyemomi Ibrahim³.

¹Adetunji OMONIJO* (MBCHB, FMC-FM), Consultant Family Physician, Federal Teaching Hospital, Ido-Eki, Ekiti State, Nigeria. Email: sundayomonijo97@gmail.com .

² Paul OLOWOYO (MBCHB, FWACP), Consultant Neurologist, AFE BABALOLA UNIVERSITY, ADO-EKITI/Federal Teaching Hospital, Ido-Eki, Ekiti State, Nigeria. Email: paulolowoyo@gmail.com.

³Azeez Oyemomi IBRAHIM (MBCHB, FMC-FM), Consultant Family Physician, Federal Teaching Hospital, Ido-Eki, Ekiti State, Nigeria. Email: ibrahimazeez1434@gmail.com.

Summary:

Background:

Spirituality has been strongly associated with good blood pressure control as it forms a strong coping mechanism in hypertensive patients.

Purpose:

This hospital-based cross-sectional study was done to determine the relationship between spirituality and blood pressure control among adult hypertensive patients in rural Southwestern Nigeria with a view to achieving good blood pressure control.

Method:

The selection was done by systematic random sampling technique. Socio-demographic and clinical information were obtained through semi-structured interviewer-administered questionnaires. The level of spirituality was assessed using the Spiritual Perspective Scale. Data were analysed using the Statistical Package for Social Sciences version 20.0. Statistical significance was set at $p \leq 0.05$.

Results:

The mean age of the respondents was 61.1 ± 11.1 years. More than half (52.6%) had high level of spirituality and more than two-thirds (67.1%) of respondents had controlled blood pressure. Respondents with high level of spirituality were 4.76 times more likely to have a good blood pressure control { $p < 0.001$, 95%CI (1.05-14.99)} than those with low level of spirituality.

Conclusion:

Proper understanding and effective utilization of this relationship will assist health professionals and researchers in the appropriate integration of this concept into patients' holistic care with the aim of achieving better blood pressure control among hypertensive patients.

Keywords: Spirituality, Blood pressure, Blood pressure control, Hypertension

Introduction:

Hypertension and its associated complications have become a common and important global public health challenge of the 21st century.¹⁻³ Hypertension is a major risk factor and a powerful predictor of cardiovascular morbidity and mortality.^{1,4}

The global prevalence of hypertension has been increasing. For instance, in the year 2000, the world was estimated to have close to 1 billion people with hypertension with a prevalence rate of 26.4% and this has been projected to increase to 1.56 billion affected individuals with a prevalence rate of 29.2% in 2025, the population burden being greater in developing countries.^{2,5} In Africa, blood pressure control rates were uniformly low and often does not exceed 45% of the hypertensive population.⁷ An hospital-based study in Port Harcourt, Nigeria shows that good hypertension control could only be achieved for just 24.2% of the patients seen in the clinic.⁶ According to a community-based study among an urban population in Zaria, Nigeria shows that hypertension control rate was only 12.4% among patients who have been on antihypertensive medications for over 3months.⁶ These gloomy statistics call for urgent action, especially as the WHO has projected a further 24% increase in the prevalence of non-communicable diseases in Nigeria, in the next 10 years.⁷

Worldwide, there has been a systematic introduction of the concept of spirituality into the medical field with a growing interest in the possible health benefits as a result of having a spiritual belief and following a religious lifestyle.⁸

Spirituality can be defined as “the aspect of humanity that refers to the way individuals seek and express meaning and purpose and the way they experience their connectedness to the moment, to self, to others, to nature, and to the significant or sacred”⁹ Spirituality has been found to play vital roles in health through disease detection and treatment compliance, beliefs that conflict with medical care, medical decision making and spiritual struggles that create stress and impair health outcome.¹⁰

Specifically, it has been noted as an essential factor influencing the health-related quality of life of chronically ill patients due to its integration into positive thoughts and health behaviours, as well as healthy coping styles which alleviate the perceived overwhelming persistent situation.^{11,12}

Spirituality has been specifically associated with good blood pressure control in hypertensive patients.¹³ The importance of addressing spirituality in hypertension like in other chronic diseases is indicated in several studies.^{8,12,14} Spirituality has been shown to be important to hypertensive patients, and some patients have indicated that they use their spirituality for coping and disease management.¹⁵

In a study conducted by Kretchy et al.,⁸ findings revealed that participants with a high level of spirituality and religious participation had significantly lower diastolic and systolic blood pressure than their counterparts with a low level of spirituality (77.8 vs. 84.7 mmHg and 137.2 vs. 149.5 mmHg, respectively) after adjusting for demographic, sociocultural, and psychological variables.¹⁶ Studies have also revealed that people who engaged in spiritual activities had lower levels of cortisol (a biological marker of stress), lipid profiles, glucose control and immune function.¹⁶⁻¹⁸

The relationship between spirituality and hypertension has been studied by several researchers in Africa. Some of them revealed a positive correlation between spirituality, medication adherence and

blood pressure control while others did not.⁸ However; there is a paucity of studies on the relationship between spirituality and hypertension in Nigeria.

This study is designed to determine the role of spirituality in the control of blood pressure among adult hypertensive patients. It also seeks to answer the following research questions:

1. What is the level of spirituality among adult hypertensive patients?
2. What is the level of blood pressure control among the respondents?
3. What is the relationship between the level of spirituality and the socio-demographic characteristics of respondents?
4. What is the relationship between the level of spirituality and blood pressure control?

MATERIALS AND METHOD

STUDY AREA/STUDY DESIGN/POPULATION

The hospital-based descriptive cross-sectional study was conducted among adult hypertensive patients aged 18 years and above attending the Family Medicine Clinic of Federal Teaching Hospital, Ido-Ekiti (FETHI) in rural Southwest Nigeria. The study was conducted over sixteen (16) weeks from April to July 2017 during which a total of 1,986 hypertensive patients were encountered.

SAMPLE SIZE DETERMINATION

The sample size was calculated using the formula for estimating proportions in a cross-sectional study: $n = z^2 pq / d^2$ (For population >10,000)¹⁹ and $nf = n/1 + (n/N)$ (For population <10,000).¹⁹

Where:

n = the minimum sample size when the population is more than 10,000.

nf = the minimum sample size when the population is less than 10,000.

N = the estimate of study population size in a given year, i.e. adult hypertensive patients who attended the Family Medicine Clinic of Federal Teaching Hospital in 2015. ($N = 4,800$).

z_{α} = the level of confidence (The standard normal deviation set at 1.96 which corresponds to the 95% confidence level obtained from the standard statistical table of normal distribution.

p = Estimated prevalence of medication adherence among hypertensive patients in a given population, (using 25.6%, which was the prevalence of medication adherence among hypertensive patients at a tertiary hospital, in South-Western Nigeria).²⁰

$q = 1 - p$.

d = Degree of accuracy desired usually set at 0.05

The minimum sample size was 276 respondents. However, to allow for non-responses during recruitment, a non-response value of 10% (28) was added to the minimum sample size (nf). This gave a minimum sample size of 304 respondents. Hence, 304 respondents were used for the study.

A sampling frame was built, by sorting out the case files of hypertensive patients attending the clinic for the day.

The selected hypertensive patients attending the clinic for the day were selected from the general patient pool, and directed to another convenient room within the family medicine complex. The potential benefits of the study were explained to them and verbal consents were obtained. The consenting patients were subjected to the inclusion and exclusion criteria. The number of consenting hypertensive patients who met the inclusion and exclusion criteria daily were between 19 and 21 out of an average of 25 participants and a total of 1,628 hypertensive patients out of the 1,986 encountered patients both consented and met the inclusion and exclusion criteria during the 16 weeks of the study.

Participants who met the inclusion and exclusion criteria on each clinic day were assigned number from 1 up to the last number for the day.

The sampling interval (K) was calculated using the formula:

$$K = N/n^{19}$$

(Where: N = total number of patients, n = calculated minimum sample size and K = sampling interval).

Hence $K = 1628/304 = 5.36$ which is approximately 5.

Systematic random sampling technique was then used to select the participants on each clinic day; hence, every 6th consenting hypertensive patient who met the inclusion criteria was enrolled for this research. Therefore, an average of four (4) hypertensive patients was recruited per day.

To ensure that a patient was not selected twice, a register for all patients that participated in the study was opened, using their hospital numbers and a colour tag on their file jackets.

INCLUSION CRITERIA

1. All consenting adult hypertensive patients aged 18 years and above.
2. Patients who have been diagnosed and on treatments for hypertension for at least 3 months in the clinic and had recorded a minimum of three (3) consecutive clinic visits in the last six months before recruitment to ensure familiarity with diagnosis and prescribed treatment modalities.

EXCLUSION CRITERIA

1. Patients with co-morbidity (Chronic Kidney Disease, Diabetes)
2. Critically ill patient as at the time of the study (Hypertensive emergency, Congestive cardiac failure, Cerebrovascular accident).
3. Pregnant hypertensive patients because they belong entirely to another class of hypertension.

DATA COLLECTION AND INSTRUMENTS

Socio-demographic variables were obtained using a semi-structured interviewer-administered questionnaire. Clinical parameters (including BP) of respondents were measured, recorded and graded by the researcher.

Assessment of patients' level of Spirituality was done using the Spiritual perspective scale (SPS). It is a 10-item tool designed to measure the extent to which individuals hold certain spiritual beliefs and engage in spiritually related behaviours.^{16,148} It was scored using the mean.^{8,21} The mean spirituality score for this study was 40.9 ± 15.2 . The scoring and determination of the level of

spirituality are found below (section 3.9.3.3.1).²¹ Scores within 10-35 was indicated a low level of spirituality and those ranging between 36-60 indicated a high level of spirituality. Using Cronbach's alpha, reliability has consistently been rated above 0.90 and average inter-item correlations are 0.878.^{8,21,22} The SPS has been used successfully in prior African studies.^{8,22}

Scoring and Determination of the Level of Spirituality²³

Ten items are measuring the beliefs and behaviours of respondents on a 6 point Likert scale.

The minimum and maximum obtainable scores were 10 and 60 respectively.

Range = minimum possible score – maximum possible score

Number of expected outcome = Respondents' level of spirituality dichotomised into high or low .i.e. 2 divisions

For this study, respondents' level of spirituality is dichotomised into high or low.

This gives a range of $60 - 10 = 50$

Interval = $\frac{\text{Range}}{\text{Number of expected outcomes}} (50)$

The number of expected outcomes (2). Interval is 25

Therefore

10 to 35 (minimum score + interval) is a low level of spirituality

36 to 60 is a high level of spirituality

Assessment of Blood Pressure was measured with an Accoson® brand mercury sphygmomanometer.

For this study, a cut-off level of both < 140mmHg and < 90mmHg for SBP and DBP respectively were considered Controlled BP. Uncontrolled BP was considered when the SBP and DBP were $\geq 140\text{mmHg}$ and $\geq 90\text{mmHg}$ respectively using the recommendation of eight Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC-VIII) criteria.²⁴

DATA ANALYSIS

All data collected were analyzed using the Statistical Package for Social Sciences (SPSS) for Windows software version 20 (SPSS Inc., Chicago, IL, USA).²⁵ The data were presented in tabular form, graphs and charts as appropriate. Frequency tables and charts were generated for relevant variables. Means, standard deviations, and percentages were determined as appropriate. The means, median and standard deviation were calculated for continuous variables while categorical variables were analyzed using proportions. The Pearson's chi-square test was used to assess the bivariate association between spirituality with the respondents' socio-demographic characteristics and blood pressure control and the Fisher's exact test was used in analyzing variables with cells less than 5 counts. Binary logistic regression analysis was used to assess the association between the independent variable (spirituality) and dependent variable (blood pressure control) while adjusting for the socio-demographic characteristics. A P-value of equal or less than 0.05 was taken to be statistically significant.

ETHICAL CLEARANCE AND CONSENT

Ethical approval was obtained from the Ethics and Research Committee of Federal Teaching Hospital, Ido-Ekiti (Protocol Number: ERC/2016/02/25/08A). Informed verbal consent was obtained from the willing participants. To maintain confidentiality, no names appeared on the questionnaires, but only numbers were used as identifiers. The reporting of this study conforms to the strengthening the Reporting of observational studies in Epidemiology (STROBE) statement²⁶. All patients' details have been de-identified.




Figure 1: Study flowchart

RESULTS

4.1 Characteristics of Respondents

A total of 304 respondents participated in the study. The respondent's ages ranged between 18 and 82 years, their mean age was 61.1 ± 11.1 years and the majority (57.9%) were older adults (65-82 years). The majority of respondents were female (53.6%) and were married (76.0%), Christians (78.6%) and rural dwellers (64.5%) from the Yoruba tribe (74.7%). Most respondents were traders (37.5%) who had a primary level of education (31.9%) and earned incomes below the poverty line (57.7%). The details of the respondent's socio-demographic characteristics were as shown in Table 1 below.

Table 2 showed the Spiritual perspective scale based on the spiritual behaviours and beliefs of respondents. Spiritual behaviours that contributed significantly to the high level of spirituality in this study were: private prayers/meditations ($\chi^2 = 3.42$, SD = 0.97) and spiritual discussions with family and friends ($\chi^2 = 3.19$, SD = 0.92), while Spiritual beliefs that contributed significantly to the high level of spirituality in this study were: forgiveness ($\chi^2 = 3.27$, SD = 0.74), seeking spiritual guidance before making decisions ($\chi^2 = 3.26$, SD = 0.87) and regarding spirituality as a means of finding live meaning and solving life's puzzle ($\chi^2 = 3.26$, SD = 0.64).

Table 3 showed the level of spirituality among the respondents. More than half (52.6%) of the respondents had a high level of spirituality while 47.4% had a low level of spirituality. The mean spirituality score obtained was 40.9 ± 15.2 .

Figure 2: A bar chart showing the level of blood pressure control among the respondents. More than two-thirds (67.1%) of the respondents had controlled blood pressure while 32.9% had uncontrolled blood pressure

Table 4 showed the relationship between spirituality and blood pressure among the respondents. It showed that Blood pressure control improves with increasing spirituality. More than two-thirds (70.1%) of those with a high level of spirituality also had good blood pressure control. The relationship was statistically significant as $p < 0.05$.

Table 5 showed the relationships between the socio-demographic characteristics of respondents and their level of spirituality. Older adult (64.4%), female (67.5%), married (58.4%), Christian (56.5%), Yoruba (63.0%) and urban dwelling (55.1%) respondents had high level of spirituality. Also, respondents that had a tertiary level of education (70.4%) who

are retired (66.7%) and earned below the poverty line (75.4%) had a high level of spirituality. All variables were statistically significant with $p < 0.05$ except the area of domicile.

Table 6 explains how some socio-demographic characteristics predicted the level of spirituality. Respondents who were middle-aged adults (1.54 times), Married (7.54 times), Christians (3.38 times), Yoruba (2.23 times), Postgraduate (2.70 times), Professionals (3.54), Less than 5 years on antihypertensive drug (2.72), Controlled blood pressure (4.76 times) and were on 2 drugs or less (2.80 times) were more likely to have a high level of spirituality. However, respondents who were male gender (0.71 times), farmers (0.78) and earned income below the poverty line (0.74 times) were less likely to have high medication adherence.

DISCUSSION

SOCIO-DEMOGRAPHIC CHARACTERISTICS OF RESPONDENTS

In Table 1, the socio-demographic profile of the respondents showed that the mean age of patients with hypertension in this study was 61.1 ± 11.1 years with the majority of the respondents being above 65 years. This was higher than 50.9 ± 9.1 years reported in a similar hospital-based study in Port-Harcourt by Akpa *et al*²⁸ and 50.6 ± 1.5 years reported in a study by Azinge *et al* in Lagos²⁹. It was however lower than the mean age of 63.1 ± 5.7 years found in an international survey.³⁰ The reason for the higher preponderance of hypertension in the older age group found in this study compared to other studies might be because this study was done in a rural population as this usually consists of older age group due to rural to urban migration of the young age groups. However, Ido-Ekiti town is a non-industrialised semi-urban setting surrounded by rural settlements.

The majority of the respondents {163 (53.6%)} in this study were females. This aligned with the findings of Iloh *et al.* (65.2%)³¹ who also reported a higher prevalence of hypertension in females. It is worthy of note that women are protected from most cardiovascular events compared to men, until after menopause, during which women are at increased risk of cardiovascular complications compared to premenopausal women.^{32,33} Gender differences in

blood pressure is inconsistent in many population studies, while some studies quoted a higher prevalence of hypertension among males,^{34,35} others reported no significant gender difference.^{28,36} These inconsistencies could be as a result of different age populations or the methodology used in these studies.

In this study about two-thirds {231 (76%)} of all the respondents were married. This was similar to findings by Marfo *et al.*³⁷ who reported 74% married hypertensive patients.³⁷ However, findings in this study was higher than the 66.4% reported in Ibadan by Ekore *et al.*³⁰ This difference may be due to the variation in marriage culture in different states and regions among the study populations in Nigeria. The percentage of respondents who did not have any formal education was 12.5%. It could be inferred that 87.5% had formal education in this study. This might be due to the study area which is located in the South-Western part of the country where there is a high literacy level which might influence patients' health-seeking behaviour in terms of them presenting in the hospital for the management of their disease. This study confirms the rating of Ekiti State as an educationally advantaged state. The reported level of literacy was much higher in a hospital-based study done among young adults in Ibadan, South-western Nigeria where about 79.1% of respondents had a minimum of secondary school education³⁶ and 40.0% reported in Lagos among market women.³⁸ These differences may be due to the different study populations among whom the studies were conducted.

LEVEL OF SPIRITUALITY AMONG RESPONDENTS

In Table 3, this study showed that the majority of the respondents (52.6%) had a high level of spirituality. The finding of the majority of the respondents being highly spiritual in this study may not be unconnected with the cultural practices which were inherent in Nigeria. This was similar to a study by Ibraheem *et al.*³⁹ in the relationship between self-reported health status and spirituality among adult patients in Ibadan, Nigeria who found a 51% level of spirituality.³⁹ In contrast, Jeri and Lynda reported 100% spirituality level among Nigerians

living in America.⁴⁰ This is higher than that obtained in the present study, and the difference may be due to the different study populations among whom the studies were conducted and the method used in the study.

In Table 2. Spiritual behaviours that contributed significantly to the high level of spirituality in this study according to Table 3 were: private prayers/meditations ($\chi^2 = 3.42$, SD = 0.97) and spiritual discussions with family and friends ($\chi^2 = 3.19$, SD = 0.92). Also, Spiritual beliefs that contributed significantly to the high level of spirituality in this study were: forgiveness ($\chi^2 = 3.27$, SD = 0.74), seeking spiritual guidance before making decisions ($\chi^2 = 3.26$, SD = 0.87) and regarding spirituality as a means of finding meaning to live and solving life's puzzle ($\chi^2 = 3.26$, SD = 0.64).

5.8. RELATIONSHIP BETWEEN SPIRITUALITY AND BLOOD PRESSURE CONTROL

In Table 4, this study has shown a high spirituality level of 52.6% and a blood pressure control level of 67.1%. The Chi-square test showed a significant relationship between them with $p < 0.001$. This is similar to the finding in a study done on spirituality and blood pressure control by Kretchy et al.,⁸ in Umuahia, Nigeria, which showed participants with a high level of spirituality and religious participation had significantly lower diastolic and systolic blood pressure than their counterparts with a low level of spirituality (77.8 vs. 84.7 mmHg and 137.2 vs. 149.5 mmHg, respectively) after adjusting for demographic, sociocultural, and psychological variables.⁸

However, the findings in this study are higher than what was observed by Quingtao et al.⁴¹ in a study conducted in china who found that 53.4% of those who had a high level of spirituality also had good blood pressure control.

Also, logistic regression done in this study according to Table 6, showed that for a single unit increase in the level of spirituality, hypertensive patients were 4.76 times more likely to achieve good blood pressure control (OR= 4.76, 95%CI = 1.05-14.99, $p < 0.001$). Hence, a

high level of spirituality is a significant predictor of good blood pressure control among hypertensive patients.

RELATIONSHIP BETWEEN SPIRITUALITY AND SOCIO-DEMOGRAPHIC CHARACTERISTICS OF RESPONDENTS

In Table 5, this study found that spirituality levels increased with age and specifically, more than half (64.4%) of the older aged group has a higher level of spirituality. This could be as a result of the fact that, as people get older they tend to find solace and meaning to life in the transcendent and hence tend to have more inclination towards spirituality.⁴² The above finding is in agreement with hospital-based research conducted by Ibraheem et al (83.9%)³⁹ who also reported an increased level of spirituality with age.

This study established the fact that women have higher spirituality than men (67.5%). This is possibly due to the vulnerability of women in this part of the world and women tend to find solace in their spiritual and religious practices. This is also similar to what Jeri et al. (63%)⁴⁰ found in his study.

In this study, a higher spirituality score was found among the respondents who were married (58.4%) when compared with the respondents who were single or divorced and this difference was statistically significant ($p=0.001$). This might be due to the possible encouragement by the spouse towards spirituality, which is not found among the single and the divorced. This trend was in agreement with the finding of Ibraheem et al.³⁹ in his study on the relationship between self-reported health status and spirituality among adult patients attending the General outpatient clinic of a tertiary hospital in Ibadan where he reported a higher spirituality level among married people (68%) compared with those who were not married.³⁹

It was observed in this study that spirituality increased with an increase in the level of literacy and this was statistically significant ($p<0.001$). This might be due to the high literacy level found in a large proportion of the respondents in this study. This is different from the

previous finding of Silva et al.⁴³ who reported that people of low socio-economic class tend to be more spiritual and that the prevalence of spiritual commitment tends to drop off among higher-income categories.⁴³

5.10. CONCLUSION

This study has shown the relationship between spirituality and blood pressure control among adult hypertensive patients in rural Southwestern, Nigeria.

It had shown 52.6% high spirituality level and 67.1% good blood pressure control.

The study found that more than two-thirds of respondents with a high level of spirituality had good blood pressure control, while more than three-quarters of respondents with a low level of spirituality had poor blood pressure control. These relationships were statistically significant.

These associations provide direction for future studies that will aid in understanding how health professionals can use this information to provide culturally sensitive and patient-centred care that will improve blood pressure control among patients with hypertension.

Several studies emphasized the importance of spirituality in overall health and some behaviour of chronically ill persons; however, spirituality has been largely overlooked in the study of blood pressure control. This study addresses this gap in the literature regarding the association between spirituality and blood pressure control in adult hypertensive patients.

5.11. RECOMMENDATIONS

The study has highlighted the significance of the relationship between spirituality and blood pressure control among patients with hypertension; hence, Physicians should take appropriate spiritual history while managing patients with hypertension to identify how spirituality can positively or negatively affect blood pressure management and control.

Also, the Physicians should encourage the health facilities to have chaplains and Imams that patients can be referred to for spiritual assistance when needed.

It is also important that the Physicians understand their limits in the process of integrating spirituality into their patient's care.

5.12. LIMITATIONS OF THE STUDY

1. The study is a cross-sectional study done in a tertiary health institution, the findings may not apply to the general population.
2. Subjects' ambulatory or home blood pressure readings in out-of-clinic settings were not monitored in this study, and thus the determination of optimal blood pressure control was based on clinic-based measurement alone instead of an average over a period of time.

Acknowledgements: The authors would like to appreciate nurses and resident doctors of family medicine department and the management of FETHI where the study was conducted.

Conflicts Of Interest: The authors declare that they have no conflicts of interest.

Funding: The researcher received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Availability of data and materials: The datasets for this study would be made available from the correspondence author on a reasonable request.

Authors' contribution:

1. AO - Conceptualization of the study, designed the study protocol, data acquisition and analysis and drafted the initial manuscript
2. PO - Literature review, data analysis and review of manuscript for intellectual content.
3. OMS - Critically revised the protocol for methodological and intellectual content.
4. AOI. Literature review, review of manuscript for intellectual content.

All authors have read and approved the final version of the manuscript prior to submission.

REFERENCES

1. Ogah OS, Okpechi I, Chukwuonye II, Akinyemi OJ, Onwubere JC, Falase OA. Blood pressure, prevalence of hypertension and hypertension related complications in Nigerian Africans: A review. *World J Cardiol* 2012; 4(12): 327-340.
2. Quraishi UM, Begum S, Agarwal A, Chandrasekhar S, Shamji U, Naeem A. A Study to Know the Prevalence and Awareness of Hypertension in Market Population of Ras Al Khaimah, UAE. *J. Basic. Appl. Sci. Res* 2013; 3(2): 458-462.
3. Abdullahi AA, Amzat J. Knowledge of hypertension among the staff of University of Ibadan. *J. Public Health Epidemiol.* 2011; 3:204-209.

4. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014. Evidence-based guideline for the management of high blood pressure in adults: Report from the panel members in appointed to the Eight Joint National Committee (JNC 8). JAMA. 2014; 311: 507-520.
5. Chythra RR, Veena GK, Avinash S, Asha K. High Blood Pressure Prevalence and Significant Correlates: A Quantitative Analysis from Coastal Karnataka, India. ISRN Preventive Medicine 2013: 1-6.
6. Oyati AI, Orogade AA, Danbauchi SS, Azuh PC. Awareness, Treatment and Control of Hypertension among Hypertensives in Zaria. Journal of Medicine in the Tropics 2011; 13(2): 139-144.
7. World Health Organisation (WHO). Facing the facts: The impact of chronic disease in Nigeria. Geneva: WHO; 2005. Available from: http://www.who.int/chp/chronic_disease_report/media/nigeria.pdf. Accessed 17th of November 2015.
8. Kretchy I, Owusu-Daaku F, Samuel D. Spiritual and religious beliefs: do they matter in the medication adherence behaviour of hypertensive patients? Journal of BioPsychoSocial Medicine 2013; 7:15-21.
9. Prasad DS, Zubair k, Das BC. Prevalence and predictors of adult hypertension in an urban eastern Indian population. 2012; 4(1): 49-52.
10. Thomas R M, Min D. Spirituality and medicine Ethics in medicine. 2014; 27(5): 1218-1220.
11. Kremer H, Ironson G. Everything Changed: Spiritual Transformation in People with HIV. International Journal of Psychiatry in Medicine. 2009; 39(3): 243–262.
12. Lewis LM. Medication adherence and spiritual perspectives among african american older women with hypertension. A qualitative study. J Gerontol Nurs. Jun 2011; 37(6):34-41.

13. Wyatt SB, Akyzbekova EL, Wofford MR. Prevalence, awareness, treatment, and control of hypertension in the Jackson Heart Study. *Hypertension*. 2008;51(3):650-656.
14. Buck AC, Williams DR, Musick MA, Sternthal MJ. An examination of the relationship between multiple dimensions of religiosity, blood pressure, and hypertension. *Social Science and Medicine*. 2009; 68(2): 314–322.
15. Giaquinto S, Spiridigliozzi C. Possible influence of spiritual and religious beliefs on hypertension. *Clin Exp Hypertens*. 2008;29(7):457-464.
16. Kharitonov SA. Religious and Spiritual Biomarkers in both Health and Disease. *Religions*. 2012; 3(2):467-497.
17. Kreikebaum S, Guarneri E, Talavera G, Madanat H, Smith T. “Evaluation of a holistic cardiac rehabilitation in the reduction of biopsychosocial risk factors among patients with coronary heart disease.” *Psychol.health Med*. 2011; 16:276-290.
18. Holt-Lunstad J, Steffen PR, Sandberg J, Jensen B. “Understanding the connection between spiritual well-being and physical health: An examination of ambulatory blood pressure, inflammation, blood lipids and fasting glucose.” *J. Behav. Med*. 2011; 34:477-488.
19. Araoye MO. Sample size determination. *Research Methodology with Statistics for Health and Social Sciences*, Ilorin: Nathadex Publishers. 2004; 115–121.
20. Ordinioha B. The prevalence of hypertension and its modifiable risk factors among lecturers of a medical school in Port Harcourt, south-south Nigeria: Implications for control effort. *Nigerian Journal of Clinical Practice* 2013; 16(1): 1 – 11.
21. Fitchett G, Powell LH. Daily spiritual experiences, systolic blood pressure, and hypertension among midlife women in SWAN. *Ann Behav Med*. 2009;37(3):257-267.

22. Reed PG. Spirituality and well-being in terminally ill hospitalized adults. *Research in Nursing and Health*. 1987; 10:335–344.
23. Petrie A, Sabin C. *Medical Statistics at a Glance*. 3rd ed. London: Wiley-Blackwell; 2009. Chapter 6: Describing data: the “spread”; p.19-24.
24. Gee ME, Campbell NR, Sarrafzadegan N, Jafar T, Khalsa TH, Mangat B, et al. Standards for the uniform reporting of hypertension in adults using population survey data: recommendations from the World Hypertension League expert committee. *The Journal of Clinical Hypertension*. 2014; 16: 773-781.
25. SPSS Inc. *Predictive Analytics Software Statistics for Windows, Version 20.0*. Chicago: SPSS Inc. 2009.
26. Von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP, et al. The strengthening the Reporting of observational studies in Epidemiology (STROBE) statement; guidelines for reporting observational studies. *Ann Intern Med*. 2007; 147: 573-7.
27. Poverty | Data [Internet]. [Data.worldbank.org](http://data.worldbank.org). 2017 [cited 5 July, 2017]. Available from: <http://data.worldbank.org/topic/poverty#>.
28. Akpa MR, Alasia DD, Emem-Chioma PC. An appraisal of hospital based blood pressure control in Port Harcourt, Nigeria. *The Nigerian Health Journal*. 2008; 8(1 - 2): 27-30.
29. Azinge EC, Sofola OA, Silva BO. Relationship between Salt intake, Salt-taste threshold and blood pressure in Nigerians. *West Afr J Med*. 2011;30(5):373-376.
30. Thoenes M, Bramlage P, Zhong S, Shang S, Volpe M, Spirk D. Hypertension control and cardio-metabolic risk: a regional perspective. *Cardiology research and practice*. 2012; 2012:1-10.
31. Iloh GU, Amadi AN, Okafor GO, Ikwudinma AO, Odu FU, Godswill-Uko EU. Adherence to lifestyle modifications among adult hypertensive Nigerians with essential hypertension in a primary care clinic of a tertiary hospital in resource-poor

- environment of Eastern Nigeria. *British Journal of Medicine and Medical Research*. 2014;4(18):3478-3490.
32. Shaw BH, Protheroe CL. Sex, drugs and blood pressure control: the impact of age and gender on sympathetic regulation of arterial pressure. *The Journal of physiology*. 2012;590(12):2841-2843.
33. Maranon R, Reckelhoff JF. Sex and gender differences in control of blood pressure. *Clin Sci (Lond)*. 2013;125(7):311-318.
34. Dzudie A, Kengne AP, Muna WF, Ba H, Menanga A, et al. Prevalence, awareness, treatment and control of hypertension in a self-selected sub-Saharan African urban population: a cross-sectional study. *BMJ open*. 2012; 2(4):1136-1217.
35. Ekwunife OI, Aguwa CN. A meta-analysis of prevalence rate of hypertension in Nigerian populations. *Journal of Public Health and epidemiology*. 2011;3(13):604-607.
36. Ekore RI, Ajayi IO, Arije A. Case finding for hypertension in young adult patients attending a missionary hospital in Nigeria. *African health sciences*. 2009;9(3):193-199.
37. Marfo AFA, Owusu-Daaku FT, Addo MO, Saana II. Ghanaian hypertensive patients understanding of their medicines and life style modification for managing hypertension. *Int J Pharm Pharm Sci*. 2014; 6(4): 165-170.
38. Busari A, Olayemi S, Oreagba I, Alabidun A. Educational intervention as a strategy for improving blood pressure status of market women in Lagos, Nigeria. *The Internet Journal of Health*. 2010; 11(1):771-777.
39. Ibraheem AB, Ibraheem WA, Adebusoye L. The relationship between self-reported health status and spirituality among adult patients attending General Outpatient Clinic of Tertiary Hospital in Ibadan. *Ann Ib Postgrad Med*. 2014;12(1):31-37.

40. Jeri D, Linda LD. Spiritual perspectives and health: A random survey in a Southern State. *Online J Rural Nurs Health Care*. 2009;9(2):23-31.
41. Qingtao M, Ying Xu, Rufeng Shi, Xin Z, Si W, Kai L, Xiaoping C. Effect of religion on hypertension in adult Buddhists and residents in China: A cross-sectional study. *Sci Rep*. 2018; 8: 8203- 8210.
42. Oluwole DA. Connection of self disclosure, sexual efficacy and spirituality among Nigerian women. *Pak J Soc Sci*. 2008;5(5):464-469.
43. Silva CF, Borges FR, Avelino CCV, Miarelli AC, Vieira GA, Goyata ST. Spirituality and Religiosity in patients with systemic arterial hypertension. *Rev bioet (impr)*. 2016; 24(2):332-343.

Table1: Socio-demographic Characteristics of Respondents

Variable	Frequency (n = 304)	Percentage (%)
Age (years)	<i>61.1 ± 11.1</i>	
Young Adult (18-35)	6	2.0
Middle-Aged Adult (36-64)	122	40.1
Older Adult (65-82)	176	57.9
Sex		
Male	141	46.4
Female	163	53.6
Marital Status		
Single	10	3.3

Married	231	76.0
Divorced/Separated	43	14.1
Widowed	20	6.6
Religion		
Christianity	239	78.6
Islam	59	19.4
Traditionalist	6	2.0
Domicile		
Urban	108	35.5
Rural	196	64.5
Ethnicity		
Yoruba	227	74.7
Igbo	46	15.1
Hausa	26	8.6
Others	5	1.6
Educational Level		
None	38	12.5
Primary	97	31.9
Secondary	63	20.7
Tertiary	88	29.0
Post Graduate	18	5.9
Occupation		
Professionals	61	20.1
Trading	114	37.5
Farming	67	22.0
Retiree	32	10.5
Self – Employed	30	9.9
Income*		
< Poverty Line (<₦20,520)	175	57.7
≥Poverty Line (≥₦20,520)	129	42.3

₦20,520 is the expected minimum income per month equivalent of \$1.90 per day at \$1 equals to ₦360 (as of July 5, 2017)²⁷

Table 2a: Spiritual Perspective Scale

S/N	SPS (Spiritual Behaviours)	Not at all	Less than once a year	About once a year	About once a month	About once a week	About once a day	ΣF	Mean	Standard Deviation
1	In talking with family and friends, how often do you mention spirituality?	4	10	16	35	102	137	304	3.19	0.92
2	How often do you share with others the problems and joys of living according to your spiritual beliefs?	10	11	10	54	102	117	304	3.08	1.10
3	How often do you read spiritually-related material?	14	15	20	50	92	113	304	2.95	1.11
4	How often do you engage in private prayer and meditation?	9	10	6	36	48	195	304	3.42	0.97

Table 2b: Spiritual Perspective Scale

S/N	SPS (Spiritual Beliefs)	Strongly Disagree	Disagree	Disagree more than agree	Agree more than disagree	Agree	Strongly Agree	ΣF	Mean	Standard Deviation
5	Forgiveness is an important part of my spirituality.	5	7	10	11	154	117	304	3.27	0.74
6	I seek spiritual guidance in making decisions in my everyday life.	5	19	20	24	127	109	304	3.26	0.87
7	My spirituality is a significant part of my life.	5	20	27	31	120	101	304	3.25	0.88
8	I frequently feel very close to God or a “higher power” in prayer, during public worship or at important moments in my daily life.	2	19	32	19	132	100	304	3.24	0.80
9	My spiritual views have had an influence upon influenced my life.	5	10	33	25	130	101	304	3.30	0.77
10	My spirituality is especially important to me because it answers many questions about the meaning of life.	2	7	21	31	151	92	304	3.26	0.64

Table 3: Assessment of the Level of Spirituality among the Respondents

Variable	Frequency n= 304	Percentage (%)
Spirituality Perspective Scale		
Low (10-35)	144	47.4
High (36-60)	160	52.6
Total	304	100.0
<i>Mean Spirituality Score ± SD</i>	<i>40.9 ± 15.2</i>	




Figure 2: A graphical representation of the level of Blood Pressure Control among Respondents

Table 4: Relationships between Blood Pressure Control and Spirituality among the Respondents

Variable	Spirituality			χ^2	p-value
	Low (10-35)	High (36-60)	Total		
	(%) n = 144	(%) n = 160	N=304		
Blood Pressure Control					
Uncontrolled ($\geq 140/90$ mmHg)	83 (83.0)	17 (17.0)	100 (100)	75.88	<0.001
Controlled (< 140/90mmHg)	61 (29.9)	143 (70.1)	204 (100)		

Table 5: Relationships between the Level of Spirituality and Sociodemographic Characteristics of Respondents

Variable	Spirituality			χ^2	p-value
	Low (%) n = 144	High (%) n = 160	Total N=304		
Age (years)					
Young Adult (18-35)	4 (66.7)	2 (33.3)	6 (100)		0.004[‡]
Middle-Aged Adult (36-64)	77 (63.6)	44 (36.4)	121 (100)		
Older Adult (65-82)	63 (35.6)	114(64.4)	177 (100)		
Sex					
Male	91 (64.5)	50 (35.5)	141 (100)	31.09	<0.001
Female	53 (32.5)	110 (67.5)	163 (100)		
Marital Status					
Single	10 (100.0)	0 (0.0)	10 (100)		0.001[‡]
Married	96 (41.6)	135 (58.4)	231 (100)		
Divorced	15 (75.0)	5 (25.0)	20 (100)		
Separated	14 (60.9)	9 (39.1)	23 (100)		
Widowed	9 (45.0)	11 (55.0)	20 (100)		
Religion					
Christianity	104 (43.5)	135 (56.5)	239 (100)		0.018[‡]
Islam	37 (62.7)	22 (37.3)	59 (100)		
Traditionalist	4 (66.7)	2 (33.3)	6 (100)		
Domicile					
Urban	88 (44.9)	108 (55.1)	196 (100)	1.35	0.245
Rural	56 (51.9)	52 (48.1)	108 (100)		
Ethnicity					
Yoruba	84 (37.0)	143 (63.0)	227 (100)		<0.001[‡]
Igbo	39 (84.8)	7 (15.2)	46 (100)		
Hausa	18 (69.2)	8 (30.8)	26 (100)		
Others	3 (60.0)	2 (40.0)	5 (100)		
Educational Status					
None	24 (63.2)	14 (36.8)	38 (100)	21.57	<0.001
Primary	57 (58.8)	40 (41.2)	97 (100)		
Secondary	31 (49.2)	32 (50.8)	63 (100)		
Tertiary	26 (29.6)	62 (70.4)	88 (100)		
Post Graduate	6 (33.3)	12 (66.7)	18 (100)		
Occupation					
Professionals	24 (39.3)	37 (60.7)	61 (100)	13.43	0.009
Trading	49 (43.0)	65 (57.0)	114 (100)		
Farming	43 (64.2)	24 (35.8)	67 (100)		
Self Employed	18 (56.3)	14 (43.7)	32 (100)		
Retired	10 (33.3)	20 (66.7)	30 (100)		
Income					
<Poverty Line	43 (24.6)	132 (75.4)	175 (100)	85.96	<0.001
≥Poverty Line	101 (78.3)	28 (21.7)	129 (100)		

[‡] = Fisher's Exact Test

Table 6: Logistics Regression for predictors of Spirituality among the Respondents

Variable		OR	95% CI		p-value
			Lower	Upper	
Age group	Young Adult (18-35)	0.54	0.53	2.68	0.051
	Middle-Aged Adult (36-65)	1.54	1.05	8.42	0.003
	Older Adult (>65)	1.0 (RC)			
Sex	Male	0.71	0.04	0.90	0.021
	Female	1.0 (RC)			
Religion	Christianity	3.38	1.22	10.81	0.004
	Islam	1.11	1.08	3.29	0.031
	Traditionalist	1.0 (RC)			
Ethnicity	Yoruba	2.23	1.19	9.41	0.011
	Igbo	1.53	1.04	2.92	0.016
	Hausa	1.10	0.03	3.05	0.062
	Others	1.0 (RC)			
Education	None	1.0 (RC)			
	Primary	1.55	0.13	6.93	0.051
	Secondary	1.11	1.04	7.42	0.012
	Tertiary	2.13	1.67	10.54	0.037
	Postgraduate	2.70	1.08	9.93	0.030
Occupation	Professionals	3.54	1.04	9.47	0.022
	Trading	1.21	1.02	6.21	0.004
	Farming	0.78	0.09	0.99	0.016
	Self – Employed	3.02	1.98	9.83	0.039
	Retired	1.0 (RC)			
Income	< Poverty Line	0.74	0.10	0.97	0.006
	>Poverty Line	1.0 (RC)			