

Sociodemographic inequity in COVID-19 vaccine uptake among youth in Zimbabwe

Leyla Larsson¹, Chido Dziva Chikwari^{1,2}, Victoria Simms^{1,2}, Mandikudza Tembo^{1,3}, Agnes Mahomva⁴, Owen Mugurungi⁵, Richard Hayes², Constance Mackworth-Young^{1,6}, Sarah Bernays^{3,7}, Constancia Mavodza^{1,2}, Tinotenda Taruvinga^{1,8}, Tsitsi Bandason¹, Ethel Dauya¹, Rashida A Ferrand^{1,8}, Katharina Kranzer^{1,8,9}

Author affiliations

¹The Health Research Unit Zimbabwe, Biomedical Research and Training Institute, Harare, Zimbabwe

²MRC International Statistics and Epidemiology Group, Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, UK

³Faculty of Public Health and Policy, London School of Hygiene and Tropical Medicine, London, UK

⁴National Response to the COVID-19 Pandemic, Office of the President and Cabinet, Harare, Zimbabwe

⁵Ministry of Health and Child Care, Harare, Zimbabwe

⁶Department of Global Health and Development, London School of Hygiene and Tropical Medicine, London, UK

⁷Faculty of Medicine and Health, University of Sydney, Sydney, Australia

⁸Department of Clinical Research, London School of Hygiene and Tropical Medicine, London, UK

⁹Department of Infectious Diseases and Tropical Medicine, Ludwig-Maximilians-Universität, Munich, Germany

Corresponding author

Leyla Larsson
The Health Research Unit Zimbabwe (THRU ZIM)
Biomedical Research and Training Institute
10, Seagrave Road
Harare, Zimbabwe
E-Mail: llarsson3@outlook.com

Acknowledgements

The CHIEDZA study is funded by the Wellcome Trust (Senior Fellowship to RAF: 206316/Z/17/Z)

Abstract

Introduction

COVID-19 vaccine acceptance research has mostly originated from high-income countries and reasons why youth may not get vaccinated may differ in low-income settings. Understanding vaccination coverage across different population groups and the sociocultural influences in healthcare delivery is important to inform targeted vaccination campaigns.

Methods

A population-based survey was conducted in 24 communities across three provinces (Harare, Bulawayo and Mashonaland East) in Zimbabwe between October 2021 and June 2022. Youth aged 18 - 24 years were recruited using random sampling. Data on sociodemographic information and COVID-19 vaccination uptake and reasons for non-uptake were collected.

Results

A total of 17,682 youth were recruited (n=10,743, 60.8% female). The median age of survey participants was 20 (IQR: 19 – 22) years. Almost two thirds (n=10,651, 60.2%) of participants reported receiving at least one dose of COVID-19 vaccine. A higher proportion of men than women had been vaccinated (68.9% vs 54.7%), and vaccination prevalence increased with age (<19 years: 57.5%, 20-22: 61.5%, >23: 62.2%). Lack of time to get vaccinated, belief that the vaccine was unsafe and anxiety about side effects (particularly infertility) were the main reasons for not getting vaccinated. Factors associated with vaccination were male sex

(OR=1.69, 95%CI:1.58-1.80), increasing age (>22 years: OR=1.12, 95%CI:1.04-1.21), education level (post-secondary: OR=4.34, 95%CI:3.27-5.76), and socioeconomic status (least poor: OR=1.32, 95%CI:1.20-1.47).

Conclusion

This study found vaccine inequity across age, sex, educational attainment and socioeconomic status among youth. Strategies should address these inequities by understanding concerns and tailoring vaccine campaigns to specific groups.

What is already known on this topic

Many countries have faced challenges when rolling out COVID-19 vaccines. Infrastructure, logistics, misinformation and vaccine hesitancy have been barriers to vaccine access and uptake globally. Vaccine nationalism by high-income countries has particularly affected countries in Africa and Asia, resulting in inequity between countries and regions.

What this study adds

Vaccine uptake among youth in Zimbabwe was more than 50% across all age-groups. Men, those with more education and those living under less socially deprived socioeconomic conditions were more like to be vaccinated. Fear of side effects and myths circulating on social media were barriers. Religion was less of a barrier than other studies reported, likely due to religious institutions' collaborations in COVID-19 vaccination efforts.

How this study might affect research, practice or policy

Vaccination campaigns should actively address specific concerns of communities, especially concerns around fertility and early death, and provide vaccines in easy-access and convenient locations. Involving community leaders in both education and vaccination efforts is pivotal given the trust and influence they have.

Introduction

By February 2023, more than 750 million SARS-CoV2 infections and 6.8 million COVID-19 associated deaths had been reported globally [1]. The development of vaccines against SARS-CoV2, which primarily protect against severe disease, has however greatly reduced both COVID-19-related mortality and morbidity [2,3]. Global vaccination programmes were fast-tracked under the COVID-19 Vaccines Global Access (COVAX) initiative at the beginning of 2021 [4]. Vaccination rates have, however, failed to meet the targets of vaccinating 70% of the world's population against COVID-19 by mid-2022 set by the World Health Organization (WHO), especially in the African continent [5].

By February 3rd 2023, Africa has reported a vaccination coverage (receiving at least 1 dose of a COVID-19 vaccine) of eligible population of 46% and of total population of 27%, with coverage varying greatly from <6% in Madagascar, Cameroon and the Democratic Republic of Congo to more than two thirds of the eligible population in Rwanda and Liberia [6,7]. Differences are partly explained by availability of vaccines, infrastructural constraints, and COVID-19 vaccine nationalism in high income countries (HIC) leading to insufficient donations of vaccine to low- and middle-income countries (LMICs) [8]. In addition, vaccine hesitancy has been reported to be one of the main barriers towards meeting global vaccine coverage targets [9-11]. Studies conducted both in HICs and LMICs have reported widespread hesitancy towards receiving the COVID-19 vaccine, despite most of the population having previously received vaccines for other viruses [12,13]

Zimbabwe recorded its first case of COVID-19 in March 2020 and was subsequently among the first African countries to implement vaccination, with the first vaccine administered on 18 February 2021 [14]. In contrast other countries in the region, Zimbabwe procured COVID-19 vaccines (SinoPharm and SinoVac) through a bilateral agreement with China [15,16]. While non-availability of vaccines was a major barrier for vaccine uptake in other African countries, this was not the case in Zimbabwe [16]. Initially, healthcare workers and people working at borders were targeted for COVID-19 vaccination, followed by those with chronic conditions and essential workers such as teachers [17]. With the availability of more vaccine doses, eligibility was rapidly extended to the rest of the adult population (people aged 18 and over). In November 2021, vaccines were also made available to adolescents aged 16-18 years and in March 2022 to children aged 12 years and older [16-18]. The initial vaccine schedule was two doses of either SinoPharm or SinoVac three weeks apart. A third booster dose was introduced in January 2022 [19-21].

The Zimbabwe vaccination campaign was administered through vaccination centres established in hospitals and clinics and outreach services. In addition, the vaccination campaign included educational programmes in schools, national mobilisation of frontline workers to assist in getting to hard-to-reach populations, and nationwide health education broadcasting [18,22-23]. Once vaccination eligibility was extended to adolescents the government partnered with organisations such as United Nations Children's Fund (UNICEF) to generate vaccination messaging targeting children and adolescents [18,25-27]. As of February 3rd 2023, the Zimbabwean Government estimates that 44% of the total population have received at least one COVID-19 vaccine dose. With approximately 60% of the population being eligible (12+ years), coverage among those eligible was 77% [1,2].

Despite the tremendous efforts put into the vaccination campaign, a considerable proportion of the general population remains unvaccinated and more recently vaccination coverage has stagnated [14]. While a Partnership for Evidence-Based Response to COVID-19 survey in September 2021 demonstrated that 82% of respondents in Zimbabwe were satisfied with the government's response, there was still some vaccine hesitancy [28]. Findings from a survey in Zimbabwe in January 2022 indicated that half of the population displayed some vaccine hesitancy, mostly driven by social media misinformation and lack of trust in official information [29]. Furthermore, there is a lack of evidence in the literature regarding uptake of COVID-19 vaccination among young people and potential causes of vaccine hesitancy in Zimbabwe.

This study investigates self-reported COVID-19 vaccine uptake among 17,862 young people aged 18-24 years across three provinces (Harare, Bulawayo, and Mashonaland East) in Zimbabwe and explores sociodemographic and -economic factors associated with uptake and reasons for possible vaccine hesitancy.

Methods

Enrolment

This study used data from a population-based survey which was conducted to ascertain the outcome of the CHIEDZA trial. CHIEDZA is a cluster randomised controlled trial conducted in three provinces (Harare, Bulawayo, and Mashonaland East) in Zimbabwe investigating the impact of providing community-based integrated HIV and sexual and reproductive health services to young people aged 16-24 years on population-level HIV outcomes

(NCT03719521) [30]. Taking advantage of a large population-based survey being undertaken as the COVID-19 vaccination campaign was being rolled out, we sought to understand COVID-19 vaccination coverage among youth to inform future vaccination strategies. Youth were randomly selected using geographic information system (GIS) methods. The survey was conducted in Harare (October– December 2021), Bulawayo (January– March 2022) and Mashonaland East (April– June 2022) aiming to recruit 16,800 18-24 year-olds (5,600 per province). COVID-19 vaccines became eligible for youth (aged 18 and above) in March/April 2021 and for 16-18 year olds in November 2021 [17]. Sociodemographic data, self-reported COVID-19 vaccination, and reasons for not being vaccinated were collected using interviewer-administered questionnaires. Participants could select multiple reasons for non-vaccination from a predefined list, or give additional reasons which were recorded as free text.

Data collection

Participants viewed an information video about the study on a tablet screen prior to providing electronic consent. Survey data were collected onto electronic tablets using SurveyCTO (Cambridge, USA). Data were stored on a server at the Biomedical Research and Training Institute (BRTI) in Zimbabwe and no identifying information was collected. This study was performed in accordance with the study protocol, the Declaration of Helsinki, as well as national and other regulatory guidelines.

Data analysis

Data analysis was conducted using Stata v16.1 (StataCorp, USA). A descriptive analysis was performed using proportions for categorical data, medians and means for continuous data,

followed by a univariable logistic regression analysis to investigate the association between sociodemographic characteristics (sex, age, educational attainment, socioeconomic status) and self-reported COVID-19 vaccine uptake. The outcome was self-report as having received at least one dose of a COVID-19 vaccine. Factors associated with the outcome in the univariable models were built into a multivariable logistic regression model, using only respondents with no missing data. To control for time, month of data collection was included in the model a priori. A Likelihood Ratio Test (LRT) was conducted for each variable to assess strength of evidence of association and reported as a *p*-value. The socioeconomic status variable was created using a principal component analysis of ownership of assets (refrigerator, bicycle, car, tv, radio, microwave, cell phone, and computer) and was then reported in quintiles. Analysis was performed for all provinces and then performed separately for each province as enrolment was conducted sequentially. Reasons for not taking up the vaccine were categorised according to the “5C’s” as described by Razai et al.: i) confidence (safety and efficacy of the vaccine), ii) complacency (perception of low risk and disease severity), iii) convenience (barriers and access), iv) communication (sources of information), and v) context (sociodemographic characteristics) [31].

Ethics

The study received ethical approval from the Medical Research Council of Zimbabwe (MRCZ) (MRCZ/A/2387), the BRTI Institutional Review Board (IRB) (AP149/2018) and the London School of Hygiene and Tropical Medicine (LSHTM) ethics committee (16124).

Patient and public involvement

The information video about the study was co-designed with and piloted among youth. The study questionnaire was also piloted with youth. The study had a Youth Advisory Board that provided guidance on study design and conduct. An extensive public engagement programme was undertaken alongside the study that included a national crowdsourcing competition to a) elicit young people's perceptions about health issues in their communities and b) train youth as researchers through a mentored programme termed Youth Researchers Academy [32].

Results

Of the 18,682 randomly sampled eligible youth in the study communities, 17,682 (94.6%) provided consent to participate. Of these, 5,849 (33.1%) were recruited in Harare, 5,969 (33.8%) in Bulawayo, and 5,864 (33.2%) in Mashonaland East. The median age of survey participants was 20 years (IQR: 19 – 22), and 10,742 (60.8%) were women. This higher proportion of women reflected the community composition.

Overall, 10,652 (60.3%) participants self-reported having received at least one dose of a COVID-19 vaccine, with 8,316 (78.1%) having received two doses (Figure 1). The proportion vaccinated was higher among males (n=4,779, 68.9%) compared to females (n=5,872, 54.7%) and increased with age (<20: 57.5%, 20-22: 61.5%, >22: 62.2%) (Table 1). Overall, Harare province, where data collection preceded the other provinces by 3 (Bulawayo) and 6 (Mashonaland East) months, had the lowest proportion vaccinated (42.6%) compared to Bulawayo (69.6%) and Mashonaland East (68.3%) (Figure 1). The proportion vaccinated and double-vaccinated increased with each subsequent month during the survey in Harare and Bulawayo, but not in Mashonaland East.

The major reasons given for not being vaccinated were lack of time, belief that the vaccine was unsafe, and anxiety about side effects (Figure 2). Men reported a lack of time as the main reason for not getting vaccinated. Women reported concerns related to side effects and safety as reasons for not getting vaccinated more frequently compared to men, especially infertility (10.7%). 755 (4.3%) women in the survey were pregnant at the time of recruitment. Of these women, 297 (39.3%) reported having received at least one dose of a COVID-19 vaccine. Of note, participants frequently reported that they were “afraid to die within 2 years of receiving the vaccine”, 17.7%, 15.6% and 8.2% in Harare, Mashonaland East and Bulawayo respectively. On the other hand, religious belief was a less commonly (< 5.0%) mentioned reason for getting vaccinated. While those living under more deprived socioeconomic circumstances had lower vaccine uptake, reasons for not getting vaccinated were similar across all socioeconomic quintiles (Supplemental Table 1).

Univariable analysis (Supplemental Table 2) showed an association between vaccine uptake age, sex, educational attainment and socioeconomic status. The association between vaccine uptake and predictors remained in the multivariable analysis (Table 2). Men (OR 1.69, 95%CI 1.59-1.80), older youth (20-22: OR 1.06, 95%CI 0.98-1.15, >22: OR 1.12, 95%CI 1.04-1.21), those with higher educational attainment (form 4: OR 1.79, 95%CI 1.39-2.30, form 6: OR 3.56, 95%CI 2.72-4.66, post-secondary: OR 4.34, 95%CI 3.27-5.76), and higher socioeconomic status (2nd quintile: OR 1.06, 95%CI 0.96-1.17, 3rd quintile: OR 1.12, 95%CI 1.01-1.23, 4th quintile: OR 1.17, 95%CI 1.06-1.29, least poor 20%: OR 1.32, 95%CI 1.20-1.47), were more likely to be vaccinated. Overall results and results stratified by province were comparable.

Discussion

This study found a COVID-19 vaccine coverage among young people aged 18-24 years of 69.6% in Bulawayo, 68.3% in Mashonaland East, and 42.6% in Harare. Despite not being a high-risk group, the national vaccination campaign reached them effectively. Vaccine uptake was however inequitable. Those who were male, older, more educated, and of higher socioeconomic status were more likely to report COVID-19 vaccination, which is in-line with studies conducted in HICs, though the specific reasons may be different. Education attainment and male sex were the strongest predictors.

The difference in COVID-19 vaccine coverage among youth across the provinces is partly explained by the staggered timing of the survey. The national vaccination campaign started in February 2021 prioritising front-line workers. Vaccination eligibility was extended to all adults in March/April 2021 and 16-18-year-olds became eligible in November 2021 (one month after study recruitment started in Harare). At the time the survey was completed in each province, the provincial vaccine coverage in Zimbabwe was 25.3% (Harare), 31.7% (Bulawayo), and 38.6% (Mashonaland East) [33]. Importantly, in the context of COVID-19 vaccination, coverage has been defined as the percentage of the total population that is vaccinated and includes children even though they may not be eligible [34]. At the end of the survey, vaccine coverage using the total eligible population (16+ years) as denominator was 73.8% nationally, and thus comparable to coverage among youth in Mashonaland East [2].

In the full adjusted model, men had 1.69 times the odds to have received the vaccine compared to women. Other studies have found higher proportions of men compared to

women reporting intention to getting the COVID-19 vaccine [35-37]. These studies have also highlighted that the difference in intention to get vaccinated against COVID-19 between men and women is less about increased health-seeking behaviour in men and more about decreased health-seeking behaviour among women, which is consistent with existing literature on general vaccine hesitancy [38,39]. This difference is likely due to specific gender differences in risk aversion and potential side effects. Reasons for not getting vaccinated in our study were mainly related to confidence, i.e. regarding safety and efficacy of the vaccine, especially among women. In this survey, 20.1% of women who were not vaccinated said they were afraid of side effects in general and 10.7% said they were afraid of infertility specifically. Furthermore, 66/458 (14.4%) of unvaccinated pregnant women reported fertility-related fears as a barrier, despite the WHO having recommended the use of SinoPharm in pregnant women [35-37,40]. This disparity may be due to confusing communication regarding pregnancy and breastfeeding at the start of the vaccination campaign, including information by official sources and on social media [41,42]. The infodemic about vaccines spread through social media has undoubtedly played an important role in Zimbabwe given the high proportion of individuals (both men and women) who reported being afraid of dying within 2 years of vaccination. This myth relates to a widely and globally circulated text message meme claiming that French virologist Luc Montagnier had said all vaccinated people will “die within two years” [42].

Fear of side effects was less frequent among unvaccinated men compared to women, and a higher proportion of men did not feel at risk (felt ‘young and healthy’) or said they were too busy for vaccination. This is despite a vaccination campaign that tried to bring vaccines to

the population by offering transport incentives such as transport money, decentralising vaccination to polyclinics, and providing vaccine outreach services [16].

Vaccination rates varied by socioeconomic status and particularly educational attainment, pointing towards health inequity. Other studies have reported that poorer and less educated people experience more barriers to vaccination [44,45]. Those with lower educational attainment are less likely to have access to accurate information and the vaccine information itself might be inaccessible in terms of language, content, and format, especially if this is provided in a written format [46]. Public health information may also include jargon which discriminates against those of lower education level. Those of lower socioeconomic status were also more likely to experience adverse events such as a cut in household income, inability to access food, higher disease risk, or loss of work than those in higher quintiles, which affects people's ability to receive a vaccine [47].

In this study only a small proportion of individuals said they were not vaccinated because of their religious beliefs. This is in contrast to the results from studies conducted in other countries in Africa where religious beliefs were among the most frequent reasons given for vaccine hesitancy [10-12]. In Zimbabwe the Apostolic church, a Pentecostal Christian denomination, has an estimated 3.5 million followers mainly among poorer and rural households. Children of members of the Apostolic church have been found to have low childhood vaccination coverage [48]. Religious affiliation was not collected in the survey, but membership of the Apostolic church is less common in urban and peri-urban settings, which may explain some of our findings. Also, the Ministry of Health and Child Care, supported by UNICEF, actively reached out to churches including the Apostolic Church to support vaccine

education and public engagement [49]. The Apostolic Women's Empowerment Trust started a COVID-19 awareness programmes in 2021 with the aim to address vaccine hesitancy among those of Apostolic faith [50].

Strengths of this study include a large representative sample of young people from three provinces at the time of the COVID-19 vaccine roll-out. Participation in the survey was high and results were comparable across all three provinces. The main limitation of this study is that COVID-19 vaccination status was self-reported and social desirability bias may have resulted in overestimation of vaccination coverage. However, the survey included a range of potentially sensitive questions, on topics including sexual and reproductive health, and substantial efforts were invested in training the survey team. The survey was conducted in urban and peri-urban settings only and vaccination coverage may be very different among young people living in rural areas. There were no questions specifying the type of vaccine received or the date of vaccination. Another limitation was that the timing of the survey relative to the COVID-19 vaccination campaign differed between the three provinces. Finally, while we asked young people to provide reasons for not being vaccinated, more detailed questions on beliefs, myths, and sources of information might have provided better understanding. Further qualitative research investigating the barriers towards vaccination, especially among women, would strongly contribute to the findings reported in this study.

Conclusion

This study showed inequitable COVID-19 vaccine coverage among young people aged 18-24 years in three provinces in Zimbabwe (Harare, Bulawayo, and Mashonaland East). Vaccination rates were lower among women, people with lower educational attainment and

among those living in more deprived socioeconomic circumstances. Fear around death and infertility fueled by myth and misinformation were among the main reasons for vaccine hesitancy. Therefore, we recommend that national vaccination campaigns should include a major focus on health education, especially for women and those with less education, and particularly around infertility and death. Information that is clear and consistent provided by trusted sources is crucial, and in this way, social media can be used in a positive manner to combat misinformation. Furthermore, campaigns should focus on community and religious leaders as they have a strong impact on their respective groups. No single programme is likely to address vaccine hesitancy in this population and thus strategies should address these inequities by understanding concerns and tailoring vaccine campaigns to specific groups instead of a one-size-fits-all approach.

References

1. World Health Organization (WHO) - WHO Coronavirus (COVID-19) Dashboard 2023. 17 February 2023. Accessed 17 February, 2023. Available from: <https://covid19.who.int/>
2. Statista. Zimbabwe: Age structure from 2011 to 2021. 20 January 2023. Accessed 17 February, 2023. Available from: <https://www.statista.com/statistics/455307/age-structure-in-zimbabwe/>
3. Ullrich F, Hanoun C, Turki AT, et al. Early report on the severity of COVID-19 in hematologic patients infected with the SARS-CoV2 omicron variant. *European Journal of Haematology*. 2022;109(4):364-372.
4. Rahmani K, Shavaleh R, Forouhi M, et al. The effectiveness of COVID-19 vaccines in reducing the incidence, hospitalization, and mortality from COVID-19: A systematic review and meta-analysis. *Frontiers in public health*. 2022;10:873596-873596.
5. World Health Organization (WHO). Strategy to achieve global COVID-19 vaccination by mid-2022, 2021. 2022.
6. CGTN Africa. Zimbabwean president, government officials receive COVID-19 booster shots. CGTN Africa. 5 January 2022. Accessed November 7, 2022. Available from: <https://africa.cgtn.com/2022/01/05/zimbabwean-president-government-officials-receive-covid-19-booster-shots/>
7. CDC A. Africa CDC COVID-19 Vaccine Dashboard 2023 [updated 11 January 2023. Available from: <https://africacdc.org/covid-19-vaccination/> accessed 11 January 2023.
8. Our world in data. Coronavirus (COVID-19) vaccinations. Our world in data. November 7, 2022. Accessed November 7, 2022. <https://ourworldindata.org/covid-vaccinations>
9. Njoga EO, Awoyomi OJ, Onwumere-Idolor OS, Awoyomi PO, Ugochukwu IC, Ozioko SN. Persisting Vaccine Hesitancy in Africa: The Whys, Global Public Health Consequences and Ways-Out—COVID-19 Vaccination Acceptance Rates as Case-in-Point. *Vaccines*. 2022;10(11):1934
10. Ackah BB, Woo M, Stallwood L, et al. COVID-19 vaccine hesitancy in Africa: a scoping review. *Global Health Research and Policy*. 2022;7(1):1-20
11. Kahn K, Pettifor A, Mataboge P, et al. COVID-19 vaccine hesitancy in rural South Africa: Deepening understanding to increase uptake and access. *Journal of Global Health*. 2022;12
12. Larson HJ, Gakidou E, Murray CJL. The Vaccine-Hesitant Moment. *New England Journal of Medicine*. 2022;387(1):58-65. doi:10.1056/NEJMra2106441
13. World Health Organization (WHO). 6 July 2022. Immunization Zimbabwe 2022 country profile. 6 July 2022. Accessed 3 February, 2023. Available from: <https://www.who.int/publications/m/item/immunization-zwe-2022-country-profile>

14. Chitungo I, Dzinamarira T, Tungwarara N, et al. COVID-19 Response in Zimbabwe: The Need for a Paradigm Shift? *COVID*. 2022;2(7):895-906
15. Sadeghalvad M, Mansourabadi AH, Noori M, et al. Recent developments in SARS-CoV-2 vaccines: A systematic review of the current studies. *Reviews in Medical Virology* 2023;33(1):e2359
16. Murewanhema G, Musuka G, Denhere K, et al. The Landscape of COVID-19 Vaccination in Zimbabwe: A Narrative Review and Analysis of the Strengths, Weaknesses, Opportunities and Threats of the Programme. *Vaccines* 2022;10(2):262.
17. Murewanhema G, Burukai TV, Chireka B, et al. Implementing national COVID-19 vaccination programmes in sub-Saharan Africa- early lessons from Zimbabwe: a descriptive cross-sectional study. *Pan Afr Med J* 2021;40:180. doi: 10.11604/pamj.2021.40.180.30824 [published Online First: 20211124]
18. UNICEF. Children's vaccination campaign gathers momentum in Zimbabwe. June 9, 2022. Accessed November 7, 2022. <https://www.unicef.org/zimbabwe/stories/childrens-vaccination-campaign-gathers-momentum-zimbabwe>
19. Xinhua. Zimbabwe receives latest vaccine donation from China amid 4th wave of COVID-19 pandemic. XINHUANET. December 21, 2021. Accessed November 7, 2022. http://www.news.cn/english/africa/2021-12/21/c_1310384569.htm
20. Zimbabwe COVID-19 SitRep 15/10/2022. Harare: Ministry of Health and ChildCare, 2022. <https://twitter.com/MoHCCZim/status/1581528130157543425> last accessed 15.10.2022
21. allAfrica. Zimbabwe: Experts Advocate for Covid Booster Jab. allAfrica. 6 January 2022. Accessed November 7, 2022. <https://allafrica.com/stories/202201060203.html>
22. Dzinamarira T, Nachipo B, Nyathi A, et al. The case to scale up edutainment as an effective public health communication intervention to combat the COVID-19 pandemic in Zimbabwe. *Health Promotion Perspectives*. 2022;12(1):34.
23. The Herald. Zimbabwe: Govt Launches Covid-19 Vaccination Blitz. allAfrica. March, 17 2022. Accessed November 7, 2022. <https://allafrica.com/stories/202203170495.html>
24. Sam-Agudu NA, Quakyi NK, Masekela R, Zumla A, Nachege JB. Children and adolescents in African countries should also be vaccinated for COVID-19. *BMJ Global Health*. 2022;7(2):e008315. doi:10.1136/bmjgh-2021-008315
25. UNICEF. Youth advocates make the difference in Zimbabwe's COVID hotspots. July 22, 2022. Accessed February 14, 2023. <https://www.unicef.org/zimbabwe/stories/youth-advocates-make-difference-zimbabwes-covid-hotspots>
26. UNICEF. Murals a game changer in Zimbabwe's battle against COVID-19. August 15, 2022. Accessed February 14, 2023. <https://www.unicef.org/zimbabwe/stories/murals-game-changer-zimbabwes-battle-against-covid-19>

27. UNICEF. UNICEF partners with Youth Advocates Zimbabwe in youth-led COVID-19 campaign. May 13, 2022. Accessed February 14, 2023. <https://www.unicef.org/zimbabwe/stories/unicef-partners-youth-advocates-zimbabwe-youth-led-covid-19-campaign>
28. PERC. Finding the Balance: Public health and social measures in Zimbabwe. October 28, 2021. Accessed January 18, 2023. https://preventepidemics.org/wp-content/uploads/2021/10/zimbabwe_en_20211028_1236.pdf
29. Kugarakuripi J, Ndoma S. Lack of trust in government, reliance on social media may drive vaccine hesitancy in Zimbabwe. January, 3 2022. Accessed January 18, 2023. <https://www.afrobarometer.org/publication/ad500-lack-trust-government-reliance-social-media-may-drive-vaccine-hesitancy-zimbabwe/>
30. Dziva Chikwari C, Dauya E, Bandason T, et al. The impact of community-based integrated HIV and sexual and reproductive health services for youth on population-level HIV viral load and sexually transmitted infections in Zimbabwe: protocol for the CHIEDZA cluster-randomised trial [version 1; peer review: awaiting peer review]. *Wellcome Open Research* 2022;7(54) doi: 10.12688/wellcomeopenres.17530.1
31. Razai MS, Oakeshott P, Esmail A, Wiysonge CS, Viswanath K, Mills MC. COVID-19 vaccine hesitancy: the five Cs to tackle behavioural and sociodemographic factors. *J R Soc Med*. Jun 2021;114(6):295-298. doi:10.1177/01410768211018951
32. Tembo M, Mackworth-Young C, Kranzer K, et al. Youth researchers academy: a report on an innovative research training programme for young people in Zimbabwe. *BMJ Innovations* 2022;8(3)
33. Our world in data. Coronavirus (COVID-19) vaccinations. Our world in data. November 7, 2022. Accessed November 7, 2022. <https://ourworldindata.org/covid-vaccinations>
34. Cutts FT, Claquin P, Danovaro-Holliday MC, Rhoda DA. Monitoring vaccination coverage: Defining the role of surveys. *Vaccine*. Jul 29 2016;34(35):4103-4109. doi:10.1016/j.vaccine.2016.06.053
35. Jensen A, Stromme M, Moyassari S, et al. COVID-19 vaccines: Considering sex differences in efficacy and safety. *Contemp Clin Trials*. Apr 2022;115:106700. doi:10.1016/j.cct.2022.106700
36. Zintel S, Flock C, Arbogast AL, Forster A, von Wagner C, Sieverding M. Gender differences in the intention to get vaccinated against COVID-19: a systematic review and meta-analysis. *Z Gesundh Wiss*. Jan 7 2022:1-25. doi:10.1007/s10389-021-01677-w
37. Afrifa-Anane GF, Larbi RT, Addo B, et al. Facilitators and barriers to COVID-19 vaccine uptake among women in two regions of Ghana: A qualitative study. *PLoS One*. 2022;17(8):e0272876. doi:10.1371/journal.pone.0272876

38. Borga LG, Clark AE, D'Ambrosio C, et al. Characteristics associated with COVID-19 vaccine hesitancy. *Scientific Reports* 2022;12(1):1-9.
39. Khan H, Dabla-Norris ME, Lima F, et al. Who doesn't want to be vaccinated? Determinants of vaccine hesitancy during COVID-19: International Monetary Fund 2021.
40. World Health Organization (WHO). The Sinopharm COVID-19 vaccine: What you need to know. 2021
41. Srivastava KC, Shrivastava D, Chhabra KG, et al. Facade of media and social media during COVID-19: A review. *Int J Res Pharm Sci* 2020:142-49
42. Rocha YM, de Moura GA, Desidério GA, et al. The impact of fake news on social media and its influence on health during the COVID-19 pandemic: A systematic review. *Journal of Public Health* 2021:1-10
43. CRUX; 2021. Did French Virologist Luc Montagnier Actually Say Vaccination Will Kill Us & Create More Variants? <https://www.youtube.com/watch?v=wkZKRND05Ks>. Accessed December 22, 2022.
44. Humer E, Jesser A, Plener PL, Probst T, Pieh C. Education level and COVID-19 vaccination willingness in adolescents. *European Child & Adolescent Psychiatry*. 2021/09/22 2021;doi:10.1007/s00787-021-01878-4
45. Shifa M, Gordon D, Leibbrandt M, Zhang M. Socioeconomic-Related Inequalities in COVID-19 Vulnerability in South Africa. *International Journal of Environmental Research and Public Health*. 2022;19(17):10480.
46. Razai MS, Chaudhry UA, Doerholt K, et al. Covid-19 vaccination hesitancy. *Bmj* 2021;373
47. Wright L, Steptoe A, Fancourt D. Are we all in this together? Longitudinal assessment of cumulative adversities by socioeconomic position in the first 3 weeks of lockdown in the UK. *J Epidemiol Community Health* 2020;74(9):683-88
48. Machekanyanga Z, Ndiaye S, Gerebe R, et al. Qualitative Assessment of Vaccination Hesitancy Among Members of the Apostolic Church of Zimbabwe: A Case Study. *J Relig Health*. Oct 2017;56(5):1683-1691. doi:10.1007/s10943-017-0428-7
49. Kholwani Nyathi (UNICEF). Zimbabwe: Experts Advocate for Covid Booster Jab. UNICEF. 10 November 2021. Accessed December 21, 2022. <https://www.unicef.org/zimbabwe/stories/religious-groups-warm-covid-19-vaccines-zimbabwe>
50. Maiden James (UNICEF). Zimbabwe's religious leaders increase efforts to tackle COVID-19 and support vaccines. UNICEF. 05 May 2021. Accessed December 21, 2022. <https://www.unicef.org/zimbabwe/press-releases/zimbabwes-religious-leaders-increase-efforts-tackle-covid-19-and-support-vaccines>

Tables and Figures

Table 1: Sociodemographic characteristics and reported COVID-19 vaccination

	n	Vaccinated
Sex		
Male	6,940	4,779 (68.9%)
Female	10,742	5,872 (54.7%)
Age		
< 20	6,809	3,916 (57.5%)
20 – 22	4,702	2,894 (61.5%)
> 22	6,171	3,841 (62.2%)
Province		
Harare	5,849	2,492 (42.6%)
Bulawayo	5,969	4,155 (69.6%)
Mashonaland East	5,864	4,004 (68.3%)
Education		
Primary/None	3,254	1,452 (44.6%)
Form 4	10,784	6,352 (58.9%)
Form 6	2,233	1,702 (76.2%)
Post-secondary	1,411	1,145 (81.2%)
Employment		
Student	4,963	3,479 (70.1%)
Employed (formal)	835	585 (70.1%)
Employed (informal)	3,158	1,821 (57.7%)
None	8,726	4,766 (54.6%)
Marital status		
Married	3,559	1,511 (42.5%)
Single	13,324	8,732 (65.5%)
Divorced or widowed	799	408 (51.1%)

Table 2: Multivariable logistic regression analysis of the association between COVID-19 vaccination and sociodemographic variables

	Overall OR (95% CI)	Harare OR (95% CI)	Bulawayo OR (95% CI)	Mashonaland East OR (95% CI)
Sex				
Female	1 (p < 0.0001)	1 (p < 0.0001)	1 (p < 0.0001)	1 (p < 0.0001)
Male	1.69 (1.59 – 1.80)	1.45 (1.29 – 1.64)	1.43 (1.27 – 1.61)	1.84 (1.63 – 2.08)
Age				
< 20	1 (p = 0.0123)	1 (p < 0.0001)	1 (p = 0.0002)	1 (p = 0.6537)
20-22	1.06 (0.98 – 1.15)	1.23 (1.06 – 1.42)	1.24 (1.08 – 1.43)	0.94 (0.81 – 1.08)
> 22	1.12 (1.04 – 1.21)	1.43 (1.25 – 1.63)	1.31 (1.14 – 1.50)	0.99 (0.86 – 1.14)
Education				
Primary	1 (p < 0.0001)	1 (p < 0.0001)	1 (p < 0.0001)	1 (p < 0.0001)
Form 4	1.79 (1.39 – 2.30)	1.81 (1.15 – 2.86)	1.83 (1.15 – 2.93)	1.39 (0.88 – 2.20)
Form 6	3.56 (2.72 – 4.66)	4.09 (2.55 – 6.57)	3.50 (2.13 – 5.75)	2.84 (1.73 – 4.66)
Secondary and above	4.34 (3.27 – 5.76)	5.20 (3.19 – 8.47)	4.44 (2.60 – 7.58)	3.51 (2.08 – 5.92)
Socioeconomic status				
Poorest	1 (p < 0.0001)	1 (p < 0.0001)	1 (p = 0.3776)	1 (p = 0.0010)
2	1.06 (0.96 – 1.17)	1.26 (1.05 – 1.50)	1.04 (0.89 – 1.22)	0.95 (0.81 – 1.13)
3	1.12 (1.01 – 1.23)	1.38 (1.16 – 1.64)	0.99 (0.78 – 1.25)	0.99 (0.83 – 1.18)
4	1.17 (1.06 – 1.29)	1.70 (1.43 – 2.03)	1.10 (0.92 – 1.32)	1.01 (0.83 – 1.23)

Least poor	1.32 (1.20 – 1.47)	1.93 (1.61 – 2.31)	1.09 (0.90 – 1.31)	1.32 (1.09 – 1.61)
------------	--------------------	--------------------	--------------------	--------------------

Analysis was adjusted for month of recruitment, OR=odds ratio, 95%CI=95% confidence interval.




Figure 1: Proportion of young people aged 18-24 who reported being vaccinated by month of the CHIEDZA prevalence survey stratified by province.




Figure 2: Reasons for not receiving the COVID-19 vaccine stratified by sex and grouped based on the 5C's of vaccine hesitancy delineated by Razai et al. shown as proportions (%) [31].

Participants could provide multiple reasons and could only answer the question if they were not vaccinated. These reasons were read out or shown to the participant for them to choose from