

**TITLE:** How long is the long COVID? a retrospective analysis of football players in two major European Championships.

Sandra Miccinilli<sup>1\*</sup>, Marco Bravi<sup>1\*</sup>, Giorgio Conti<sup>1</sup>, Federica Bressi<sup>1</sup>, Silvia Sterzi<sup>1</sup>, Fabio Santacaterina<sup>1\*</sup>, and Massimo Ciccozzi<sup>2</sup>

<sup>1</sup> Department of Physical and Rehabilitation Medicine, Università Campus Bio-Medico di Roma, Via Alvaro del Portillo, 200, 00128 Rome, Italy

<sup>2</sup> Unit of Medical Statistics and Molecular Epidemiology, Università Campus Bio-Medico di Roma, Via Alvaro del Portillo, 200, 00128 Rome, Italy

\*: Corresponding Author - [f.santacaterina@policlinicocampus.it](mailto:f.santacaterina@policlinicocampus.it)

†: the authors Marco Bravi and Sandra Miccinilli contributed equally to this paper.

**Keywords:** SARS-CoV-2; muscle injury; soccer; muscle strain

## ABSTRACT

**Objectives,** the goal of this study was to investigate the correlation between SARS-CoV-2 infection and muscle injuries among a large sample of professional soccer players.

**Methods,** A retrospective cohort study was conducted on professional soccer players from the Serie A and LaLiga leagues during the 2019-2020 and 2020-2021 football seasons. The players were divided into two groups based on whether they contracted the Sars-CoV-2 infection (C+) or not (C-) during the 2020/2021 season. Data collection was conducted using the Transfermarkt24 site.

**Results,** In the 2019-2020 both championships showed non-significant differences in the average number of muscular injuries between the C+ group and the C- group (Serie A:  $p=0.194$ ; 95%CI: -0.044 to 0.215, LaLiga  $p=0.915$ ; 95%CI: -0.123 to 0.137). In the 2020-2021 the C+ group had a significantly higher number of muscular injuries compared to the C- group in both championships (Serie A:  $p<0.001$ ; 95%CI 0.731 to 1.038; LaLiga:  $p<0.001$ ; 95%CI: 0.773 to 1.054). Multiple linear regression analysis confirmed that belonging to C+ in the season 2020/2021 was the variable that most strongly influenced the probability of having a muscle injury in both championships. Survival analysis revealed a hazard ratio of 3.73 (95%CI 3.018 to 4.628) and of 5.14 (95% CI 3.200 to 8.254) for Serie A and LaLiga respectively.

## Conclusions

This retrospective cohort study revealed a significant association between SARS-CoV-2 infection and increased risk of muscle injury, emphasizing the importance of carefully considering the infection in the decision-making process for determining athletes' readiness to return to sport.

## Introduction

In late December 2019, The World Health Organization (WHO) office in China was informed about novel cases of pneumonia of unknown aetiology detected in the city of Wuhan, Hubei province [1]. Afterwards, a new type of coronavirus, named SARS-CoV-2, was isolated and identified by the Chinese authorities. The coronavirus disease 2019 (COVID-19) caused by SARS-CoV-2 was classified as a pandemic on 11 of March 2020 [2] by the World Health Organization (WHO). All countries were involved in surveillance of the disease and all sports that included close contact were temporarily suspended.

Football (soccer) is a physically demanding sport that requires a high level of neuromuscular readiness, including strength, reactivity, and muscular power [3,4]. The characteristics of this sport could therefore justify the increases in muscle injury rates associated with the increase in the number of official seasonal competitions and weekly training sessions [5,6]. However, the outbreak of the SARS-CoV-2 virus has had a significant impact on professional athletes, even in mild cases, leading to muscle weakness, poor tolerance to physical exercise, period of inactivity and absence from

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

sporting practice, with a considerable impact also on bio-psychosocial context [7–9]. COVID-19 is known to cause severe inflammatory responses, respiratory failure, acute respiratory distress syndrome (ARDS), and bilateral pneumonia [10]; in some cases these symptoms can continue causing what is called long COVID. It can also affect the musculoskeletal system, causing myalgia and sarcopenia in COVID-19 positive patients [11]. Therefore, the question arises whether COVID-19 may have influenced the number of muscle injuries in elite athletes. Previous studies in the literature have observed that the approximately three months of sports activity suspension due to the lockdown caused an increase in the incidence of injuries in the main European football leagues, including the Bundesliga (GER) [12], LaLiga (SPA) [13] and Premier League (ENG) [14]. The only exception is represented by Italian first division (Serie A) [15], where a non-statistically significant difference was observed between pre- and post-lockdown injuries. However, these studies did not show whether contracting SARS-CoV-2 infection can be considered an additional risk factor for injury occurrence. Only the recent prospective study by Wezenbeek et al. [16] involving three Belgian professional male football teams during the first half of the 2020-2021 season, reported a five-fold higher risk of developing a muscle strain after a SARS-CoV-2 infection. The goal of this study is to further investigate the findings reported by Wezenbeek et al. [16] and to retrospectively examine, on a larger sample size, the correlation between SARS-CoV-2 infection and muscle injuries among professional football players from two different championships, Serie A and LaLiga. These two championships were chosen because Serie A was an exception in terms of post-lockdown injury increase, while LaLiga had the most active collaboration in identifying and communicating positive cases and its results were more discordant between pre- and post-lockdown.

However, these studies did not show whether having contracted SARS-CoV-2 infection can be considered an additional risk factor for occurrence of injuries. Recently only the prospective study by Wezenbeek et al., involving three Belgian professional male football teams, during the first half of the 2020–2021 season reported a five times higher risk of developing a muscle strain after a SARS-CoV-2 infection.

The goal of this study is to deepen the results reported by Wezenbeek et al. and to retrospectively verify, on a larger sample, the correlation between SARS-CoV-2 infection and muscle injuries among professional footballers of two different championship Serie A and LaLiga. The choice of these two different championships was because Serie A represented the exception of post-lockdown injuries increase and both Serie A and LaLiga had the most active collaboration in identifying and communicating positive cases [13]. Our hypothesis is that there is a relationship between muscle injury and previous SARS-CoV-2 infection.

## **Materials and methods**

### **Study design**

A retrospective cohort study was conducted according to the STROBE guidelines [17]. The aim was to identify a potential correlation between SARS-CoV-2 infection and muscle injuries in professional soccer players from the Serie A and LaLiga leagues during the 2019-2020 and 2020-2021 football seasons. The study included all players from Serie A and LaLiga during the two seasons, as well as those who changed league levels during this time period (e.g. from a minor league such as Serie B in Italy to the top national league, Serie A) and those who moved to a foreign team. Athletes were divided into two groups: COVID-positive (C+) and COVID-negative (C-).

The grouping was performed as follows: 1) all football players infected during the 2020-2021 season were classified as “C+”, while all players who did not contract the infection during this time period were classified as “C-”; 2) players who contracted the infection during the 2019-2020 season or during off-season periods (e.g. season summer suspension) were excluded from the study (Serie A  $n = 29$ ; LaLiga  $n = 38$ ) and their data were removed from both seasons. This created a database with pre-COVID (2019-2020 season) and post-COVID (2020-2021 season) data.

## Patient and public involvement

This study investigating the association between SARS-CoV-2 infection and the occurrence of muscle injuries during the sports season is the result of the involvement of medical staff and young athletes from some sports clubs who were the first to notice this possible correlation. These have also been involved in the discussion of the results of the study.

## Data collection

Data collection was conducted using the Transfermarkt24 site (<https://www.transfermarkt.co.uk/>) a website founded in Germany in May 2000 which contains a wealth of football information from leagues around the world, including data on the type and severity of injuries. This methodology is consistent with previous studies [18–21]. The data collected are specified in Table 1.

**Table 1: data collected and relative definitions used in the study**

Birth date	
Age at the end of the 2020-2021 season	
Team	
Muscle injuries season 2019-2020	Total number of muscular injury sustained by a player that resulted from a in-season football match or football training
Muscle injuries season 2020-2021	
Matches played in the 2019-2020	Total number of matches played during the season
Matches played in the 2020-2021	
Minutes played season 2019-2020	Total number of minutes played during the season
Minutes played season 2020-2021	
Date of SARS-CoV-2 infection	Date of in-season SARS-CoV-2 infection
Date of muscle injuries season 2019-2020	Date of each muscle injury occurred during in-season period
Date of muscle injuries season 2020-2021	
Season 2019-2020 start and end date	First and last match date of the season
Season 2020-2021 start and end date	

## Statistical analysis

Statistical analysis was performed according to the CHAMP statement [22] using MedCalc software (Version 20, MedCalc Software Ltd). The normal distribution of the data was verified using the Kolmogorov-Smirnov test. The comparison of mean injuries between the two groups in the 2019-2020 and 2020-2021 seasons was performed using a parametric t-Student test. A multiple linear regression analysis was carried out in both seasons to analyse the relationship between the dependent variable, muscle injuries, and the independent variables of age, SARS-CoV-2 infection, minutes played, and matches played. A multiple linear regression analysis was carried out in both seasons to analyse the relationship between the dependent variable, muscle injuries, and the independent variables of age, SARS-CoV-2 infection, minutes played, and matches played. Additionally, a Kaplan-Meier survival analysis was performed, using the elapsed time (in days) from the start of each championship to the day on which the injury occurred during the in-season period as the "survival time." If no injuries occurred, the total duration of the season in days was used as the survival time. The Kaplan-Meier curves were analysed using the Logrank Test in both seasons.

## Results

The study included 634 players from Serie A and 649 players from LaLiga, for a total of 1283 elite football players. Of the Italian championship players, 171 (27.3%) were infected with SARS-CoV-2, with 29 excluded due to being infected during the 2019-2020 season. In LaLiga, a total of 165

(25.4%) players were infected with SARS-CoV-2, with 38 excluded as they were infected during the 2019-2020 season (Figure 1).

In the 2019-2020 Serie A championship, non-significant differences ( $p=0.194$ ; 95%CI:  $-0.044$  to  $0.215$ ) were found in the average number of muscular injuries between the C+ group ( $0.52\pm0.75$ ) and the C- group ( $0.43\pm0.68$ ). In the 2020-2021 season, the C+ group had a significantly higher number of muscular injuries ( $1.44\pm0.96$ ) compared to the C- group ( $0.56\pm0.78$ ) ( $p<0.001$ ; 95%CI  $0.731$  to  $1.038$ ) (figure 2). The multiple linear regression analysis revealed that, in the 2019-2020 season, muscular injuries were mainly related to age, minutes played, and matches played by the athletes (table 2). In the 2020-2021 season, in addition to age, belonging to the C+ group was found to be the variable that most strongly influenced the probability of having a muscle injury, unlike the previous season (table 2).

Table 2: Multiple linear regression analysis of Serie A football players. The factors described, relating to the 2019-2020 and 2020/2021 season, represent the dependent variables capable of influencing or not the number of muscle injuries within the same season.

Independent variables	Coefficient	Std. Error	t	P
<b>2019/2020 season</b>				
(Constant)	-0,123			
Age	0,023	0,006	3,900	<b>&lt;0,001</b>
Minutes played	-0,000	0,000	-3,679	<b>&lt;0,001</b>
Matches played	0,025	0,007	3,613	<b>&lt;0,001</b>
Group (C+/C-)	-0,108	0,065	-1,658	0,098
<b>2020/2021 season</b>				
(Constant)	1,770			
Age	0,020	0,007	2,814	<b>0,005</b>
Minutes played	-0,000	0,000	-1,269	0,205
Matches played	0,006	0,007	0,910	0,363
Group (C+/C-)	-0,889	0,078	-11,371	<b>&lt;0,001</b>

The survival analysis based on the Kaplan-Meier curves (figure 3) revealed that in both groups, the injury rates during the 2019-2020 season did not differ significantly ( $p=0.401$ ). However, during the 2020-2021 season, the difference in injury rates between the two groups became statistically significant ( $p<0.001$ ), with a hazard ratio of 3.73 (95%CI 3.018 to 4.628).

In LaLiga, the results indicated that during the 2019-2020 season, there was no significant difference in the average number of muscular injuries between the C+ and C- groups, with  $0.40\pm0.67$  and  $0.39\pm0.65$  injuries, respectively ( $p=0.915$ ; 95%CI:  $-0.123$  to  $0.137$ ). However, during the 2020-2021 season, the average number of injuries was significantly higher ( $p<0.001$ ; 95%CI:  $0.773$  to  $1.054$ ) in the C+ group ( $1.23\pm1.04$ ) compared to the C- group ( $0.32\pm0.59$ ) (figure 2). The multiple linear regression analysis indicated that in the 2019-2020 season, age was the main factor related to muscle injuries ( $p=0.044$ ), while games played and minutes played by athletes did not significantly influence the occurrence of injuries. However, in the 2020-2021 season, the group variable had a significant influence on the muscle injury rate (table 3).

Table 3: Multiple linear regression analysis of LaLiga football players. The factors described, relating to the 2019-2020 and 2020/2021 season, represent the dependent variables capable of influencing or not the number of muscle injuries within the same season.

Independent variables	Coefficient	Std. Error	t	P
<b>2019/2020 season</b>				
(Constant)	0,110			
Age	0,0129	0,006411	2,022	<b>0,044</b>

Minutes played	-0,000	0,00006924	-0,426	0,670
Matches played	0,002	0,006129	0,337	0,736
Group (C+/C-)	-0,042	0,06376	-0,663	0,508
<b>2020/2021 season</b>				
(Constant)	2,141			
Age	0,000	0,007	0,087	0,931
Minutes played	0,000	0,000	0,697	0,486
Matches played	-0,004	0,006	-0,749	0,454
Group (C+/C-)	-0,902	0,069	-12,916	<b>&lt;0,001</b>

Also in the Spanish championship, the survival analysis (figure 2) showed how the injuries rates in the 2019-2020 season did not differ significantly ( $p=0.246$ ), while in the season 2020-2021 the difference becomes statistically significant ( $p< 0.001$ ), with a hazard ratio of 5.14 (95% CI 3.200 to 8.254).

## Discussion

The main aim of this study was to determine whether professional soccer players who contracted SARS-CoV-2 had an increased risk of muscular injuries. The findings of this study reported a three to five times increased risk of muscle injury after SARS-CoV-2 infection.

Over the years, UEFA has conducted numerous studies on the match/minute ratio played by the elite athletes of the teams participating in its international tournaments to define a "normal" value for the incidence of injuries in football [23–25]. In particular, Hägglund et al.'s 11-year retrospective study [23] observed the incidence of injuries in more than 1000 hours of play and found an incidence of 2 injuries per year per individual footballer or an average of 50 injuries if we take into consideration the team group (list of 25 athletes). If we consider only muscular injuries, the incidence drops to 15 injuries out of a shortlist of 25 footballers, with an average per athlete of 0.6 injuries [25]. This data is essential for commenting on the results of our study. Our results showed that in both leagues, players who did not contract COVID-19 had injury rates lower than the average of 0.6 reported by Bengtsson et al. [25] in both seasons. Instead, regarding players infected with SARS-CoV-2, they showed in the pre-covid season an injury rate in line with the data of Bengtsson et al. [25], while in the 2020/2021 season, after the infection, it increased significantly in both leagues (1.44 for Serie A and 1.23 for LaLiga).

Our results indicate that SARS-CoV-2 infection could be a risk factor for muscular injury, in line with the recent study by Wezenbeek et al. [16]. Multiple linear regression analysis showed that in the 2019-2020 season, the risk of injury was significantly influenced by age, according to a previous study by Green et al. [26]. However, in the 2020-2021 season, SARS-CoV-2 infection was the variable that highly influenced the risk of muscle injury. Therefore, belonging to the C+ group in the 2020-2021 season completely distorts the correlation between the various variables and the muscle injury rates, becoming the main factor capable of influencing the latter. In other words, while older players with a high number of games and playing time were at greater risk of incurring muscle injuries in the 2019-2020 season, having contracted COVID-19 was the primary and most important risk factor in the following season. Similarly, in LaLiga, age was the main risk factor for muscle injury in the 2019-2020 season, while in the 2020-2021 season, the only variable that significantly influenced muscle injuries was the SARS-CoV-2 infection. The survival analysis showed that during the 2019-

2020 season, the two groups had a similar risk of muscular injury. In the 2020/2021 season, the hazard risk ratio was 3.7 to 5.1 times greater among those athletes infected by SARS-CoV-2, in line with the results of Wezenbeek et al. [16], which reported a five-time higher hazard rate to develop a muscle strain after SARS-CoV-2 infection.

As discussed by Wezenbeek et al. [16], there are two possible explanations for the correlation between SARS-CoV-2 infection and an increased risk of muscular injury. The first hypothesis is that strict quarantine rules implemented during the 2020/2021 season led to prolonged periods of abstention from training and sports participation, resulting in muscular detraining [27,28] and subsequent loss of muscle strength [29]. Muscle strength has been shown to be a protective factor for muscular injury [30], and the loss of muscle strength is more significant in highly trained athletes with greater initial muscle mass [31]. The second explanation is related to the direct biological effects of the virus. SARS-CoV-2 infection causes hyperinflammation, an increase in inflammatory markers, an increase in the neutrophil/lymphocyte ratio, and possible depletion of circulating T cells [32]. Furthermore, the virus may cause direct damage to muscle tissue by targeting the ACE 2 receptor [18], which is widely present on muscle tissue [33]. In addition, the virus can cause disturbances in blood flow and oxygen transport, leading to reduced muscle oxygenation during exercise. Lower muscle oxygen saturation [34] and  $\text{VO}_2$  max [35] are possible risk factors for muscle injury.

### Clinical Implications

These results emphasize the importance of carefully considering the infection in the decision-making process for determining athletes' readiness to return to sport (RTS). In fact, as already described by Elliot et al. [36] in addition to the known complications, it is necessary to consider that COVID-19 can be responsible for musculoskeletal complications. Therefore, our opinion is that the decision regarding the RTS should not only take into account the remission of COVID-19 related symptoms, rest and the cessation of post-COVID-19 drug therapy, but should include specific assessment and training programs, generally used after a musculoskeletal injury [37] and therefore consider COVID-19 not only a risk factor but a real injury.

### Limitations

One possible limitation of this study is that the muscle injury events were collected from a single database, which may have resulted in some injuries being overlooked, omitted, or interpreted differently, leading to over- or underestimation. Another limitation of this study is that it was not possible to control for some factors that may have influenced the increased risk of injury, such as previous muscle injuries in unexamined seasons and severity of COVID-19-related symptoms. Nevertheless, the study's strength lies in its design, with a large homogeneous population group of male professional soccer players followed prospectively with a standardized methodology.

### Conclusion

This retrospective cohort study, conducted on a large sample of professional male football players from two main European leagues, has revealed a significant association between SARS-CoV-2 infection and increased risk of muscular injury. The results of this study show that the risk of muscular injury is 3.7 to 5.1 times higher in football players who have been infected with SARS-CoV-2. Overall, our study highlights the need for further investigation into the effects of SARS-CoV-2 on athletes and the development of tailored rehabilitation protocols to ensure safe return to play.

### Competing Interests

The authors declare that they have no conflict of interest.

**Patient and public involvement:** patients and/or the public were involved in this research. Refer to the Methods section for further details.

## Funding

This research received no external funding.

## Contributorship

Contributors MB, GC, and MC contributed to the conception of the study. SM, MB, GC, FB, SS, FS and MC contributed to the design of the study. GC, MB and FS completed the acquisition of the data. MB GC and MC performed the data analysis. All authors assisted with the interpretation. MB and SM were the principal authors of the manuscript. All authors contributed to the drafting and revision of the final article. All authors approved the final submitted version of the manuscript.

## Ethical approval information

Not applicable

## Acknowledgements

Not applicable

## Data availability

Data are available on reasonable request to the corresponding author.

## References

- 1 Huang C, Wang Y, Li X, *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;**395**:497–506. doi:10.1016/S0140-6736(20)30183-5
- 2 Cucinotta D, Vanelli M. WHO Declares COVID-19 a Pandemic. *Acta Bio Medica : Atenei Parmensis* 2020;**91**:157. doi:10.23750/ABM.V91I1.9397
- 3 Manzi V, Annino G, Savoia C, *et al.* Relationship between aerobic fitness and metabolic power metrics in elite male soccer players. *Biol Sport* 2021;**39**:599–606. doi:10.5114/BIOLSPORT.2022.106389
- 4 Krstrup P, Mohr M, Steensberg A, *et al.* Muscle and blood metabolites during a soccer game: Implications for sprint performance. *Med Sci Sports Exerc* 2006;**38**:1165–74. doi:10.1249/01.MSS.0000222845.89262.CD
- 5 Ekstrand J, Waldén M, Hägglund M. Hamstring injuries have increased by 4% annually in men's professional football, since 2001: a 13-year longitudinal analysis of the UEFA Elite Club injury study. *Br J Sports Med* 2016;**50**:731–7. doi:10.1136/BJSPORTS-2015-095359
- 6 Ekstrand J, Bengtsson H, Waldén M, *et al.* Hamstring injury rates have increased during recent seasons and now constitute 24% of all injuries in men's professional football: the UEFA Elite Club Injury Study from 2001/02 to 2021/22. *Br J Sports Med* 2023;**57**:292–8. doi:10.1136/BJSPORTS-2021-105407
- 7 Serviente C, Decker ST, Layec G. Long-Term Recovery from SARS-CoV-2 (COVID-19): From heart to muscle: pathophysiological mechanisms underlying long-term physical sequelae from SARS-CoV-2 infection. *J Appl Physiol* 2022;**132**:581. doi:10.1152/JAPPLPHYSIOL.00734.2021
- 8 Jia L, Carter M v., Cusano A, *et al.* The Effect of the COVID-19 Pandemic on the Mental and Emotional Health of Athletes: A Systematic Review. *Am J Sports Med* Published Online First: 2022. doi:10.1177/03635465221087473

- 9 Haan R, Ali Alblooshi ME, Syed DH, *et al.* Health and Well-Being of Athletes During the Coronavirus Pandemic: A Scoping Review. *Front Public Health* 2021;**9**:255. doi:10.3389/FPUBH.2021.641392/BIBTEX
- 10 Huang C, Wang Y, Li X, *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;**395**:497. doi:10.1016/S0140-6736(20)30183-5
- 11 Montes-Ibarra M, Oliveira CLP, Orsso CE, *et al.* The Impact of Long COVID-19 on Muscle Health. *Clin Geriatr Med* 2022;**38**:545. doi:10.1016/J.CGER.2022.03.004
- 12 Seshadri DR, Thom ML, Harlow ER, *et al.* Case Report: Return to Sport Following the COVID-19 Lockdown and Its Impact on Injury Rates in the German Soccer League. *Front Sports Act Living* 2021;**3**:604226. doi:10.3389/FSPOR.2021.604226/FULL
- 13 García-Calvo T, Fernandez-Navarro J, Díaz-García J, *et al.* The impact of COVID-19 lockdown on soccer positional and physical demands in the Spanish La Liga. <https://doi.org/101080/2473393820222055784> Published Online First: 2022. doi:10.1080/24733938.2022.2055784
- 14 Mannino BJ, Yedikian T, Mojica ES, *et al.* The COVID lockdown and its effects on soft tissue injuries in Premier League Athletes. <https://doi.org/101080/0091384720211980746> Published Online First: 2021. doi:10.1080/00913847.2021.1980746
- 15 Marotta N, de Sire A, Gimigliano A, *et al.* Impact of COVID-19 lockdown on the epidemiology of soccer muscle injuries in Italian Serie A professional football players. *Journal of Sports Medicine and Physical Fitness* 2022;**62**:356–60. doi:10.23736/S0022-4707.21.12903-2
- 16 Wezenbeek E, Denolf S, Willems TM, *et al.* Association between SARS-COV-2 infection and muscle strain injury occurrence in elite male football players: a prospective study of 29 weeks including three teams from the Belgian professional football league. *Br J Sports Med* 2022;**56**:818–23. doi:10.1136/BJSPORTS-2021-104595
- 17 Vandembroucke JP, von Elm E, Altman DG, *et al.* Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *PLoS Med* 2007;**4**:1628–54. doi:10.1371/JOURNAL.PMED.0040297
- 18 Annino G, Manzi V, Alashram AR, *et al.* COVID-19 as a Potential Cause of Muscle Injuries in Professional Italian Serie A Soccer Players: A Retrospective Observational Study. *International Journal of Environmental Research and Public Health* 2022, Vol 19, Page 11117 2022;**19**:11117. doi:10.3390/IJERPH191711117
- 19 Wand T. Analysis of the Football Transfer Market Network. *J Stat Phys* 2022;**187**:1–12. doi:10.1007/S10955-022-02919-1/FIGURES/4
- 20 Grassi A, Macchiarola L, Filippini M, *et al.* Epidemiology of Anterior Cruciate Ligament Injury in Italian First Division Soccer Players. *Sports Health* 2020;**12**:279. doi:10.1177/1941738119885642
- 21 Schiffner E, Latz D, Grassmann JP, *et al.* Fractures in German elite male soccer players. *Journal of Sports Medicine and Physical Fitness* 2019;**59**:110–5. doi:10.23736/S0022-4707.17.07901-4
- 22 Mansournia MA, Collins GS, Nielsen RO, *et al.* A Checklist for statistical Assessment of Medical Papers (the CHAMP statement): explanation and elaboration. *Br J Sports Med* 2021;**55**:1009–17. doi:10.1136/bjsports-2020-103652



- 23 Hägglund M, Waldén M, Magnusson H, *et al.* Injuries affect team performance negatively in professional football: an 11-year follow-up of the UEFA Champions League injury study. *Br J Sports Med* 2013;**47**:738–42. doi:10.1136/BJSPORTS-2013-092215
- 24 Waldén M, Hägglund M, Ekstrand J. UEFA Champions League study: a prospective study of injuries in professional football during the 2001–2002 season. *Br J Sports Med* 2005;**39**:542–6. doi:10.1136/BJSM.2004.014571
- 25 Bengtsson H, Ekstrand J, Hägglund M. Muscle injury rates in professional football increase with fixture congestion: an 11-year follow-up of the UEFA Champions League injury study. *Br J Sports Med* 2013;**47**:743–7. doi:10.1136/BJSPORTS-2013-092383
- 26 Green B, Bourne MN, van Dyk N, *et al.* Recalibrating the risk of hamstring strain injury (HSI): A 2020 systematic review and meta-analysis of risk factors for index and recurrent hamstring strain injury in sport. *Br J Sports Med* 2020;**54**:1081–8. doi:10.1136/BJSPORTS-2019-100983
- 27 Sarto F, Impellizzeri FM, Spörri J, *et al.* Impact of Potential Physiological Changes due to COVID-19 Home Confinement on Athlete Health Protection in Elite Sports: a Call for Awareness in Sports Programming. *Sports Medicine* 2020;**50**:1417–9. doi:10.1007/S40279-020-01297-6/METRICS
- 28 Heo SJ, Park SK, Jee YS. Detraining Effects of COVID-19 Pandemic on Physical Fitness, Cytokines, C-Reactive Protein and Immunocytes in Men of Various Age Groups. *Int J Environ Res Public Health* 2022;**19**:1845. doi:10.3390/IJERPH19031845
- 29 de Boer MD, Maganaris CN, Seynnes OR, *et al.* Time course of muscular, neural and tendinous adaptations to 23 day unilateral lower-limb suspension in young men. *J Physiol* 2007;**583**:1079–91. doi:10.1113/JPHYSIOL.2007.135392
- 30 Lauersen JB, Andersen TE, Andersen LB. Strength training as superior, dose-dependent and safe prevention of acute and overuse sports injuries: a systematic review, qualitative analysis and meta-analysis. *Br J Sports Med* 2018;**52**:1557–63. doi:10.1136/BJSPORTS-2018-099078
- 31 Miles MP, Heil DP, Larson KR, *et al.* Prior resistance training and sex influence muscle responses to arm suspension. *Med Sci Sports Exerc* 2005;**37**:1983–9. doi:10.1249/01.MSS.0000176302.99185.BE
- 32 Tan LY, Komarasamy TV, RMT Balasubramaniam V. Hyperinflammatory Immune Response and COVID-19: A Double Edged Sword. *Front Immunol* 2021;**12**:3981. doi:10.3389/FIMMU.2021.742941/BIBTEX
- 33 Hasan LK, Deadwiler B, Haratian A, *et al.* Effects of COVID-19 on the Musculoskeletal System: Clinician’s Guide. *Orthop Res Rev* 2021;**13**:141. doi:10.2147/ORR.S321884
- 34 Vasquez-Bonilla AA, Brazo-Sayavera J, Timón R, *et al.* Monitoring Muscle Oxygen Asymmetry as a Strategy to Prevent Injuries in Footballers. <https://doi.org/10.1080/0270136720222026865> Published Online First: 2022. doi:10.1080/02701367.2022.2026865
- 35 Watson A, Brindle J, Brickson S, *et al.* Preseason aerobic capacity is an independent predictor of in-season injury in collegiate soccer players. *Clinical Journal of Sport Medicine* 2017;**27**:302–7. doi:10.1097/JSM.0000000000000331


- 36 Elliott N, Martin R, Heron N, *et al.* Infographic. Graduated return to play guidance following COVID-19 infection. *Br J Sports Med* 2020;**54**:1174–5. doi:10.1136/BJSPORTS-2020-102637
- 37 Martin RRL, Cibulka MT, Bolgla LA, *et al.* Hamstring Strain Injury in Athletes. *Journal of Orthopaedic and Sports Physical Therapy* 2022;**52**:CPG1–44. doi:10.2519/JOSPT.2022.0301/ASSET/IMAGES/LARGE/JOSPT-CPG1-FIG003.JPEG

## Figure legends


Figure 1: Participants enrolment flowchart

Figure 2: Average number of muscle injuries in the 2019/2020 and 2020/2021 seasons. The dashed blue line represents the average number of injuries per season identified by Bengtsson et al [25].

Figure 3: Kaplan-Meier curves relating to muscular injuries events of Serie A and LaLiga in the 2019-2020 and 2020-2021 seasons.



Serie A



LaLiga

