

Predicting High-Risk Fetal Cardiac Disease Anticipated to Need Immediate Postnatal Stabilization and Intervention with Planned Pediatric Cardiac Operating Room Delivery

Cardiac OR Delivery for High Acuity Fetal CHD

Amol Moray MD^{1*}, Proscovia M. Mugaba MD^{1*}, Chloe Joynt MD², Angela McBrien MD¹, Luke Eckersley MD, PhD¹, Ernest Philipos MD², Paula Holinski MD^{3,4}, Lindsay Ryerson MD³, James Y Coe MD⁵, Sujata Chandra MD⁶, Amanda Aiken MD⁶, Billy Wong MD⁶, Michele Derbyshire MD⁷, Maria Lefebvre BScN², Mohamed Al Aklabi MD⁸, and Lisa K Hornberger MD^{1,6}.

*contributed equally to the development of this manuscript

Affiliations: ¹Fetal & Neonatal Cardiology Program, Division of Cardiology, Department of Pediatrics; ²Division of Neonatology, Department of Pediatrics; ³Division of Critical Care, ⁴Department of Anesthesia; ⁵Interventional Cardiology, Division of Cardiology, Department of Pediatrics; ⁶Department of Obstetrics & Gynecology; ⁷Stollery Pediatric and Mazankowski Cardiac Operating Rooms; ⁸Division of Pediatric Cardiovascular Surgery, Department of Surgery; The Mazankowski Alberta Heart Institute and Women's & Children's Health Research Institute, University of Alberta

For Correspondence: Lisa K Hornberger MD, Pediatric Cardiology, Stollery Children's Hospital, WMC 4.26 8440 112 Street, Edmonton, Alberta; lisa.hornberger@ahs.ca; (780)407-3963 and fax (780)407-3954

Word Count: 5950

ABSTRACT

Background: Distances between delivery centers and cardiac services can make the care of fetuses with cardiac disease(CD) at risk of acute cardiorespiratory instability(ACRI) at birth a challenge. In 2013 we implemented a fetal echocardiography(FE)-based algorithm targeting fetuses considered high-risk for ACRI at ≤ 2 hours of birth for Caesarian section(CS) delivery in our pediatric cardiac operating room(PCOR) of our children's hospital. We examine the experience and outcomes of affected newborns.

Methods: We reviewed maternal and postnatal medical records of all fetuses with CD at high-risk for ACRI encountered January 2013-March 2022. Secondary analysis was performed including all fetuses with diagnoses of d-transposition of the great arteries/intact ventricular septum(d-TGA/IVS) and hypoplastic left heart syndrome(HLHS) encountered over the study period.

Results: Forty fetuses were considered high-risk for ACRI: 15 d-TGA/IVS and 7 HLHS with restrictive atrial septum(RAS), 4 absent pulmonary valve syndrome, 3 obstructed anomalous pulmonary veins, 2 severe Ebstein anomaly, 2 thoracic/intracardiac tumors and 7 others. PCOR delivery occurred for 33 but not for 7 (5 d-TGA/IVS, 2 HLHS with RAS). For high-risk cases, FE had a positive predictive value of 50% for intervention/ECMO/death at ≤ 2 hours and 70% at ≤ 24 hours. Of "low-risk" cases, 6/46 with d-TGA/IVS and 0/45 with HLHS required intervention at ≤ 2 hours. FE predicted intervention/ECMO/death at ≤ 2 hours with a sensitivity of 67%, specificity 93%, and positive and negative predictive values of 87% and 87%, respectively, for d-TGA/IVS, and 100%, 95%, 71%, and 100% for HLHS, respectively.

Conclusions: FE predicts need for urgent intervention in majority with d-TGA/IVS and HLHS, and in half of the entire spectrum of high-risk CD.

Non-standard Abbreviations and Acronyms

ACRI: acute cardiorespiratory instability

ASO: arterial switch operation

BAS: balloon atrial septostomy

CD: cardiac disease

CS Cesarean section

d-TGA/IVS -dtransposition of the great arteries/intact ventricular septum

ECMO: extracorporeal membrane oxygenation

FE: fetal echocardiography

HLHS: hypoplastic left heart syndrome

LOC: level of care

PCOR: pediatric cardiac operating room

RAS: restrictive atrial septum

TAPVD: total anomalous pulmonary venous drainage

TOF/APV: tetralogy of Fallor with absent pulmonary valve syndrome

INTRODUCTION

Advances in fetal echocardiography (FE) have made it possible to accurately evaluate fetal cardiac anatomy and circulatory physiology in a significant proportion of fetuses with cardiac disease (CD).^{1,2} This has enabled the prediction of the intrauterine and perinatal course of most fetal CD,^{2,3} which, in turn has led to improved perinatal care and the development of intrauterine interventions preventing the evolution of more severe secondary cardiac pathology.²⁻⁷

In more recent years, FE has increasingly facilitated recognition of a subset of critical CD with potential for acute cardiorespiratory instability (ACRI) soon after birth. Prenatal predictors of ACRI are evolving, particularly for d-transposition of the great arteries with intact ventricular septum (d-TGA/IVS)⁸⁻¹¹ and hypoplastic left heart syndrome (HLHS) with a restrictive or intact atrial septum (RAS).^{12,13} Other critical CD lesions have also been recognized as potentially life-threatening at birth.¹⁴⁻¹⁸ With this knowledge, FE-based algorithms have evolved to guide perinatal management, including mode and location of delivery. Only two existing studies report promising results of how this approach could ultimately improve the clinical outcomes of these challenging patients^{3,19}; however further data are needed that explore approaches with different types of multidisciplinary infrastructure and geographical challenges faced by many pediatric cardiac programs internationally.

It is not uncommon for obstetrical services to be separated by a short, but important, distance from the pediatric cardiac surgical or interventional centers. This renders the ex-utero transport of a critically ill newborn to the cardiac center a challenge, and may reduce the benefit of a prenatal diagnosis. There has been a move by some centers towards delivery of such fetuses under controlled circumstances in a pediatric cardiac operating room (PCOR) with a full cardiac

team who can stabilize the newborn by performing cardiac intervention within minutes of birth.^{3,19} There is a paucity of data on the efficacy of this approach and its risks in the published literature.

In 2013, our Western Canadian centralized pediatric cardiac program initiated a FE-based algorithm using a multi-disciplinary team approach to risk-stratify and triage fetuses for specialized delivery room care, 5.4 kilometers from high-risk obstetrical services. The current study reviews our experience and assesses the accuracy of FE in predicting the need for urgent postnatal intervention for a spectrum of high-risk pathologies. We further focus on the more common lesions of d-TGA/IVS and HLHS with RAS to examine the sensitivity and specificity of current prenatal predictors.

METHODS

This retrospective cohort study was approved of by the Human Research Ethics Committee of the University of Alberta. Pregnancies complicated by major structural and functional fetal CD, including life-threatening arrhythmias, evaluated and managed in our fetal and neonatal cardiology program from January 2013 to March 2022 were identified through our institutional fetal cardiology database. Firstly, those complicated by fetal CD assigned to delivery room level of care (LOC) 4 as per the American Heart Association Fetal Cardiology Guidelines²⁰ were analyzed. We then performed analysed all pregnancies complicated by a fetal diagnosis of d-TGA/IVS and HLHS evaluated and delivered at our institution over the time same period.

Fetal Echocardiography-Based Algorithm: In 2013, given the distance between our quaternary obstetrical hospital, the Lois Hole Hospital for Women, and the pediatric cardiac program at the Stollery Children's Hospital, our institution evolved a multi-disciplinary team approach guided by FE to risk-stratify CD²⁰: infants having non-ductus arteriosus dependent CD remained at the obstetrical hospital (LOC 1-2), those with ductus arteriosus-dependent CD were transferred to the Stollery Children's Hospital within hours of birth (LOC 2-3), and CD cases deemed at high-risk of ACRI were delivered in our PCOR by Caesarian section (CS) (LOC 4). FE evaluation in each case entailed assessing for lesion-specific criteria to evaluate CD severity using published data, whenever available, and/or clinical experience with multidisciplinary team consensus. In d-TGA/IVS, we assessed for RAS and a small/restrictive ductus arteriosus,⁸⁻¹¹ which include tethered, fixed position of septum primum, small atrial flow orifice (≤ 3 mm), ductus arteriosus constriction, and increased ductal flow velocities suggestive of obstruction. Serial studies were performed to assess for evolving features with ≥ 2 exams after 35 weeks for most. In HLHS, we assessed for features of RAS including foramen ovale size < 3 mm, thick atrial septum and pulmonary vein Doppler flow forward/reverse velocity-time integral (FR VTI) ratio, with a ratio of < 3 near term suggestive of severe left atrial hypertension necessitating emergent intervention.¹² Maternal hyperoxygenation was performed as previously described.²¹

For other prenatally diagnosed CD at risk of ACRI, we assessed for general and lesion-specific features as elaborated in Table 1, including presence/absence of hydrops, cardiac size, combined cardiac output, and evidence of biventricular dysfunction as previously reported.^{2-4,14-18} Where available, noncardiac data was acquired through ultrasound and fetal magnetic resonance imaging for extracardiac pathology and features of airway obstruction (TOF/APV)¹⁶ or lung

parenchyma suggestive of ‘nutmeg lung’ (HLHS with RAS and obstructed total anomalous pulmonary venous drainage (TAPVD)).²²

Pediatric Cardiac Operating Room Delivery Planning: For those fetuses deemed at highest-risk of ACRI (LOC 4), expected to require an intervention at ≤ 2 hours and/or considered at exceptionally high-risk during inter-hospital transport, CS delivery in the PCOR was recommended with the cardiac team on standby to carry out acute neonatal care. CS delivery was based on the predicted need for urgent postnatal interventions dependent on a pre-assembled multidisciplinary team with specific technical skills. For delivery in the PCOR, the fetus had to have no major extracardiac pathology including chromosomal/genetic anomaly that would make aggressive management unlikely to alter a poor prognosis. The mother could not be deemed at high-risk given the lack of obstetrical support in the PC hospital. Our management algorithm is presented in Figure 1.

Once a pregnancy was identified as potentially meeting criteria for PCOR delivery, and the patient agreed, clinical details were reviewed by a multidisciplinary team, and, with the group’s consensus, the delivery plan was made, taking into consideration the ideal gestational age, date, needs at delivery for acute neonatal care and disposition of the newborn (surgical neonatal intensive care unit versus pediatric cardiac intensive care unit) depending on likelihood for needing extracorporeal membrane oxygenation (ECMO). The teams involved in the consensus included obstetrical, neonatology, and fetal cardiology specialists consistently and other pediatric cardiology subspecialists as applicable, cardiac surgeons, pediatric cardiac critical care, pediatric cardiac anesthesia and both obstetrical and neonatal nursing. Delivery was planned for 38-39 weeks of gestation unless there was a high-risk of earlier delivery or acute fetal

compromise/death. On the morning of delivery, both maternal and neonatal multidisciplinary teams assembled prior to the delivery in preparation for enacting the perinatal care plan. Delivery occurred in one PCOR room and the baby was transported at birth to an adjoining PCOR where the cardiac care team was available for resuscitation. Following CS, the mother was monitored by an obstetrical nurse for 3-5 hours in the recovery room to ensure clinical stability prior to being transported back to the obstetrical hospital for further care. It was understood, should the mother go into spontaneous labor, delivery would occur in the obstetrical hospital with early notification of the multidisciplinary team and urgent neonatal transport protocols enacted.

Data Collection and Analysis: Demographic and clinical data were obtained through a chart review of fetal/postnatal echocardiograms and medical records. We extracted data on the fetal CD and findings suggesting risk of ACRI, delivery room and postnatal care plan, delivery information, postnatal findings and management and their timing from birth, and postnatal outcomes. For those in whom CS in the PCOR occurred, we analyzed maternal postnatal outcomes documented in the same admission. We compared the fetal diagnosis and delivery room and postnatal care plans with the postnatal diagnosis, clinical course and treatment/interventions performed and we used this to assess the utility of using FE in identifying high-risk fetal CD. We further defined its accuracy in predicting postnatal ACRI (positive predictive value), for purposes of this review defined as a combined outcome of intervention/ECMO/death occurring at ≤ 2 hours and ≤ 24 hours after birth. We performed a sub-analysis of all fetal d-TGA/IVS and HLHS diagnosed prenatally in our program and managed at the Stollery during the study period. We determined sensitivity, specificity, positive and negative predictive values of FE for fetuses at risk of postnatal ACRI shortly after birth.

RESULTS

Over the study period, 670 fetuses with CD or persistent brady or tachyarrhythmias were encountered in our program. Of these, 40(6.1%) were identified for delivery in the PCOR, including 37 singleton and 3 twin pregnancies (one affected twin each). Figure 2 summarizes the CD diagnoses and outcomes of this cohort, all confirmed to have no major discrepancies between the prenatal and postnatal cardiac diagnoses after birth. The largest CD subgroups included 15 with d-TGA/IVS and 7 with HLHS with RAS. The 18 others included 4 with TOF/APV, 3 with TAPVD with obstruction, 2 with severe Ebstein anomaly, 2 with persistent, life-threatening arrhythmias and 7 with other CD (Table 1). Figures 3-5 demonstrate examples of ACRI cases.

Of the 40 fetuses, 33(82.5%) were delivered in the PCOR as planned at a median gestation of 37.4(range 32.7-39.9)weeks and median birth weight of 3.0(range 2.1-4.4)kg. Of these 33, 8 (24.2%) were delivered at <37 weeks' gestation: 3 with twin gestations (2 dichorionic, 1 monochorionic at 35.0-36.0 weeks), 4 for evolving compromise/hydrops and/or non-reassuring obstetrical surveillance (32.7-36.7 weeks), 1 with spontaneous labor with rapid transport to the PCOR (33.6 weeks). Seven others of the 40 were delivered out of the PCOR at a median gestation of 38.4(range 37.4-40.0) weeks and median birth weight 3.3(range 2.8-3.7)kg, including 5 with d-TGA/IVS with early spontaneous labor and 2 with HLHS and RAS whose mothers declined a CS.

Overall, for the high-risk cohort, FE had a positive predictive value of 50% for predicting the combined outcome of intervention/ECMO/death at ≤ 2 hours and 70% within the first 24 hours.

There was no mortality at delivery for those delivered in the PCOR, and survival to hospital discharge was 82%.

The thirty-three mothers (median age 31, range 19-36 years) with a PCOR CS delivery had spinal anesthesia. Eight had a known concomitant medical condition: 3 gestational diabetes, 1 chronic hepatitis B infection with cholestasis of pregnancy, 1 systemic lupus, 1 gestational hypertension, 1 asthma and 1 chronic kidney disease (G2A2 stage). The delivery represented a primary CS in the majority (28/33, 85%), including 13 primiparous mothers. All were successfully transferred back to the obstetrical hospital in stable condition within 3-5 hours of CS. Overall, there were no major maternal complications (e.g. postpartum hemorrhage, Caesarian hysterectomy, or infection), and total length of hospital stay for the mothers was a median of 2(2-4) days. Minor complications occurred in 5 following hospital discharge (1 seroma, 1 wound pain, 1 spinal headache, 1 mild dehydration, 1 mild superficial wound infection), and were treated as outpatients with mostly supportive treatment.

d-TGA/IVS Delivery & Outcomes: Of 61 fetuses diagnosed with d-TGA/IVS during the study period, 15 were deemed high-risk (LOC4) and anticipated to require urgent balloon atrial septostomy (BAS) after birth. Ten of the 15 had a PCOR delivery (median gestation 38.6, 37.9-39.9weeks). Of these 10, 6 underwent urgent septostomy at ≤ 2 hours after birth for significant cyanosis despite intubation, medications and supplemental oxygen. In a 7th, an intact atrial septum necessitated emergent arterial switch operation (ASO) in the first hour. Two others with an atrial septal defect of < 3 mm underwent BAS at 6 and 9 hours, respectively, due to progressive hypoxemia. The 10th neonate with a redundant septum primum and 3-4mm interatrial

communication, deemed challenging for BAS, underwent ASO at 32 hours post-delivery for hypoxemia.

With early spontaneous labor, 5 of the 15 d-TGA/IVS planned for PCOR delivery were born vaginally at the obstetrical hospital (median gestation 38.2, 37.6-38.7weeks). All 5 were transported to the Stollery Children's Hospital: 4 underwent immediate successful BAS at ≤ 2 hours after birth for severe hypoxemia, and the 5th arrested on arrival requiring ECMO for stabilization. All 15 cases underwent ASO at a median age of 5(0-11) days, with all but one surviving to hospital discharge.

All 46 d-TGA/IVS cases classified as lower-risk (LOC 2/3) and encountered within the study period were delivered in our obstetrical hospital, 14(30%) by CS, and were safely transferred for specialized cardiac care. Six(13%), underwent successful BAS at ≤ 2 hours after birth for severe hypoxemia. An additional 9 underwent BAS at ≤ 24 hours (range 3-14 hours) after birth, and one preterm infant required a BAS on day 11. All underwent ASO and survived to hospital discharge.

For the entire cohort with d-TGA/IVS, FE had a sensitivity of 67%, specificity of 93%, and positive and negative predictive values of 80% and 87%, respectively, for predicting the outcome of cardiac intervention/ECMO/death at ≤ 2 hours after birth.

HLHS Delivery & Outcomes: Of 52 fetuses diagnosed with HLHS during the study period, 7 were classified as high-risk for ACRI, requiring urgent atrial septoplasty. Five had PCOR deliveries at a median gestational age of 37(35-39)weeks and median birth weight 3.0 (2.5-4.0)kg. Three underwent atrial septoplasty with stenting at ≤ 2 hours of delivery. One of the 3, a

twin, who delivered at 35 weeks with fetal MRI features of “nutmeg lung”, required ECMO after atrial septoplasty and demised at <24 due to disseminated intravascular coagulopathy with persistently high lactates and intraventricular hemorrhage. A 4th developed florid pulmonary congestion necessitating urgent atrial septoplasty at 6 hours, and a 5th case with a redundant atrial septum, initially deemed not to require urgent atrial septoplasty, underwent early Norwood-Sano surgery at 36 hours given evolving hypoxemia and pulmonary venous congestion. Two of the 7 high-risk HLHS were delivered vaginally: one in a local hospital to a mother lost to follow-up from our service after the plan for PCOR delivery was made. This baby demised within minutes of delivery. Another mother who declined CS, had spontaneous labor and delivered vaginally at our obstetrical hospital. On arrival to our PC hospital, with profound hypoxemia the neonate required ECMO. Care was withdrawn at <24 hours due to severe end organ dysfunction. Of the surviving 4 of 7 high-risk HLHS infants, 3 underwent Norwood-Sano surgery at a median age of 5 (2 -9) days and 1 a hybrid procedure at 20 days. All four survived to hospital discharge. Survival of high-risk HLHS who had a PCOR delivery was 80% versus 0% without PCOR delivery.

All 45 low-risk fetal HLHS cases encountered within the study period were delivered at the obstetrical hospital, 16(36%) by CS, and were safely transferred for specialized cardiac treatment. None required intervention prior to Norwood-Sano or hybrid procedure and all survived to hospital discharge. For the entire cohort of fetuses with HLHS, FE had a sensitivity of 100%, specificity of 95%, and a positive and negative predictive values of 71% and 100%, respectively, in predicting the combined outcome of cardiac intervention/ECMO/death at ≤ 2 hours of delivery.

Other than the “low-risk” d-TGA/IVS cases requiring urgent BAS, no other prenatally diagnosed CD case encountered over the study period with intent-to-treat and not deemed at risk for ACRI required emergent intervention/ECMO or died at ≤ 2 hours after birth.

DISCUSSION

A small but important subset of fetuses with major CD are at risk of ACRI shortly after birth and require specialized delivery planning to provide immediate stabilization with an intervention and/or medical treatment. With increasing experience in fetal CD diagnosis, the benchmark of a successful fetal cardiology program has shifted from diagnostic accuracy to predicting perinatal/neonatal presentation with detailed delivery care planning in an effort to optimize perinatal outcomes. Our results suggest that FE can be used to effectively predict most high-risk cases of d-TGA/IVS and HLHS with RAS, and the need for very early intervention in at least 50% of other cardiovascular pathology through a multidisciplinary team approach. We further describe a unique algorithm of an organized PCOR delivery remote from our quaternary obstetrical hospital, an approach to date that has received little attention.

Ideally, for the highest-risk fetal CD, delivery should occur in or within close proximity to the PC hospital to access the necessary infrastructure for resuscitation. Unfortunately, many programs internationally have a significant distance separating the delivery room from the PC hospital, making the ex-utero transport of a critically unstable neonate challenging. Even transporting such newborns from an adjoining building or between floors can, at times, be difficult. Alternatively, neonatal cardiac resuscitation can be orchestrated where the delivery occurs, but for many centers, this would be remote from other necessary equipment and

personnel. Determining where delivery of high-risk cardiac newborns should occur is part of a complex decision-making process involving collaboration between multiple teams, and requires consideration of infrastructure, including facilities, equipment and personnel, within any program. It requires communication algorithms which take into consideration factors such as availability of expertise on site to offer both maternal and neonatal care, treatment options, the distance from the delivery room to the intensive care or interventional/surgical unit, and practical aspects of transporting a sick newborn safely and efficiently. An additional factor is availability of appropriate delivery and resuscitation space on the right day and time. The PCOR delivery approach used at our institution represents an important move towards minimizing cardiac mortality and morbidity of the highest risk fetal CD diagnoses by reducing crucial time to a life-saving intervention or stabilization. Arrival of the neonate in the intervention suite without prior significant cardiorespiratory compromise should have a positive impact on the success rate of intervention with a reduction in complications, and ultimately lead to improved short and long-term outcomes. Parents are key decision makers, and this type of perinatal care planning requires parental commitment (especially for the mother) to consent to this strategy of care.

Our study shows successful implementation of an algorithm of high acuity perinatal/neonatal care through a multidisciplinary, cardiac program-wide approach. We witnessed no important maternal morbidity beyond the CS, and mitigated mortality and morbidity risks for many affected newborns. We used a combined outcome of intervention/ECMO/death at ≤ 2 hours of delivery to evaluate our institutional experience of predicting such cases. The distance from our obstetrical to PC hospitals and the terrain covered during transport, including a heavily trafficked areas and dependence on a bridge, warranted this time cut-off for intervention. However, this

only takes into consideration newborns requiring intervention; whereas, it did not capture risks of transporting a fragile newborn with potential for cardiopulmonary extremis and collapse in transit. Without a randomized trial, it is difficult to fully ascertain the benefits of avoiding transportation. However, the two cases requiring emergent atrial septoplasty, both arresting on arrival to our cardiac center necessitating ECMO and others requiring immediate intervention following PCOR delivery, emphasize the critical nature of such newborns and the need to balance risks of late prematurity as well as maternal CS outside of the obstetrical hospital versus, for some, potential arrest and even death. We attempted to mitigate maternal risks by only considering largely lower-risk mothers, moving the smaller but fully complemented obstetrical team for the procedure, and following a brief period of monitoring relocating the mother to the obstetrical hospital for ongoing postpartum care.

In a high-volume pediatric cardiac program, HLHS and d-TGA/IVS with RAS/IAS comprise the bulk of fetuses requiring cardiac interventions at birth as witnessed in our experience. It was reassuring that previously established FE-based parameters in retrospectively examined cohorts⁸⁻¹³ were fairly accurate in guiding delivery planning for most with these two lesions. For HLHS with RAS we found FE to be highly accurate (100% sensitivity and 95% specificity) in predicting postnatal pulmonary venous hypertension requiring decompression of the left atrium. Facilitated by improving prenatal detection of d-TGA in our province,²³ we prospectively predicted a need for BAS at ≤ 2 hours with a sensitivity of 67% and specificity of 93% and failed to predict the need for urgent BAS in only 13% of the others. As has been reported previously,²⁴ there are still limitations in predicting d-TGA/IVS with RAS with resultant profound hypoxemia at birth, especially in late 3rd trimester pregnancies. One advantage of our approach is that the

infant with d-TGA/IVS is delivered at a time when personnel and equipment for BAS are immediately available for a day-time procedure, often performed during a short “honeymoon period” when the baby has not as yet become profoundly hypoxemic and acidotic as a consequence. Such interventions, under less duress, with earlier optimized mixing should contribute to less morbidity and mortality previously reported,²⁵ and ultimately improved longer-term outcomes. Furthermore, immediate availability of other services, should a rare catastrophic event occur, provides further benefit of this approach.

For the other lesions in our cohort with limited published data to guide delivery planning, we formulated PCOR care plans based on anticipated pathophysiology and fragility in the perinatal transition with expert multidisciplinary consensus. With experience, we have modified our approach further, particularly for Ebstein anomaly and TOF/APV, including more routine use of fetal MRI to assess airways in the latter, where most infants may have been sufficiently stabilized for transport. For TAPVD, although we correctly predicted obstruction, we found it challenging to predict the exact timing of intervention. Low velocity non-pulsatile pulmonary vein flow aids in predicting obstruction, but in our cohort, one baby required acute intervention with pulmonary vein flow patterns more akin to RAS in HLHS. Another required ECMO by four hours and still another had surgical repair several hours after birth following necessary axial imaging. In the latter 2 cases, PCOR delivery provided additional time of relative stability for further imaging prior to repair. In a few cases we had missed opportunities and could have been more proactive in instituting the interventions without waiting for decompensation. Every institution has a learning curve when a new process is employed, and we would approach such

cases more aggressively in the current era. Timely post-delivery review of individual cases has allowed us to continue to refine our algorithm.

Limitations: The current study was a single-center experience over nearly a decade. The study was limited by its small sample size, particularly for more rare lesions, making it difficult to draw strong conclusions regarding the accuracy of FE for all CD. There was also a potential for intervention bias introduced by having a predetermined delivery and postnatal care plan, mitigated, to some extent, by relying on the newborn's clinical status following the initial perinatal transition and a multidisciplinary team decision as to whether to proceed. Thus, if the baby could be stabilized, a planned early intervention did not occur. It is acknowledged that the transition norms of oxygen saturation, blood pressure and heart rates of many CD in the first 10-15 minutes after birth is an emerging area of study and that ongoing experience may alter our interpretation in the future regarding need for and timing of intervention.²⁶ The study did not explore the potential long-term benefits of early restoration of adequate cardiac output and/or oxygenation, the impact on morbidity and mortality after further definitive surgical intervention or the longer-term impact, including on neurodevelopment. Additionally, the limited number of mothers undergoing CS in the PCOR may not truly reflect the potential short- and long-term implications of obstetrical surgery performed in a children's hospital.

CONCLUSION

FE predicted need for urgent intervention in majority with d-TGA/IVS and HLHS, and in half of the entire spectrum considered high-risk. Based on this experience, FE can be used to predict ACRI, especially for d-TGA/IVS and HLHS with RAS, thus allowing for planned perinatal and

acute neonatal specialty delivery room care. Delivery-planning needs to be tailored to a given institution's geographical factors and available infrastructure to take full advantage of the prenatal diagnosis. For some, such as our own, a well-organized algorithm which includes delivery within the PCOR of the highest acuity fetal CD cases may be warranted and can be successfully implemented with multidisciplinary involvement for the care of the mother and baby.

Acknowledgements: We want to acknowledge the critical contributions of our maternal-fetal-medicine nurses, operating room and obstetrical managers as well as other staff who have made the execution of these deliveries possible.

Source of Funding: Drs Moray and Mugaba were funded through the Stollery Children's Hospital Foundation and the Department of Pediatrics at the University of Alberta.

Disclosures: None

REFERENCES

1. Maulik D, Nanda NC, Maulik D, Vilchez G. A brief history of fetal echocardiography and its impact on the management of congenital heart disease. *Echocardiography* 2017;34(12):1760-1767.
2. Sanapo L, Pruetz JD, Słodki M, Goens MB, Moon-Grady AJ, Donofrio MT. Fetal echocardiography for planning perinatal and delivery room care of neonates with congenital heart disease. *Echocardiography*. 2017;34(12):1804-1821.
3. Donofrio MT. Predicting the Future: Delivery Room Planning of Congenital Heart Disease Diagnosed by Fetal Echocardiography. *Am J Perinatol*. 2018;35(6):549-552.
4. Donofrio MT, Levy RJ, Schuette JJ, Skurow-Todd K, Sten MB, Stallings C, Pike JI, Krishnan A, Ratnayaka K, Sinha P, et al. Specialized delivery room planning for fetuses with critical congenital heart disease. *Am J Cardiol*. 2013;111(5):737-47.
5. Holland BJ, Myers JA, Woods CR. Prenatal diagnosis of critical congenital heart disease reduces risk of death from cardiovascular compromise prior to planned neonatal cardiac surgery: a meta-analysis. *Ultrasound Obstet Gynecol*. 2015;45(6):631-8.
6. Moon-Grady AJ, Morris SA, Belfort M, Chmait R, Dangel J, Devlieger R, Emery S, Frommelt M, Galindo A, Gelehrter S, et al. International Fetal Cardiac Intervention Registry. International Fetal Cardiac Intervention Registry: A Worldwide Collaborative Description and Preliminary Outcomes. *J Am Coll Cardiol* 2015;66(4):388-99.
7. Freud LR, Tworetzky W. Fetal interventions for congenital heart disease. *Curr Opin Pediatr*. 2016;28(2):156-62.
8. Maeno YV, Kamenir S, Sinclair B, van der Velde ME, Smallhorn JF, Hornberger LK. Prenatal Features of Ductus Arteriosus Constriction and Restrictive Foramen Ovale in d-Transposition of the Great Arteries. *Circulation* 1999; 99:1209-1214.
9. Punn R, Silverman NH. Fetal predictors of urgent balloon atrial septostomy in neonates with complete transposition. *J Am Soc Echocardiogr* 2011;24(4):425-30.
10. Vigneswaran TV, Zidere V, Miller OI, Simpson JM, Sharland GK. Usefulness of the Prenatal Echocardiogram in Fetuses With Isolated Transposition of the Great Arteries to Predict the Need for Balloon Atrial Septostomy. *Am J Cardiol*. 2017;119(9):1463-1467.
11. Słodki M, Axt-Flidner R, Zych-Krekora K, Wolter A, Kawecki A, Enzensberger C, Gulczyńska E, Respondek-Liberska M; International Prenatal Cardiology Collaboration Group.

New method to predict need for Rashkind procedure in fetuses with dextro-transposition of the great arteries. *Ultrasound Obstet Gynecol.* 2018;51(4):531-536.

12. Divanović A, Hor K, Cnota J, Hirsch R, Kinsel-Ziter M, Michelfelder E. Prediction and perinatal management of severely restrictive atrial septum in fetuses with critical left heart obstruction: clinical experience using pulmonary venous Doppler analysis. *J Thorac Cardiovasc Surg* 2011;141(4):988-94.

13. Sathanandam SK, Philip R, Gamboa D, Van Bergen A, Ilbawi MN, Knott-Craig C, Waller BR, Javois AJ, Cuneo BF, Management of hypoplastic left heart syndrome with intact atrial septum: a two-centre experience. *Cardiol Young* 2016; 26:1072–1081.

14. Freud LR, Escobar-Diaz MC, Kalish BT, Komarlu R, Puchalski MD, Jaeggi ET, Szwast AL, Freire G, Levasseur SM, Kavanaugh-McHugh A, et al. Outcomes and Predictors of Perinatal Mortality in Fetuses with Ebstein Anomaly or Tricuspid Valve Dysplasia in the Current Era: A Multicenter Study. *Circulation* 2015;132(6):481-9.

15. Wertaschnigg D, Jaeggi M, Chitayat D, Shannon P, Ryan G, Thompson M, Yoo SJ, Jaeggi E. Prenatal diagnosis and outcome of absent pulmonary valve syndrome: contemporary single-center experience and review of the literature. *Ultrasound Obstet Gynecol.* 2013;41(2):162-7.

16. Chelliah A, Moon-Grady AJ, Peyvandi S, Chiu JS, Bost JE, Schidlow D, Carroll SJ, Davey B, Divanovic A, Hornberger L, et al. Contemporary Outcomes in Tetralogy of Fallot With Absent Pulmonary Valve After Fetal Diagnosis. *J Am Heart Assoc* 2021;10(12): e019713.

17. Sridharan S, Sullivan I, Tomek V, Wolfenden J, Škovránek J, Yates R, Janoušek J, Dominguez TE, Marek J. Flecainide versus digoxin for fetal supraventricular tachycardia: Comparison of two drug treatment protocols. *Heart Rhythm* 2016;13(9):1913-9.

18. Weber R, Kantor P, Chitayat D, Friedberg MK, Golding F, Mertens L, Nield LE, Ryan G, Seed M, Yoo SJ, Manlhiot C, Jaeggi E. Spectrum and outcome of primary cardiomyopathies diagnosed during fetal life. *JACC Heart Fail* 2014;2(4):403-11.

19. Berkley EM, Goens MB, Karr S, Rappaport V. Utility of fetal echocardiography in postnatal management of infants with prenatally diagnosed congenital heart disease. *Prenat Diagn.* 2009 Jul;29(7):654-8.

20. Donofrio MT, Moon-Grady AJ, Hornberger LK, Copel JA, Sklansky MS, Abuhamad A, Cuneo BF, Huhta JC, Jonas RA, Krishnan A, et al. Diagnosis and treatment of fetal cardiac disease: a scientific statement from the American Heart Association. *Circulation* 2014;129(21):2183-242.

21. Szwast A, Tian Z, McCann M, Donaghue D, Rychik J. Vasoreactive response to maternal hyperoxygenation in the fetus with hypoplastic left heart syndrome. *Circ Cardiovasc Imaging*. 2010;3(2):172-8.
22. Saul D, Degenhardt K, Iyoob SD, Surrey LF, Johnson AM, Johnson MP, Rychik J, Victoria T. Hypoplastic left heart syndrome and the nutmeg lung pattern in utero: a cause and effect relationship or prognostic indicator? *Pediatr Radiol*. 2016;46(4):483-9.
23. Ravi P, Mills L, Fruitman D, Savard W, Colen T, Khoo NS, Serrano-Lomelin J, Hornberger LK. Population Trends in Prenatal Detection of Transposition of Great Arteries In Alberta: Impact of Obstetrical Screening Ultrasound Guidelines. *Ultrasound Obstet Gynecol*. 2018;51(5):659-664.
24. Donofrio MT, Skurow-Todd K, Berger JT, McCarter R, Fulgium A, Krishnan A, Sable CA. Risk-stratified postnatal care of newborns with congenital heart disease determined by fetal echocardiography. *J Am Soc Echocardiogr*. 2015;28(11):1339-49.
25. Petit CJ, Rome JJ, Wernovsky G, Mason SE, Shera DM, Nicolson SC, Montenegro LM, Tabbutt S, Zimmerman RA, Licht DJ. Preoperative brain injury in transposition of the great arteries is associated with oxygenations and time to surgery, not balloon atrial septostomy. *Circulation* 2009; 119:709-716.
26. Thomas A, Ma AL, Weinberg DD, Huber M, Ades A, Rychik J, Foglia EE. Delivery room oxygen physiology and respiratory interventions for newborns with cyanotic congenital heart disease. *J Perinatol* 2021; 41:2309–231.

Figure Legends

Figure 1. Algorithm for High Risk Fetal Cardiac Disease with Delivery in the PCOR.

CD-cardiac disease; CV-cardiovascular; MFM/1^oOB-maternal-fetal medicine specialist/primary obstetrician; NICU-neonatal intensive care unit/service; OB-obstetrical; OR-operating room; PCOR-pediatric cardiac operating room.

Figure 2. Outcomes of the Fetuses Considered at High Risk of Acute Cardiovascular

Compromise. AS-aortic stenosis; ASO-arterial switch operation; BAS-balloon atrial septostomy; BAV-balloon aortic valvuloplasty; ECMO-extracorporeal membrane oxygenation; IAS-interatrial stent; LPA-left pulmonary artery; CM-cardiomyopathy; PM-pacemaker; TAPVD-total anomalous pulmonary venous drainage; TOF/APV-Tetralogy of Fallot with absent pulmonary valve; TV-tricuspid valve

Figure 3. d-TGA/IVS with Restrictive Atrial Septum. (A) Cross-sectional 4-chamber and (B)

sagittal (B) fetal echo images showing a tethered atrial septum with an aneurysmal foramen ovale flap (arrow A, * B) bowing to the left and tethered to the superior portion of septum secundum. An atrial communication could not be demonstrated. (C) 2D image of the ductus arteriosus demonstrating a discretely narrowed lumen (arrow). (D) Doppler interrogation of the ductus arteriosus showed increased systolic and diastolic velocities suggestive of ductal constriction. (E) Subcostal view at postnatal echocardiography revealed no demonstrable atrial communication, further suggested when (F) the septostomy catheter (arrow) could only tent and not cross the septum. With severe cyanosis and metabolic acidosis, the newborn underwent an arterial switch operation at <1 hour. LA-left atrium; RA-right atrium, RV-right ventricle.

Figure 4. Supracardiac TAPVD with Obstruction. (A) 4-chamber image at 36 weeks showing right(RV) >left(LV) ventricular asymmetry and an increased distance (arrow) between the left atrium (LA) and descending aorta (dAo). Increased echogenicity of the lung parenchyma was suggestive of pulmonary edema. RA-right atrium. (B) In the 3-vessel view, the superior vena cava (S) and aorta (A) were of comparable in size, and there was a structure (V) to the left of the pulmonary artery (P). (C) By color Doppler the vertical vein (red flow and V) coursed over the left pulmonary artery (LPA) to join the left innominate vein. Where the vertical vein coursed over the LPA there was obstruction by color and (D) pulsed Doppler, with a mean gradient of 5mmHg. (E) Pulmonary vein flow was low-velocity and nonphasic flow in keeping with obstruction. This infant required ECMO at 4 hours for progressive cyanosis.

Figure 5. Cardiac Compression by a Large Mediastinal Tumor. (A) Cross-sectional and (B) sagittal fetal echo images at 23 weeks demonstrated a tumor in the superior mediastinum with cystic(C) and solid(S) components abutting the pulmonary artery (PA) and mild cardiac dextroposition. Rapid growth of the tumor from 30 weeks led to development of hydrops. (C) At 33 weeks the now massive tumor occupied much of the fetal chest with significant cardiac compression. These findings prompted preterm delivery. With progressive cardiorespiratory embarrassment, the tumor (D) was removed surgically 7 hours after birth. Pathology assessment confirmed a germ cell tumor, and an acute hemorrhage into the tumor was the cause of its rapid increase in size in utero.

Table 1 – Fetal Cardiac Pathology Recommended for Delivery in the Pediatric Cardiac Operating Room

FE diagnosis	FE indication for PC OR delivery	GA at delivery	Neonatal Course	Outcome
Critical AS	RAS, severely dilated and hypokinetic LV and dilated LA, ascites	37	IAS, BAV ≤ 2hours and stabilized. With severe MS Stage 1 Norwood D14	Fontan completion, stable
Complete heart block	Severe bradycardia, HR low 50s, LV EFE and early onset hydrops	37	HR 50s with acidosis. PPM inserted at ~10 hours	Alive, stable
Large LMCA to coronary sinus fistula	Continuous run-off in aortic arch, potential risk of myocardial ischemia	39	No intervention/treatment required, stable	Fistula resolved by 1 year, stable
Persistent SVT	Reduced BiV function with hydrops	34	AVRT, early initiation of antiarrhythmic drugs	Controlled on drugs, stable
TOF/APV	LMB compression, cardiac shift, polyhydramnios	37	Intubated at birth with prone position and inotropes, TOF repair D7 due to frequent profound desaturations	Alive, stable
TOF/APV	Compression of both main bronchi	39	No ventilatory support required	TOF repair at 3 mos
TOF/APV	Compression of trachea and bronchi	37	Immediate intubation, extubated D2	TOF repair at 4 mos
TOF/APV	RMB severely compressed or atretic	38	CPAP only but weaned at D2	TOF repair at 4.5 mos
Cardiomyopathy	Severely reduced BiV function		LVAD on D14, had heart transplant D27	Alive, stable
Cardiac tumour	Critical RVOT obstruction and LV compression by tumour, with LCO	37	Immediate tumour aspiration; further debulking twice	Pulmonary embolism death at 2 mos
Mediastinal tumour	Acute enlargement of tumor with evolving hydrops	33	Immediate intubation, surgical resection at ~7hours, followed by chemotherapy	Alive, stable
Severe Ebstein TV	Severe cardiomegaly, LCO	36	Cone repair D3	Died on D11
Severe Ebstein TV	Severe cardiomegaly, severe PR LCO	37	ECMO at 12 hours; Starnes procedure D4	Remained hospitalized death at 4 mos
RAI with Supracardiac TAPVD, uAVSD, PAtr	Obstructed pulmonary veins	39	TAPVD repair + central shunt at ~11 hours	Remained hospitalised, death at 3 mos
Supracardiac TAPVD	Obstructed pulmonary veins	37	ECMO at ~4 hours; TAPVD repair D2	Alive, stable
Supracardiac	Obstructed pulmonary veins	34	Emergent TAPVD repair at 1 hour	Alive, PHT on

TAPVD				medication
LPA sling in monochoronic twins	Severe tracheal compression & polyhydramnios	36	Immediate intubation, LPA re-implantation and tracheoplasty D15	Alive, stable
Truncus arteriosus	Severe truncal stenosis with decreased BiV function and severe hypertrophy	35	Truncus repair on D5	Died on D26

Legend: **AS**, aortic stenosis; **BAV**, balloon aortic valvuloplasty; **BiV**, biventricular function; **CS**, Caesarian section; **D**, day; **ECMO**, extracorporeal membrane oxygenation; **FE**, fetal echocardiography; **GA**, gestational age; **IAS**, intact atrial septum; **LA**, left atrium; **LCO**, low cardiac output; **LMCA**, left main coronary artery; **LMB**, left main bronchus; **LPA**, left pulmonary artery **LV**, left ventricle; **mos**, months; **MS**, mitral stenosis; **PA**, pulmonary artery; **PAtr**, pulmonary atresia; **PHT**, pulmonary hypertension; **RAI**, right atrial isomerism; **RAS**, restrictive atrial septum; **RMB**, right main bronchus; **RVOT**, right ventricular outflow tract; **SV**, single ventricle; **SVT**, supraventricular tachycardia; **TAPVD**, total anomalous pulmonary venous drainage; **TOF/APV**, tetralogy of Fallot/absent pulmonary valve syndrome; **uAVSD**, unbalanced atrioventricular septal defect









