

1 **Prevalence and predictors of HIV testing among young men in**
2 **Papua New Guinea: a cross-sectional analysis of a nationally**
3 **representative sample**

4
5 McKenzie Maviso^{1*}

6
7 ¹ Division of Public Health, School of Medicine and Health Sciences, University of Papua New
8 Guinea, Port Moresby, Papua New Guinea

9
10 *Corresponding author

11 Email: mckenzie.maviso@upng.ac.pg

12

13

14

15

16

17

18

19

20

21

22

23

24

25 **Prevalence and predictors of HIV testing among young men in**
26 **Papua New Guinea: a cross-sectional study of a nationally**
27 **representative sample**

28

29 **Abstract**

30 **Background**

31 HIV testing is an important component of HIV prevention and serves as a gateway to other HIV-related
32 services. However, the uptake remains suboptimal among young people, particularly in highly
33 prevalent settings such as Papua New Guinea (PNG). This study used nationally representative data to
34 assess the prevalence and determine predictors of HIV testing uptake among young men aged 15–24
35 years in PNG.

36 **Methods**

37 The 2016–2018 PNG Demographic and Health Survey (PNGDHS) data was used. A total of 1,275
38 young men aged 15–24 years were included in the final analysis. Descriptive, bivariate, and
39 multivariable logistic regression analyses were performed to determine independent predictors of HIV
40 testing. Adjusted odds ratios (AORs) with 95% confidence intervals (CIs) were reported. All analyses
41 were adjusted using survey weights to account for unequal sampling probabilities.

42 **Results**

43 The overall prevalence of HIV testing among young men was 17.2% (95% CI: 15–19). Of those who
44 were tested for HIV, most lived in rural areas (73%), about one-third had a sexual debut at age <15
45 years (32.9%), and 33.9% inconsistently used condoms during sex. In multivariable analysis, men aged
46 20–24 years (AOR 1.18, 95% CI: 1.00–2.31), who own mobile phones (AOR 1.43, 95% CI: 1.00–
47 2.55), who were aware that consistent condom use during sex can reduce HIV risk (AOR 2.18, 95% CI:

48 1.18–4.04), who had paid for sex (AOR 1.75, 95% CI: 1.01–5.83), and who had two or more sexual
49 partners (AOR 1.37, 95% CI: 1.01–3.14) had increased odds of HIV testing. However, decreased odds
50 of HIV testing were found among men who were never married (AOR 0.51, 95% CI: 0.29–0.88), lived
51 in rural areas (AOR 0.54, 95% CI: 0.32–0.92), and consistently used condoms during sex (AOR 0.59,
52 95% CI: 0.34–1.01).

53 **Conclusion**

54 The findings show that HIV testing is low among young men in PNG. To increase HIV testing uptake
55 among young men, it is crucial to implement comprehensive youth-friendly HIV/STI education and
56 tailored sensitization programs and enable more accessible and affordable HIV testing services. Also,
57 outreach and community-based testing programs for young men in prioritized areas requiring urgent
58 prevention intervention are feasible options in PNG.

60 **Introduction**

61 Considerable progress has been made in responding to HIV/AIDS under the Sustainable Development
62 Goals framework to end the epidemic by 2030 [1]. Since 2010, new HIV infection rates have decreased
63 by 14%, and 78% of people living with HIV know their status [2]. However, in the Asia and Pacific
64 region, the HIV response is impacted by marked disparities and varied epidemic trends, notably
65 affecting young people aged 15–24 years, who represent a growing proportion of people who
66 disproportionately endure the burden of HIV. In 2022, an estimated 440,000 young people were living
67 with HIV, accounting for about 25% of new HIV infections in the region [2].

68 The United Nations' 95-95-95 targets to end the HIV epidemic by 2030 called for 95% of
69 people living with HIV to know about their serostatus [3,4]. HIV testing is a key strategic entry point to
70 prevention, treatment, care, and support services. It confers multiple benefits for individuals who test

71 positive and those who test negative and encourages preventive behaviors [5,6]. Despite the significant
72 benefits, young people remain the least likely to get an HIV test and know their HIV status [15,27].
73 Studies have shown that not all, particularly men willingly seek HIV testing services as a way to
74 prevent HIV transmission and/or acquisition due to various barriers, such as individual (poor
75 HIV/AIDS knowledge, testing availability, and unwillingness to test), social (stigma, discrimination,
76 and lack of social support), and health service utilization (loss of trust or confidence in health workers
77 and lack of medication, including ART) [7–9]. These barriers undermine efforts to increase HIV testing
78 and highlight the critical need to develop appropriate HIV prevention strategies.

79 Among young people, their sexual behaviors and susceptibility to HIV infection differ widely
80 by social context, as well as increased individual autonomy and a lack of social control [10,11]. For
81 example, high-risk behaviors such as early sexual debut, substance use (alcohol consumption and drug
82 use), inconsistent condom use, and multiple and concurrent sexual relationships, as well as poor HIV
83 knowledge, low-risk perception, and sensation-seeking behaviors, increase HIV risks and transmissions
84 among young people [12–15]. In addition, biological factors such as low rates of male circumcision
85 [16,17] and manifestations of other sexually transmitted infections (STIs) (chlamydia, gonorrhea,
86 syphilis, and trichomoniasis) [18,19] contribute to some of the increased HIV risk and burden among
87 this population. Despite young people’s vulnerability to HIV infection, they are frequently overlooked
88 in the design and implementation of national AIDS strategies [20,21]. They remain an important target
89 group for HIV prevention and STI surveillance due to their unique behavioral and social-related
90 vulnerability.

91 PNG is one of the culturally diverse countries in the Western Pacific region and is grappling
92 with a concentrated-generalized HIV epidemic, indicating significant disparities in HIV prevalence
93 between key populations (e.g., sex workers and men who have sex with men) and the general
94 population [22,23]. HIV prevalence is 1% in the general adult population [24]. Among young people,

95 the prevalence of HIV is estimated at 0.2% for males and 0.5% for females [24], demonstrating the
96 high vulnerability of this priority group in the country. Although HIV testing services are widely
97 available throughout PNG [25], the current testing rate among young people is suboptimal, especially
98 among those most at risk of HIV [26,27]. Furthermore, a major obstacle in the country continues to be
99 young men's participation in HIV programs and their access to sexual and reproductive health care
100 [28,29]. Evidence concerning HIV knowledge and risk reduction behaviors (e.g., consistent use of
101 condoms and having only one faithful sexual partner) among young men is scarce; although a recent
102 study among young women in PNG found that early sexual debut, having one sexual partner and not
103 having an STI were associated with not testing for HIV [26]. However, young men's knowledge of
104 HIV and sexual relationships is established through a complex interplay of sociocultural norms,
105 expectations, attitudes, and information gleaned from various sources [30].

106 Despite the established profile of risky sexual behaviors among young people [26,31,32], little
107 is known about the prevalence and correlates of HIV testing uptake in young men in PNG. Moreover,
108 evidence concerning predictors influencing the uptake of HIV testing among young men is scarce. This
109 creates a critical knowledge gap that must be filled to inform national-level policy development and
110 targeted interventions. In this study, data from a nationally representative survey was used to determine
111 the prevalence and predictors of HIV testing uptake among young men aged 15–24 years in PNG.

112 **Methods and materials**

113 **Data sources**

114 Data for the current study were derived from the 2016–2018 PNG Demographic and Health Survey
115 (PNGDHS), which used a stratified, multistage cluster sampling method to ensure a sample
116 representative of the population. The survey was conducted between October 2016 and December 2018
117 and collected data on the country's important demographic, socioeconomic, and health indicators. The

118 survey used the list of census units (CUs) from the 2011 PNG National Population and Housing Census
119 [33] as the sampling frame, which contains information on CU location, type of residence (urban or
120 rural), estimated number of residential households, and population by sex. Administratively, the
121 country is divided into four (4) main regions (Southern, Highlands, Momase, and Islands) comprising
122 22 provinces, and each province is subdivided into urban and rural areas. The PNG National Statistical
123 Office (NSO) with technical support and assistance from the Inner City Fund (ICF) through the DHS
124 Program conducted the study. The governments of PNG and Australia, the United Nations Population
125 Fund (UNFPA), and the Children’s Fund of the United Nations (UNICEF) funded the survey [34].

126 **Sampling procedures**

127 The 2016–2018 PNGDHS sample stratification and selection were achieved in two strata: urban
128 and rural areas, except for the National Capital District, which has no rural areas. In each stratum,
129 samples of households were selected in two stages, with CU constituting the primary sampling unit and
130 households as the secondary sampling unit. The first stage involved selecting 800 CUs using a
131 probability proportional to the CU size. In the second stage, 24 households were selected from each
132 cluster using probability sampling, yielding a total sample size of approximately 19,200 households. A
133 total of 7,333 men aged 15–49 years participated in the survey. Young men who never had sex and
134 aged 25 or more were excluded from this study. A total weighted sample of 1,275 young men aged 15–
135 24 years who ever had sex with variables of interest were included in this study. Details about the
136 sampling technique, survey team training, household selection, survey questionnaires, and validation
137 procedures are available in the final report [34].

138

139

140

141 **Variables and measurements**

142 The primary outcome variable of this study was “ever tested for HIV?” and was dichotomized as “0”
143 for “no” or “1” for “yes.” The explanatory variables were selected based on the literature and their
144 relevance [18–20] and were divided into three categories: (1) Sociodemographic factors include age,
145 marital status, educational level, employment status, wealth index, region, and place of residence. Also,
146 household factors measured included exposure to mass media (listening to the radio and watching
147 television), access to the internet, and mobile phone ownership. (2) HIV-related knowledge includes
148 ever heard of AIDS or STIs, can a healthy person have HIV? Can a person get HIV by sharing food
149 with an infected person? Can a person get HIV through witchcraft? Consistent condom use can reduce
150 HIV risk and knowing an HIV testing facility. (3) Sexual risk behaviors include age of sexual debut,
151 number of sexual partners (including spouse), having ever paid for sex, and consistent condom use
152 during sex, and having had an STI (unspecific) in the last 12 months.

153 **Statistical analysis**

154 Data extraction, recoding, and analysis were performed using IBM Statistical Package for the Social
155 Sciences (SPSS), Version 26.0 (Armonk, NY: IBM Corp.). The sample was weighted using the
156 primary sampling unit variable, stratification variable, and weight variable to restore its
157 representativeness and obtain a better estimate throughout the analysis. Descriptive statistics were used
158 to explore the characteristics of the sample and were presented as weighted frequencies (n) and
159 percentages (%). A chi-square test for independence was performed to assess the strength of the
160 association between the dependent variable and each independent variable. Variables that were
161 significant and those with $p \leq 0.25$ in the bivariate analysis were included in a multivariable analysis to
162 determine their collective associations with HIV testing [38]. Since the PNGDHS used a two-stage
163 stratified sampling technique, a complex samples analysis technique was employed to determine

164 independent predictors of HIV testing. Adjusted odds ratios (AORs) with 95% confidence intervals
165 (CIs) were reported. A $p \leq 0.05$ was used to assess statistical significance.

166 **Ethical Considerations**

167 Permission to use data was obtained from the DHS program (<https://dhsprogram.com>) and was only
168 used for this study. The Institutional Review Board (IRB) of Inner City Fund (ICF) International and
169 the IRB committee of PNG examined and approved the 2016–2018 PNGDHS protocols. All the
170 respondents had provided verbal informed consent before each interview. There were no ethical
171 considerations on the researcher's part as the data were anonymized entirely, with no identifiable
172 information on the survey participants when analyzed.

173 **Results**

174 **Sociodemographic characteristics of young men**

175 Table 1 presents the participants' characteristics. Overall, 1,275 sexually active young men aged 15–24
176 years enrolled in the study. The mean age was 20.84 (\pm 2.31) years, while the majority (70.2%) were
177 aged 20–24 years and lived in rural areas (82%). Nearly half had attained primary (44.3%) and
178 secondary (45.5%) education, and about two-thirds (62%) were unemployed. Regarding mass media
179 exposure, more than two-thirds of young men listened to the radio (71.7%), while less than half
180 watched television (47.1%) and had internet access (33.6%). More than half (57.8%) of young men
181 owned mobile phones.

182

183

184

185

Table 1. Sociodemographic characteristics of participants (N = 1,275)

Characteristics	Frequency	
	<i>n</i>	%
<i>Sociodemographic</i>		
Age (years)		
15–19	380	29.8
20–24	895	70.2
Marital status (missing, <i>n</i> = 26)		
Never married	940	75.2
Married	310	24.8
Educational level		
No formal education	73	5.7
Primary	565	44.3
Secondary	580	45.5
Tertiary	58	4.5
Employment status		
Not employed	791	62.0
Employed	485	38.0
Wealth Index		
Poorest	120	9.4
Poorer	119	9.3
Middle	189	14.8
Richer	294	23.0
Richest	553	43.4
Place of residence		
Urban	229	18.0
Rural	1,046	82.0
Region		
Southern	300	23.6
Highlands	455	35.7
Momase	373	29.2
Islands	147	11.5
Read newspaper/magazine		
No	369	28.9
Yes	907	71.1
Listen to the radio		

No	361	28.3
Yes	914	71.7
Watch television		
No	674	52.9
Yes	601	47.1
Internet access		
No	847	66.4
Yes	428	33.6
Own a mobile phone		
No	537	42.2
Yes	738	57.8

Note: Weighted frequencies (*n*) and percentages (%)

186

187 **The prevalence of having ever been tested for HIV**

188 Overall, 17.1% (95% CI: 0.15, 0.19) of the young men had ever been tested for HIV (Fig 1).

189

190 **Fig 1.** The rates of HIV testing among young men in PNG.

191

192 **HIV knowledge and sexual risk behaviors**

193 Table 2 shows the HIV-related knowledge and sexual risk behaviors of young men. The vast majority
 194 of young men had adequate knowledge that a healthy person can have HIV (90.8%) and that using
 195 consistent condom use during sex can reduce the risk of HIV transmission (80.2%). Most knew of an
 196 HIV testing facility (70.8%). Regarding young men's sexual risk behaviors, over a quarter were
 197 reported to have initiated sex at an early age (<15 years) (28.2%) and inconsistently used condoms
 198 during sex (27.7%). The majority of young men had a single sexual partner in the last 12 months
 199 (88.7%).

200

Table 2. HIV knowledge and sexual risk behaviors of young men (N = 1,275)

Characteristics	Frequency	
	<i>n</i>	%
<i>HIV Knowledge</i>		
Ever heard of AIDS		
No	92	7.2
Yes	1,183	92.8
Ever heard of STIs		
No	74	5.8
Yes	1,201	94.2
A healthy person can have HIV (missing, <i>n</i> = 326)		
No	88	9.2
Yes	861	90.8
A person can get HIV by sharing food with an infected person (missing, <i>n</i> = 298)		
No	822	84.1
Yes	155	15.9
A person can get HIV through witchcraft (missing, <i>n</i> = 297)		
No	908	92.8
Yes	70	7.2
Consistent condom use during sex can reduce HIV risk (missing, <i>n</i> = 311)		
No	191	19.8
Yes	773	80.2
Know HIV testing facility (missing, <i>n</i> = 93)		
No	346	29.2
Yes	836	70.8
<i>Sexual Risk Behavior</i>		
Age of sexual debut (years)		
<15	360	28.2
>15	916	71.8
Number of sexual partners, including spouse (last 12 months)		
0–1	1,130	88.7
2 or more	144	11.3
Ever paid for sex (missing, <i>n</i> = 49)		
No	1,164	94.9

Yes	63	5.1
Consistent condom use during sex (missing, $n = 95$)		
No	326	27.7
Yes	854	72.3
Had any STIs (last 12 months) [†]		
No	1,211	94.9
Yes	65	5.1

The type of STI was not specified in the dataset.

Note: Weighted frequencies (n) and percentages (%)

201

202 **Bivariate analysis of predictors of HIV testing among young men**

203 Table 2 presents the relationship between sociodemographic characteristics, HIV knowledge, sexual
204 risk behaviors, and HIV testing uptake among young men. Overall, only 17.1% ($n = 222$) of young men
205 have ever been tested for HIV across sociodemographic, HIV knowledge, and sexual risk behavior
206 strata. Having ever been tested for HIV was high among men aged 20–24 years (78.8%), those who
207 were married (66.5%), had attained secondary education (58.1%), were unemployed (53.6%), and lived
208 in rural areas (73%). Similarly, HIV testing remained high among those who read a newspaper or
209 magazine (81.5%), listened to the radio (79.2%), and watched television (53.2%). Regarding HIV
210 knowledge, having ever heard of AIDS and STIs ($p < 0.001$), being aware that consistent use of
211 condoms can reduce HIV risk ($p = 0.011$), and knowing an HIV testing facility ($p < 0.001$) were
212 significantly associated with HIV testing uptake. Furthermore, there was a statistically significant
213 association between HIV testing and the following sexual risk behavior factors: age of sexual debut (p
214 = 0.049), ever paid for sex ($p < 0.001$), consistent condom use ($p = 0.02$), and having an STI in the past
215 12 months ($p < 0.001$).

216

217

218

Table 3. Bivariate analysis of sociodemographic, HIV knowledge, behavior, and HIV testing (N = 1,275)

Characteristics	Ever been tested for HIV				P-value
	No (n = 1,053)	%	Yes (n = 222)	%	
<i>Sociodemographics</i>					
Age (years)					0.002*
15–19	333	31.6	47	21.1	
20–24	720	68.4	175	78.8	
Marital status (missing, n = 26)					0.001**
Never married	238	23.0	72	33.5	
Married	796	77.0	143	66.5	
Educational level					0.000**
No formal education	68	6.5	5	2.3	
Primary	490	46.5	75	33.8	
Secondary	450	42.7	129	58.1	
Tertiary	45	4.3	13	5.9	
Employment status					0.005*
Not employed	671	63.7	119	53.6	
Employed	382	36.3	103	46.4	
Wealth Index					0.271
Poorest	96	9.1	24	10.9	
Poorer	94	8.9	25	11.3	
Middle	162	15.4	27	12.2	
Richer	235	22.3	58	26.2	
Richest	466	44.3	87	39.4	
Place of residence					0.000**
Urban	169	16.1	60	27.0	
Rural	883	83.9	162	73.0	
Region					0.000**
Southern	248	23.6	53	23.8	
Highlands	355	33.7	100	44.8	
Momase	332	31.5	41	18.4	
Islands	118	11.2	29	13.0	
Read newspaper/magazine					0.000**
No	328	31.1	41	18.5	
Yes	725	68.9	181	81.5	
Listen to the radio					0.006*

No	315	29.9	46	20.8	
Yes	738	70.1	175	79.2	
Watch television					0.048*
No	570	54.1	104	46.8	
Yes	483	45.9	118	53.2	
Internet access					0.000**
No	730	69.3	117	52.7	
Yes	323	30.7	105	47.3	
Own a mobile phone					0.000**
No	475	45.1	62	28.1	
Yes	578	54.9	159	71.9	
<i>HIV Knowledge</i>					
Ever heard of AIDS					0.000**
No	92	8.7	0	0	
Yes	961	91.3	222	100	
Ever heard of STIs					0.000**
No	47	7.0	0	0	
Yes	979	93.0	222	100	
A healthy person can have HIV (missing, $n = 326$)					0.170
No	77	9.9	11	6.5	
Yes	702	90.1	158	93.5	
A person can get HIV by sharing food with an infected person (missing, $n = 298$)					0.248
No	673	83.5	148	87.1	
Yes	133	16.5	22	12.9	
A person can get HIV through witchcraft (missing, $n = 297$)					0.376
No	739	92.5	168	94.4	
Yes	60	7.5	10	5.6	
Consistent condom use during sex can reduce HIV risk (missing, $n = 311$)					0.011*
No	170	21.3	21	12.7	
Yes	627	78.7	145	87.3	
Know HIV testing facility (missing, $n = 93$)					
No	346	36.0	0	0	0.000**
Yes	614	64.0	222	100	
<i>Sexual Risk Behaviors</i>					

Age of sexual debut (years)					0.049*
<15	287	27.3	73	32.9	
>15	766	72.7	149	67.1	
Number of sexual partners, including spouse (in the last 12 months)					0.101
0–1	941	89.4	189	85.5	
2 or more	112	10.6	32	14.5	
Ever paid for sex (missing, <i>n</i> = 49)					0.000**
No	971	96.7	192	88.1	
Yes	37	3.7	26	11.9	
Consistent condom use during sex (missing, <i>n</i> = 95)					0.020*
No	251	26.2	75	33.9	
Yes	707	73.8	146	66.1	
Had any STIs (last 12 months) [†]					0.000**
No	1,010	95.9	200	90.1	
Yes	43	4.1	22	9.9	

* $p \leq 0.05$

** $p \leq 0.001$

[†]The type of STI was not specified in the dataset.

219

220 **Multivariable analysis of predictors of HIV testing among young men**

221 Table 3 shows the results of the multivariable analysis examining the association between HIV testing
 222 uptake and sociodemographic, HIV-related knowledge, and sexual risk behavior variables. Age, marital
 223 status, place of residence, being aware that consistent condom use during sex can reduce HIV risk, ever
 224 paid for sex, number of sexual partners, and consistent condom use during sex were statistically
 225 significant predictors of HIV testing uptake among young men. In the multivariable analysis, men aged
 226 20–24 years (AOR 1.18, 95% CI: 1.00–2.31), who own mobile phones (AOR 1.43, 95% CI: 1.00–
 227 2.55), who were aware that consistent condom use during sex can reduce HIV risk (AOR 2.18, 95% CI:
 228 1.18–4.04), who had paid for sex (AOR 1.75, 95% CI: 1.01–5.83), and who had two or more sexual
 229 partners (AOR 1.37, 95% CI: 1.01–3.14) had increased odds of HIV testing. However, decreased odds

230 of HIV testing uptake were found among men who were never married (AOR 0.51, 95% CI: 0.29–
231 0.88), lived in rural areas (AOR 0.54, 95% CI: 0.32–0.92), and consistently used condoms during sex
232 (AOR 0.59, 95% CI: 0.34–1.01).

233 **Discussion**

234 This study assessed the prevalence of HIV testing and determined important sociodemographic factors,
235 HIV-related knowledge, and sexual risk behaviors associated with HIV testing uptake among young
236 men aged 15–24 years using the PNGDHS data. Findings from this study revealed that only 17.1% of
237 young men in this study had ever been tested for HIV, which is lower than that of research in other
238 countries [37,39,40]. The differences could be attributed to differences in HIV knowledge and
239 awareness and the availability and accessibility of HIV testing services in these countries. There is
240 evidence that young men’s low uptake of HIV testing may be attributed to a lack of access to HIV
241 testing services and a lack of trust in these services [35,41]. In addition, fear of stigmatization, fear of
242 being diagnosed positive for HIV, the perceived risk of sexual exposure, poor attitudes by healthcare
243 professionals, and the perceived psychological burden have been found to impede HIV testing uptake
244 [35]. In PNG, HIV testing services have been more focused on the prevention of parent-to-child
245 transmission [42–44] and key populations (e.g., sex workers) [45,46], while such progress is yet to be
246 achieved with high-risk young populations. Given the importance of early HIV diagnosis and linking
247 access to treatment and care, age-specific policy development is required to increase HIV testing
248 uptake among this priority group.

249

Table 3. Multivariable analysis of predictors of HIV testing among young men (N = 1,275)

Characteristics	Ever been tested for HIV				p-value
	COR	95% CI	AOR	95% CI	
Age (years)					0.026*
15–19	Ref		Ref		
20–24	1.73	1.09–2.76	1.18	1.00–2.31	
Marital status					0.016*
Never married	0.59	0.39–0.89	0.51	0.29–0.88	
Married	Ref		Ref		
Educational level					0.749
No formal education	0.25	0.05–1.09	0.83	0.13–5.33	
Primary	0.53	0.18–1.58	1.23	0.34–4.49	
Secondary	0.98	0.31–3.09	1.55	0.42–5.69	
Tertiary	Ref		Ref		
Employment status					0.380
Not employed	0.66	0.45–0.97	0.78	0.45–1.36	
Employed	Ref		Ref		
Place of residence					0.023*
Urban	Ref		Ref		
Rural	0.52	0.35–0.77	0.54	0.32–0.92	
Region					0.313
Southern	0.87	0.52–1.46	1.19	1.61–2.32	
Highlands	1.15	0.71–1.86	1.15	0.60–2.18	
Momase	0.49	0.25–1.02	0.61	0.28–1.34	
Islands	Ref		Ref		
Read newspaper/magazine					0.437

No	Ref		Ref		
Yes	1.99	1.32–3.01	1.27	0.69–2.35	
Listen to the radio					0.349
No	Ref		Ref		
Yes	1.61	1.05–2.47	1.38	0.70–2.72	
Watch television					0.291
No	Ref		Ref		
Yes	1.34	0.86–2.09	0.69	0.36–1.36	
Internet access					0.250
No	Ref		Ref		
Yes	2.03	1.26–3.25	1.41	0.79–2.52	
Own a mobile phone					0.026*
No	Ref		Ref		
Yes	2.09	1.38–3.19	1.43	1.00–2.55	
A healthy person can have HIV					0.145
No	Ref		Ref		
Yes	1.56	0.75–3.22	1.95	0.79–4.77	
A person can get HIV by sharing food with an infected person					0.340
No	Ref		Ref		
Yes	0.76	0.42–1.38	0.69	0.33–1/47	
Consistent condom use during sex can reduce HIV risk					0.013*
No	Ref		Ref		
Yes	1.86	1.04–3.33	2.18	1.18–4.04	
Age at first sex					0.509
<15	Ref		Ref		
>15	0.77	0.51–1.17	0.84	0.49–1.43	
Ever paid for sex					0.036*

No	Ref		Ref		
Yes	3.56	1.85–6.84	1.75	1.01–5.83	
Number of sexual partners, including spouse (last 12 months)					0.046*
0–1	Ref		Ref		
2 or more	1.45	0.87–2.40	1.37	1.00–3.14	
Consistent condom use during sex					0.050*
No	Ref		Ref		
Yes	0.69	0.45–1.05	0.59	0.34–1.01	
Had any STIs (last 12 months)					0.639
No	Ref		Ref		
Yes	2.63	1.36–5.09	0.75	0.22–2.54	

* $p \leq 0.05$

COR, crude odds ratio. AOR, adjusted odds ratio. CI, confidence interval. Ref, reference category.

Note: Variables such as ever heard of AIDS and STIs had no responses for “ever tested for HIV” and were excluded in the final analysis.

250 Consistent with previous studies [35,36,47,48], this study indicated that demographic factors,
251 such as age, marital status, and place of residence, were significant predictors of HIV testing uptake.
252 HIV testing uptake tends to be significantly higher with increasing age, with men aged 20–24 years
253 being 1.18 times more likely to be tested than adolescent men (aged 15–19 years). A plausible
254 explanation could be that men aged 15–19 have a lower self-perceived likelihood of HIV compared to
255 those of increased age, who are more likely to be sexually active and knowledgeable about HIV and its
256 related risks [36,48,49]. It is also possible that they are more likely to have married and have more
257 lifetime exposure to HIV testing services, such as during antenatal care [36,50]. Health education and
258 awareness, including school-based interventions to increase HIV knowledge and risk perceptions, are
259 feasible options that could promote positive sexual risk behaviors and increase HIV testing uptake
260 among young men.

261 There was a negative association between HIV testing uptake and marital status. The odds of
262 HIV testing remained lower among unmarried young men than those who were married. The result
263 corroborated recent studies in Sub-Saharan Africa [36,51,52]. One possible explanation for this could
264 be that young men frequently encounter challenges accessing available HIV testing services, are
265 dissatisfied with the services provided, and have confidentiality issues [35,48]. Another plausible
266 reason could be that young men have a low-risk perception and are unaware of the physical signs of
267 HIV and its adverse consequences [53]. Evidence has shown that young men with low-risk perceptions
268 who do not have access to HIV testing services are less likely to have received an HIV test in their
269 lifetime [54,55]. Youth-friendly health services can help close the HIV testing gap among young men
270 by addressing access barriers, providing a more accessible, equitable, and effective social environment,
271 and linking to HIV treatment and care.

272 The results further revealed that the place of residence was negatively associated with HIV
273 testing. Young men who were from rural areas were less likely to be tested for HIV, consistent with

274 previous studies elsewhere [52,56]. The possible reason for this could be due to the availability and
275 accessibility of HIV testing services in urban areas rather than in rural areas, where young men would
276 have fewer opportunities to be tested for HIV. This disparity may also reflect a lack of HIV services, a
277 lack of trained HIV-trained health workers, and inequitable outreach programs to reach young men in
278 rural areas, as previously reported [57,58]. While prioritizing areas that require tailored and youth-
279 specific interventions to improve HIV testing access and uptake, there is a need to further expand HIV
280 services to rural areas. Mobile and rural health outreach programs are critical for reaching young men
281 lacking health care access and increasing HIV testing uptake, and should not be neglected.

282 In agreement with studies in Kenya and Ethiopia [59,60], the current study found that mobile
283 phone ownership among young men increases the uptake of HIV testing compared to non-mobile
284 owners. Mobile phone interventions in healthcare (mHealth) have shown success in improving access
285 to care and treatment, especially in settings where health disparities are more pronounced [61,62].
286 Evidence is emerging that the use of mobile technologies remains an effective strategy to reach young
287 people improve access to treatment and retention in care and provide information on HIV services and
288 prevention interventions, including risk reduction [59,63,64]. Furthermore, mobile phones have been
289 shown to address barriers of provider prejudice, stigmatization, discrimination, fear of refusal, lack of
290 privacy, and confidentiality [65,66]. Using mHealth technology provides a potentially promising
291 implementation strategy for interventions to remedy disparities and barriers to HIV testing and the care
292 continuum among young men. The study underscores the significance of preliminary, formative
293 research on population disparities in accessibility, effectiveness, willingness, and use of mobile phones
294 in HIV prevention programs.

295 In this study, young men who knew that consistent condom use during sex could reduce HIV
296 risk were 2.18 times more likely to be tested for HIV than their counterparts. The results were similar
297 to previous studies [67–69], indicating that having a better knowledge of condom efficacy and

298 effectiveness and sexual modes of HIV transmission were associated with condom use. This study also
299 established that young men who consistently use condoms during sex were less likely to be tested for
300 HIV. This is consistent with evidence suggesting that increased knowledge of HIV and risk perceptions
301 were associated with a positive attitude toward condom use [67,68]. It is also possible that discussion
302 of HIV with a sexual partner, knowing the partner's HIV status, and being in a steady relationship has
303 been shown to influence the consistent use of condoms [70,71]. Regular health education and
304 information dissemination on condom self-efficacy and risk reduction are crucial for preventing HIV
305 infection among this priority group.

306 Regarding sexual risk behaviors, young men who had paid for sex were more likely to be tested
307 for HIV than their counterparts. These findings are consistent with previous studies elsewhere [72,73],
308 which suggest that young men who engage in risky sexual behaviors such as sex work and
309 inconsistently use condoms appropriately perceive their risks of HIV acquisition and transmission and,
310 as a result, undertake HIV testing. The fact that they engage in such risky behaviors may imply that
311 their overall risk perception differs from that of those who do not engage in risky behaviors [74].
312 Strategies for young men who transact sex should not only concentrate on HIV testing and treatment
313 approaches but also on the social contexts in which they engage in transacting sex. Similarly, having
314 multiple sexual partners was significantly associated with HIV testing uptake. This finding
315 corroborates a study in Canada [75] that found an association between HIV testing and having
316 concurrent and multiple sexual partners among men. Having concurrent and multiple sexual partners
317 and risky sexual behaviors are predisposed to HIV and other STIs. Evidence shows that young people
318 who have multiple sex partners and are knowledgeable about HIV and its risks are more likely to
319 utilize HIV testing services [75,76]. The positive association between having multiple partners and the
320 uptake of HIV testing implies that there is a need for developing targeted programs for behavior change
321 and risk reduction for young people in this context.

322 Contrary to other studies [35,36], no association between the uptake of HIV testing and
323 educational level, employment status, wealth index, mass media exposure, age of sexual debut, or
324 having an STI was found in this study. However, this study has provided useful findings that have the
325 potential to inform the strengthening of current HIV prevention programs to target young men in PNG.
326 Improving HIV testing uptake among young people will play a vital role in reducing the risk of
327 acquiring HIV among young men, hence optimizing their health and well-being [53]. Targeted and
328 coordinated programs aimed at increasing HIV testing services among this priority group remain
329 imperative to achieve the national targets.

330 **Study strengths and limitations**

331 The study is based on nationwide population data with a nationally representative sample. Also,
332 appropriate estimation adjustments, such as weighting, were employed. However, the study has
333 limitations, and findings should be interpreted cautiously. Given that the data are from a cross-sectional
334 study, it is impossible to prove the temporal relationship. Since the study's outcome depended on self-
335 reporting, social desirability bias is highly probable, which might have led to under-reporting by the
336 participants. In addition, the dataset did not indicate whether young men were in same-sex relationships
337 (such as men who have sex with men) or not, which could have influenced the uptake of HIV testing.

338 **Conclusion**

339 Findings from this study showed that one in six young men had ever tested for HIV in PNG, which is
340 well below the national HIV/STI targets. Men aged 20–24 years who owned mobile phones, were
341 aware that consistent condom use reduces HIV risks, had ever paid for sex, and had multiple sexual
342 partners were significant predictors of HIV testing. To increase HIV testing uptake among young men,
343 it is crucial to implement targeted and comprehensive youth-friendly HIV/STI education and

344 sensitization programs and enable more accessible and affordable HIV testing services. Moreover,
345 concerted efforts for outreach and community-based testing programs, especially in rural and
346 prioritized areas where HIV testing services are inaccessible, are feasible options for young people in
347 PNG.

348

349 **Supporting information**

350 S1 Table. STROBE Checklist.

351 S1 Dataset. HIV testing-young men dataset.

352

353 **Acknowledgments**

354 The author would like to thank the DHS program and its partners for the approval to analyze the data.

355 Also, the author is greatly indebted to Dr. Fatch Welcome Kalembo from Curtin University, who
356 provided constructive feedback and suggestions to strengthen the quality of this manuscript.

357

358 **Author contributions**

359 Conceptualization: McKenzie Maviso

360 Data curation: McKenzie Maviso

361 Formal analysis: McKenzie Maviso

362 Methodology: McKenzie Maviso

363 Software: McKenzie Maviso

364 Validation: McKenzie Maviso

365 Writing – original draft: McKenzie Maviso

366 Writing – review & editing: McKenzie Maviso

367

368 **References**

- 369 1. Assefa Y, Gilks CF. Ending the epidemic of HIV/AIDS by 2030: Will there be an endgame to
370 HIV, or an endemic HIV requiring an integrated health systems response in many countries?
371 *International Journal of Infectious Diseases*. 2020;100: 273–277. doi:10.1016/j.ijid.2020.09.011
- 372 2. UNAIDS. The path that ends AIDS: UNAIDS Global AIDS Update 2023. Geneva, Switzerland;
373 2023. Available: <https://thepath.unaids.org/>
- 374 3. Frescura L, Godfrey-Faussett P, Feizzadeh A. A, El-Sadr W, Syarif O, Ghys PD, et al. Achieving
375 the 95 95 95 targets for all: A pathway to ending AIDS. *PLOS ONE*. 2022;17: e0272405.
376 doi:10.1371/journal.pone.0272405
- 377 4. UNAIDS. Understanding Fast-Track: Accelerating action to end the AIDS Epidemic by 2030.
378 Geneva, Switzerland; 2015. Available:
379 [https://www.unaids.org/sites/default/files/media_asset/201506_JC2743_Understanding_FastTrack](https://www.unaids.org/sites/default/files/media_asset/201506_JC2743_Understanding_FastTrack_en.pdf)
380 [_en.pdf](https://www.unaids.org/sites/default/files/media_asset/201506_JC2743_Understanding_FastTrack_en.pdf)
- 381 5. Delavande A, Kohler H-P. The Impact of HIV Testing on Subjective Expectations and Risky
382 Behavior in Malawi. *Demography*. 2012;49: 1011–1036. doi:[https://doi.org/10.1007/s13524-012-](https://doi.org/10.1007/s13524-012-0119-7)
383 [0119-7](https://doi.org/10.1007/s13524-012-0119-7)
- 384 6. Scott-Sheldon LA, Carey MP, Carey KB, Cain D, Simbayi LC, Mehlomakhulu V, et al. HIV
385 testing is associated with increased knowledge and reductions in sexual risk behaviours among
386 men in Cape Town, South Africa. *African Journal of AIDS Research*. 2013;12: 195–201.
387 doi:<https://doi.org/10.2989/16085906.2013.863219>
- 388 7. Fauk NK, Sukmawati AS, Berek PAL, Ernawati, Kristanti E, Wardoyo SSI, et al. Barriers to HIV
389 testing among male clients of female sex workers in Indonesia. *International Journal for Equity in*
390 *Health*. 2018;17: 68. doi:10.1186/s12939-018-0782-4
- 391 8. Hlongwa M, Mashamba-Thompson T, Makhunga S, Hlongwana K. Barriers to HIV testing uptake
392 among men in sub-Saharan Africa: a scoping review. *African Journal of AIDS Research*. 2020;19:
393 13–23. doi:10.2989/16085906.2020.1725071
- 394 9. Tetteh JK, Frimpong JB, Budu E, Adu C, Mohammed A, Ahinkorah BO, et al. Comprehensive
395 HIV/AIDS knowledge and HIV testing among men in sub-Saharan Africa: a multilevel
396 modelling. *Journal of Biosocial Science*. 2021/11/05 ed. 2022;54: 975–990.
397 doi:10.1017/S0021932021000560
- 398 10. Caltabiano M, Castiglioni M, De-Rose A. Changes in the sexual behaviour of young people:
399 introduction. *Genus*. 2020;76: 38. doi:10.1186/s41118-020-00107-1

- 400 11. Hittner JB, Owens EC, Swickert RJ. Influence of Social Settings on Risky Sexual Behavior.
401 SAGE Open. 2016;6: 2158244016629187. doi:10.1177/2158244016629187
- 402 12. Cho H-S, Yang Y. Relationship Between Alcohol Consumption and Risky Sexual Behaviors
403 Among Adolescents and Young Adults: A Meta-Analysis. International Journal of Public Health.
404 2023;68. doi:<https://doi.org/10.3389/ijph.2023.1605669>
- 405 13. Harris B, McCredie MN, Truong T, Regan T, Thompson CG, Leach W, et al. Relations Between
406 Adolescent Sensation Seeking and Risky Sexual Behaviors Across Sex, Race, and Age: A Meta-
407 Analysis. Archives of Sexual Behavior. 2023;52: 191–204. doi:<https://doi.org/10.1007/s10508-022-02384-7>
- 409 14. Manu A, Ogum-Alangea D, Azilaku JC, Anaba EA, Torpey K. Risky sexual behaviours and HIV
410 testing among young people in Ghana: evidence from the 2017/2018 Multiple Indicator Cluster
411 Survey. Reproductive Health. 2022;19: 125. doi:<https://doi.org/10.1186/s12978-022-01439-1>
- 412 15. Bolarinwa OA, Ajayi KV, Sah RK. Association between knowledge of Human Immunodeficiency
413 Virus transmission and consistent condom use among sexually active men in Nigeria: An analysis
414 of 2018 Nigeria Demographic Health Survey. Jilo GK, editor. PLOS Glob Public Health. 2022;2:
415 e0000223. doi:<https://dx.plos.org/10.1371/journal.pgph.0000223>
- 416 16. Wilcken A, Miiro-Nakayima F, Hizaamu RN, Keil T, Balaba-Byansi D. Male circumcision for
417 HIV prevention - a cross-sectional study on awareness among young people and adults in rural
418 Uganda. BMC Public Health. 2010;10: 209. doi:10.1186/1471-2458-10-209
- 419 17. Benders A, Nakamura K, Seino K, Al-Sobaihi S. Factors Associated with Low Uptake of Medical
420 Male Circumcision Among Adolescent Boys in Tanzania: A Multinomial Logistic Regression
421 Modeling. HIV/AIDS - Research and Palliative Care. 2022;14: 565–575.
422 doi:10.2147/HIV.S387380
- 423 18. Folayan MO, Sam-Agudu NA, Harrison A. Exploring the why: risk factors for HIV and barriers
424 to sexual and reproductive health service access among adolescents in Nigeria. BMC Health
425 Services Research. 2022;22: 1198. doi:10.1186/s12913-022-08551-9
- 426 19. Alexis Sentís, Mario Martin-Sanchez, Maider Arando, Martí Vall, María Jesus Barbera, Inma
427 Ocaña, et al. Sexually transmitted infections in young people and factors associated with HIV
428 coinfection: an observational study in a large city. BMJ Open. 2019;9: e027245.
429 doi:10.1136/bmjopen-2018-027245
- 430 20. Pettifor A, Bekker L-G, Hosek S, DiClemente R, Rosenberg M, Bull SS, et al. Preventing HIV
431 among young people: research priorities for the future. J Acquir Immune Defic Syndr. 2013;63
432 Suppl 2: S155–S160. doi:10.1097/QAI.0b013e31829871fb
- 433 21. Pettifor A, Stoner M, Pike C, Bekker L-G. Adolescent lives matter: preventing HIV in
434 adolescents. Curr Opin HIV AIDS. 2018;13: 265–273. doi:10.1097/COH.0000000000000453
- 435 22. Hakim AJ, Coy K, Badman SG, Willie B, Narokobi R, Gabuzzi J, et al. One size does not fit all:
436 HIV prevalence and correlates of risk for men who have sex with men, transgender women in

- 437 multiple cities in Papua New Guinea. *BMC Public Health*. 2019;19: 623.
438 doi:<https://doi.org/10.1186/s12889-019-6942-7>
- 439 23. Kelly-Hanku A, Willie B, Weikum D, Boli Neo R, Kupul M, Coy K. Kauntim mi tu: Multi-site
440 summary report from the key population integrated bio-behavioural survey, Papua New Guinea.
441 Goroka: Papua New Guinea Institute of Medical Research and Kirby Institute, UNSW Sydney.
442 2018. Available: [https://www.kirby.unsw.edu.au/research/reports/kauntim-mi-tu-port-moresby-](https://www.kirby.unsw.edu.au/research/reports/kauntim-mi-tu-port-moresby-2017)
443 2017
- 444 24. UNAIDS. Country facts - Papua New Guinea 2022. UNAIDS; 2022. Available:
445 <https://www.unaids.org/en/regionscountries/countries/papuanewguinea>
- 446 25. Grundy J, Dakulala P, Wai K, Maalsen A, Whittaker M. Independent State of Papua New Guinea
447 Health System Review. World Health Organization. Regional Office for South-East Asia; 2019.
448 Available: <https://iris.who.int/handle/10665/280088>
- 449 26. Maviso M, Kalembo FW. Prevalence and determinants of not testing for HIV among young adult
450 women in Papua New Guinea: findings from the Demographic and Health Survey, 2016–2018.
451 *BMJ Open*. 2024;14: e075424. doi:10.1136/bmjopen-2023-075424
- 452 27. Worth H. HIV Prevention in Papua New Guinea: Is It Working or Not? *World Journal of AIDS*.
453 2012;2: 117.
- 454 28. Baigry MI, Ray R, Lindsay D, Kelly-Hanku A, Redman-MacLaren M. Barriers and enablers to
455 young people accessing sexual and reproductive health services in Pacific Island Countries and
456 Territories: A scoping review. *PLOS ONE*. 2023;18: e0280667.
457 doi:<https://doi.org/10.1371/journal.pone.0280667>
- 458 29. Gray N, Azzopardi P, Kennedy E, Willersdorf E, Creati M. Improving Adolescent Reproductive
459 Health in Asia and the Pacific: Do We Have the Data? A Review of DHS and MICS Surveys in
460 Nine Countries. *Asia Pac J Public Health*. 2013;25: 134–144. doi:10.1177/1010539511417423
- 461 30. Kelly A, Worth H, Akuani F, Kepa B, Kupul M, Walizopa L, et al. Gendered talk about sex,
462 sexual relationships and HIV among young people in Papua New Guinea. *Culture, Health &*
463 *Sexuality*. 2010;12: 221–232. doi:<https://doi.org/10.1080/13691050903181107>
- 464 31. Kuzma J, Himata R. A pilot study of the knowledge, behaviour and attitudes connected with
465 HIV/AIDS among youth in PNG. *Contemporary PNG Studies*. 2005;3: 1–12.
- 466 32. Yakam LT. Factors associated with alcohol consumption among students in Divine Word
467 University, Madang Province, Papua New Guinea. *Contemporary PNG Studies*. 2021;34: 17–27.
- 468 33. National Statistical Office (NSO). National Population and Housing Census 2011. Port Moresby,
469 Papua New Guinea: National Statistical Office; 2015. Available: [https://png-](https://png-data.sprep.org/resource/papua-new-guinea-2011-national-report)
470 [data.sprep.org/resource/papua-new-guinea-2011-national-report](https://png-data.sprep.org/resource/papua-new-guinea-2011-national-report)
- 471 34. National Statistical Office (NSO) [Papua New Guinea], ICF. Papua New Guinea Demographic
472 and Health Survey 2016-18. Port Moresby, Papua New Guinea, and Rockville, Maryland, USA:
473 NSO and ICF; 2019. Available: <https://www.dhsprogram.com/pubs/pdf/FR364/FR364.pdf>

- 474 35. Ajayi AI, Awopegba OE, Adeagbo OA, Ushie BA. Low coverage of HIV testing among
475 adolescents and young adults in Nigeria: Implication for achieving the UNAIDS first 95. PLOS
476 ONE. 2020;15: e0233368. doi:<https://doi.org/10.1371/journal.pone.0233368>
- 477 36. Sonko I, Chung M-H, Hou W-H, Chen W-T, Chang P-C. Predictors of HIV testing among youth
478 aged 15–24 years in The Gambia. PLOS ONE. 2022;17: e0263720.
479 doi:<https://doi.org/10.1371/journal.pone.0263720>
- 480 37. Asaolu IO, Gunn JK, Center KE, Koss MP, Iwelunmor JI, Ehiri JE. Predictors of HIV Testing
481 among Youth in Sub-Saharan Africa: A Cross-Sectional Study. PLOS ONE. 2016;11: e0164052.
482 doi:<https://doi.org/10.1371/journal.pone.0164052>
- 483 38. Zhang Z. Model building strategy for logistic regression: purposeful selection. *Annals of*
484 *Translational Medicine*; Vol 4, No 6 (March 25, 2016): *Annals of Translational Medicine*
485 (Toward Precision Medicine in Neurological Diseases). 2016 [cited 31 Dec 2015].
486 doi:<http://dx.doi.org/10.21037/atm.2016.02.15>
- 487 39. Ijaiya MA, Anibi A, Abubakar MM, Obanubi C, Anjorin S, Uthman OA. A multilevel analysis of
488 the determinants of HIV testing among men in Sub-Saharan Africa: Evidence from Demographic
489 and Health Surveys across 10 African countries. PLOS Global Public Health. 2024;4: e0003159.
490 doi:[10.1371/journal.pgph.0003159](https://doi.org/10.1371/journal.pgph.0003159)
- 491 40. Olakunde BO, Adeyinka DA, Olawepo JO, Pharr JR. HIV testing among men in Nigeria: a
492 comparative analysis between young people and adults. *AIDS Care*. 2020;32: 155–162.
493 doi:[10.1080/09540121.2019.1622642](https://doi.org/10.1080/09540121.2019.1622642)
- 494 41. Aluzimbi G, Lubwama G, Muyonga M, Hladik W. HIV Testing and Risk Perceptions: A
495 Qualitative Analysis of Secondary School Students in Kampala, Uganda. *J Public Health Afr*.
496 2017;8: 577. doi:[10.4081/jphia.2017.577](https://doi.org/10.4081/jphia.2017.577)
- 497 42. Carmone A, Bomai K, Bongsi W, Frank TD, Dalepa H, Loifa B, et al. Partner testing, linkage to
498 care, and HIV-free survival in a program to prevent parent-to-child transmission of HIV in the
499 Highlands of Papua New Guinea. *Global Health Action*. 2014;7: 24995.
500 doi:<https://doi.org/10.3402/gha.v7.24995>
- 501 43. Badman SG, Valley LM, Toliman P, Kariwiga G, Lote B, Pomat W, et al. A novel point-of-care
502 testing strategy for sexually transmitted infections among pregnant women in high-burden
503 settings: results of a feasibility study in Papua New Guinea. *BMC Infectious Diseases*. 2016;16:
504 250. doi:<https://doi.org/10.1186/s12879-016-1573-4>
- 505 44. Saweri OP, Batura N, Pulford J, Khan MM, Hou X, Pomat WS, et al. Investigating health service
506 availability and readiness for antenatal testing and treatment for HIV and syphilis in Papua New
507 Guinea. *BMC Pregnancy and Childbirth*. 2022;22: 780. doi:<https://doi.org/10.1186/s12884-022-05097-w>
- 509 45. Hakim AJ, Badman SG, Weikum D, Amos A, Willie B, Narokobi R, et al. Considerable distance
510 to reach 90-90-90 targets among female sex workers, men who have sex with men and
511 transgender women in Port Moresby, Papua New Guinea: findings from a cross-sectional

- 512 respondent-driven sampling survey. *Sex Transm Infect.* 2020;96: 143.
513 doi:<http://dx.doi.org/10.1136/sextrans-2019-053961>
- 514 46. Mitchell E, Hakim A, Nosi S, Kupul M, Boli-Neo R, Aeno H, et al. A socio-ecological analysis of
515 factors influencing HIV treatment initiation and adherence among key populations in Papua New
516 Guinea. *BMC Public Health.* 2021;21: 2003. doi:<https://doi.org/10.1186/s12889-021-12077-w>
- 517 47. Ajiboye AS, Eshetu F, Lulseged S, Getaneh Y, Tademe N, Kifle T, et al. Predictors of HIV testing
518 among youth 15–24 years in urban Ethiopia, 2017–2018 Ethiopia population-based HIV impact
519 assessment. *PLOS ONE.* 2023;18: e0265710. doi:10.1371/journal.pone.0265710
- 520 48. Alem AZ, Liyew AM, Guadie HA. Spatial pattern and associated factors of HIV testing and
521 counselling among youths (15–24 years) in Ethiopia. *BMC Public Health.* 2021;21: 644.
522 doi:10.1186/s12889-021-10677-0
- 523 49. Wang Y, Kinsler JJ, Kiwuwa-Muyingo S. Factors associated with HIV testing among youth in
524 Tanzania based on the 2016–2017 Tanzania HIV Impact Survey (THIS). *PLOS Global Public
525 Health.* 2022;2: e0000536. doi:10.1371/journal.pgph.0000536
- 526 50. Kalibbala D, Mpungu SK, Ssuna B, Muzeyi W, Mberesero H, Semitala FC, et al. Determinants of
527 testing for HIV among young people in Uganda. A nested, explanatory-sequential study. *PLOS
528 Global Public Health.* 2022;2: e0000870. doi:10.1371/journal.pgph.0000870
- 529 51. Nshimirimana C, Vuylsteke B, Smekens T, Benova L. HIV testing uptake and determinants
530 among adolescents and young people in Burundi: a cross-sectional analysis of the Demographic
531 and Health Survey 2016–2017. *BMJ Open.* 2022;12: e064052.
532 doi:<http://dx.doi.org/10.1136/bmjopen-2022-064052>
- 533 52. Adugna DG, Worku MG. HIV testing and associated factors among men (15–64 years) in
534 Eastern Africa: a multilevel analysis using the recent demographic and health survey. *BMC Public
535 Health.* 2022;22: 2170. doi:10.1186/s12889-022-14588-6
- 536 53. Musonda E, Phiri M, Shasha L, Bwalya C, Musemangezhi S, Ishimwe SMC, et al. Prevalence of
537 HIV testing uptake among the never-married young men (15–24) in sub-Saharan Africa: An
538 analysis of demographic and health survey data (2015–2020). *PLOS ONE.* 2023;18: e0292182.
539 doi:10.1371/journal.pone.0292182
- 540 54. Jung MS, Dlamini NS, Cui X, Cha K. Prevalence of HIV testing and associated factors among
541 young adolescents in Eswatini: a secondary data analysis. *BMC Pediatrics.* 2022;22: 659.
542 doi:<https://doi.org/10.1186/s12887-022-03698-0>
- 543 55. Muravha T, Hoffmann CJ, Botha C, Maruma W, Charalambous S, Chetty-Makkan CM. Exploring
544 perceptions of low risk behaviour and drivers to test for HIV among South African youth. *PLOS
545 ONE.* 2021;16: e0245542. doi:10.1371/journal.pone.0245542
- 546 56. Mabaso M, Maseko G, Sewpaul R, Naidoo I, Jooste S, Takatshana S, et al. Trends and correlates
547 of HIV prevalence among adolescents in South Africa: evidence from the 2008, 2012 and 2017
548 South African National HIV Prevalence, Incidence and Behaviour surveys. *AIDS Research and
549 Therapy.* 2021;18: 97. doi:<https://doi.org/10.1186/s12981-021-00422-3>

- 550 57. Sanga ES, Mukumbang FC, Mushi AK, Lerebo W, Zarowsky C. Understanding factors
551 influencing linkage to HIV care in a rural setting, Mbeya, Tanzania: qualitative findings of a
552 mixed methods study. *BMC Public Health*. 2019;19: 383. doi:10.1186/s12889-019-6691-7
- 553 58. Pellowski JA. Barriers to care for rural people living with HIV: a review of domestic research and
554 health care models. *J Assoc Nurses AIDS Care*. 2013;24: 422–437.
555 doi:<https://doi.org/10.1016/j.jana.2012.08.007>
- 556 59. Jennings Mayo-Wilson L, Kang B-A, Mathai M, Mak'anyengo MO, Ssewamala FM. Mobile
557 phone access, willingness, and usage for HIV-related services among young adults living in
558 informal urban settlements in Kenya: A cross-sectional analysis. *International Journal of Medical
559 Informatics*. 2022;161: 104728. doi:<https://doi.org/10.1016/j.ijmedinf.2022.104728>
- 560 60. Nigatu M, Kabeta T, Taye A, Belina M. HIV voluntary counseling and testing uptake and
561 associated factors among Ethiopian youths: evidence from the 2016 EDHS using multilevel
562 modeling. *BMC Infectious Diseases*. 2021;21: 334. doi:10.1186/s12879-021-06021-x
- 563 61. Hall CS, Fottrell E, Wilkinson S, Byass P. Assessing the impact of mHealth interventions in low-
564 and middle-income countries – what has been shown to work? *Global Health Action*. 2014;7:
565 25606. doi:10.3402/gha.v7.25606
- 566 62. Nwaozuru U, Obiezu-Umeh C, Shato T, Uzoaru F, Mason S, Carter V, et al. Mobile health
567 interventions for HIV/STI prevention among youth in low- and middle-income countries
568 (LMICs): a systematic review of studies reporting implementation outcomes. *Implement Sci
569 Commun*. 2021;2: 126–126. doi:10.1186/s43058-021-00230-w
- 570 63. Garg PR, Uppal L, Mehra S, Mehra D. Mobile Health App for Self-Learning on HIV Prevention
571 Knowledge and Services Among a Young Indonesian Key Population: Cohort Study. *JMIR
572 Mhealth Uhealth*. 2020;8: e17646–e17646. doi:<https://doi.org/10.2196/17646>
- 573 64. Ippoliti NB, L'Engle K. Meet us on the phone: mobile phone programs for adolescent sexual and
574 reproductive health in low-to-middle income countries. *Reproductive Health*. 2017;14: 11.
575 doi:10.1186/s12978-016-0276-z
- 576 65. Mulawa MI, Rosengren AL, Amico KR, Hightow-Weidman LB, Muessig KE. mHealth to reduce
577 HIV-related stigma among youth in the United States: a scoping review. *Mhealth*. 2021;7: 35.
578 doi:<http://dx.doi.org/10.21037/mhealth-20-68>
- 579 66. Feroz AS, Ali NA, Khoja A, Asad A, Saleem S. Using mobile phones to improve young people
580 sexual and reproductive health in low and middle-income countries: a systematic review to
581 identify barriers, facilitators, and range of mHealth solutions. *Reprod Health*. 2021;18: 9–9.
582 doi:10.1186/s12978-020-01059-7
- 583 67. Hosseini Hooshyar S, Karamouzian M, Mirzazadeh A, Haghdoost AA, Sharifi H, Shokoohi M.
584 Condom Use and its Associated Factors Among Iranian Youth: Results From a Population-Based
585 Study. *Int J Health Policy Manag*. 2018;7: 1007–1014. doi:10.15171/ijhpm.2018.65

- 586 68. Yosef T, Nigussie T. Behavioral Profiles and Attitude toward Condom Use among College
587 Students in Southwest Ethiopia. *BioMed Research International*. 2020;2020: 1–6.
588 doi:10.1155/2020/9582139
- 589 69. Masoda M, Govender I. Knowledge and attitudes about and practices of condom use for reducing
590 HIV infection among Goma University students in the Democratic Republic of Congo. *Southern*
591 *African Journal of Epidemiology and Infection*. 2013;28: 61–68.
592 doi:10.1080/10158782.2013.11441521
- 593 70. Ajayi AI, Ismail KO, Akpan W. Factors associated with consistent condom use: a cross-sectional
594 survey of two Nigerian universities. *BMC Public Health*. 2019;19: 1207.
595 doi:<https://doi.org/10.1186/s12889-019-7543-1>
- 596 71. Ajayi OA, Ogunsola OO, Akinro Y, Adamu-Oyegun S, Wudiri K, Ojo TO, et al. Consistent
597 condom use and associated factors among HIV positive women of reproductive age on anti-
598 retroviral treatment in Ogun State, Nigeria. *Pan Afr Med J*. 2022;43: 101.
599 doi:10.11604/pamj.2022.43.101.32806
- 600 72. Hodgins C, Stannah J, Kuchukhidze S, Zembe L, Eaton JW, Boily M-C, et al. Population sizes,
601 HIV prevalence, and HIV prevention among men who paid for sex in sub-Saharan Africa (2000–
602 2020): A meta-analysis of 87 population-based surveys. *PLOS Medicine*. 2022;19: e1003861.
603 doi:10.1371/journal.pmed.1003861
- 604 73. Gebresilassie F, Ayele B, Hadgu T, Gebretnsae H, Negash D, Demoz Ghebremdhin K, et al.
605 Predictors of Condom Use Among Youth of the Rural Tigray, Northern Ethiopia: Community-
606 Based Cross-Sectional Study. *HIV/AIDS - Research and Palliative Care*. 2023;15: 377–385.
607 doi:<https://doi.org/10.2147/HIV.S412337>
- 608 74. Dizechi S, Brody C, Tuot S, Chhea C, Saphonn V, Yung K, et al. Youth paying for sex: what are
609 the associated factors? Findings from a cross-sectional study in Cambodia. *BMC Public Health*.
610 2018;18: 113. doi:<https://doi.org/10.1186/s12889-017-4999-8>
- 611 75. Luginaah NA, Konkor I, Lawson ES, Mkandawire P, Husbands W, Omorodion F, et al.
612 Concurrent sexual partnerships and HIV testing among heterosexual Black men in Ontario,
613 Canada: findings from the weSpeak study. *Ethn Health*. 2022;27: 1825–1840.
614 doi:10.1080/13557858.2021.1976395
- 615 76. Wilson Chialepeh N, Sathiyasusuman A. Associated Risk Factors of STIs and Multiple Sexual
616 Relationships among Youths in Malawi. *PLOS ONE*. 2015;10: e0134286.
617 doi:10.1371/journal.pone.0134286

618
619

Ever tested for HIV among young men in PNG

