

1 **Gut microbiome predicts atopic diseases in an infant cohort with reduced bacterial**
2 **exposure due to social distancing**

3

4 Katri Korpela^{a1,2}, Sadhbh Hurley^{a3,4}, Sinead Ahearn Ford², Ruth Franklin³, Susan Byrne^{3,4},
5 Nonhlanhla Lunjani², Brian Forde^{2,5}, Ujjwal Neogi⁶, Carina Venter⁷, Jens Walter^{2,5,8},
6 Jonathan Hourihane^{b3,4}, Liam O'Mahony^{b2,5,8}

7 CORAL Study Group: Naomi MCCallion RCSI and Rotunda Hospital Dublin, Aideen Byrne
8 CHI and TCD, Dublin, Martin White Coombe Hopsital and RCSI, John Fitzsimons CHI and
9 RCSI

10 ^aBoth share first authorship

11 ^bBoth share senior authorship

12

13 ¹Human Microbiome Research Program, Faculty of Medicine, University of Helsinki,
14 Helsinki, Finland

15 ²APC Microbiome Ireland, University College Cork, Cork, Ireland

16 ³Paediatrics and Child Health, Royal College of Surgeons in Ireland, Dublin, Ireland.

17 ⁴Children's Health Ireland, Dublin, Ireland.

18 ⁵School of Microbiology, University College Cork, Cork, Ireland

19 ⁶The Systems Virology Lab, Division of Clinical Microbiology, Department of Laboratory
20 Medicine, Karolinska Institute, ANA Futura, Campus Flemingsberg, Stockholm, Sweden.

21 ⁷Section of Allergy & Immunology, Department of Pediatrics, Children's Hospital Colorado,
22 University of Colorado School of Medicine, Aurora, Colorado; Children's Hospital
23 Colorado, Aurora, Colorado, USA.

24 ⁸Dept. of Medicine, University College Cork, Cork, Ireland

25

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

26 **Abstract**

27 Several hypotheses link altered microbial exposure in affluent societies to increased
28 prevalence of allergies, but none have been experimentally tested in humans. Here we
29 capitalize on the opportunity to study a cohort of infants (CORAL) raised during COVID-19
30 associated social distancing measures to test the interactions between bacterial exposure
31 and fecal microbiome composition with atopic outcomes. We show that fecal *Clostridia*
32 levels were significantly lower in CORAL infants and correlated with a microbial exposure
33 index. Microbiota composition was the most significant component of regression models
34 predicting risk of atopic dermatitis (AUC 0.86) or food allergen sensitization (AUC 0.98)
35 and mediated the effects of multiple environment factors on disease risk. Although diet
36 had a larger effect on microbiota composition than environmental factors linked to
37 dispersal, most effects were mediated through the microbiota. This study provides critical
38 information to refine existing hypothesis on the importance of the gut microbiota to immune
39 development.

40 **Introduction**

41 Human mucosal surfaces and body cavities harbor diverse communities of
42 commensal microbes that play essential roles in regulation of host metabolic responses,
43 epithelial barrier function, immune education, and immune regulation (1-6). Humans
44 evolved in an environmental and social context that enabled reliable transmission and
45 dispersal of human adapted symbionts, accompanied by diets rich in substrates that
46 provide nutritional support for the gut microbiota (7,8). Hallmarks of these evolutionary
47 relationships can be observed in the microbiomes of non-industrialized human populations
48 and non-human primates, whose microbiomes are shaped by dispersal and social
49 behavior (9-11). Given that the gut microbiota provides essential, and evolution routed
50 benefits to human development, alterations in microbial transmission through lifestyle
51 factors linked to industrialization and western dietary habits provide a possible mechanism
52 for the rise in chronic, non-communicable diseases in affluent societies.

53 There is an “immunological window of opportunity” early in life when the immune
54 system is particularly responsive to environmental exposures (including infections,
55 nutrition, and microbiome) that help establish the thresholds, patterns of reactivity, and
56 functional trajectories that can have long term consequences including altered risk of
57 immune mediated diseases (12-15). The immune response to commensal microbes is not
58 simply a form of host defense but represents an intimate and sophisticated bidirectional
59 dialogue that ensures a stable microenvironment is maintained with important symbiotic
60 physiological effects on the host. These microbe-host interactions promote appropriate
61 immune responses that effectively defend against infection, with limited collateral damage
62 to host tissue, and without any subsequent aberrant inflammatory or allergic reactions (16-
63 20).

64 Epidemiological factors linking immune-mediated diseases with living environment,
65 lifestyle and social interactions (e.g. household size, absence of older siblings, antibiotics,
66 mode of delivery, breast feeding rates, rural versus urban living, dietary intake, etc.) are
67 also known to impact early-life gut microbiota assembly (21-24). These environmental
68 factors alter ecological processes that shape community assembly, such as dispersal
69 (household size, siblings, c-sections) and selection (antibiotics, breast feeding, diet).
70 Metacommunity theory predicts that the composition and dynamics within any local
71 community (e.g. within a human individual) are governed both by processes that occur
72 within the community (dietary resources, antibiotics) and by the process of dispersal
73 (siblings, hygiene, family size) which links communities together (25-27). These theoretical
74 frameworks led us to hypothesize that allergies, and the increase in allergy with
75 industrialization, arise from modern lifestyle-factors disrupting human gut metacommunity
76 composition and function.

77 The importance of environmental factors that influence metacommunity dynamics
78 through dispersal (family size, siblings, social interactions) can usually only be examined in
79 observational studies. However, the enforced social distancing measures that were
80 employed during sustained periods of mass lockdowns to prevent spread of the SARS-
81 CoV-2 virus provided a real-world unique opportunity to experimentally test the
82 metacommunity hypothesis of atopic disease. The CORAL (Impact of **COR**ona Virus
83 Pandemic on **A**llergic and Autoimmune Dysregulation in Infants Born During Lockdown)
84 study examined effects of the social isolation restrictions imposed during the early stages
85 of the COVID-19 pandemic on infants that spent most of their early life in social isolation.
86 Compared with a historical Irish cohort, infants in the CORAL cohort had higher rates of
87 atopic dermatitis (25% versus 17%) and food allergen sensitization (7% versus 4%) at 12
88 months of age (28). Furthermore, some deficits in social communication were evident at

89 their 12-month assessment (29). Here we extend this work, hypothesizing that
90 environmental factors that alter metacommunity processes affected microbiome
91 composition, and that these interactions might mediate allergic outcomes, which are some
92 of the earliest manifestations of inappropriate immune reactivity. To test this hypothesis,
93 we performed 16S rRNA tag sequencing on fecal DNA and applied an ecological modeling
94 approach to determine the relative importance of the environmental drivers of bacterial
95 community assembly. We then determined to what degree interactions between bacterial
96 taxa and dietary factors could predict levels of atopic dermatitis and food allergen
97 sensitization in this population.

98 **Results**

99 *CORAL infant cohort characteristics*

100 CORAL demographic details have been previously published (28, 30). In summary
101 55% were male, 96% Caucasian, 46% first born infants, 38% second born or later infants,
102 65% were born by spontaneous vaginal delivery. Despite multiple waves of COVID-19 in
103 the community during this time, social isolation was effective in minimizing exposure to
104 SARS-CoV-2 as only 3% of infants had a detectable antibody response to the virus at 12
105 months of age. In addition, only 17% of infants required antibiotics by 12 months
106 (compared to 80% in a pre-pandemic UK birth cohort) further indicating reduced exposure
107 to infectious agents (31).

108 Families of infants in the CORAL cohort had a median of one social contact outside
109 the home at birth, increasing to four when the infants reached 6 months of age. The
110 median number of people who had kissed the infant in the first 6 months was 3 (including
111 parents). In addition, social isolation restrictions led to 25% of infants not having met a
112 child their own age by their first birthday (32).

113

114 *Fecal bacterial community composition*

115 We sequenced fecal samples from all infants for which background questionnaire
116 data was available at 6 (n=351) and 12 months (n=343), using 16S rRNA tag sequencing.
117 Consistent with previous studies, infant microbiota composition was strongly associated
118 with age and birth mode. Age explained 26% of the variation in composition between
119 samples (PERMANOVA, p=0.001) and birth mode explained 3% at 6 months
120 (PERMANOVA, p=0.001) and 1.5% at 12 months (p=0.004) (**Supplementary Fig 1a**). Age
121 was significantly associated with the relative abundances of 29 (33%) of the 89 detected
122 bacterial families (FDR-corrected p<0.05). At the order level, *Bifidobacteriales*,


123 *Lactobacillales*, *Enterobacteriales*, and *Veillonellales* decreased from 6 to 12 months of
124 age, while *Verrucomicrobiales*, *Erysipelotrichales*, *Clostridiales*, *Bacteroidales*, and
125 *Coriobacteriales* increased from 6 to 12 months of age (**Supplementary Fig 1b**).

126 Also consistent with previous studies, C-section (CS) delivery or exposure to
127 intrapartum antibiotics during vaginal birth (VAB) were associated with increased relative
128 abundance of the genera *Sarcina* (*Clostridiaceae*), *Flavonifractor* (*Ruminococcaceae*), and
129 *Erysipelatoclostridium* (*Erysipelatoclostridiaceae*) and reduced relative abundance of
130 *Collinsella* (*Coriobacteriaceae*) at 6 months of age. In addition, CS delivery was associated
131 with increased relative abundance of *Enterobacter* (*Enterobacteriaceae*), *Amedibacterium*
132 (*Erysipelotrichaceae*), *Blautia*, and *Roseburia* (*Lachnospiraceae*), and reduced relative
133 abundance of *Bacteroides* (*Bacteroidaceae*) and *Parabacteroides* (*Porphyromonadaceae*)
134 (FDR-corrected $p < 0.05$). At 12 months, most birth-mode related differences were no
135 longer significant, except for *Bacteroides* that was still significantly reduced in the CS-
136 delivered infants, and *Faecalibacillus* (*Erysipelotrichaceae*) in the VAB group (FDR-
137 corrected $p < 0.05$).

138

139 *CORAL microbiota comparison with historical cohorts*

140 We compared the microbiota composition in the CORAL cohort to the general
141 pattern of microbiota development observed across multiple previously published cohorts,
142 including data from 34 studies, 3825 infants, and 5732 fecal samples (33). Similar time-
143 dependent abundance of *Bacilli*, *Proteobacteria* and *Bacteroidia* were observed in the
144 CORAL and previous cohorts (**Fig. 1**). However, CORAL infants had significantly lower
145 levels of *Clostridia* at both 6 months and 12 months of age ($p < 0.001$). In addition, the
146 relative abundance of *Bifidobacterium* (class *Actinomycetia*) was elevated at 12 months in
147 comparison to other cohorts ($p < 0.0001$).



148

149 **Fig 1.** Comparison to the general pattern of microbiota development in other cohorts. The
 150 relative abundances of the most abundant bacterial taxa (phylum Proteobacteria, classes
 151 *Bacilli*, *Bacteroidia*, *Clostridia*, and family *Bifidobacteriaceae*). The trend lines are calculated
 152 from a third-degree polynomial regression model weighted by the sample size of each
 153 study. The boxplots represent the relative abundances in the CORAL cohort. Significance
 154 of the differences is marked with asterisks (***) $p < 0.001$.


155

156 We also compared CORAL microbiota data with a recently published Irish infant
 157 cohort (INFANTMET) that was studied prior to the pandemic (34, 35). *Clostridia* were also
 158 significantly less abundant in the CORAL cohort compared to the INFANTMET cohort at
 159 both 6 ($p < 0.001$) and 12 ($p = 0.02$) months of age, while *Bifidobacteria* were more abundant
 160 in the CORAL cohort at 12 months of age, ($p < 0.0001$) after adjusting for birth mode,
 161 breastfeeding, and antibiotic use. The *Clostridia* pattern was mainly driven by the reduced
 162 relative abundance of the genera *Sarcina* (Clostridiaceae, $p < 0.0001$) and *Eubacterium*
 163 (Eubacteriaceae, $p < 0.0001$) at 6 months, *Enterocloster* (*Ruminococcaceae*, $p < 0.001$) and
 164 *Hungatella* (Lachnospiraceae, $p < 0.0001$) at both time points, and additionally family
 165 *Ruminococcaceae* at 12 months ($p < 0.001$) and the genus *Flavonifractor*
 166 (*Ruminococcaceae*, $p < 0.0001$) at 12 months. At 6 months, the CORAL cohort harbored a

167 higher microbial richness than the pre-pandemic cohort ($p < 0.0001$), but the pattern had
168 reversed by 12 months due to the lower rate of increase in richness in CORAL ($p < 0.0001$).

169 In order to determine if reduced abundance of *Clostridia* could be related to
170 differences in microbial exposures, we examined correlations with a microbial exposure
171 index generated by combining answers to questionnaire data relating to potential sources
172 of microbial exposures (i.e. having a family member classified as an essential worker who
173 was allowed to work outside the home, number of siblings, number of adults in the
174 household, attending daycare outside of home, having pets, and living in a house as
175 compared to an apartment). The relative abundance of *Clostridia* was associated with the
176 exposure index ($p = 0.039$) at 6 months of age (**Fig. 2**), supporting the hypothesis that
177 reduced exposure to external sources of microbes was linked with the lower-than-
178 expected relative abundance of *Clostridia*. At the genus level, all *Clostridial* genera
179 showed a positive trend with external exposure, which was statistically significant ($p < 0.05$)
180 in 19/38 (50%) of the genera detected at 6 months of age (**Fig. 2**). At 6 months, the
181 exposure index was also positively associated with microbial richness ($p = 0.01$). The
182 recorded exposures were not significantly associated with *Clostridia* ($p = 0.73$) or richness
183 ($p = 0.08$) by 12 months of age.

184



185

186 **Fig. 2.** Association of the class Clostridia (first panel), and Clostridial genera with external
 187 exposure index. Trendlines, 95% confidence intervals, and p-values are calculated based
 188 on a third-degree polynomial regression model. Significant associations are highlighted
 189 with pink.

190

191 *Relative contribution of environmental factors on microbiome development in the CORAL*
 192 *cohort*


193

194 We analyzed the contribution of multiple potential contributing factors on the gut
 195 microbiota composition at 6 and 12 months, using the envfit function in R package vegan.
 196 At both time points, diet was the main determinant of microbiota composition, exceeding
 197 even the effect of birth mode (**Fig. 3a**). The measured variables explained 32% and 31%
 198 of the variation in microbiota composition at 6 and 12 months, respectively. The impact of
 medical interventions, primary caesarean delivery, and intrapartum antibiotics were higher

214 *Bacteroides* (**Fig. 4c and Fig. 4d**). Cow's milk formula consumption, on the other hand,
215 had the opposite effect to breastfeeding and vaginal birth at 12 months, shifting the
216 microbiota away from the *Bifidobacterium/Bacteroides*-dominated composition.

217 The impact of factors linked to human exposure dramatically increased from 6 to 12
218 months with effect of the number of siblings increasing from 1% to 9% of the variation.
219 Pets and siblings at 6 and 12 months, respectively, shifted the microbiota from the infant-
220 type *Bacilli-Enterobacteria*-composition towards a more mature composition with increased
221 relative abundance of members of *Ruminococcaceae* and *Lachnospiraceae* (**Fig. 4e**).
222 Pets was the strongest environmental factor at 6 months, while at 12 months, the living
223 environment (rural versus urban, house versus apartment) became more significant, with
224 exposure to a rural living environment shifting the microbiota towards increased relative
225 abundance of the *Clostridial* families *Ruminococcaceae*, *Lachnospiraceae*, and
226 *Peptostreptococcaceae* (**Fig. 4f**).

227



228

229 **Fig. 4.** Association between bacterial genera and food categories (a,b), most significant
230 variables (c,d), and external exposures (e,f) at 6 months (a,c,e) and 12 months (b,d,f).
231 Heat maps (a,b,e,f) represent the correlations between log-transformed relative
232 abundances of bacteria, coloured by class, and dietary/exposure variables. Panels (c,d)
233 show the significant associations ($p < 0.05$) of exposure variables with the first two principal
234 coordinates as well as the bacterial genera most strongly associated with the coordinates,
235 with samples coloured by birth mode.

236

237 *Microbiome mediates the effects of environmental factors on atopic dermatitis and food*
238 *allergen sensitization*


239 Virtually all the environmental factors identified above as predictors of microbiome
240 variation (diet, breastfeeding, siblings, pets, etc.) have also been described as
241 epidemiological factors that influence atopic dermatitis and food allergen sensitization. To
242 determine the extent to which the microbiota mediates the effects of epidemiological
243 factors, we compared the estimated impact of exposures on disease outcomes with and
244 without adjusting for gut microbiota composition using logistic regression models, adjusting
245 for parental allergic diseases and ethnicity.

246 The atopic dermatitis prediction model based purely on the environmental factors
247 without gut microbiota data achieved an AUC of 0.82 (95% confidence interval 0.76-0.88).
248 We then selected the most significantly associated microbial genera measured at 6 and 12
249 months and added those into the model. Adding the gut microbiota data as additional
250 explanatory variables in the model increased the AUC to 0.87 (95% CI 0.82-0.92)
251 (**Supplementary Fig 2**). The gut microbiota composition at both 6 months and 12 months
252 was strongly predictive of atopic dermatitis diagnosis at 12 months (**Fig. 5a**). The atopic
253 dermatitis-associated microbes, apart from *Akkermansia* (*Akkermansiaceae*) and

254 *Amedibacterium* (*Erysipelotrichaceae*), were mostly Clostridia, with all but *Sarcina*
255 (*Clostridiaceae*) at 6 months having a positive association with atopic dermatitis. In
256 particular *Enterocloster* (*Lachnospiraceae*), was associated at both timepoints with
257 increased risk of atopic dermatitis. Regarding exposures, atopic dermatitis was positively
258 associated with attending daycare, and smoking in the household, and negatively
259 associated with living in a rural area and with antibiotics use between 6 and 12 months of
260 age. The effect of smoking was mainly direct, rather than microbiota mediated, since
261 including the gut microbiota in the model did not reduce the estimated effect size.
262 Interestingly, our analysis revealed that rural living had a protective direct effect on atopic
263 dermatitis, but also a predisposing microbiota-mediated effect through increased relative
264 abundances of *Intestinibacter* and *Eisenbergiella* (**Fig. 5b**). The effects of daycare and
265 antibiotics were microbiota mediated, evidenced by the decrease in effect size when
266 microbiota was included in the model. Attending daycare was associated with reduced
267 relative abundance of *Akkermansia*, and antibiotics reduced the relative abundances of
268 *Intestinibacter* and *Enterocloster* (**Fig. 5b**).

269 In terms of diet, atopic dermatitis was positively associated with the frequent infant
270 consumption of starchy vegetables, fish, peanuts and pistachios, and negatively with the
271 consumption of grain, pine nuts, and kimchi (**Fig. 5a**). Many foods had both direct and
272 microbiota-mediated effects on atopic dermatitis risk, with the direct and microbiota-
273 mediated effects partly countering each other (**Fig. 5b**). Starchy vegetables and peanuts
274 had a direct predisposing association, but they were also associated with reduced
275 abundances of atopic dermatitis-associated microbes, such as *Akkermansia*. In contrast,
276 the protective effects of kimchi and pine nuts were almost fully explained by their negative
277 impact on *Akkermansia*, *Intestinibacter*, *Ruthenibacterium*, and *Enterocloster*.
278 Breastfeeding, siblings, birth mode and antibiotics in the first 6 months did not have a

279 direct association with atopic dermatitis but appeared to modulate risk through their effects
 280 on the microbiota.
 281



282
 283 **Fig. 5.** Predictors of atopic dermatitis (a-b) and food sensitization (c-d) based on logistic
 284 regression models with and without microbiota. In panels a and c, negative deviance
 285 indicates a negative association. Panels b and d show the direct and microbiota-mediated
 286 associations between exposures and health outcomes at 1 year. The width of the
 287 connecting lines indicates strength of the association (deviance explained).
 288

289 The AUC for the food allergen sensitization model without gut microbiota was 0.88
 290 (95% CI 0.82-0.94) and adding the gut microbiota increased the AUC to 0.98 (95% CI

291 0.97-1.00) (**Supplementary Fig 3**). Gut microbiota composition at 6 months emerged as
292 the strongest predictor of food allergen sensitization at 12 months of age, and it appeared
293 to both mediate the effects of several dietary components and exposures, and act as an
294 important confounder for the effects of others (**Fig. 5c**). Food sensitization was primarily
295 associated with the 6-month gut microbiota composition, the predictive microbes being
296 members of Clostridia except for *Bacillus* (Bacilli, *Bacillaceae*) and *Odoribacter*
297 (*Bacteroidia*, *Odoribacteraceae*). Regarding exposures, having more than 2 adults in the
298 household, or an essential worker, or a pet (adult/pet exposure) had a protective
299 association with food sensitization that was mostly mediated through decreased relative
300 abundances of *Bacillus* and *Peptoniphilus* (**Fig. 5d**). Maternal smoking had a direct
301 influence on food sensitization and an influence mediated by *Peptoniphilus*, while passive
302 smoking only had a direct predisposing effect. In addition, having siblings and C-section
303 birth were associated with reduced relative abundances of food sensitization-associated
304 microbes, while antibiotics in the first 6 months had the opposite effect.

305 Of the dietary components, tempeh had a direct positive association and egg had a
306 direct negative association with food sensitization (**Fig. 5c-d**). Cheese, orange/red
307 vegetables, and walnuts had a direct positive association, as well as microbiota-mediated
308 effects that were partly protective. The predisposing effect of beef was fully explained by
309 its positive impact on *Peptoniphilus*. Grains and sesame had a direct negative association
310 with food sensitization as well as microbiota-mediated protective effects through reduced
311 relative abundances of *Monoglobus*, *Oscillibacter*, and *Bacillus*.

312

313 Discussion

314 In this study, we establish that social isolation due to the COVID-19 pandemic
315 altered the early life assembly of the gut microbiota in infants, which was associated with
316 risk of atopic diseases. We discovered that the primary factors that impact early-life
317 microbiome assembly include specific dietary factors and exposures to other humans.
318 Furthermore, we established that known epidemiological links between breast feeding,
319 siblings, antibiotics, and diet with risk of atopy are partly mediated via the microbiome.
320 Overall, our findings provide direct support of the 'metacommunity hypothesis' of allergic
321 disease whereby lifestyle factors affecting ecological processes that shape the human gut
322 metacommunity at local and wider scales predispose humans to immunological deficits
323 that contribute to allergy risk.

324 Neonatal gut microbiota is shaped by maternal gut microbes obtained at birth, but
325 the relative contributions of the initial vertical colonization and later horizontal exposures to
326 microbiota maturation in infancy has not previously been well established (36,37). Despite
327 the strict social distancing measures, which were largely complied with in Ireland, the
328 overall gut microbiota composition and development followed closely the expected natural
329 pattern, indicating that the most important source of gut microbes in infants is the mother
330 and other family members (38). During the first months of life, the infant microbiota is
331 dominated by strains from *Actinobacteria* and *Bacteroidia*, which are maternally derived in
332 vaginally born neonates (36). In contrast, maternal strains of *Clostridia* were never
333 observed in neonates suggesting that *Clostridia* are transmitted from environmental
334 sources, perhaps related to lack of persistence previously described for sporulating
335 microbes (39). This supports our findings that environmentally derived taxa, mainly
336 *Clostridia*, are reduced in socially isolated infants, while levels of maternally derived taxa
337 are consistent with historical cohorts. Some species from the *Clostridia* family are well

338 recognized as important immunomodulatory microbes, and altered *Clostridia* levels were
339 previously observed early in life in allergic children (40-43). Our results show that some
340 members of *Clostridia* that were reduced in the CORAL cohort, were associated with
341 increased risk of atopic outcomes, suggesting potentially protective effects of delayed
342 colonization. This highlights the need to better define the optimal timings for microbiota
343 maturation during infancy.

344 Higher *Bifidobacterium* levels at 12 months might be related to a reduction in
345 antibiotic use and a high breastfeeding rate in this cohort. Antibiotic use in infancy and
346 later childhood causes a dramatic, long-term reduction in bifidobacteria (44,45). The
347 relative importance of undisturbed gut microbiota development in the first year of life
348 versus delayed *Clostridia* colonization on the long-term health of infants is an important
349 remaining question with implications for infant care practices and decisions, such as entry
350 to daycare and early social contacts. There is likely an optimal age when the benefits of
351 increasing social contacts begin to outweigh the risks of infection and consequent
352 antibiotic exposure. The effect may also vary between outcomes: early-life antibiotic use
353 was associated with a microbiota-mediated risk of obesity in later childhood (45), which is
354 perhaps reduced in the socially distancing cohort, while the risk of immune-related
355 diseases may be increased.

356 Following acquisition of potential colonizers, our results suggest that the impact of
357 diet on infant gut microbiota development clearly outweighs the effects of social contacts
358 and environment. During breastfeeding, human milk oligosaccharides support specialized
359 microbes, such as bifidobacteria, that protect against infection and support immune
360 development via multiple mechanisms including secretion of metabolites (e.g. SCFAs,
361 indole-3-lactic acid) (46). The introduction of complementary foods leads to a rapid
362 expansion in microbiota diversity and a shift to taxa that can utilize dietary fibers instead of

363 human milk oligosaccharides, accompanied by changes in production of known
364 immunomodulatory microbial metabolites, such as butyrate (47-49). In the CORAL cohort,
365 plant based dietary sources (including legumes, tree nuts and sesame seeds) had the
366 most significant effects on microbiota composition at 6 and 12 months. Of note, the early
367 introduction of potentially allergenic plants foods such as peanut and tree nuts is
368 suggested to protect against food allergy development (50,51). Our study suggests that
369 these foods are utilized by the early life microbiota, which may play a role in their allergy
370 protective effects complementing mechanisms associated with direct sampling by the
371 immune system. It was remarkable to what degree our findings aligned with what is
372 currently considered healthy eating, and foods that support healthy host-microbe
373 interactions (52). Specifically, plant-based foods in general had similar effects as
374 breastfeeding on microbiota composition, which was not observed for animal-based foods
375 within this age group, potentially due to fibers present within the plant-based foods (52,53).
376 This supports current practice to encourage plant based dietary sources during weaning.
377 The effect sizes of dietary components and other measured exposures were greater than
378 that expected from previous studies (54,55), perhaps due to the reduced overall
379 complexity of exposures contributing to the CORAL infant's microbiota thereby allowing
380 more accurate identification of effects on specific taxa. In addition, the greatly reduced use
381 of antibiotics in this cohort removed an important confounding factor to gut microbiota
382 development.

383 Pets and the presence of essential workers who could leave the family home
384 increased the relative abundance of spore-forming, environmentally transmitted bacteria.
385 House type and living in a rural environment also contributed to microbiota composition at
386 12 months of age. While exposure to biodiverse environments is important to support
387 assembly and maturation of a diverse microbiome, human-adapted symbionts might only

388 be acquired from contact with other humans. Indeed, while the effect of siblings on allergic
389 disease risk were gut microbiota mediated, the protective effect of living in a rural
390 environment was not, indicating that exposure to green environments exerts its protective
391 effects through other mechanisms, but has limited benefits in terms of gut microbiota
392 development.

393 Social distancing measures designed to limit spread of SARS-CoV-2 were
394 undoubtedly instrumental in preventing excessive morbidity and mortality in Ireland prior to
395 the availability of vaccines. However, it is important to understand more fully how these
396 practices affected the microbiome, and then, in response, to develop public measures and
397 practices that can, if appropriate, increase exposure to beneficial microbes and
398 simultaneously reduce risk of transmission of infectious agents. This cohort of infants born
399 during the pandemic also allowed us to assess the impact of dietary, environmental, and
400 social exposures on development of early life gut metacommunity processes over the first
401 year of life, and their association with immune-mediated disorders such as atopic
402 dermatitis and food allergen sensitization. Whether these differences will be short lived,
403 with rapid recovery post-lockdown, or whether there will be longer term effects remains to
404 be seen.

405 **Methods**

406 *Study Outline*

407 The CORAL (Impact of **C**orona Virus Pandemic on **A**llergic and Autoimmune
408 Dysregulation in Infants Born During Lockdown) Study is a longitudinal prospective
409 observational study of allergy, immune function and neurodevelopment in a population of
410 term Irish infants born during the first 3 months of the COVID-19 pandemic, March–May
411 2020. A total of 3773 infants were born in the two participating major maternity hospitals in
412 Dublin from March to May 2020. Invitations were sent to the families of 3065 term babies
413 who were eligible for inclusion. Exclusion criteria were pre-birth PCR-proven SARS-CoV-2
414 infection in a parent or co-dwelling person, intravenous antibiotics in the neonatal period,
415 multiple births or major congenital anomaly. A total of 351 infants were recruited
416 postnatally to the CORAL Study and mothers did not need to be first-time parents. Babies
417 had a clinical review along with lateral flow COVID-19 antibody screening at 6 and
418 12 months of age. Detailed epidemiological information was collected at recruitment and at
419 each 6-month review, with a dietary questionnaire performed at 12 months of age.

420 At recruitment and at each 6-month review, families completed an online questionnaire
421 which collected evolving epidemiological information under three main categories. Firstly,
422 the infant's home environment including household number, presence of siblings, urban or
423 rural location, type of dwelling, childcare outside the home, household pets, and maternal
424 and household smoking status. Secondly, infant health and healthcare utilisation including
425 infant weaning, infant illnesses, antibiotic use, SARS-CoV-2 contacts, testing and positivity,
426 infant hospitalisation, regular medication use and vaccination status. Lastly information was
427 collected regarding symptoms and signs of allergic conditions including suspected adverse
428 food reactions, atopic dermatitis and allergic airways disease.

429 At 12 month clinical review infants were assessed for signs of atopic dermatitis and
430 severity scoring was completed if present. Skin prick testing was completed to the 3 most
431 common food allergens (peanut, egg and milk) and aeroallergens (house dust mite, grass
432 pollen and cat dander) in all infants.

433 The food frequency questionnaire measured frequency of intake of baby foods, fruits,
434 vegetables, fermented foods, pulses/legumes, meat, poultry, grains, dairy, eggs, fish and
435 seafoods, peanuts, tree nuts and seeds, soy, sugary and fatty foods. We also enquired about
436 intake of multivitamins, prebiotics and probiotics. Frequencies included: Avoid, Never, Given
437 and then stopped, Rarely: less than once a month, at least once a month, at least once a
438 week, once a day, more than once a day.

439 This study involves human participants and ethical permission was granted by the
440 National COVID-19 Ethics Committee (20-NREC-COV-067). Parent/guardian gave
441 informed consent to participate in the study before taking part. Ireland spent the majority of
442 period March 2020–April 2021 in maximal (level 5; no home visitors, stay at home except
443 for essential purposes, exercise within 2–5 km of home, essential retail only, unable to sit
444 in restaurants/cafes, work from home unless essential worker) lockdown and so babies
445 spent most of the first year of their lives in social isolation.

446

447 *Microbiota sampling and sequencing*

448 Fecal samples were collected at home using a stool sampling kit with preservative
449 (Norgen Biotek), at 6 and 12 months of age. Stabilized samples were returned via post
450 and stored at -80 C until analysis. Microbial DNA was isolated using the DNeasy
451 PowerSoil Pro Kit (Qiagen) and MiSeq sequencing of the v3 16S rRNA generated
452 approximately 20,000 reads per sample (University of Minnesota Genomics Center).

453

454 *Bioinformatic processing*

455 The forward and reverse reads were merged, and the merged reads were quality
456 and chimera filtered using vsearch version 2.14.2 (56). Unique reads occurring only once
457 in the total dataset were removed. Each read was annotated against the RDP 11.5
458 database using the syntax command in vsearch with 90 % confidence cut-off. The
459 annotated reads were summarized at different taxonomic levels.

460

461 *Statistical analysis*

462 The statistical analyses were conducted in R. Principal coordinates analysis
463 (function capscale in R package vegan) using Pearson correlation distances was used to
464 depict the overall gut microbiota composition and its determinants. Functions envfit and
465 adonis in package vegan was used to assess the importance of the background variables
466 on gut microbiota composition. Heat maps were created using the heatmap.2 function in
467 package gplots. Negative binomial models (function glm.nb in package MASS) with the
468 sample-specific read count as an offset were used to identify the factors associated with
469 individual bacterial taxa (57). Logistic regression was used to assess the effects of gut
470 microbiota and diet on the risk of atopic dermatitis and food sensitization. Area under the
471 curve (AUC) was used to assess the predictive ability of the logistic regression models,
472 using the function roc in package pROC (58).

473 **Acknowledgements**

474 *Ethical approval:* National Research Ethics Committee for COVID-19-related Research
475 (NREC COVID-19) Approval number 20-NREC-COV-067.

476

477 *Funding:* This work was supported by Royal College of Surgeons in Ireland, Temple
478 Street Hospital Foundation in Dublin, Ireland (Approval number RPAC 20-02), and the
479 Clemens Von Pirquet Foundation in Geneva, Switzerland. The authors are supported by a
480 Science Foundation Ireland research center grant 12/RC/2273_P2. We would like to thank
481 Dr Yvonne d'Art for initial conceptualization and co-design of the study and Prof. Catherine
482 Stanton and Dr. Kiera Healy for sharing INFANTMET 16S rRNA sequencing data.

483

484 *Disclosures:* C. Venter reports grants from Reckitt Benckiser, Food Allergy Research and
485 Education, and National Peanut Board, and personal fees from Reckitt Benckiser, Nestle
486 Nutrition Institute, Danone, Abbott Nutrition, Else Nutrition, and Before Brands. L.
487 O'Mahony reports consultancy with Precision BioticsAlimentary Health, grants from
488 GlaxoSmithKline and Chiesi, and participation in speaker bureau for Nestle, Yakult, Reckitt
489 and Abbott. J. Hourihane is a board member of Clemens Von Pirquet Foundation. No
490 other conflicts were declared for the remaining authors.

491 **References**

- 492 1. Valdes AM, Walter J, Segal E, Spector TD. Role of the gut microbiota in nutrition and
493 health. *BMJ*. 2018 Jun 13;361:k2179.
- 494 2. Lunjani N, Ahearn-Ford S, Dube FS, Hlela C, O'Mahony L. Mechanisms of microbe-
495 immune system dialogue within the skin. *Genes Immun* 2021 Oct;22(5-6):276-288.
- 496 3. Akdis CA. Does the epithelial barrier hypothesis explain the increase in allergy,
497 autoimmunity and other chronic conditions? *Nat Rev Immunol*. 2021 Apr 12.
- 498 4. Lunjani N, Walsh LJ, Venter C, Power M, MacSharry J, Murphy DM, O'Mahony L.
499 Environmental influences on childhood asthma - The effect of diet and microbiome on
500 asthma. *Pediatr Allergy Immunol*. 2022 Dec; e13892.
- 501 5. Wilmes P, Martin-Gallausiaux C, Ostaszewski M, Aho VTE, Novikova PV, Laczny CC,
502 Schneider JG. The gut microbiome molecular complex in human health and disease. *Cell*
503 *Host Microbe*. 2022 Sep 14;30(9):1201-1206.
- 504 6. Forde B, Yao L, Shaha R, Murphy S, Lunjani N, O'Mahony L. Immunomodulation by
505 foods and microbes: Unravelling the molecular tango. *Allergy*. 2022 Dec;77(12):3513-
506 3526.
- 507 7. Sonnenburg ED, Sonnenburg JL. Starving our microbial self: the deleterious
508 consequences of a diet deficient in microbiota-accessible carbohydrates. *Cell Metab*. 2014
509 Nov 4;20(5):779-786.
- 510 8. Walter J, O'Mahony L. The importance of social networks-An ecological and
511 evolutionary framework to explain the role of microbes in the aetiology of allergy and
512 asthma. *Allergy* 2019; 74(11):2248-2251.
- 513 9. Johnson Katerina V-A, Watson Karli K, Dunbar Robin IM, Burnet Philip WJ. Sociability
514 in a non-captive macaque population is associated with beneficial gut bacteria. *Frontiers in*
515 *Microbiology* 2022; 13.
- 516 10. Martínez I, Stegen JC, Maldonado-Gómez MX, Eren AM, Siba PM, Greenhill AR,
517 Walter J. The gut microbiota of rural papua new guineans: composition, diversity patterns,
518 and ecological processes. *Cell Rep* 2015;11:527-538.
- 519 11. Moeller AH, Foerster S, Wilson ML, Pusey AE, Hahn BH, Ochman H. Social behavior
520 shapes the chimpanzee pan-microbiome. *Sci Adv* 2016;2:e1500997.
- 521 12. Fujimura KE, Sitarik AR, Havstad S, Lin DL, Levan S, Fadrosch D, et al. 2016. Neonatal
522 gut microbiota associates with childhood multisensitized atopy and T cell differentiation.
523 *Nat Med* 22: 1187–1191.

- 524 13. Arrieta M-C, Stiemsma LT, Dimitriu PA, Thorson L, Russell S, Yurist-Doutsch S, et al.
525 2015. Early infancy microbial and metabolic alterations affect risk of childhood asthma. *Sci*
526 *Transl Med* 7: 307ra152.
- 527 14. Olin A, Acevedo N, Lakshmikanth T, Chen Y, Johansson C, Alm J, Scheynius A,
528 Brodin P. Longitudinal analyses of development of the immune system during the first five
529 years of life in relation to lifestyle. *Allergy*. 2022 May;77(5):1583-1595.
- 530 15. Lunjani N, Tan G, Dreher A, Sokolowska M, Groeger D, Warwyzniak M, et al.
531 Environment-dependent alterations of immune mediators in urban and rural South African
532 children with atopic dermatitis. *Allergy*. 2022 Feb;77(2):569-581.
- 533 16. Brodin P. Immune-microbe interactions early in life: A determinant of health and
534 disease long term. *Science*. 2022 May 27;376(6596):945-950.
- 535 17. Groeger D, Schiavi E, Grant R, Kurnik-Łucka M, Michalovich D, Williamson R, et al.
536 Intranasal *Bifidobacterium longum* protects against viral-induced lung inflammation and
537 injury in a murine model of lethal influenza infection. *EBioMedicine*. 2020 Oct;60:102981.
- 538 18. Schiavi E, Gleinser M, Molloy E, Groeger D, Frei R, Ferstl R, et al. The Surface-
539 Associated Exopolysaccharide of *Bifidobacterium longum* 35624 Plays an Essential Role
540 in Dampening Host Proinflammatory Responses and Repressing Local TH17 Responses.
541 *Appl Environ Microbiol*. 2016 Nov 21;82(24):7185-7196.
- 542 19. Konieczna P, Groeger D, Ziegler M, Frei R, Ferstl R, Shanahan F, et al.
543 *Bifidobacterium infantis* 35624 administration induces Foxp3 T regulatory cells in human
544 peripheral blood: potential role for myeloid and plasmacytoid dendritic cells. *Gut*. 2012
545 Mar;61(3):354-66.
- 546 20. Lyons A, O'Mahony D, O'Brien F, MacSharry J, Sheil B, Ceddia M, et al. Bacterial
547 strain-specific induction of Foxp3+ T regulatory cells is protective in murine allergy models.
548 *Clin Exp Allergy*. 2010 May;40(5):811-9.
- 549 21. Strachan DP. Hay fever, hygiene, and household size. *BMJ*. 1989 Nov 18;
550 299(6710):1259-60.
- 551 22. von Mutius E, Martinez FD, Fritzsche C, Nicolai T, Reitmeir P, Thiemann HH. Skin test
552 reactivity and number of siblings. *BMJ*. 1994 Mar 12;308(6930):692-5.
- 553 23. Lunjani N, Satitsuksanoa P, Lukasik Z, Sokolowska M, Eiwegger T, O'Mahony L.
554 Recent developments and highlights in mechanisms of allergic diseases: Microbiome.
555 *Allergy*. 2018 Dec;73(12):2314-2327.
- 556 24. Huang YJ, Marsland BJ, Bunyavanich S, O'Mahony L, Leung DY, Muraro A, Fleisher
557 TA. The microbiome in allergic disease: Current understanding and future opportunities-

- 558 2017 PRACTALL document of the American Academy of Allergy, Asthma & Immunology
559 and the European Academy of Allergy and Clinical Immunology. *J Allergy Clin Immunol.*
560 2017 Apr;139(4):1099-1110.
- 561 25. Nemergut DR, Schmidt SK, Fukami T, O'Neill SP, Bilinski TM, Stanish LF, et al.
562 Patterns and processes of microbial community assembly. *Microbiol Mol Biol Rev.* 2013
563 Sep;77(3):342-56.
- 564 26. Kennedy KM, de Goffau MC, Perez-Muñoz ME, Arrieta MC, Bäckhed F, Bork P, et al.
565 Questioning the fetal microbiome illustrates pitfalls of low-biomass microbial studies.
566 *Nature.* 2023 Jan;613(7945):639-649.
- 567 27. Miller ET, Svanbäck R, Bohannan BJM. Microbiomes as Metacommunities:
568 Understanding Host-Associated Microbes through Metacommunity Ecology. *Trends Ecol*
569 *Evol.* 2018 Dec;33(12):926-935.
- 570 28. Hurley S, Franklin R, McCallion N, et al. Allergy-related outcomes at 12 months in the
571 CORAL birth cohort of Irish children born during the first COVID 19 lockdown. *Pediatr*
572 *Allergy Immunol.* 2022;33:e13766.
- 573 29. Byrne S, Sledge H, Franklin R, Boland F, Murray DM, Hourihane [JO'B](#); CORAL Study
574 group. Social communication skill attainment in babies born during the COVID-19
575 pandemic: a birth cohort study. *Arch Dis Child.* 2023 Jan;108(1):20-24.
- 576 30. Lawler M, Franklin R, McCallion N, et al.. The impact of COVID-19 lockdown on
577 infants' coronavirus exposure and routine healthcare access in Ireland: The CORAL birth
578 cohort study at 6 months. *Pediatr Allergy Immunol* 2021;32:1876–9.
- 579 31. Harris JM, Mills P, White C, Moffat S, Newman Taylor AJ, Cullinan P. Recorded
580 infections and antibiotics in early life: associations with allergy in UK children and their
581 parents. *Thorax.* 2007;62(7):631-637.
- 582 32. Sledge H, Lawler M, Hourihane [JO'B](#), et al.. Parenting a newborn baby during the
583 COVID-19 pandemic: a qualitative survey. *BMJ Paediatr Open* 2022;6:e001348.
- 584 33. Korpela K, de Vos WM. Early life colonization of the human gut: microbes matter
585 everywhere. *Curr Opin Microbiol.* 2018 Aug;44:70-78.
- 586 34. Fouhy F, Watkins C, Hill CJ, O'Shea CA, Nagle B, Dempsey EM, et al. Perinatal
587 factors affect the gut microbiota up to four years after birth. *Nat Commun.* 2019 Apr
588 3;10(1):1517.
- 589 35. Hill CJ, Lynch DB, Murphy K, Ulaszewska M, Jeffery IB, O'Shea CA, et al. Evolution of
590 gut microbiota composition from birth to 24 weeks in the INFANTMET Cohort. *Microbiome.*
591 2017 Jan 17;5(1):4.

- 592 36. Korpela K, Costea P, Coelho LP, Kandels-Lewis S, Willemsen G, Boomsma DI,
593 Segata N, Bork P. Selective maternal seeding and environment shape the human gut
594 microbiome. *Genome Res.* 2018 Apr;28(4):561-568.
- 595 37. Korpela K, Helve O, Kolho KL, Saisto T, Skogberg K, Dikareva E, et al. Maternal Fecal
596 Microbiota Transplantation in Cesarean-Born Infants Rapidly Restores Normal Gut
597 Microbial Development: A Proof-of-Concept Study. *Cell.* 2020 Oct 15;183(2):324-334.e5.
- 598 38. Valles-Colomer M, Blanco-Míguez A, Manghi P, Asnicar F, Dubois L, Golzato D, et al.
599 The person-to-person transmission landscape of the gut and oral microbiomes. *Nature.*
600 2023 Feb;614(7946):125-135.
- 601 39. Hildebrand F, Gossmann TI, Frioux C, Özkurt E, Myers PN, Ferretti P, et al. Dispersal
602 strategies shape persistence and evolution of human gut bacteria. *Cell Host Microbe.*
603 2021 Jul 14;29(7):1167-1176.e9.
- 604 40. Atarashi K, Tanoue T, Oshima K, Suda W, Nagano Y, Nishikawa H, et al. Treg
605 induction by a rationally selected mixture of Clostridia strains from the human microbiota.
606 *Nature.* 2013 Aug 8;500(7461):232-6. doi: 10.1038/nature12331.
- 607 41. Savage JH, Lee-Sarwar KA, Sordillo J, Bunyavanich S, Zhou Y, O'Connor G, et al. A
608 prospective microbiome-wide association study of food sensitization and food allergy in
609 early childhood. *Allergy.* 2018 Jan;73(1):145-152. doi: 10.1111/all.13232.
- 610 42. Bunyavanich S, Shen N, Grishin A, Wood R, Burks W, Dawson P, et al. Early-life gut
611 microbiome composition and milk allergy resolution. *J Allergy Clin Immunol.* 2016
612 Oct;138(4):1122-1130. doi: 10.1016/j.jaci.2016.03.041.
- 613 43. Feehley T, Plunkett CH, Bao R, Choi Hong SM, Culleen E, Belda-Ferre P, et al.
614 Healthy infants harbor intestinal bacteria that protect against food allergy. *Nat Med.* 2019
615 Mar;25(3):448-453.
- 616 44. Korpela K, Salonen A, Virta LJ, Kekkonen RA, Forslund K, Bork P, de Vos WM.
617 Intestinal microbiome is related to lifetime antibiotic use in Finnish pre-school children. *Nat*
618 *Commun.* 2016 Jan 26;7:10410.
- 619 45. Korpela K, Salonen A, Virta LJ, Kekkonen RA, de Vos WM. Association of Early-Life
620 Antibiotic Use and Protective Effects of Breastfeeding: Role of the Intestinal Microbiota.
621 *JAMA Pediatr.* 2016 Aug 1;170(8):750-7.
- 622 46. Henrick BM, Rodriguez L, Lakshmikanth T, Pou C, Henckel E, Arzoomand A, et al.
623 Bifidobacteria-mediated immune system imprinting early in life. *Cell.* 2021 Jul
624 22;184(15):3884-3898.e11.

- 625 47. Laursen MF, Andersen LB, Michaelsen KF, Mølgaard C, Trolle E, Bahl MI, Licht TR.
626 Infant Gut Microbiota Development Is Driven by Transition to Family Foods Independent of
627 Maternal Obesity. *mSphere*. 2016 Feb 10;1(1):e00069-15.
- 628 48. Tsukuda, N., Yahagi, K., Hara, T. et al. Key bacterial taxa and metabolic pathways
629 affecting gut short-chain fatty acid profiles in early life. *ISME J* 15, 2574–2590 (2021).
- 630 49. Roduit C, Frei R, Ferstl R, Loeliger S, Westermann P, Rhyner C, et al. High levels of
631 butyrate and propionate in early life are associated with protection against atopy. *Allergy*.
632 2019 Apr;74(4):799-809.
- 633 50. Logan K, Bahnson HT, Ylescupidez A, Beyer K, Bellach J, Campbell DE, et al. Early
634 introduction of peanut reduces peanut allergy across risk groups in pooled and causal
635 inference analyses. *Allergy*. 2022 Nov 27. doi: 10.1111/all.15597.
- 636 51. de Silva D, Halken S, Singh C, Muraro A, Angier E, Arasi S, et al. Preventing food
637 allergy in infancy and childhood: Systematic review of randomised controlled trials. *Pediatr*
638 *Allergy Immunol*. 2020 Oct;31(7):813-826. doi: 10.1111/pai.13273.
- 639 52. Armet AM, Deehan EC, O'Sullivan AF, Mota JF, Field CJ, Prado CM, Lucey AJ, Walter
640 J. Rethinking healthy eating in light of the gut microbiome. *Cell Host Microbe*. 2022 Jun
641 8;30(6):764-785.
- 642 53. Venter C, Meyer RW, Greenhawt M, Pali-Schöll I, Nwaru B, Roduit C, et al. Role of
643 dietary fiber in promoting immune health-An EAACI position paper. *Allergy*. 2022
644 Nov;77(11):3185-3198.
- 645 54. Stewart CJ, Ajami NJ, O'Brien JL, Hutchinson DS, Smith DP, Wong MC, et al.
646 Temporal development of the gut microbiome in early childhood from the TEDDY study.
647 *Nature*. 2018 Oct;562(7728):583-588.
- 648 55. Beller L, Deboutte W, Falony G, Vieira-Silva S, Tito RY, Valles-Colomer M, et al.
649 Successional Stages in Infant Gut Microbiota Maturation. *mBio*. 2021 Dec
650 21;12(6):e0185721.
- 651 56. Rognes T, Flouri T, Nichols B, Quince C, Mahé F. VSEARCH: a versatile open source
652 tool for metagenomics. *PeerJ*. 2016 Oct 18;4:e2584.
- 653 57. Venables, W. N. & Ripley, B. D. (2002) *Modern Applied Statistics with S*. Fourth
654 Edition. Springer, New York. ISBN 0-387-95457-0.
- 655 58. Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez JC, Müller M. pROC: an
656 open-source package for R and S+ to analyze and compare ROC curves. *BMC*
657 *Bioinformatics*. 2011 Mar 17;12:77.