# Study of the Effect of Intestinal Microbes on Obesity: A Bibliometric Analysis

## 1 Zehao Su<sup>1,2</sup>, Chenyu Tian<sup>1,2</sup>, Guan Wang<sup>1\*</sup>, Jingjing Guo<sup>1,2\*</sup>, Xiaoyan Yang<sup>1,2\*</sup>

- <sup>2</sup> <sup>1</sup> Department of Endocrinology & Metabolism and West China Biomedical Big Data Center, West
- 3 China Hospital, Sichuan University/West China School of Nursing, Chengdu 610041, Sichuan,
- 4 China.
- <sup>5</sup> <sup>2</sup>Med-X Center for Informatics, Sichuan University, Chengdu 610041, China.

# Keywords: obesity<sub>1</sub>, intestinal microbiota<sub>2</sub>, bibliometric analysis<sub>3</sub>, visualization<sub>4</sub>, co-occurrence analysis<sub>5</sub>.

8 Abstract

9 Background: Obesity is a serious public health problem. According to statistics, there are millions obese people worldwide. The intestinal microbiota is the microbiota that lives in the gastrointestinal 10 11 tract of the human body and can interact with the external environment through diet. Probing and 12 summarizing the relationship between intestinal microbes and obesity has important guiding significance for the accurate control of the research direction and expanding the choice of obesity 13 14 treatment methods. We aimed to use bibliometrics to qualitatively and quantitatively analyze the 15 published literature and to reveal the research hotspots and development trends in the effects of intestinal microbes on obesity research. 16

Methods: We obtained documents from the core collection of Web of Science (WoSCC) from 2013– 2022 on February 2nd, 2023. Microsoft Excel, Biblioshiny R language software packages and CiteSpace were employed to collect publication data, analyze publication trends, and visualize relevant results.

21 **Results:** We identified 8888 original research articles on the effects of intestinal microbes on obesity published between 2013 and 2022. China made the highest number of publications on the topic, with 22 23 3036 publications (34.16%), and the United States with 87065 citations made the greatest contribution. 24 University of Copenhagen was the most prolific institution (209 publications). Patrice D. Cani 25 published the most related articles (97 publications), whereas Peter J. Turnbaugh was cited the most frequently (3307 citations). Nutrients was the journal with the most studies (399 publications, 4.49%), 26 27 and Nature was the most commonly co-cited journal (6446 citations). The burst keywords "obese 28 patient" and "serum" recently appeared as research frontiers.

Conclusion: Although China has the largest number of publications, the United States, depending on its largest citations, has become the leader in the effects of intestinal microbes on obesity research. Current research hotspots focus on related mechanisms of the effects of intestinal microbes on obesity and therapeutic methods for obesity with intestinal microbes. Through diet, probiotic preparations and regulation of the intestinal flora, fecal microbe transplantation may have a significant impact on reducing the obesity epidemic. Considering the great potential and application prospects, intestinal microbe applications in obesity therapy will remain a research hotspot in the future.

#### 36 1 Introduction

37 Obesity is a serious public health problem. According to statistics, there are approximately 600 38 million obese people and 1.9 billion overweight individuals worldwide (El-Sayed Moustafa and 39 Froguel, 2013). It is predicted that the number of obese people in the world will reach 1.12 billion by 40 2030 (KellyYang et al., 2008). As far as we know, obesity is not only a change in appearance but also 41 associated with disorders of lipid and glucose metabolism, chronic inflammation, oxidative stress and 42 increased risk of a variety of diseases. According to research, obesity is one of the nonnegligible 43 reasons for cardiovascular disease, diabetes and malignant tumors (BoccellinoDi Domenico et al., 2018; 44 Scherer and Hill, 2016), and it is estimated that more than 80% of the global burden of diabetes and 45 cardiovascular disease will occur in developing countries such as China and India in 2025 46 (MohanGupta et al., 2016). However, the etiology of obesity is not yet completely clear. The increasing 47 prevalence of obesity will not be explained by changes in energy balance alone. Therefore, it is vital 48 for the whole world to determine the formation mechanism of obesity and then explore effective 49 methods to treat it and reduce the incidence and mortality of obesity-related diseases.

50 The intestinal microbiota is a microbiota that lives in the gastrointestinal tract of the human body 51 and can interact with the external environment through diet (AlessandroMarcello et al., 2017). These 52 microbes colonize the intestinal tract through mother-to-child transmission from the prenatal period. 53 The colonization of human intestinal microbes continues after birth and is regulated by factors 54 including gestational age, mode of delivery (natural delivery or cesarean section), diet (breastfeeding 55 or baby food), hygiene and antibiotics. The environment and diet of the first three years of life are 56 essential for the acquisition of adult-like microbiota and the establishment of bacteria-host symbiosis 57 that affect immune and nervous system development. At the age from 2 to 5, the intestinal microbiota will gradually stabilize, and there is no significant difference with intestinal microbiota in adults 58 59 (RodríguezMurphy et al., 2015). Under normal circumstances, the intestinal microbiota maintains a 60 symbiotic relationship with the human body, and a good symbiotic relationship between them is very 61 important for the maintenance of human health (JandhyalaTalukdar et al., 2015), but instantaneous 62 changes in the intestinal ecosystem can lead to the destruction of the symbiotic relationship between 63 microorganisms and hosts, such as aging, obesity, sedentary lifestyles, and dietary patterns, and the 64 effects of antibiotics can change the intestinal microbiota (NazliYang et al., 2004). Because of the 65 important role of the intestinal ecosystem in maintaining host physiology, its changes can also cause a 66 variety of physiological disorders in the human body, including low-grade inflammation, metabolic 67 disorders, excessive lipid accumulation and loss of insulin sensitivity, in turn leading to changes in the composition of the intestinal microbiota and decreasing the diversity of flora and metabolic pathways 68 69 (Halmos and Suba, 2016; Sidhu and van der Poorten, 2017), thus increasing the risk of metabolic 70 diseases such as obesity and diabetes (TangKitai et al., 2017). Probing and summarizing the 71 relationship between intestinal microbes and obesity has important guiding significance for the 72 accurate control of the research direction and expanding the choice of obesity treatment methods.

73 Bibliometrics is a discipline that uses mathematics and statistics to quantitatively analyze literature 74 and information. Through quantitative analysis of the subject literature, we can construct its knowledge 75 structure and explore its development trend. At present, it has been widely used in many research fields 76 (GaoDou et al., 2021; LiuGao et al., 2022; WangMa et al., 2020). Overall, this analysis method and 77 the traditional literature review are both based on previous research, summarize the current situation 78 and shortcomings of the research, and finally guide further research. Literature review emphasizes the 79 content of the article, that is, to summarize the aspects and shortcomings of the existing research, and 80 most of them quote the representative papers in the literature according to a preset research context.

81 On the other hand, bibliometric analysis places more emphasis on the analysis of "quantity". There is

82 no need to analyze the research content of each document in detail but to analyze the number of 83 published documents, the distribution of authors and the relationship of citations in the target field.

84 Most of the included documents are highly quoted. Compared with the traditional literature review,

85 bibliometric analysis can make a more intuitive systematic analysis of all the literature in this field in

86 a visual way, which is helpful for researchers entering a new field to grasp the overall trend of the field.

87 There have been many review articles in the field of intestinal microbes and obesity, but the quantitative

analysis in the field is insufficient. Therefore, in addition to briefly summarizing the hotpots in review 88

89 articles, we use bibliometrics to qualitatively and quantitatively analyze the published literature and

90 intends to reveal the research hotspots and development trends in this field.

#### 91 2 Materials and methods

#### 92 2.1 Data sources and search strategies

Data analysis and graph acquisition

93 Data were obtained from the core collection of Web of Science (WoSCC), a database of Clarivate 94 Analytics. The following search strategies were presented: ((overweight OR obesity OR obes\* OR 95 over\$weight) AND (gastrointestin\* OR gastro-intestin\* OR gut OR intestin\*) AND (bacteria OR 96 prebiotic OR probiotic OR microbiot\* OR microbiome\* OR flora OR microflora)). In the end, a total 97 of 8888 original articles in English were retrieved. All the documents were downloaded as "Full Record and Cited References" and saved as "Bibtex" and "plain text file". The above data were obtained on 98 99 2nd February 2023.

# 100

2.2

#### 101 The corresponding files retrieved from WoSCC were imported into Microsoft Excel (16.66.1), 102 Biblioshiny and CiteSpace (6.1. R6) to perform a bibliometric and visual analysis. Annual publications, 103 country annual publications, and publisher annual publications were analyzed with Microsoft Excel. 104 The Biblioshiny platform provides a web-based graphical interface in the Bibliometrix package, which 105 was used to conduct total citation, average article cited, publications of journal and country 106 collaboration map. CiteSpace is interactive visualization analysis software developed by Professor 107 Chen Chaomei to conduct different types of network analyses, such as country collaboration networks, 108 institution collaboration networks, author collaboration networks, co-cited author collaboration 109 networks, co-cited journal collaboration networks, and keyword co-occurrence, which aid in visually 110 analyzing the knowledge domain and emerging trends. A dual-map overlay was constructed for 111 journals. The parameters of CiteSpace were as follows: time slicing (2013 to 2022), years per slice (1), 112 links (strength: cosine, scope: within slices), selection criteria (g-index: k=5 in country, institution, 113 author collaboration network analysis and keyword co-occurrence), pruning (pathfinder, pruning sliced 114 networks). Cluster labels used keywords by log likelihood ratio (LLR). To identify emerging topics,

115 we detected keywords and references with a strong citation burst.

116 Ethical approval was waived by the local ethical board because the data came from public databases, 117 and no human or animal subjects were involved.

#### 118 3 Results

#### 119 3.1 An overview of the annual growth trend

120 As shown in Figure 1A, from 2013 to 2022, a total of 8888 English-language articles met our 121 inclusion criteria. The number of annual publications began to exceed 500 in 2016, and only after 3

122 years did the annual publications exceed 1000. The annual number of publications increased from 267 123 in 2013 to 1768 in 2022, with an amazing increase of over sixfold. We further analyzed the annual 124 national output of the 10 most productive countries and regions (Figure 1B). The different colors represent different countries, and the slope represents the speed of annual publication number increase. 125 126 China ranked first in the number of annual publications, followed by the United States and Spain. The 127 top 10 publishers according to their contribution to the total number of articles about the effect of 128 intestinal microbes on obesity are shown in Figure 1C. The different colors represent different years' 129 publications. Regarding the number of publications, Elsevier (1685) and Springer Nature (1388) far 130 surpassed other publishers, confirming their unsurpassable international position in the publishing 131 industry.

#### 132 **3.2** Country influence and collaboration

133 In the ten years from 2013-2022, 124 countries or regions participated in the domain of intestinal 134 microbes and obesity. China ranked first with 3036 publications, accounting for 34.16% of all 135 publications, followed by the United States (2356, 26.51%) and Spain (480, 5.40%) (Figure 2A, Table 136 1). Nevertheless, the United States tops the ranking of the most cited countries with a frequency 137 (number of documents) of 87065, followed by China (65250), France (13809), and Canada (13496) 138 (Supplementary Table S1). The country collaboration map shows the overall perspective of countries' 139 and regions' academic cooperation (Supplementary Figure S1). The shade of blue represents the 140 number of articles published, dark blue indicates more publications, and as the color becomes lighter, 141 the publications decreased. The red line represents that there is academic collaboration between the 142 linked two countries. The thicker the line is, the more cooperation two countries have. The United 143 States had the broadest academic associations with other countries or regions, while its collaboration 144 with China was the tightest, followed by Canada. The United States and China both had a frequent 145 connection with European countries. Moreover, among European countries, there was also tight 146 collaboration, but communication among the other countries or regions still needs to be strengthened.

147 There were 179 institutions included in the results of institution analysis. The top 10 institutions 148 mostly came from China (5) and the USA (2), followed by Denmark (1), Belgium (1), and Spain (1). 149 Chinese academic of sciences, the University of Copenhagen and Shanghai Jiao Tong University had 150 the highest total link strength, revealing close cooperation with other institutions (Figure 2B, Table 2).

#### 151 **3.3** Distribution and co-authorship of authors

152 In this analysis, 16 authors had published at least 21 publications. The 10 most productive authors 153 are listed in Table 3. Most of the authors were from Europe, with Belgium (n=3), followed by China 154 (n=3), Canada (n=2), Sweden (n=1), and France (n=1). Patrice D. Cani (97 publications) and Nathalie 155 M. Delzenne (73 publications), both from Université catholique de Louvain, contributed the top two 156 most publications, followed by Reimer, Raylene A (51 publications) from the University of Calgary. 157 The top two authors maintained the most contact with other research labs, and the cooperation between 158 Patrice D. Cani and Nathalie M. Delzenne was close (Figure 3A). Co-cited authors are shown in Figure 159 3B and Table 4. Peter J. Turnbaugh, and Patrice D. Cani rank the top two, which manifests their 160 centrality in the research field.

#### 161 **3.4** Analysis of journals and co-cited academic journals

162 As shown in Table 5, the top 10 most productive journals published 2155 publications, accounting 163 for 24.25% of all publications. Nutrients (impact factor (IF) 2021, 6.706) published the most articles 164 (399 publications, 4.49%), followed by Scientific Reports (IF 2021, 4.997; 339 publications, 3.81%) and Food & Function (IF 2021, 6.317; 281 publications, 3.16%). The top three highest co-cited journals 165 166 were Nature (IF 2021, 69.504; 6446 citations), PLOS ONE (IF 2021, 3.752; 6247 citations), and 167 Proceedings of the National Academy of Sciences of the United States of America (IF 2021, 12.779; 168 5329 citations). The co-citation relationship among different journals was visualized in a co-citation 169 network (Figure 4, Table 6). Nature had the highest centrality (0.18), followed by PLOS ONE (0.17), 170 which means that the published articles in the two journals were widely recognized and cited by 171 researchers.

172 Supplementary Figure S2 shows the dual-map overlay of journals. The yellow, green, and orange 173 spline waves represent citations made by the source articles. Each spline curve starts with the citing 174 map on the left and points to the cited map on the right. The label represents the subject covered by the 175 journal. The dual-map overlay shows seven main citation paths. The published articles were mostly 176 focused on journals in the fields of medicine, medical, clinical, molecular, biology, and immunology 177 and partly focused on veterinary, animal and science. Cited articles were mostly published in journals 178 in the fields of molecular biology and genetics, and many articles were published in journals in the 179 fields of health, nursing, medicine, environmental, toxicology and nutrition.

#### 180 **3.5** Analysis of keywords

#### 181 **3.5.1 Co-occurrence analysis**

Keywords could accurately reflect the main research point of an article, which has a high condensation in a research field and can directly point to the center of the text. Therefore, a high frequency of keywords represents hot issues in a research field and research hotspots. The keyword cooccurrence graph for the effects of intestinal microbes on obesity is shown in Figure 5. The density value was 0.0457. The ten most frequent keywords were gut microbiota, obesity, inflammation, insulin resistance, intestinal microbiota, health, metabolism, diet, gut microbiome, and metabolic syndrome (Table 7).

#### 189 **3.5.2 Cluster analysis**

The clustering analysis of the keywords yielded 288 nodes and 1083 links, with a Q value of 0.4123 (>0.3) and an S value of 0.7178 (>0.5), which are two important indicators to evaluate the significance of the clustering effect. There were 8 clusters, including diversity, insulin resistance, short-chain fatty acids, activation, food intake, bariatric surgery, body composition, and nonalcoholic fatty liver disease (Figure 6). "diversity" # 0 was the largest cluster, followed by "insulin resistance" # 1 and "short-chain fatty acids" # 2. Timeline View analysis was conducted to further analyze the keywords of the effects of intestinal microbes on obesity (Supplementary Figure S3).

#### 197 **3.5.3 Burst detection.**

198 CiteSpace was further used to perform burst detection with high frequency and therefore revealed 199 research frontiers and hotspot trends over a period. As shown in Supplementary Table S2, the first 25 200 keywords were sorted by strengths of burst to discover the research hotspots about the effects of 201 intestinal microbes on obesity. Keywords such as "ecology" (strength 28.33), "human gut microbiota" 202 (strength 27.34), "microbiota" (strength 23.44), "diet induced obesity" (strength 21.12), and "irritable

bowel disease" (strength 16) were the strongest keywords leading the research boom from 2013-2018.
Keywords such as "obese" (strength 9.3) and "serum" (9.69) were the most recent trending keywords.
Supplementary Table S3 shows the top 25 references with the strongest citation bursts, which were

206 recognized as a critical milestone and led the development direction of the field for a while.

#### 207 4 Discussion

In this study, we performed a bibliometric analysis of the effects of intestinal microbes on obesityrelated studies from 2013 to 2022 using the core collection of WoSCC to comprehensively understand global research trends and hotspots and provide references for researchers in this field or those who want to become involved in the field. Based on the above analysis, it is certain that obesity is closely influenced by intestinal microbes.

#### 213 4.1 Global trends in effects of intestinal microbes on obesity

214 The amount of annual scientific production is an important indicator of development in an academic 215 field. In our study, a total of 8888 original articles from 1392 journals met the inclusion criteria. The 216 article annual growth rate reached 22.93%, and the annual growth rate from 2019 to 2020 was the 217 highest. An article published by Lawrence A. David et al., which was the most co-cited, found that an 218 altered diet could lead to a rapid gut microbiome response. The results showed that short-term 219 consumption of diets composed entirely of animal or plant products alters microbial community 220 structure and overwhelms interindividual differences in microbial gene expression (DavidMaurice et 221 al., 2014). The second most cited article was published by Amandine Everard et al. They isolated a 222 mucin-degrading bacterium that resides in the mucus layer, named Akkermansia muciniphila. The 223 results showed that the abundance of A. muciniphila decreased in obese and type 2 diabetic mice, but 224 prebiotic feeding could normalize A. muciniphila abundance. Moreover, high-fat diet-induced 225 metabolic disorders, including fat-mass gain, metabolic endotoxemia, adipose tissue inflammation, and 226 insulin resistance, could be reversed by A. muciniphila treatment (EverardBelzer et al., 2013). Although 227 these two articles are about different research, they both proved the correlations between intestinal microbes and the host, which have greatly driven the development of the field. 228

229 Among the 10 most influential journals in effects of intestinal microbes on obesity research, four 230 were also among the top 10 co-cited journals (Nature, Proceedings of the national academy of sciences 231 (PNAS), Gut, Science). Among these, the IF value of Nature was 69.504 (2021), followed by Science 232 (63.832), Gut (31.795) and PNAS (12.779), indicating their outstanding contributions in the field of 233 effects of intestinal microbes on obesity research. The most frequently co-cited journals were Nature, PLOS ONE, PNAS, and Gut. Notably, three of the top 10 co-cited references were published in Nature, 234 235 one of them was published in PNAS, and one of them was published in Gut. The main research hotspots 236 were about the mechanism of dietary intervention or human genetics to intestinal microbes. Moreover, 237 richness or kinds of intestinal microbes can also modulate the host's metabolism, such as obesity or type 2 diabetes. These findings have laid the foundation for a deeper understanding of the regulatory 238 239 relationship between intestinal microbes and obesity or type 2 diabetes.

#### 240 4.2 Co-authorship networks in effects of intestinal microbes on obesity research

241 The closeness of collaboration between countries/regions, institutions, and authors was assessed, 242 which can help find the laws of scientific research cooperation, guiding more effective scientific

243 research activities, and promoting potential collaborative opportunities for other groups. According to 244 Table 1, China, with 3036 articles, ranks first among the top 10 total publication countries, followed 245 by the United States, with 2356 publications. Notably, the total number of publications in the USA (87065) was greater than that in China (65250) and far greater than that in France (13809). Canada 246 247 (13496) and the other top 10 most cited countries. The total citations of the United States and China 248 are far more than several other top 10 most cited countries citations combined. Without doubt, the 249 United States and China demonstrate their position as leaders in the field. Notably, Israel with 49 250 publications was the most cited country per article (average article citations, 124.18), suggesting that 251 the quality of research on effects of intestinal microbes on obesity in Israel is very high, the similar 252 story is unfolding in many European countries such as Belgium (average article citations, 107.85), 253 Sweden (average article citations, 103.95), etc., which sets a model for other countries/regions in this 254 research field (Supplementary Table S1, S4). To improve the quality of articles, the reliability of the 255 data source and the rigor of the experimental design may be the first factors affecting the quality of 256 publications, and more attention should be given to these aspects in future studies. According to Figure 257 2B, the bottom right is most dominated by universities from China, while the upper left is most 258 dominated by schools in North America or Europe. Nevertheless, more cooperation between the two 259 clusters could be conducted to catalyze breakthrough progress in research on the effects of intestinal 260 microbes on obesity.

261 Four of the top 10 institutions are based in China, followed by the United States with three, which 262 are thereby maximizing regional advantages and demonstrating the dominance of the United States 263 and China in the field. This may partly explain why China and the United States consistently maintain 264 a high quantity of publications. University of Copenhagen in Denmark was the most productive 265 institution worldwide, followed by Chinese academic of sciences in China, indicating that these two 266 institutions participated in the most collaborations with other institutions worldwide. Although Spain 267 and Canada ranked third and fourth in terms of total publications, only one of the Spanish research 268 institutions ranked in the top 10, indicating a lack of institutions with professional and research stature 269 in terms of the effects of intestinal microbes on obesity research (Table 1-2). The most effective 270 organizations and groups are leading the trends in the effects of intestinal microbes on obesity research; 271 thus, further study at these institutions will ensure continuous future development in this field.

#### 272 4.3 Basic knowledge and hotspots in effects of intestinal microbes on obesity research

#### 273 4.3.1 Basic knowledge

274 The most frequently cited references possess pivotal academic position in the field. Either positive 275 or negative conclusions from the documents influence others' research direction. In our study, the top 276 co-cited references were used to investigate the knowledge base for the effects of intestinal microbes 277 on obesity research. Among the top 10 co-cited references, we found that second-generation 278 sequencing was a vital technology that relies on its high-throughput characteristic, making it easy to 279 sequence the transcriptome or genome of a species (Schuster, 2008). With the development of 280 sequencing technology, a large amount of data is generated. To make more accurate and full use of 281 sequencing data, software and data analysis platforms have also been developed. An open-source 282 software package named DADA2 is used for modeling and correcting Illumina-sequenced amplicon 283 errors. With the help of this software, researchers could accurately reconstruct amplicon-sequenced 284 communities at the highest resolution, which ensured the accuracy of the research to the greatest extent 285 (CallahanMcMurdie et al., 2016). Another utility tool named QIIME 2 could serve not only as a 286 marker-gene analysis tool but also as a multidimensional and powerful data science platform that can 287 be rapidly adapted to analyze diverse microbiome features (BolyenRideout et al., 2019). These tools 288 help drive rapid development in microbiome research. With the help of sequencing, recent research

289 found that the richness of the human gut microbiome correlated with human metabolic markers. 290 Individuals with a low bacterial richness were characterized by more marked overall adiposity, insulin 291 resistance and dyslipidemia and a more pronounced inflammatory phenotype when compared with 292 high bacterial richness individuals (Le ChatelierNielsen et al., 2013). Correspondingly, with higher gut 293 microbiome gene richness and A. muciniphila abundance could exhibit the healthiest metabolic status, 294 particularly in fasting plasma glucose, plasma triglycerides and body fat distribution (DaoEverard et 295 al., 2016). The obesity-associated gut microbial species Bacteroides thetaiotaomicron, a glutamate-296 fermenting commensal, is linked to changes in circulating metabolites, and the abundance of 297 Bacteroides thetaiotaomicron is markedly decreased in obese individuals and inversely correlated with 298 serum glutamate concentration (LiuHong et al., 2017). Human obesity is a heterogeneous condition in 299 the context of pathogenesis, pathophysiology and therapeutic responsiveness. Studies of alterations in 300 the genome-the microbial gut metagenome-may define subsets of adult individuals with different 301 metabolic risk profiles, which could contribute to resolving some of the heterogeneity associated with 302 adiposity-related phenotypes (Le ChatelierNielsen et al., 2013). All these findings suggest that gut 303 microbiome richness is a key factor in maintaining the homeostasis of human health.

304 The gut microbiome can rapidly respond to an altered diet, and a dynamic balance can be achieved 305 between intestinal microbes and diet immediately, indicating that not only a long-term diet but also a 306 short-term diet can influence the gut microbiome (DavidMaurice et al., 2014). Weight-loss intervention by bariatric surgery partially reversed obesity-associated microbial and metabolic alterations in obese 307 308 individuals (LiuHong et al., 2017). This means that external interventions can affect intestinal microbes. 309 Research also proved that transmissible and modifiable interactions between diet and microbiota 310 influence host biology. The transformation correlated with invasion of members of *Bacteroidales* from 311 Ln into Ob microbiota that prevented development of increased adiposity and body mass in Ob cage 312 mates and transformed their microbiota's metabolic profile to a leanlike state. Therefore, it may be 313 possible to intervene in obesity by targeting the gut microbiota (RidauraFaith et al., 2013), and there is 314 already evidence proving this speculation. Akkermansia muciniphila is a mucin-degrading bacterium 315 that resides in the mucus layer. Research found that A. muciniphila decreased in obese and type 2 316 diabetic mice, and A. muciniphila treatment reversed high-fat diet-induced metabolic disorders, 317 including fat-mass gain, metabolic endotoxemia, adipose tissue inflammation, and insulin resistance 318 (EverardBelzer et al., 2013). Pasteurized A. muciniphila could enhance its capacity to reduce fat mass 319 development, insulin resistance and dyslipidemia. Moreover, Amuc 1100, a purified membrane 320 protein from Akkermansia muciniphila, could improve the gut barrier and partly recapitulate the 321 beneficial effects of the bacterium (PlovierEverard et al., 2017). All these findings are of great driving 322 significance, indicating that it is feasible to use microbes to treat obesity, but more research is needed 323 to support this hypothesis.

324 In addition to the underlying effect of intestinal microbes on obesity, substantial evidence has suggested that intestinal microbes also have an effect on type 2 diabetes, which is also a major public 325 326 health issue throughout the world. A metagenome-wide association study (MGWAS) was used to 327 identify and validate type-2-diabetes-associated markers and to establish the concept of a metagenomic 328 linkage group, which enabled taxonomic species-level analyses. Type 2 diabetes is accompanied by a 329 decrease in the abundance of some universal butyrate-producing bacteria and an increase in various 330 opportunistic pathogens (QinLi et al., 2012). Conversely, high bacterial richness was related to a 331 reduction in insulin resistance (Le ChatelierNielsen et al., 2013), and A. muciniphila treatment could 332 also reverse insulin resistance (EverardBelzer et al., 2013) (Table 8).

#### 333 4.3.2 Research hotspots

334 Keywords with high frequency are usually used to accurately reveal the main research interests and hotspots in the field within a certain period, which help researchers to precisely catch up to the research 335 336 trends from numerous studies. By analyzing the co-occurrence keywords, "gut microbiota" and 337 "obesity" had the highest frequency and centrality ranking the top two, indicating that the main research 338 interests focused on the relationship between "gut microbiota" and "obesity". The effects of intestinal 339 microbes on obesity have been supported by several trials, which have all shown that intestinal 340 microbes modulate the metabolism of obesity (EverardBelzer et al., 2013; Le ChatelierNielsen et al., 341 2013; RidauraFaith et al., 2013). The clustering function was used to show the whole network map. 342 The 8 clusters of keywords presented in this paper illustrate 8 research topics, which to some extent 343 represent 8 significant aspects of the research. Studies in the same cluster might have similar research themes. As shown in Figure 6, the top two largest clusters, "diversity" # 0 and "insulin resistance" # 1, 344 345 showed that the two topics were still the main themes in the effect of intestinal microbes on obesity 346 research. Simultaneously, a timeline view analysis was conducted to further observe the evolution pace 347 of each cluster and recognize the pivotal research direction through a microperspective. We can clearly 348 observe in Supplementary Figure S3 that the research focus has shifted with time; by 2020, the research on several topics was more balanced. After 2020, the research in the "shot-chain fatty acids" #2 and 349 350 "activation" #3 directions gradually weakened. The results indicated that gut microbiome research and 351 the treatment of obesity and obesity-related diseases have become the main themes.

352 Furthermore, burst detection was an analytic tool that could effectively capture the dramatic 353 increases in references or keywords in one research field within a specified period; therefore, it served 354 as an important indicator of research hotspots or research frontiers over time. The evolution of the burst 355 keywords over the last 10 years is shown in Supplementary Table S2. According to the ranking by 356 burstiness strength, we observed that the top 15 keywords started to burst in 2013, and "ecology" remained the longest with an end time in 2018. More importantly, the "obese patient" and "serum" 357 358 burst separately in 2019 and 2020 and continue to burst until now, which indicated that the two topics 359 have received continuous attention in recent years and might be the main trends of research on the 360 effect of intestinal microbes on obesity. Notably, there was a sudden and rapid growth in the number 361 of articles in the field, which might be related to the term "serum", which was only a new burst in that 362 vear.

363 Bariatric surgery is considered the only effective and sustainable weight loss method for obese 364 patients. There is a 50-70% decrease in body weight and fat mass after surgical procedures such as 365 Roux-Y gastric bypass (RYGB) and sleeve gastrectomy (SG) (PaganelliLuyer et al., 2019). 366 Nevertheless, obese patients undergoing bariatric surgery may experience overgrowth of small 367 intestinal bacteria (SabateCoupaye et al., 2017), a condition that snags with weight loss and increases 368 the risk of micronutrient deficiency that appears to be harmful for the configuration and composition 369 of intestinal microbiota (HibberdWu et al., 2017; Mach and Clark, 2017). To maintain the effect of 370 surgery and avoid weight rebound, it is important for obese patients to correct the microbial balance 371 and improve microbiota-host interactions with specific interventions (Aron-WisnewskyPrifti et al., 372 2019). Research has also shown that the elevated pH resulting from RYGB could ensure the survival 373 of probiotic bacteria, making it possible for surgical patients to receive probiotic therapy (80). 374 Probiotics are a kind of active microorganism beneficial to the host that colonizes the human intestinal 375 tract and reproductive system, and they can improve the host microecological balance and play a 376 beneficial role. Bariatric surgery is often followed by an increase in Streptococcaceae and a decline in 377 Bifidobacteriaceae (PaganelliLuyer et al., 2019). Proper supplementation with probiotics can 378 compensate for the intestinal microbial imbalance caused by surgery. Moreover, recent research 379 showed that probiotic intervention could increase the levels of peptide YY and GLP-1 in mice (TianLi 380 et al., 2016), reduce the level of intestinal inflammation (TsaiLin et al., 2019), and induce the

381 production of anti-inflammatory cytokines (FolignéParayre et al., 2016). Meanwhile, intestinal peptide 382 signals activate the gut-brain axis, which, in turn, exerts endocrine effects on other organ systems, 383 especially the brain, regulating appetite, metabolism, and other dietary behaviors(Bliss and Whiteside, 384 2018: Hussain and Bloom. 2013: NevrinckVan Hée et al., 2012: RastelliVan Hul et al., 2020). 385 Consequently, weight loss and reduced insulin resistance occur. However, due to the heterogeneity of 386 experimental techniques, methods and objects, the effectiveness and safety of microbial product 387 intervention in improving obesity against intestinal microbiota need to be further verified, and the 388 mechanism is not clear. Few studies have focused on the development of new functional 389 microbiological products, and most of them have been carried out in mice. Before microbial products 390 are reasonably and effectively treated for obesity, a large number of studies are needed, especially 391 randomized controlled clinical trials.

392 Serum samples are a less invasive biomatrix that reflects the dynamic changes in the metabolome 393 of the whole organism (DunnBroadhurst et al., 2011). The serum metabolome has emerged as a 394 technique that focuses on defining the functional status of host-microbial relationships in biological 395 specimens, which can reflect the dynamic changes in metabolites and explore disease-related 396 metabolites or dysregulated metabolic pathways (metabolome analysis for investigating host-gut 397 microbiota interactions). A top 10 co-cited reference showed the gut microbiome and serum 398 metabolome alterations in obesity that patients after bariatric surgery not only appeared partially 399 reversed obesity-associated microbial but also accompanied by metabolic alterations, including the 400 decreased abundance of B. thetaiotaomicron and the elevated serum glutamate concentration, 401 suggesting that it may be possible to intervene in obesity by targeting the gut microbiota (LiuHong et 402 al., 2017). There are complex interactions between the host and intestinal microbes in carbohydrate. 403 amino acid, lipid and nucleic acid metabolism. Intestinal microbes can use their respective metabolites 404 to maintain intestinal viability while affecting the development, homeostasis and function of the host 405 immune system through nutrition- and metabolite-dependent mechanisms (Brestoff and Artis, 2013). 406 Small molecules, such as vitamins, fatty acids, amino acids, and bile acids, regulate host-intestinal 407 metabolic homeostasis by binding to specific host membranes or nuclear receptors (HustedTrauelsen 408 et al., 2017; PutignaniDel Chierico et al., 2016). The references with the highest citation bursts are 409 unique contributions to the field of study. We found that articles related to bile acids have become the 410 most cited articles in recent years (KohDe Vadder et al., 2016; WangLiao et al., 2019). Bile acids 411 directly and rapidly affect the metabolism of bacteria, including membrane damage and disruption of 412 amino acid, nucleotide and carbohydrate metabolism, and short-term exposure to bile acids 413 significantly affects host metabolism by altering the bacterial community structure (TianGui et al., 414 2020). Another study showed that the gut microbiota also produced many other metabolites with 415 important functions, such as short-chain fatty acids (SCFAs) and bile acids (KrishnanDing et al., 2018). 416 Short-chain fatty acids can directly activate G-coupled receptors, inhibit histone deacetylases, and 417 serve as energy substrates (KohDe Vadder et al., 2016). They thus affect various physiological 418 processes and may contribute to health and disease. According to the above analysis, although notable 419 advances in the effects of intestinal microbes on obesity have been made, our understanding of the 420 interrelationships between them remains descriptive, and we still have numerous gaps to fill.

#### 421 **4.4 Strengths and Limitations**

422 The bibliometric study conducted a systematic analysis of the basic situation, research hotspots, 423 and trends in effects of intestinal microbes on obesity from a visualization perspective. The results of 424 the bibliometric study were objective and accurate, which could provide a comprehensive guide for

425 academics who are already or wish to work in this field. Nevertheless, there are still some limitations 426 in our study. First, owing to the nature of the CiteSpace software, our data are filtered only from the 427 WoSCC database, which is not sufficiently comprehensive and may lead to data omission. Second, our 428 results were processed by CiteSpace software with certain algorithms, which could lead to bias in some 429 of the results. Finally, only English articles were included from the database and analysis, potentially 430 leading to a source bias.

# 431 **5** Conclusion

432 Intestinal microbes possess essential research value and application prospects in obesity therapy. 433 Compared with other review articles, the contribution of the study is evident in its visualized ways to 434 reveal the effect of intestinal microbes on obesity. Research on the effects of intestinal microbes on obesity is increasing rapidly, highlighted by the outstanding contributions of China, the United States 435 436 and Spain to the development of this field. Current research hotspots focus on related mechanisms of 437 the effects of intestinal microbes on obesity and therapeutic methods for obesity with intestinal microbes. Diet, probiotic preparations and regulation of the intestinal flora, fecal microbe 438 439 transplantation may have significant impacts on reducing the obesity epidemic. "Obese patient" and 440 "serum" have become the latest burst that have attracted attention in recent years and will serve as the 441 focus of future research. More cooperation and communication between countries and institutions 442 should be strengthened to promote development in this field and benefit more patients with 443 obesity.Data availability statement

444 The original contributions presented in the study are included in the article, and further inquiries 445 can be directed to the corresponding author.

## 446 Author Contributions

ZS and JG conceived the study. XY guided the methodological process. ZS, CT participated in
data collection and analysis. ZS wrote the manuscript. GW, JG and XY revised the manuscript. All
authors contributed to the article and approved the submitted version.

autions contributed to the article and approved the subm

## 450 Funding

451 This study was funded by West China Hospital, Sichuan University (Project No. 339180262).

## 452 Acknowledgments

The authors are indebted to Prof. Zhiyong Zong, who contributed his time, knowledge, and energy to developing this document.

## 455 **Conflict of interest**

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## 458 **Publisher's note**

459 All claims expressed in this article are solely those of the authors and do not necessarily represent 460 those of their affiliated organizations or those of the publisher, the editors and the reviewers. Any 461 product that may be evaluated in this article or claim that may be made by its manufacturer is not

462 guaranteed or endorsed by the publisher.

#### 463 Supplementary Material

464 The Supplementary Material for this article can be found online at:


#### 465 **References**

- 466 Alessandro, F., Marcello, D., Rosa, D. S., Valentina, G., and Luca, M. 2017. Gut microbiota, obesity and metabolic
- 467 disorders. *Minerva gastroenterologica e dietologica* 63 (4), 337-44. doi: 10.23736/S1121-421X.17.02376-5.
- Aron-Wisnewsky, J., Prifti, E., Belda, E., Ichou, F., Kayser, B. D., Dao, M. C., et al. 2019. Major microbiota dysbiosis in
  severe obesity: fate after bariatric surgery. *Gut* 68 (1), 70-82. doi: 10.1136/gutjnl-2018-316103.
- 470 Bliss, E. S., and Whiteside, E. 2018. The Gut-Brain Axis, the Human Gut Microbiota and Their Integration in the
- 471 Development of Obesity. Front Physiol 9, 900. doi: 10.3389/fphys.2018.00900.
- 472 Boccellino, M., Di Domenico, M., Donniacuo, M., Bitti, G., Gritti, G., Ambrosio, P., et al. 2018. AT1-receptor blockade:
- 473 Protective effects of irbesartan in cardiomyocytes under hypoxic stress. *PLoS One* 13 (10), e0202297. doi:
- 474 10.1371/journal.pone.0202297.
- 475 Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., et al. 2019. Reproducible,
- interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol* 37 (8), 852-857. doi:
  10.1038/s41587-019-0209-9.
- Brestoff, J. R., and Artis, D. 2013. Commensal bacteria at the interface of host metabolism and the immune system. *Nature immunology* 14 (7), 676-684. doi: 10.1038/ni.2640.
- Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J., and Holmes, S. P. 2016. DADA2: Highresolution sample inference from Illumina amplicon data. *Nat Methods* 13 (7), 581-3. doi: 10.1038/nmeth.3869.
- 482 Dao, M. C., Everard, A., Aron-Wisnewsky, J., Sokolovska, N., Prifti, E., Verger, E. O., et al. 2016. Akkermansia
- muciniphila and improved metabolic health during a dietary intervention in obesity: relationship with gut microbiome
   richness and ecology. *Gut* 65 (3), 426-36. doi: 10.1136/gutjnl-2014-308778.
- David, L. A., Maurice, C. F., Carmody, R. N., Gootenberg, D. B., Button, J. E., Wolfe, B. E., et al. 2014. Diet rapidly and
  reproducibly alters the human gut microbiome. *Nature* 505 (7484), 559-63. doi: 10.1038/nature12820.
- 487 Dunn, W. B., Broadhurst, D., Begley, P., Zelena, E., Francis-McIntyre, S., Anderson, N., et al. 2011. Procedures for
- 488 large-scale metabolic profiling of serum and plasma using gas chromatography and liquid chromatography coupled to
- 489 mass spectrometry. *Nature protocols* 6 (7), 1060-1083. doi: 10.1038/nprot.2011.335.
- El-Sayed Moustafa, J. S., and Froguel, P. 2013. From obesity genetics to the future of personalized obesity therapy. *Nat Rev Endocrinol* 9 (7), 402-13. doi: 10.1038/nrendo.2013.57.
- 492 Everard, A., Belzer, C., Geurts, L., Ouwerkerk, J. P., Druart, C., Bindels, L. B., et al. 2013. Cross-talk between
- Akkermansia muciniphila and intestinal epithelium controls diet-induced obesity. *Proc Natl Acad Sci U S A* 110 (22),
  9066-71. doi: 10.1073/pnas.1219451110.
- 495 Foligné, B., Parayre, S., Cheddani, R., Famelart, M. H., Madec, M. N., Plé, C., et al. 2016. Immunomodulation properties
- 496 of multi-species fermented milks. *Food Microbiol* 53 (Pt A), 60-9. doi: 10.1016/j.fm.2015.04.002.


- 497 Gao, K., Dou, Y., Lv, M., Zhu, Y., Hu, S., and Ma, P. 2021. Research hotspots and trends of microRNA in
- 498 periodontology and dental implantology: a bibliometric analysis. *Ann Transl Med* 9 (14), 1122. doi: 10.21037/atm-21499 726.
- 500 Halmos, T., and Suba, I. 2016. [Physiological patterns of intestinal microbiota. The role of dysbacteriosis in obesity,
- 501 insulin resistance, diabetes and metabolic syndrome]. Orv Hetil 157 (1), 13-22. doi: 10.1556/650.2015.30296.
- 502 Hibberd, M. C., Wu, M., Rodionov, D. A., Li, X., Cheng, J., Griffin, N. W., et al. 2017. The effects of micronutrient
- deficiencies on bacterial species from the human gut microbiota. *Science translational medicine* 9 (390), eaal4069. doi:
   10.1126/scitranslmed.aal4069.
- Hussain, S. S., and Bloom, S. R. 2013. The regulation of food intake by the gut-brain axis: implications for obesity. *Int J Obes (Lond)* 37 (5), 625-33. doi: 10.1038/ijo.2012.93.
- 507 Husted, A. S., Trauelsen, M., Rudenko, O., Hjorth, S. A., and Schwartz, T. W. 2017. GPCR-mediated signaling of 508 metabolites. *Cell metabolism* 25 (4), 777-796. doi: 10.1016/j.cmet.2017.03.008.
- 509 Jandhyala, S. M., Talukdar, R., Subramanyam, C., Vuyyuru, H., Sasikala, M., and Nageshwar Reddy, D. 2015. Role of 510 the normal gut microbiota. *World J Gastroenterol* 21 (29), 8787-803. doi: 10.3748/wjg.v21.i29.8787.
- Kelly, T., Yang, W., Chen, C. S., Reynolds, K., and He, J. 2008. Global burden of obesity in 2005 and projections to
  2030. *Int J Obes (Lond)* 32 (9), 1431-7. doi: 10.1038/ijo.2008.102.
- 513 Koh, A., De Vadder, F., Kovatcheva-Datchary, P., and Bäckhed, F. 2016. From Dietary Fiber to Host Physiology: Short-514 Chain Fatty Acids as Key Bacterial Metabolites. *Cell* 165 (6), 1332-1345. doi: 10.1016/j.cell.2016.05.041.
- 515 Krishnan, S., Ding, Y., Saedi, N., Choi, M., Sridharan, G. V., Sherr, D. H., et al. 2018. Gut Microbiota-Derived
- 516 Tryptophan Metabolites Modulate Inflammatory Response in Hepatocytes and Macrophages. *Cell Rep* 23 (4), 1099-517 1111. doi: 10.1016/j.celrep.2018.03.109.
- Le Chatelier, E., Nielsen, T., Qin, J., Prifti, E., Hildebrand, F., Falony, G., et al. 2013. Richness of human gut microbiome correlates with metabolic markers. *Nature* 500 (7464), 541-6. doi: 10.1038/nature12506.
- Liu, R., Hong, J., Xu, X., Feng, Q., Zhang, D., Gu, Y., et al. 2017. Gut microbiome and serum metabolome alterations in
  obesity and after weight-loss intervention. *Nat Med* 23 (7), 859-868. doi: 10.1038/nm.4358.
- 522 Liu, Z., Gao, K., Hai, Y., and Liu, T. 2022. Developments, Focuses, and Trends in Early-Onset Scoliosis From 2005 to
- 523 2020: A Systematic Bibliometric Analysis. *World Neurosurg* 158, e697-e710. doi: 10.1016/j.wneu.2021.11.044.
- 524 Mach, N., and Clark, A. 2017. Micronutrient deficiencies and the human gut microbiota. *Trends in microbiology* 25 (8), 525 607-610. doi: 10.1016/j.tim.2017.06.004.
- 526 Mohan, I., Gupta, R., Misra, A., Sharma, K. K., Agrawal, A., Vikram, N. K., et al. 2016. Disparities in Prevalence of
- 527 Cardiometablic Risk Factors in Rural, Urban-Poor, and Urban-Middle Class Women in India. PLoS One 11 (2),
- 528 e0149437. doi: 10.1371/journal.pone.0149437.
- 529 Nazli, A., Yang, P. C., Jury, J., Howe, K., Watson, J. L., Söderholm, J. D., et al. 2004. Epithelia under metabolic stress
- 530 perceive commensal bacteria as a threat. Am J Pathol 164 (3), 947-57. doi: 10.1016/s0002-9440(10)63182-3.
- 531 Neyrinck, A. M., Van Hée, V. F., Piront, N., De Backer, F., Toussaint, O., Cani, P. D., et al. 2012. Wheat-derived
- arabinoxylan oligosaccharides with prebiotic effect increase satietogenic gut peptides and reduce metabolic endotoxemia
   in diet-induced obese mice. *Nutr Diabetes* 2 (1), e28. doi: 10.1038/nutd.2011.24.
- 534 Paganelli, F. L., Luyer, M., Hazelbag, C. M., Uh, H. W., Rogers, M. R. C., Adriaans, D., et al. 2019. Roux-Y Gastric
- 535 Bypass and Sleeve Gastrectomy directly change gut microbiota composition independent of surgery type. Sci Rep 9 (1),
- 536 10979. doi: 10.1038/s41598-019-47332-z.

- 537 Plovier, H., Everard, A., Druart, C., Depommier, C., Van Hul, M., Geurts, L., et al. 2017. A purified membrane protein
- from Akkermansia muciniphila or the pasteurized bacterium improves metabolism in obese and diabetic mice. *Nat Med* 23 (1), 107-113. doi: 10.1038/nm.4236.
- 25 (1), 107 115. doi: 10.1050/miii. 1250.
- 540 Putignani, L., Del Chierico, F., Vernocchi, P., Cicala, M., Cucchiara, S., Dallapiccola, B., et al. 2016. Gut microbiota 541 dysbiosis as risk and premorbid factors of IBD and IBS along the childhood–adulthood transition. *Inflammatory bowel*
- dysbiosis as risk and premorbid factors of IBD and IBS along the childhood–adulthood transition. *Inflammatory bowel diseases* 22 (2), 487-504. doi: 10.1097/MIB.00000000000602.
- Qin, J., Li, Y., Cai, Z., Li, S., Zhu, J., Zhang, F., et al. 2012. A metagenome-wide association study of gut microbiota in
  type 2 diabetes. *Nature* 490 (7418), 55-60. doi: 10.1038/nature11450.
- 545 Rastelli, M., Van Hul, M., Terrasi, R., Lefort, C., Régnier, M., Beiroa, D., et al. 2020. Intestinal NAPE-PLD contributes
- to short-term regulation of food intake via gut-to-brain axis. *Am J Physiol Endocrinol Metab* 319 (3), E647-e657. doi:
- 547 10.1152/ajpendo.00146.2020.
- Ridaura, V. K., Faith, J. J., Rey, F. E., Cheng, J., Duncan, A. E., Kau, A. L., et al. 2013. Gut microbiota from twins
  discordant for obesity modulate metabolism in mice. *Science* 341 (6150), 1241214. doi: 10.1126/science.1241214.
- 550 Rodríguez, J. M., Murphy, K., Stanton, C., Ross, R. P., Kober, O. I., Juge, N., et al. 2015. The composition of the gut
- 551 microbiota throughout life, with an emphasis on early life. *Microb Ecol Health Dis* 26, 26050. doi:
- 552 10.3402/mehd.v26.26050.
- 553 Sabate, J.-M., Coupaye, M., Ledoux, S., Castel, B., Msika, S., Coffin, B., et al. 2017. Consequences of small intestinal
- bacterial overgrowth in obese patients before and after bariatric surgery. *Obesity surgery* 27, 599-605. doi:
- Scherer, P. E., and Hill, J. A. 2016. Obesity, Diabetes, and Cardiovascular Diseases: A Compendium. *Circ Res* 118 (11),
   1703-5. doi: 10.1161/circresaha.116.308999.
- Schuster, S. C. 2008. Next-generation sequencing transforms today's biology. *Nat Methods* 5 (1), 16-8. doi:
  10.1038/nmeth1156.
- Sidhu, M., and van der Poorten, D. 2017. The gut microbiome. *Australian family physician* 46 (4), 206-211. doi:
  10.3316/informit.741870194405168.
- Tang, W. H., Kitai, T., and Hazen, S. L. 2017. Gut Microbiota in Cardiovascular Health and Disease. *Circ Res* 120 (7),
  1183-1196. doi: 10.1161/circresaha.117.309715.
- 563 Tian, P., Li, B., He, C., Song, W., Hou, A., Tian, S., et al. 2016. Antidiabetic (type 2) effects of Lactobacillus G15 and
- Q14 in rats through regulation of intestinal permeability and microbiota. *Food Funct* 7 (9), 3789-3797. doi:
  10.1039/c6fo00831c.
- Tian, Y., Gui, W., Koo, I., Smith, P. B., Allman, E. L., Nichols, R. G., et al. 2020. The microbiome modulating activity of
  bile acids. *Gut microbes* 11 (4), 979-996. doi: 10.1080/19490976.2020.1732268.
- Tsai, Y. L., Lin, T. L., Chang, C. J., Wu, T. R., Lai, W. F., Lu, C. C., et al. 2019. Probiotics, prebiotics and amelioration
  of diseases. *J Biomed Sci* 26 (1), 3. doi: 10.1186/s12929-018-0493-6.
- 570 Wang, K., Liao, M., Zhou, N., Bao, L., Ma, K., Zheng, Z., et al. 2019. Parabacteroides distasonis Alleviates Obesity and
- 571 Metabolic Dysfunctions via Production of Succinate and Secondary Bile Acids. *Cell Rep* 26 (1), 222-235.e5. doi:
- 572 10.1016/j.celrep.2018.12.028.

- 573 574 Wang, Z., Ma, D., Pang, R., Xie, F., Zhang, J., and Sun, D. 2020. Research progress and development trend of social
- media big data (smbd): Knowledge mapping analysis based on citespace. ISPRS International Journal of Geo-
- 575 Information 9 (11), 632. doi: 10.3390/ijgi9110632.




- 577 FIGURE 1 Global publication trend and country or region analysis for the effect of intestinal microbes on obesity
- 578 research. (A) Annual worldwide publication output. (B) Growth trends in publication output from the top 10 most
- 579 productive countries. (C) Top 10 publishers according to their contribution to the total number of articles on the effect of 580 intestinal microbes on obesity research.




581 FIGURE 2 CiteSpace network visualization map of countries/regions and institutions. (A) CiteSpace network

582 visualization map of countries/regions involved in the effect of intestinal microbes on obesity research (node label: by 583 citation, label font size: proportional). (B) CiteSpace network visualization map of institutions involved in the effect of


584 intestinal microbes on obesity research (node label: by citation, label font size: proportional).




585 FIGURE 3 CiteSpace network visualization map of authors and co-cited authors. (A) CiteSpace network visualization 586 map of authors involved in the effect of intestinal microbes on obesity research (node label: by citation, label font size: 587 proportional). (B) CiteSpace network visualization map of co-cited authors involved in the effect of intestinal microbes 588 on obesity research (node label: by centrality, label font size: proportional).



589 FIGURE 4 CiteSpace network visualization map of co-cited journals involved in the effect of intestinal microbes on 590 obesity research (node label: by centrality, label font size: proportional).



591 FIGURE 5 Keyword co-occurrence network for the effect of intestinal microbes on obesity research (node label: by 592 centrality, label font size: uniformed).



- 593 FIGURE 6 Keyword clusters for the effect of intestinal microbes on obesity research.
- 594
- 595 TABLE 1 Top 10 productive countries/regions related to the effect of intestinal microbes on obesity.

Rank	Countries	Articles	Centrality	Percentage(N/8888)
1	China	3036	0.02	34.16%
2	the United States	2356	0.25	26.51%
3	Spain	480	0.14	5.40%
4	Canada	466	0.07	5.24%
5	France	432	0.11	4.86%
6	Japan	406	0.02	4.57%
7	Germany	382	0.12	4.30%
8	Italy	372	0.12	4.19%
9	South Korea	364	0.05	4.10%
10	England	355	0.23	3.99%

596

597 TABLE 2 Top 10 productive institutions related to the effect of intestinal microbes on obesity.

Rank	Institutions	Articles	Centrality	Country
1	University of Copenhagen	209	0.28	Denmark

2	Chinese academic of sciences	202	0.18	China
3	Shanghai Jiao Tong university	174	0.09	China
4	China Agricultural University	129	0.04	China
5	Catholic University of Louvain	114	0.08	Belgium
6	Zhejiang University	107	0.14	China
7	Inst Salud Carlos III	94	0.10	Spain
8	University of California, Davis	93	0.11	the United States
9	University of Illinois	91	0.09	the United States
10	University of Chinese academic of sciences	90	0.02	China

598

599 TABLE 3 Top 10 productive authors related to the effect of intestinal microbes on obesity.

Rank	Name	Articles	Centrality	Country
1	Patrice D. Cani	97	0.03	Belgium
2	Nathalie M. Delzenne	73	0.03	Belgium
3	Reimer, Raylene A	51	0.00	Canada
4	Backhed, Fredrik	35	0.06	Sweden
5	Neyrinck, Audrey M	34	0.01	Belgium
6	Wang, Jing	33	0.00	China
7	Liu, Bin	32	0.00	China
8	Marette, Andre	28	0.01	Canada
9	Chen, Wei	28	0.01	China
10	Clement, Karine	27	0.01	France

600

601 TABLE 4 Top 10 co-cited authors related to the effect of intestinal microbes on obesity.

Rank	Cited authors	Articles	Centrality	Country
1	Peter J. Turnbaugh	3307	0.33	the United States
2	Patrice D. Cani	2879	0.27	Belgium
3	Ruth E. Ley	2508	0.11	the United States
4	Backhed, Fredrik	1660	0.09	Sweden
5	J. Gregory Caporaso	1448	0.1	the United States
6	Amandine Everard	1370	0.08	Belgium
7	Junjie Qin	1250	0.12	China
8	Robert C Edgar	935	0.01	the United States
9	Nicola Segata	806	0.01	Italy
10	Emmanuelle Le Chatelier	751	0.01	France

602

Sources	Articles	Percentage(N/8888)	IF
Nutrients	399	4.49%	6.706
Scientific Reports	339	3.81%	4.997
Food & Function	281	3.16%	6.317
PLOS ONE	265	2.98%	3.752
Frontiers in microbiology	226	2.54%	6.064
Journal of functional foods	171	1.92%	5.223
Molecular nutrition & food research	136	1.53%	6.575
Frontiers in nutrition	116	1.31%	6.590
Journal of agricultural and food chemistry	114	1.28%	5.895
International journal of molecular sciences	108	1.22%	6.208

604 TABLE 5 Top 10 productive journals related to the effect of intestinal microbes on obesity.

605

606 TABLE 6 Top 10 co-cited journals related to the effect of intestinal microbes on obesity.

Rank	Co-cited journals	Articles	Centrality	IF
1	Nature	6446	0.18	69.504
2	PLOS ONE	6247	0.17	3.752
3	Proceedings of the national academy of sciences (PNAS)	5329	0.02	12.779
4	Gut	4389	0.01	31.795
5	Science	4180	0.01	63.832
6	Diabetes	4032	0.02	9.305
7	Scientific reports	3979	0.01	4.997
8	Cell	3308	0.01	66.850
9	Nutrients	3250	0.05	6.706
10	Gastroenterology	3114	0.04	33.883

608 TABLE 7 Top 10 keywords related to the effect of intestinal microbes on obesity.

Rank	Keywords	Articles	Centrality
1	gut microbiota	4673	0.39
2	obesity	3587	0.23
3	inflammation	1578	0.04
4	insulin resistance	1368	0.12
5	intestinal microbiota	1038	0.01
6	health	869	0.02
7	metabolism	862	0.02

8	diet	848	0.01
9	gut microbiome	825	0.01
10	metabolic syndrome	714	0.01

#### 609

610 TABLE 8 The top 10 co-cited references related to the effect of intestinal microbes on obesity.

Rank	Title	Author	Article type	Journal	DOI	Articles
1	Diet rapidly and reproducibly alters the human gut microbiome	Lawrence A. David et al., 2013	Letter	Nature	10.1038/nature12 820	383
2	Cross-talk between Akkermansia muciniphila and intestinal epithelium controls diet-induced obesity	Amandine Everard et al.,2013	Article	PNAS	10.1073/pnas.121 9451110	367
3	Akkermansia muciniphila and improved metabolic health during a dietary intervention in obesity- relationship with gut microbiome richness and ecology	Maria Carlota Dao et al., 2016	Article	Gut	10.1136/gutjnl- 2014-308778	303
4	A metagenome-wide association study of gut microbiota in type 2 diabetes	Junjie Qin et al., 2012	Article	Nature	10.1038/nature11 450	291
5	Richness of human gut microbiome correlates with metabolic markers	Emmanuelle Le Chatelier et al., 2013	Article	Nature	10.1038/nature12 506	285
6	Gut microbiota from twins discordant for obesity modulate metabolism in mice	Vanessa K. Ridaura et al., 2013	Article	Science	10.1126/science.1 241214	276
7	A purified membrane protein from <i>Akkermansia muciniphila</i> or the pasteurized bacterium improves metabolism in obese and diabetic mice	Hubert Plovier et al., 2017	Letters	Nature Medicine	10.1038/nm.4236	275
8	DADA2: High-resolution sample inference from Illumina amplicon data	Benjamin J Callahan et al., 2016	Brief communications	Nature methods	10.1038/nmeth.38 69	273
9	Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2	Evan Bolyen et al., 2019	Correspondence	Nature biotechnology	10.1038/s41587- 019-0209-9	265
10	Gut microbiome and serum metabolome alterations in obesity and after weight-loss intervention	Ruixin Liu et al., 2017	Article	Nature medicine	10.1038/nm.4358	264