

1 **Full article title:** Musculoskeletal phenotypes in 3q29 deletion syndrome

2 **Authors:** Rebecca M Pollak¹, Jacob C Tilmon², Melissa M Murphy³, Michael J Gambello⁴,

3 Rossana Sanchez Russo⁴, John P Dormans⁵, Jennifer G Mulle^{1,6}

4 **Affiliations:** ¹Center for Advanced Biotechnology and Medicine, Robert Wood Johnson Medical

5 School, Rutgers University; ²Indiana University School of Medicine; ³Department of Pediatrics,

6 Emory University School of Medicine; ⁴Department of Human Genetics, Emory University

7 School of Medicine; ⁵Emeritus Professor of Orthopedic Surgery, University of Pennsylvania;

8 ⁶Department of Psychiatry, Robert Wood Johnson Medical School, Rutgers University

9 **Corresponding author information:** Jennifer Gladys Mulle, MHS, PhD; Center for Advanced

10 Biotechnology and Medicine and Department of Psychiatry, Robert Wood Johnson Medical

11 School, Rutgers University, 679 Hoes Lane West, Piscataway, NJ 08854; 848-445-9866;

12 jennifer.mulle@rutgers.edu

13 **Funding:** NIH R01 MH110701, NIH T32 GM0008490

14 **Competing interests:** The authors have no conflicts of interest to report.

15 **Availability of data and materials:** The datasets used and/or analyzed during the current study

16 are available from the corresponding author on reasonable request.

17 **Ethics approval and consent to participate:** This study was approved by Emory University's

18 Institutional Review Board (IRB00064133) and Rutgers University's Institutional Review Board

19 (PRO2021001360). All study subjects gave informed consent prior to participating in this study.

20 **Consent for publication:** Not applicable.

21 **Acknowledgements:** We gratefully acknowledge our study population, the 3q29 deletion

22 community, for their participation and commitment to research.

23 **Authors' contributions:** RMP performed the statistical analysis, produced all figures and tables,
24 and wrote the manuscript. JCT performed the initial literature review, designed the survey
25 instrument, collected the data, and performed the preliminary data analysis. MMM, MJG, RSR,
26 JPD, and JGM contributed to the study design, data collection, and preliminary data analysis.
27 MMM and MJG assisted in refining the survey instrument. JGM edited the manuscript and
28 provided guidance on analyzing and interpreting data. JGM was the principal investigator
29 responsible for study direction. All authors participated in commenting on the drafts and have
30 read and approved the final manuscript.

31 **Abstract**

32 3q29 deletion syndrome (3q29del) is a rare genomic disorder caused by a 1.6 Mb deletion
33 (hg19, chr3:195725000–197350000). 3q29del is associated with neurodevelopmental and
34 psychiatric phenotypes, including an astonishing >40-fold increased risk for schizophrenia, but
35 medical phenotypes are less well-described. We used the online 3q29 registry
36 (3q29deletion.org) to recruit 57 individuals with 3q29del (56.14% male) and requested
37 information about musculoskeletal phenotypes with a custom questionnaire. 85.96% of
38 participants with 3q29del reported at least one musculoskeletal phenotype. Congenital
39 anomalies were most common (70.18%), with pes planus (40.35%), pectus excavatum (22.81%),
40 and pectus carinatum (5.26%) significantly elevated relative to the pediatric general population.
41 49.12% of participants reported fatigue after 30 minutes or less of activity. Bone fractures
42 (8.77%) were significantly elevated relative to the pediatric general population, suggesting
43 3q29del impacts bone strength. Participants commonly report receiving medical care for
44 musculoskeletal complaints (71.93%), indicating that these phenotypes impact quality of life for
45 individuals with 3q29del. This is the most comprehensive description of musculoskeletal
46 phenotypes in 3q29del to date, suggests ideas for clinical evaluation, and expands our
47 understanding of the phenotypic spectrum of this syndrome.

48 **Keywords:** 3q29 deletion; musculoskeletal; orthopedic; copy number variant; genomic disorder

49 Introduction

50 The 3q29 deletion is a rare (~1:30,000) (Kendall et al., 2017; Stefansson et al., 2014) 1.6
51 Mb deletion on chromosome 3 (hg19, chr3:195725000–197350000) (Ballif et al., 2008;
52 Glassford et al., 2016; Willatt et al., 2005). Individuals with 3q29 deletion syndrome (3q29del)
53 are at substantially increased risk for a variety of neurodevelopmental and neuropsychiatric
54 phenotypes, including mild to moderate intellectual disability (ID), a 19-fold increased risk for
55 autism spectrum disorder (ASD), and a greater than 40-fold increased risk for schizophrenia (SZ)
56 (Ballif et al., 2008; Cox & Butler, 2015; Girirajan et al., 2012; Glassford et al., 2016; Itsara et al.,
57 2009; Kirov et al., 2012; Klaiman et al., 2022; Marshall et al., 2017; Mulle, 2015; Mulle et al.,
58 2010; Pollak et al., 2019; Sanchez Russo et al., 2021; Sanders et al., 2015; Szatkiewicz et al.,
59 2014; Willatt et al., 2005). Medical features of 3q29del are less well-defined, with most reports
60 focusing on the high rate of cardiac manifestations associated with the 3q29 deletion, namely
61 congenital heart defects (Cox & Butler, 2015). However, more work is needed to define the
62 spectrum of other medical phenotypes associated with the 3q29 deletion.

63 Musculoskeletal complaints, including axial, extremity, and neuromuscular phenotypes,
64 have been associated with an array of copy number variant disorders that have other
65 phenotypic similarities to 3q29del. Significant musculoskeletal phenotypes have been identified
66 in carriers of the 22q11.2 deletion, 16p11.2 reciprocal deletion and duplication, and 7q11.23
67 deletion (Williams syndrome) (Al-Kateb et al., 2014; Bassett et al., 2005; Chapman et al., 1996;
68 Cherniske et al., 2004; Copes et al., 2016; Damasceno et al., 2014; Digilio et al., 2003; Ficcadenti
69 et al., 2015; Homans et al., 2018; Kaplan et al., 1989; Martin et al., 1984; Morava et al., 2002;
70 Morris et al., 1988; Ren et al., 2020; Ricchetti et al., 2004; Stagi et al., 2010; Taeusch et al.,

71 2004; Wu et al., 2015). These phenotypes include high rates of scoliosis, joint phenotypes
72 including joint laxity and joint contractures, and reduced bone density. Similarly, some
73 musculoskeletal phenotypes have been associated with 3q29del, including chest cavity
74 deformities and finger and toe anomalies (Cox & Butler, 2015; Sanchez Russo et al., 2021). One
75 of these studies, performed by members of our team, diagnosed musculoskeletal anomalies via
76 direct clinical assessment of 3q29 deletion carriers by trained medical geneticists (Sanchez
77 Russo et al., 2021). However, musculoskeletal concerns were not the main focus of either of
78 these studies, and more detail is needed to understand the full phenotypic spectrum associated
79 with the 3q29 deletion.

80 In the present study, we report the spectrum of musculoskeletal phenotypes
81 systematically ascertained from a large cohort of individuals with 3q29del. Using a custom
82 questionnaire, we were able to assess various domains of musculoskeletal function, including
83 axial, extremity, and neuromuscular phenotypes; imaging procedures; and medical
84 interventions. These data expand our understanding of the greater phenotypic spectrum
85 associated with the 3q29 deletion and provide important insight regarding an under-reported
86 phenotypic domain.

87 **Methods**

88 *Editorial policies and ethical considerations*

89 This study was approved by Emory University's Institutional Review Board
90 (IRB00064133) and Rutgers University's Institutional Review Board (PRO2021001360).

91 *Sample*

92 57 individuals with 3q29 deletion syndrome (56.14% male) were recruited from the
93 online 3q29 deletion registry (3q29deletion.org), ranging in age from 0.75-46.5 years (mean =
94 12.02 ± 9.37 years). Inclusion criteria for the present study were membership in the 3q29
95 deletion registry as of June 2020 and fluency in English. Participants gave informed consent to
96 participate in the study via an online form. A description of the study sample can be found in
97 Table 1.

98 *Measures*

99 A custom survey instrument was designed by a member of the study team to assess
100 musculoskeletal phenotypes and was reviewed by expert clinicians. The survey consisted of 74
101 questions across 7 different domains related to musculoskeletal function, including axial and
102 extremity skeleton pathology, joint pathology, neuromuscular function, presence of congenital
103 malformations, other common pediatric musculoskeletal conditions, and any medical
104 treatments received (Supplemental Information). Demographic data was collected separately
105 as a part of the 3q29 deletion registry. Questionnaires were filled out by parents or primary
106 caregivers. The survey was hosted on a local REDCap database, and a link to the survey with
107 instructions was emailed to members of the 3q29 deletion registry between June and July of
108 2020.

109 *Analysis*

110 All analyses were performed in R version 4.0.4 (R Core Team, 2008). Data were
111 summarized as counts and proportions of the total participants that reported a given
112 phenotype. The rate of phenotypes in our study sample was compared to the pediatric general
113 population using an exact binomial test via the base R package (R Core Team, 2008). Pediatric

114 general population prevalence rates were established from previously published large studies
115 focusing on pediatric cohorts with an age range similar to that of our study subjects with
116 3q29del (García-Rodríguez et al., 1999; Konieczny et al., 2013; Naranje et al., 2016; Westphal et
117 al., 2009). Data visualization was performed using the plotly R package (Sievert et al., 2017).

118 **Results**

119 *High rates of reported musculoskeletal complaints in individuals with 3q29del*

120 Participants with 3q29del overwhelmingly reported musculoskeletal phenotypes, with
121 85.96% (n=49) of participants reporting at least one musculoskeletal complaint. The most
122 common phenotypic category was congenital anomalies, with 70.18% (n=40) of participants
123 reporting one or more (Figure 1). Excluding congenital phenotypes, 40.35% (n=23) of
124 participants reported at least one joint problem, 15.79% (n=9) of participants reported at least
125 one pain problem, and 59.65% (n=34) of respondents reported at least one neuromuscular
126 problem (Figure 1).

127 Respondents more commonly reported phenotypes in the extremities (68.42%, n=39)
128 compared to axial (36.84%, n=21) or neuromuscular (59.65%, n=34) manifestations (Table 2).
129 The most commonly reported extremity phenotype was pes planus (40.35%, n=23), followed by
130 joint stiffness (28.07%, n=16) (Table 2). The most commonly reported axial phenotype was
131 pectus excavatum (22.81%, n=13) (Table 2). The most commonly reported neuromuscular
132 phenotype was fatigue in 30 minutes or less of physical activity (49.12%, n=28), followed by gait
133 anomalies (36.84%, n=21) (Table 2).

134 *Variable presence of congenital anomalies within 3q29 deletion carriers*

135 While 68.85% of participants with 3q29del reported at least one congenital anomaly,
136 the specific phenotypes experienced by study participants varied. The most commonly reported
137 congenital anomaly was pes planus (40.35%, n=23), followed by a unique or atypical facial
138 structure (31.58%, n=18) and pectus excavatum (22.81%, n=13) (Figure 2A). To determine
139 whether the reports of congenital anomalies by our study subjects was higher than the
140 pediatric general population, we compared the prevalence of three specific phenotypes. Pes
141 planus (40.35%, n=23, $p < 2.2E-16$), pectus excavatum (22.81%, n=13, $p = 3.41E-13$), and pectus
142 carinatum (5.26%, n=3, $p = 0.007$) were all reported at significantly higher rates by our study
143 subjects compared to the pediatric general population (2.7%, 1.275%, and 0.675%, respectively)
144 (García-Rodríguez et al., 1999; Westphal et al., 2009), suggesting that the 3q29 deletion confers
145 an increased risk for congenital musculoskeletal anomalies.

146 *Joint complaints in 3q29 deletion carriers are driven by joint stiffness*

147 Joint problems were a commonly reported phenotype in our study sample. To
148 determine whether there were specific areas of concern versus an equal distribution of
149 complaints across categories, we compared the different classes of joint phenotypes included
150 on our survey instrument. We found that the majority of participants that reported joint
151 problems reported joint stiffness (28.07%, n=16) (Figure 2B). Joint laxity was not commonly
152 reported by our participants, with rates ranging from 0% (shoulder and knee laxity) to 7.02%
153 (n=4, finger laxity) (Figure 2B). These data suggest that the 3q29 deletion does not have a
154 strong phenotypic impact on joint function.

155 *High rate of fatigue in individuals with 3q29del*

156 To determine the level of musculoskeletal endurance in our study population, we asked
157 how long the individual with 3q29del can participate in physical activity before becoming
158 fatigued. 49.12% (n=28) of our study participants reported becoming fatigued after 30 minutes
159 or less of physical activity (Table 2, Figure 2C). Study participants most commonly reported
160 becoming fatigued after 15 to 30 minutes of physical activity (36.84%, n=21) (Figure 2C). These
161 data suggest that musculoskeletal endurance may be impaired by the 3q29 deletion.

162 *Orthopedic imaging procedures among 3q29del study subjects*

163 To determine the proportion of our study sample that received any medical imaging
164 procedures, we asked participants to report any X-ray, CT scan, or MRI performed. 36.84%
165 (n=21) of our study sample reported receiving at least one imaging procedure. The most
166 commonly reported imaging procedures were spinal X-ray and skull MRI (12.28%, n=7) (Table
167 3). Rates of extremity imaging procedures were relatively low; the most commonly reported
168 extremity imaging procedures were hand and hip X-rays (8.77%, n=5) (Table 3). These data
169 suggest that the musculoskeletal phenotypes associated with the 3q29 deletion may not
170 commonly require advanced imaging procedures.

171 *Elevated rate of bone fractures associated with the 3q29 deletion*

172 Coincident with the relatively low rates of diagnostic imaging received by our study
173 sample, few participants reported medical diagnoses confirmed by imaging procedures.
174 Notably, 8.77% (n=5) of our study sample reported a bone fracture confirmed by diagnostic
175 imaging (Table 4); this is significantly elevated compared to the pediatric general population
176 prevalence of 0.947% ($p=0.0002$) (Naranje et al., 2016). 7.02% (n=4) of our study sample
177 reported scoliosis confirmed by diagnostic imaging (Table 4); this is not significantly increased

178 compared to the pediatric general population prevalence of 0.47-5.2% (midpoint=2.835%,
179 p=0.07) (Konieczny et al., 2013), however it suggests that the 3q29 deletion may affect risk for
180 scoliosis. These data demonstrate that the 3q29 deletion is associated with bone fracture
181 liability. A larger sample size is needed to definitively determine whether 3q29 deletion
182 participants have scoliosis at a significantly increased rate compared to the pediatric general
183 population.

184 *Medical care for musculoskeletal complaints is common for 3q29 deletion carriers*

185 We sought to determine whether the musculoskeletal complaints present in our study
186 sample of individuals with 3q29del are associated with high rates of medical intervention.
187 Therefore, we asked study participants whether they had ever received physical therapy, an
188 orthopedic evaluation, or orthopedic surgery for their musculoskeletal complaints. 71.93%
189 (n=41) of our study participants reported receiving at least one of these medical interventions.
190 A majority of individuals with 3q29del received physical therapy (64.91%, n=37) (Figure 3).
191 26.79% (n=15) of our study sample received an orthopedic evaluation, and 7.02% (n=4)
192 underwent orthopedic surgery (Figure 3). These data suggest that some musculoskeletal
193 phenotypes associated with the 3q29 deletion are severe enough to require intervention. These
194 data further suggest that musculoskeletal complaints in 3q29 deletion syndrome may be
195 improved or corrected with less invasive medical interventions such as physical therapy, as
196 compared to surgery.

197 **Discussion**

198 This is the first study of 3q29del to systematically ascertain musculoskeletal phenotypes
199 as a primary outcome measure. We found high rates of musculoskeletal phenotypes in our

200 study sample of individuals with 3q29del, with 85.96% (n=49) of participants reporting at least
201 one complaint. Congenital anomalies (70.18%, n=40) and neuromuscular problems (59.65%,
202 n=34) were common. Respondents reported more phenotypes localized to the extremities
203 (68.42%, n=39) as compared to the axial skeleton (36.84%, n=21). Individuals with 3q29del also
204 received high rates of medical interventions for their musculoskeletal phenotypes, including
205 physical therapy (64.91%, n=37), orthopedic evaluation (26.79%, n=15), and orthopedic surgery
206 (7.02%, n=4). Together, these data suggest that the 3q29 deletion confers a previously
207 underappreciated risk for musculoskeletal phenotypes.

208 We found that the rates of several musculoskeletal phenotypes were significantly
209 increased in our study sample relative to the pediatric general population (García-Rodríguez et
210 al., 1999; Naranje et al., 2016; Westphal et al., 2009). Pes planus, pectus excavatum, pectus
211 carinatum, and bone fractures were all elevated in our 3q29del study sample. Scoliosis was not
212 significantly elevated in our study sample relative to the pediatric general population, though
213 there was an increase that is suggestive of an effect (Konieczny et al., 2013). A previous review
214 of case reports and a prior study by our team both identified high rates of chest cavity
215 deformities, and case reports have identified scoliosis in some 3q29 deletion carriers (Cox &
216 Butler, 2015; Sanchez Russo et al., 2021). Together with the present study, these data suggest
217 that the 3q29 deletion confers specific, but underappreciated, risks for musculoskeletal
218 phenotypes.

219 Musculoskeletal phenotypes have been associated with several copy number variant
220 disorders, including 22q11.2 deletion syndrome, 16p11.2 deletion and duplication syndromes,
221 and Williams syndrome. These syndromes have been associated with scoliosis, extremity

222 anomalies, joint problems, and reduced bone mineral density in isolated studies (Al-Kateb et al.,
223 2014; Bassett et al., 2005; Chapman et al., 1996; Cherniske et al., 2004; Copes et al., 2016;
224 Damasceno et al., 2014; Digilio et al., 2003; Ficcadenti et al., 2015; Homans et al., 2018; Kaplan
225 et al., 1989; Martin et al., 1984; Morava et al., 2002; Morris et al., 1988; Ren et al., 2020;
226 Ricchetti et al., 2004; Stagi et al., 2010; Taeusch et al., 2004; Wu et al., 2015). It has been
227 suggested that the more pressing medical and psychiatric concerns, including congenital heart
228 defects and elevated risk for ASD and SZ, may overshadow orthopedic concerns (Homans et al.,
229 2018). The 3q29 deletion is also associated with a complex constellation of medical,
230 neurodevelopmental, and neuropsychiatric comorbidities that may result in orthopedic
231 phenotypes receiving less attention. However, the high rate of medical care reported by our
232 population specifically for orthopedic concerns highlights the importance of assessing any
233 relevant phenotypes early and introducing appropriate interventions, to lessen the impact of
234 orthopedic problems across the lifespan.

235 Due to the variable nature of musculoskeletal phenotypes associated with the 3q29
236 deletion, clinical care should address multiple areas of risk. Based on the present study, in
237 combination with previous reports, individuals with 3q29del should receive regular orthopedic
238 exams, including periodic scoliosis screening (Cox & Butler, 2015; Sanchez Russo et al., 2021).
239 Regularly assessing spinal curvature will allow for early detection of any anomalies, which is
240 critical for maximizing the benefit of early-stage, less-intensive therapeutics, such as bracing.
241 Individuals with 3q29 deletion syndrome should also receive physical therapy assessments,
242 including the six-minute walk test (Andersen et al., 2016; Kuettel et al., 2022), to help guide
243 clinical care. Additionally, the significantly elevated incidence of bone fracture in our study

244 sample suggests that the 3q29 deletion may have an adverse impact on bone mineral density,
245 which may be compounded by documented feeding challenges within the 3q29del population
246 (Wawrzonek et al., 2022). Reduced bone mineral density has also been identified in other copy
247 number variant disorders, including 22q11.2 deletion syndrome and Williams syndrome
248 (Cherniske et al., 2004; Copes et al., 2016; Ficcadenti et al., 2015; Homans et al., 2018; Stagi et
249 al., 2010). To monitor the progression of this phenotype, periodic DEXA scans or another
250 measurement of bone mineral density, paired with basic labs including measurements of
251 calcium and vitamin D levels, will help to guide clinical care. It is important to note the relatively
252 low frequency of medical imaging procedures reported by our study participants; while this
253 may indicate that the musculoskeletal phenotypes in individuals with 3q29del do not commonly
254 require advanced imaging, it may also suggest that musculoskeletal concerns in this population
255 receive less clinical attention as compared to the significant neurodevelopmental and
256 behavioral manifestations of the syndrome. Because our population is relatively young, it is not
257 known whether the 3q29 deletion is a risk factor for late-onset phenotypes related to reduced
258 bone density, such as osteoporosis. These data highlight the importance of longitudinal follow
259 up to assess later onset conditions.

260 While this study provides critical insight into the musculoskeletal phenotypes associated
261 with 3q29del, it is not without limitations. First, the data was gathered via an optional survey
262 sent out to 3q29 deletion registry participants. It is possible that those individuals who are
263 more severely affected would be more likely to participate. This would potentially bias our
264 results, and lead to an inflation of the rates of different musculoskeletal phenotypes. However,
265 some respondents reported no phenotypes, suggesting that the entire study sample is not

266 comprised of severely affected individuals. Second, the data collected in this study is parent- or
267 self-report, rather than direct clinical observation, and can be subject to recall bias. However,
268 the elevated rates of extremity anomalies, chest wall deformities, and scoliosis mirror a large
269 review of case studies as well as a recent publication by our team that used direct assessment
270 (Cox & Butler, 2015; Sanchez Russo et al., 2021). Our sample is young (average age 12.02 ± 9.37
271 years) so our study cannot answer questions about later onset phenotypes. Finally, while this is
272 the largest study of musculoskeletal phenotypes in 3q29del to date, the sample size is still
273 limited and lacks racial and ethnic diversity. Future studies will work to increase the enrollment
274 of underrepresented minorities in the online 3q29 deletion registry.

275 The present study is the first to systematically ascertain musculoskeletal phenotypes in
276 a large sample of individuals with 3q29del. A majority of our study participants reported at least
277 one musculoskeletal phenotype. Congenital phenotypes, including pectus excavatum, pectus
278 carinatum, and pes planus, were frequently reported and were significantly elevated relative to
279 the pediatric general population (García-Rodríguez et al., 1999; Westphal et al., 2009).
280 Participants also reported a high rate of neuromuscular phenotypes, including gait anomalies
281 and fatigue in 30 minutes or less of physical activity. The rate of bone fractures was significantly
282 increased compared to the pediatric general population (Naranje et al., 2016). Medical
283 interventions, particularly physical therapy, were common in our study sample. These data
284 indicate that musculoskeletal phenotypes are a part of the phenotypic spectrum of the 3q29
285 deletion. Individuals with 3q29del should be monitored for the development of orthopedic
286 concerns, including scoliosis and reduced bone mineral density. Clinical monitoring, combined

287 with early interventions, including physical therapy, where appropriate will improve the
288 function and quality of life for individuals with 3q29del.

289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331

References

- Al-Kateb, H., Khanna, G., Filges, I., Hauser, N., Grange, D. K., Shen, J., Smyser, C. D., Kulkarni, S., & Shinawi, M. (2014). Scoliosis and vertebral anomalies: additional abnormal phenotypes associated with chromosome 16p11.2 rearrangement. *American Journal of Medical Genetics. Part A*, *164a*(5), 1118-1126. <https://doi.org/10.1002/ajmg.a.36401>
- Andersen, L. K., Knak, K. L., Witting, N., & Vissing, J. (2016). Two- and 6-minute walk tests assess walking capability equally in neuromuscular diseases. *Neurology*, *86*(5), 442-445. <https://doi.org/10.1212/wnl.0000000000002332>
- Ballif, B. C., Theisen, A., Coppinger, J., Gowans, G. C., Hersh, J. H., Madan-Khetarpal, S., Schmidt, K. R., Tervo, R., Escobar, L. F., Friedrich, C. A., McDonald, M., Campbell, L., Ming, J. E., Zackai, E. H., Bejjani, B. A., & Shaffer, L. G. (2008). Expanding the clinical phenotype of the 3q29 microdeletion syndrome and characterization of the reciprocal microduplication. *Molecular Cytogenetics*, *1*, 8. <https://doi.org/10.1186/1755-8166-1-8>
- Bassett, A. S., Chow, E. W., Husted, J., Weksberg, R., Caluseriu, O., Webb, G. D., & Gatzoulis, M. A. (2005). Clinical features of 78 adults with 22q11 Deletion Syndrome. *American Journal of Medical Genetics. Part A*, *138*(4), 307-313. <https://doi.org/10.1002/ajmg.a.30984>
- Chapman, C. A., du Plessis, A., & Pober, B. R. (1996). Neurologic findings in children and adults with Williams syndrome. *J Child Neurol*, *11*(1), 63-65. <https://doi.org/10.1177/088307389601100116>
- Cherniske, E. M., Carpenter, T. O., Klaiman, C., Young, E., Bregman, J., Insogna, K., Schultz, R. T., & Pober, B. R. (2004). Multisystem study of 20 older adults with Williams syndrome. *American Journal of Medical Genetics Part A*, *131*(3), 255-264.
- Copes, L. E., Pober, B. R., & Terilli, C. A. (2016). Description of common musculoskeletal findings in Williams Syndrome and implications for therapies. *Clin Anat*, *29*(5), 578-589. <https://doi.org/10.1002/ca.22685>
- Cox, D. M., & Butler, M. G. (2015). A clinical case report and literature review of the 3q29 microdeletion syndrome. *Clinical dysmorphology*, *24*(3), 89-94. <https://doi.org/10.1097/MCD.0000000000000077>
- Damasceno, M. L., Cristante, A. F., Marcon, R. M., & Barros Filho, T. E. (2014). Prevalence of scoliosis in Williams-Beuren syndrome patients treated at a regional reference center. *Clinics (Sao Paulo)*, *69*(7), 452-456. [https://doi.org/10.6061/clinics/2014\(07\)02](https://doi.org/10.6061/clinics/2014(07)02)
- Digilio, M. C., Angioni, A., De Santis, M., Lombardo, A., Giannotti, A., Dallapiccola, B., & Marino, B. (2003). Spectrum of clinical variability in familial deletion 22q11.2: from full manifestation to extremely mild clinical anomalies. *Clin Genet*, *63*(4), 308-313. <https://doi.org/10.1034/j.1399-0004.2003.00049.x>
- Ficcadenti, A., Zallocco, F., Neri, R., Giovannini, L., Tirabassi, G., & Balercia, G. (2015). Bone density assessment in a cohort of pediatric patients affected by 22q11DS. *J Endocrinol Invest*, *38*(10), 1093-1098. <https://doi.org/10.1007/s40618-015-0295-6>
- García-Rodríguez, A., Martín-Jiménez, F., Carnero-Varo, M., Gómez-Gracia, E., Gómez-Aracena, J., & Fernández-Crehuet, J. (1999). Flexible flat feet in children: a real problem? *Pediatrics*, *103*(6), e84. <https://doi.org/10.1542/peds.103.6.e84>

- 332 Girirajan, S., Rosenfeld, J. A., Coe, B. P., Parikh, S., Friedman, N., Goldstein, A., Filipink, R. A.,
333 McConnell, J. S., Angle, B., & Meschino, W. S. (2012). Phenotypic heterogeneity of
334 genomic disorders and rare copy-number variants. *New England Journal of Medicine*,
335 367(14), 1321-1331.
- 336 Glassford, M. R., Rosenfeld, J. A., Freedman, A. A., Zwick, M. E., Mulle, J. G., & Unique Rare
337 Chromosome Disorder Support, G. (2016). Novel features of 3q29 deletion syndrome:
338 Results from the 3q29 registry. *American Journal of Medical Genetics. Part A*, 170A(4),
339 999-1006. <https://doi.org/10.1002/ajmg.a.37537>
- 340 Homans, J. F., Tromp, I. N., Colo, D., Schlösser, T. P. C., Kruyt, M. C., Deeney, V. F. X., Crowley, T.
341 B., McDonald-McGinn, D. M., & Castelein, R. M. (2018). Orthopaedic manifestations
342 within the 22q11.2 Deletion syndrome: A systematic review. *American Journal of*
343 *Medical Genetics. Part A*, 176(10), 2104-2120. <https://doi.org/10.1002/ajmg.a.38545>
- 344 Itsara, A., Cooper, G. M., Baker, C., Girirajan, S., Li, J., Absher, D., Krauss, R. M., Myers, R. M.,
345 Ridker, P. M., Chasman, D. I., Mefford, H., Ying, P., Nickerson, D. A., & Eichler, E. E.
346 (2009). Population analysis of large copy number variants and hotspots of human
347 genetic disease. *American Journal of Human Genetics*, 84(2), 148-161.
348 <https://doi.org/10.1016/j.ajhg.2008.12.014>
- 349 Kaplan, P., Kirschner, M., Watters, G., & Costa, M. T. (1989). Contractures in patients with
350 Williams syndrome. *Pediatrics*, 84(5), 895-899.
- 351 Kendall, K. M., Rees, E., Escott-Price, V., Einon, M., Thomas, R., Hewitt, J., O'Donovan, M. C.,
352 Owen, M. J., Walters, J. T. R., & Kirov, G. (2017). Cognitive Performance Among Carriers
353 of Pathogenic Copy Number Variants: Analysis of 152,000 UK Biobank Subjects.
354 *Biological Psychiatry*, 82(2), 103-110. <https://doi.org/10.1016/j.biopsych.2016.08.014>
- 355 Kirov, G., Pocklington, A. J., Holmans, P., Ivanov, D., Ikeda, M., Ruderfer, D., Moran, J.,
356 Chambert, K., Toncheva, D., Georgieva, L., Grozeva, D., Fjodorova, M., Wollerton, R.,
357 Rees, E., Nikolov, I., van de Lagemaat, L. N., Bayés, A., Fernandez, E., Olason, P. I., . . .
358 Owen, M. J. (2012). De novo CNV analysis implicates specific abnormalities of
359 postsynaptic signalling complexes in the pathogenesis of schizophrenia. *Molecular*
360 *Psychiatry*, 17(2), 142-153. <https://doi.org/10.1038/mp.2011.154>
- 361 Klaiman, C., White, S. P., Saulnier, C., Murphy, M., Burrell, L., Cubells, J., Walker, E., & Mulle, J.
362 G. (2022). A distinct cognitive profile in individuals with 3q29 deletion syndrome. *J*
363 *Intellect Disabil Res*. <https://doi.org/10.1111/jir.12919>
- 364 Konieczny, M. R., Senyurt, H., & Krauspe, R. (2013). Epidemiology of adolescent idiopathic
365 scoliosis. *J Child Orthop*, 7(1), 3-9. <https://doi.org/10.1007/s11832-012-0457-4>
- 366 Kuettel, J., Bay, R. C., & McIsaac, T. L. (2022). Configuration variability of the six-minute walk
367 test among licensed physical therapists working with neurologic conditions: a pilot
368 survey. *Physiotherapy Theory and Practice*, 1-17.
369 <https://doi.org/10.1080/09593985.2022.2140318>
- 370 Marshall, C. R., Howrigan, D. P., Merico, D., Thiruvahindrapuram, B., Wu, W., Greer, D. S.,
371 Antaki, D., Shetty, A., Holmans, P. A., Pinto, D., Gujral, M., Brandler, W. M., Malhotra, D.,
372 Wang, Z., Fajardo, K. V. F., Maile, M. S., Ripke, S., Agartz, I., Albus, M., . . . Schizophrenia
373 Working Groups of the Psychiatric Genomics, C. (2017). Contribution of copy number
374 variants to schizophrenia from a genome-wide study of 41,321 subjects. *Nature*
375 *Genetics*, 49(1), 27-35. <https://doi.org/10.1038/ng.3725>

- 376 Martin, N., Snodgrass, G., & Cohen, R. (1984). Idiopathic infantile hypercalcaemia--a continuing
377 enigma. *Archives of Disease in Childhood*, *59*(7), 605-613.
- 378 Morava, E., Lacassie, Y., King, A., Illes, T., & Marble, M. (2002). Scoliosis in velo-cardio-facial
379 syndrome. *J Pediatr Orthop*, *22*(6), 780-783.
- 380 Morris, C. A., Demsey, S. A., Leonard, C. O., Dilts, C., & Blackburn, B. L. (1988). Natural history of
381 Williams syndrome: physical characteristics. *The Journal of Pediatrics*, *113*(2), 318-326.
- 382 Mulle, J. G. (2015). The 3q29 deletion confers >40-fold increase in risk for schizophrenia.
383 *Molecular Psychiatry*, *20*(9), 1028-1029. <https://doi.org/10.1038/mp.2015.76>
- 384 Mulle, J. G., Dodd, A. F., McGrath, J. A., Wolyniec, P. S., Mitchell, A. A., Shetty, A. C., Sobreira, N.
385 L., Valle, D., Rudd, M. K., Satten, G., Cutler, D. J., Pulver, A. E., & Warren, S. T. (2010).
386 Microdeletions of 3q29 confer high risk for schizophrenia. *American Journal of Human*
387 *Genetics*, *87*(2), 229-236. <https://doi.org/10.1016/j.ajhg.2010.07.013>
- 388 Naranje, S. M., Erali, R. A., Warner, W. C., Jr., Sawyer, J. R., & Kelly, D. M. (2016). Epidemiology
389 of Pediatric Fractures Presenting to Emergency Departments in the United States. *J*
390 *Pediatr Orthop*, *36*(4), e45-48. <https://doi.org/10.1097/bpo.0000000000000595>
- 391 Pollak, R. M., Murphy, M. M., Epstein, M. P., Zwick, M. E., Klaiman, C., Saulnier, C. A., the Emory
392 3q29 Project, & Mulle, J. G. (2019). Neuropsychiatric phenotypes and a distinct
393 constellation of ASD features in 3q29 deletion syndrome: results from the 3q29 registry.
394 *Mol Autism*, *10*(1), 30. <https://doi.org/10.1186/s13229-019-0281-5>
- 395 R Core Team. (2008). R: A language and environment for statistical computing. *R Foundation for*
396 *Statistical Computing, Vienna, Austria*. <https://www.r-project.org/>
- 397 Ren, X., Yang, N., Wu, N., Xu, X., Chen, W., Zhang, L., Li, Y., Du, R. Q., Dong, S., Zhao, S., Chen, S.,
398 Jiang, L. P., Wang, L., Zhang, J., Wu, Z., Jin, L., Qiu, G., Lupski, J. R., Shi, J., . . . Liu, P.
399 (2020). Increased TBX6 gene dosages induce congenital cervical vertebral malformations
400 in humans and mice. *Journal of Medical Genetics*, *57*(6), 371-379.
401 <https://doi.org/10.1136/jmedgenet-2019-106333>
- 402 Ricchetti, E. T., States, L., Hosalkar, H. S., Tamai, J., Maisenbacher, M., McDonald-McGinn, D.
403 M., Zackai, E. H., & Drummond, D. S. (2004). Radiographic study of the upper cervical
404 spine in the 22q11.2 deletion syndrome. *J Bone Joint Surg Am*, *86*(8), 1751-1760.
405 <https://doi.org/10.2106/00004623-200408000-00020>
- 406 Sanchez Russo, R., Gambello, M. J., Murphy, M. M., Aberizk, K., Black, E., Burrell, T. L., Carlock,
407 G., Cubells, J. F., Epstein, M. T., Espana, R., Goines, K., Guest, R. M., Klaiman, C., Koh, S.,
408 Leslie, E. J., Li, L., Novacek, D. M., Saulnier, C. A., Sefik, E., . . . Mulle, J. G. (2021). Deep
409 phenotyping in 3q29 deletion syndrome: recommendations for clinical care. *Genetics in*
410 *Medicine*, *23*(5), 872-880. <https://doi.org/10.1038/s41436-020-01053-1>
- 411 Sanders, S. J., He, X., Willsey, A. J., Ercan-Sencicek, A. G., Samocha, K. E., Cicek, A. E., Murtha, M.
412 T., Bal, V. H., Bishop, S. L., Dong, S., Goldberg, A. P., Jinlu, C., Keaney, J. F., 3rd, Klei, L.,
413 Mandell, J. D., Moreno-De-Luca, D., Poultney, C. S., Robinson, E. B., Smith, L., . . . State,
414 M. W. (2015). Insights into Autism Spectrum Disorder Genomic Architecture and Biology
415 from 71 Risk Loci. *Neuron*, *87*(6), 1215-1233.
416 <https://doi.org/10.1016/j.neuron.2015.09.016>
- 417 Sievert, C., Parmer, C., Hocking, T., Chamberlain, S., Ram, K., Corvellec, M., & Despouy, P.
418 (2017). plotly: Create Interactive Web Graphics via 'plotly.js'. *R package version 4.6.0*.
419 <https://CRAN.R-project.org/package=plotly>

- 420 Stagi, S., Lapi, E., Gambineri, E., Manoni, C., Genuardi, M., Colarusso, G., Conti, C., Chiarelli, F.,
421 de Martino, M., & Azzari, C. (2010). Bone density and metabolism in subjects with
422 microdeletion of chromosome 22q11 (del22q11). *Eur J Endocrinol*, *163*(2), 329-337.
423 <https://doi.org/10.1530/eje-10-0167>
- 424 Stefansson, H., Meyer-Lindenberg, A., Steinberg, S., Magnusdottir, B., Morgen, K., Arnarsdottir,
425 S., Bjornsdottir, G., Walters, G. B., Jonsdottir, G. A., Doyle, O. M., Tost, H., Grimm, O.,
426 Kristjansdottir, S., Snorrason, H., Davidsdottir, S. R., Gudmundsson, L. J., Jonsson, G. F.,
427 Stefansdottir, B., Helgadottir, I., . . . Stefansson, K. (2014). CNVs conferring risk of autism
428 or schizophrenia affect cognition in controls. *Nature*, *505*(7483), 361-366.
429 <https://doi.org/10.1038/nature12818>
- 430 Szatkiewicz, J. P., O'Dushlaine, C., Chen, G., Chambert, K., Moran, J. L., Neale, B. M., Fromer, M.,
431 Ruderfer, D., Akterin, S., Bergen, S. E., Kähler, A., Magnusson, P. K. E., Kim, Y., Crowley, J.
432 J., Rees, E., Kirov, G., O'Donovan, M. C., Owen, M. J., Walters, J., . . . Sullivan, P. F. (2014).
433 Copy number variation in schizophrenia in Sweden [Immediate Communication].
434 *Molecular Psychiatry*, *19*, 762. <https://doi.org/10.1038/mp.2014.40>
435 <https://www.nature.com/articles/mp201440#supplementary-information>
- 436 Taeusch, H., Ballard, R., & Gleason, C. (2004). *Avery's Diseases of the Newborn*. Saunders.
- 437 Wawrzonek, A. J., Sharp, W., Burrell, T. L., Gillespie, S. E., Pollak, R. M., Murphy, M. M., & Mülle,
438 J. G. (2022). Symptoms of Pediatric Feeding Disorders Among Individuals with 3q29
439 Deletion Syndrome: A Case-Control Study. *J Dev Behav Pediatr*, *43*(3), e170-e178.
440 <https://doi.org/10.1097/dbp.0000000000001009>
- 441 Westphal, F. L., Lima, L. C., Lima Neto, J. C., Chaves, A. R., Santos Júnior, V. L., & Ferreira, B. L.
442 (2009). Prevalence of pectus carinatum and pectus excavatum in students in the city of
443 Manaus, Brazil. *J Bras Pneumol*, *35*(3), 221-226. <https://doi.org/10.1590/s1806-37132009000300005>
- 444 Willatt, L., Cox, J., Barber, J., Cabanas, E. D., Collins, A., Donnai, D., FitzPatrick, D. R., Maher, E.,
445 Martin, H., Parnau, J., Pindar, L., Ramsay, J., Shaw-Smith, C., Sistermans, E. A.,
446 Tettenborn, M., Trump, D., de Vries, B. B. A., Walker, K., & Raymond, F. L. (2005). 3q29
447 microdeletion syndrome: clinical and molecular characterization of a new syndrome.
448 *American Journal of Human Genetics*, *77*(1), 154-160. <https://doi.org/10.1086/431653>
- 449 Wu, N., Ming, X., Xiao, J., Wu, Z., Chen, X., Shinawi, M., Shen, Y., Yu, G., Liu, J., Xie, H., Gucev, Z.
450 S., Liu, S., Yang, N., Al-Kateb, H., Chen, J., Zhang, J., Hauser, N., Zhang, T., Tasic, V., . . .
451 Zhang, F. (2015). TBX6 null variants and a common hypomorphic allele in congenital
452 scoliosis. *N Engl J Med*, *372*(4), 341-350. <https://doi.org/10.1056/NEJMoa1406829>
453
454

456 **Table 1. Demographics of study participants with 3q29del.**

	Mean	SD
Age	12.02	9.37
	N	%
Sex		
Male	32	56.14%
Female	25	43.86%
Race		
White	54	94.74%
Black	1	1.75%
Asian	2	3.51%
Ethnicity		
Not Hispanic/Latino	40	70.18%
Hispanic/Latino	7	12.28%
Unknown/not reported	10	17.54%
Informant relationship		
Biological parent	52	91.12%
Self	3	5.26%
Other	2	3.51%

457

458

459 **Table 2. Musculoskeletal phenotypes reported by participants with 3q29del.**

	N	%
Axial phenotypes		
Any axial phenotype	21	36.84%
Scoliosis	4	7.02%
Cervical pain for more than 1 month	3	5.26%
Thoracic pain for more than 1 month	1	1.75%
Lumbar pain for more than 1 month	3	5.26%
Pectus excavatum	13	22.81%
Pectus carinatum	3	5.26%
Sixth lumbar vertebra	0	0.00%
Extremity phenotypes		
Any extremity phenotype	39	68.42%
Joint stiffness	16	28.07%
Joint contracture	2	3.51%
Joint laxity	8	14.03%
Daily joint pain for more than 1 month	5	8.77%
Pes planus	23	40.35%
Long fingers	10	17.54%
Camptodactyly	2	3.51%
Nail hypoplasia	3	5.26%
Uniquely shaped fingers or toes	9	15.79%
Bone fracture	5	8.77%
Leg length discrepancy	0	0.00%
Neuromuscular phenotypes		
Any neuromuscular phenotype	34	59.65%
Gait anomalies	21	36.84%
Fatigue in 30 minutes or less of activity	28	49.12%

460

461

462 **Table 3. Imaging procedures reported by participants with 3q29del.**

Body part	Imaging procedure	N	%
Skull	X-ray	1	1.75%
	CT	4	7.02%
	MRI	7	12.28%
Neck	X-ray	1	1.75%
	CT	0	0.00%
	MRI	3	5.26%
Spine	X-ray	7	12.28%
	CT	0	0.00%
	MRI	3	5.26%
Shoulder	X-ray	2	3.51%
	CT	0	0.00%
	MRI	0	0.00%
Arm	X-ray	2	3.51%
	CT	0	0.00%
	MRI	0	0.00%
Elbow	X-ray	0	0.00%
	CT	0	0.00%
	MRI	0	0.00%
Hand	X-ray	5	8.77%
	CT	0	0.00%
	MRI	0	0.00%
Rib	X-ray	1	1.75%
	CT	0	0.00%
	MRI	0	0.00%
Pelvis	X-ray	2	3.51%
	CT	0	0.00%
	MRI	1	1.75%
Hip	X-ray	5	8.77%
	CT	0	0.00%
	MRI	0	0.00%
Leg	X-ray	1	1.75%
	CT	0	0.00%
	MRI	1	1.75%
Knee	X-ray	2	3.51%
	CT	0	0.00%
	MRI	2	3.51%

Ankle	X-ray	3	5.26%
	CT	0	0.00%
	MRI	0	0.00%
Foot	X-ray	3	5.26%
	CT	0	0.00%
	MRI	0	0.00%

463

464

465 **Table 4. Participant-reported musculoskeletal diagnoses confirmed by imaging procedures.**

Diagnosis	N	%
Bone fracture	5	8.77%
Scoliosis	4	7.02%
Leg length discrepancy	0	0.00%
Mass/tumor	0	0.00%
Sixth lumbar vertebra	0	0.00%
Other condition	4	7.02%

466

467 **Figure titles and legends**

468 **Figure 1. Musculoskeletal phenotypes reported by participants with 3q29del.** Rate of
469 musculoskeletal phenotypes across four categories reported by participants with 3q29del (n =
470 57).

471 **Figure 2. Specific musculoskeletal phenotypes reported by participants with 3q29del.** A) Rate
472 of congenital musculoskeletal phenotypes reported by participants with 3q29del (n = 57). B)
473 Rate of joint phenotypes reported by participants with 3q29del (n = 57). C) Amount of physical
474 activity until becoming fatigued reported by participants with 3q29del (n = 57).

475 **Figure 3. Medical interventions reported by participants with 3q29del.** Rate of medical
476 interventions received for musculoskeletal phenotypes reported by participants with 3q29del
477 (n = 57), showing high rates of medical care needed for musculoskeletal concerns.

100

medRxiv preprint doi: <https://doi.org/10.1101/2023.04.03.23288084>; this version posted April 3, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](#).

80

60

40

20

0


Percentage of respondents




Joint problems

Pain problems

Neuromuscular problems

Congenital anomalies





100

medRxiv preprint doi: <https://doi.org/10.1101/2023.04.03.23288084>; this version posted April 3, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](#).

80

60

40

20

0

Percentage of respondents



Orthopedic
evaluation



Orthopedic
surgery



Physical
therapy