1	Drivers of Early Childhood Vaccination Success in Nepal, Senegal, and Zambia: A
2	Multiple Case Study Analysis Using the Consolidated Framework of Implementation
3	Research
4	Authors: Cam Escoffery ^a , Emily Ogutu ^a , Zoe Sakas ^a , Kyra A. Hester ^a , Anna Ellis ^a , Katie
5	Rodriguez ^a , Chandni Jaishwal ^a , Chenmua Yang ^a , Sameer Dixit ^d , Anindya Bose ^e , Moussa
6	Sarr ^d , William Kilembe ^c , Robert A. Bednarczyk ^a , Matthew C. Freeman ^a
7	Affiliations:
8	^a Rollins School of Public Health, Emory University, Atlanta, GA, USA
9	^b Center for Molecular Dynamics Nepal, Katmandu, Nepal
10	^c Center for Family Health Research in Zambia, Lusaka, Zambia
11	^d Institut de Recherche en Santé de Surveillance Epidemiologique et de Formation
12	(IRESSEF), Dakar, Senegal
13	^e WHO Nepal Immunization Preventable Disease Division
14	* Corresponding author: Matthew C. Freeman; <u>matthew.freeman@emory.edu</u>
15	www.FreemanResearchGroup.org

16 Abstract

Introduction The fundamental components of a vaccine delivery system are welldocumented, but robust evidence is needed on *how* the related processes and implementation strategies - including the facilitators and barriers- contribute to improvements in childhood vaccination coverage. The purpose of this study was to identify critical facilitators and barriers to the implementation of common interventions across three countries that have dramatically increased coverage of early childhood

vaccination over the past 20 years, and to qualify common or divergent themes in theirsuccess.

Methods We conducted 277 key informant interviews and focus group discussions with public health leaders at the regional, district, and local levels and community members in Nepal, Senegal, and Zambia to identify intervention activities and the facilitators and barriers to implementation. We used thematic analysis grounded in the Consolidated Framework for Implementation Research (CFIR) to identify immunization program key facilitators and barriers.

Results We found that the common facilitators to program implementation across the 31 countries were the CFIR inner setting constructs of 1) networks and communications, 2) 32 goals and feedback, 3) relative priority, and 4) readiness for implementation; and outer 33 setting constructs of 4) cosmopolitanism and 5) external policies and mandates. The 34 common barriers were incentives and rewards, available resources, access to knowledge 35 and information, and patients needs and resources. Critical to the success of these 36 national immunization programs were prioritization and codification of health as a 37 human right, clear chain of command and shared ownership of immunization, 38 communication of program goals and feedback, offering of incentives at multiple levels, 39 training of staff central to vaccination education, the provision of resources to support the 40 program, key partnerships and guidance on implementation and adoption of vaccination 41 policies. 42

43 Conclusion Adequate organizational commitment, resources, communication, training,
44 and partnerships were the most critical facilitators for these countries to improve
45 childhood vaccination.

46

- 48 Keywords: childhood immunization, exemplars, implementation science, routine
- 49 immunization, immunization coverage

50 Introduction

Vaccination averts an estimated 4-5 million deaths annually; children, in particular, have 51 benefited from this protection against communicable diseases [1]. Countries have 52 reported progress in vaccine delivery over the years; however, immunization coverage has 53 varied widely among and within countries [2]. Several countries within Africa and 54 55 SouthEast Asia regions have outperformed their peers with significant increases in routine immunization coverage since 2000 – including Nepal, Senegal, and Zambia [3-56 6]. Understanding the factors that contributed to the success of these countries can add 57 to the existing literature and be adopted by other countries to improve their immunization 58 program performance. 59

To disseminate effective strategies to increase childhood immunization, there needs to be 60 61 understanding of what works (facilitators) to improve immunization performance among different countries. Identification of those factors can reduce childhood mortality, 62 medical expenditures and increase the future productivity of the country through the 63 longevity of their children. Implementation science is the study of methods to promote 64 65 the adoption and implementation of evidence-based interventions, strategies, or policies to public health and healthcare settings [7]. This provides us with a framework to help 66 understand the context of implementation, assess and improve public health 67 performance, and support and inform intervention scale-up [8]. The use of 68 implementation science frameworks to examine childhood vaccination programs in 69 countries with high immunization coverage rates provides an opportunity to identify and 70 describe factors that may have supported effective programs and childhood vaccination 71 improvements in low- and middle-income countries (LMICs). The Consolidated 72

Framework for Implementation Research (CFIR) is a widely applied framework that describes the context in which interventions and programs are delivered. CFIR is a meta theory that includes five domains: intervention characteristics, inner setting, outer setting, individual characteristics, and process (Table 1)[9]. It has been widely used since its creation and subsequent publication in 2009 [10], and can explain important factors related to the results of program delivery (e.g., implementation outcomes) [11].

A few studies have applied CFIR to examine children immunization programs 79 internationally among LMICs [12-14]. These studies used CFIR to explain facilitators and 80 barriers to implementation that impacted intervention effectiveness for vaccination 81 initiatives. Facilitators in program delivery included intervention flexibility, self-efficacy 82 of health workers, leadership support and resources (vaccine stocking), and cold chain 83 supervision (structural characteristics). Barriers included acceptability of the vaccine, 84 vaccine costs to mothers, vaccine hesitancy, inadequate cold chain infrastructure and lack 85 of incentives for health workers such as community health workers [12, 15]. A quantitative 86 evaluation study of the Global Polio Eradication Initiative (PEI) in Nigeria using CFIR 87 found that successful contributing factors of community engagement for PEI were the 88 external social environment, and political factors [14]. Application of the CFIR framework 89 may guide the design or selection of more appropriate interventions for a particular health 90 facility or community, identify critical factors at the national and subnational levels of 91 government that lead to better public health delivery. This can ultimately lead to greater 92 acceptability and adoption of vaccination protocols by healthcare providers, which can 93 contribute to increased levels of vaccine uptake [10, 16]. 94

The purpose of this study was to identify common interventions, and the critical facilitators and barriers to implementation using CFIR framework, that contributed to exemplary growth in routine early childhood vaccine coverage in Nepal, Senegal, and Zambia. Findings from this study may be used to develop actionable recommendations for improving immunization programming globally. We examined the CFIR inner setting (i.e., Ministry of Health) and outer setting (i.e, external partners and stakeholders) constructs that were related to vaccine service delivery and demand generation.

102 Methods and materials

Study design. The Exemplars in Vaccine Delivery research project focused on how Nepal, 103 Senegal, and Zambia succeeded in achieving catalytic growth in childhood routine 104 105 immunization coverage from 2000-2019. Details about the overall project and methodological approach can be found elsewhere [17, 18]. We used a multi-case study 106 design to explore critical determinants to the implementation of the national 107 immunization programs by applying CFIR, as described in Table 1. Among the five 108 domains of the CFIR framework, we focused on the relevance of the inner and outer 109 settings. 110

Our specific research questions for this paper were: 1) What are the key intervention strategies employed by the national vaccination programs? and 2) What are the facilitators and barriers related to the implementation of the vaccine programs?

Table 1. Inner and Outer Setting Domains of the Consolidated Framework for Implementation Research

Co	Consolidated Framework for Implementation Research Constructs				
Co	nstruct	Description			
II.	INNER SETTING				
A	Networks &	The nature and quality of webs of social networks and the			
	Communications	nature and quality of formal and informal communications			
		within an organization.			
В	Implementation climate	The absorptive capacity for change, shared receptivity of			
		involved individuals to an intervention, and the extent to			
		which use of that intervention will be rewarded, supported,			
		and expected within their organization.			
	B1. Organizational	Extrinsic incentives such as goal-sharing awards, performance			
Incentives & Rewards		reviews, promotions, and raises in salary, and less tangible			
		incentives such as increased stature or respect.			
	B2. Goals and Feedback	The degree to which goals are clearly communicated, acted			
		upon, and fed back to staff, and alignment of that feedback			
		with goals.			
	B3. Relative Priority	Individuals' shared perception of the importance of the			
		implementation within the organization.			
С	Readiness for	Tangible and immediate indicators of organizational			
	implementation	commitment to its decision to implement an intervention.			
	C1. Available Resources	The level of resources dedicated for implementation and on-			
		going operations, including money, training, education,			
		physical space, and time.			
	C2. Access to Knowledge &	Ease of access to digestible information and knowledge about			
	Information	the intervention and how to incorporate it into work tasks.			

I. O	UTER SETTING	
A	Patient Needs &	The extent to which patient needs, as well as facilitators and
	Resources	barriers to meet those needs, are accurately known and
		prioritized by the organization.
В	Cosmopolitanism	The degree to which an organization is networked with other
		external organizations.
С	External Policy &	A broad construct that includes external strategies to spread
	Incentives	interventions, including policy and regulations (governmental
		or other central entity), external mandates, recommendations
		and guidelines, pay-for-performance, collaboratives, and
		public or benchmark reporting.

116

Procedures. In consultation with national stakeholders and available data, we selected 117 three provinces within Nepal, Senegal, and Zambia, while considering population density, 118 vaccination coverage, and contextual factors [17]. Qualitative data were collected between 119 August 2019 and April 2021. Data were collected by the Center for Family Health 120 Research in Zambia (CFHRZ), Center for Molecular Dynamics Nepal (CMDN) in Nepal, 121 122 Institute de Recherche en Santé de Surveillance Epidémiologique et de Formation (IRESSEF) in Senegal, and the Emory University research team. The participants 123 included the Ministry of Health (MoH) staff, key partners, public health officers, local 124 organizations, and community members (Table 2). 125

Measures. Topic guides asked participants about facilitators and barriers to implementation, immunization activities and interventions, drivers for high routine immunization coverage, and key partners and their roles. Key informant interview (KII) and focus group discussion (FGD) guides were translated into local languages by research
assistants. All interview guides were piloted before use and adjusted iteratively
throughout data collection. Data was collected by a different team, other than the authors
and the names of the participants were not included. Data collection tools can be found
on Open Science Framework [19].

Data analysis. Key informant interviews and focus group discussions were audio-134 135 recorded. Recordings were transcribed verbatim into the language of the original audio 136 recording, and then translated into English. When audio recordings were in English, the recordings were transcribed directly. Data were password protected, and reviewed by 137 more than one person in the research team, to ensure accuracy and completeness in 138 deidentification. We created a codebook that was iteratively improved based on the 139 emerging themes [20, 21]. Codes were deductive based on CFIR constructs and open-140 ended questions. Participants were categorized based their role: 1) Ministry at the 141 national level, 2) Ministry at the subnational level, 3) Partners, 4) Local implementers, 142 and 5) Community members. Transcripts were coded and analyzed using MaxQDA2020 143 software (Berlin, Germany). We used the visual tool in MaxQDA to detect the frequency 144 of codes to select the major CFIR constructs within the inner and outer settings for this 145 analysis. For each theme, we described the valence of the construct to vaccination 146 147 coverage success: positive influence on implementation, negative, or mixed. At the individual transcript level, valence was determined by interviewees' accounts related to 148 the specific construct (e.g., available resources, external mandates, etc.). At the case level, 149 we considered whether the interviewees agreed with each other in terms of the constructs' 150 influence on vaccination. Finally, we assigned an overall valence across the interviewee 151 categories for constructs using methods recommended by others [22]. 152

Ethics. The study was considered exempt by the Institutional Review Board committee of 153 Emory University, Atlanta, Georgia, USA (IRB00111474) and approved by ethics 154 committees in each country; Nepal Health Research Council in Kathmandu, Nepal 155 (NHRC; Reg. no. 347/2019); the National Ethical Committee for Health Research 156 157 (CERNS; Comité National d'Ethique pour la Recherche en Santè) in Dakar, Senegal (00000174); and the University of Zambia Biomedical Research Ethics Committee 158 (Federal Assurance No. FWA00000338, REF. No. 166-2019) and the National Health 159 Research Authority in Zambia. All interviewees gave informed consent to participate in 160 the study. 161

162 **Results**

We conducted 277 activities in Nepal, Senegal, and Zambia (Table 2). Table 3 outlines key 163 interventions, programs, and policies across all three countries, organized by level of 164 implementation. At the national level, the most commonly implemented interventions 165 across the three countries were the introduction of new vaccines, presence of a national 166 immunization technical advisory board, expansion of health posts, and media campaigns. 167 At the subnational level, all countries reported use of the Reaching Every District (RED) 168 initiative [23], micro-planning, expansion of the vaccine cold chain, community health 169 worker training, monitoring and evaluation meetings, and vaccination campaigns. At the 170 local level, all employed community health workers, outreach efforts for vaccine 171 promotion, media outreach, micro-planning, defaulter tracing, community and religious 172 leadership engagement, and education of grandparents and/or parents using community 173 groups. 174

175 Table 2. Summary of countries, partner organizations, regions, districts and

176 data Collection Activities[24]

Country	Nepal Senegal		Zambia			
In Country Research Partner	Center for Dynamics,	Molecular Nepal	Institut de Recherche en Santé de Surveillance, de Surveillance Epidémiologique et de Formation (IRESSEF)		Center for Family Health Research in Zambia	
Data collection period (MM/YYYY)	8/2019 – 1	2/2019	12/2020 - 4/20	21	10/2019 - 02/2020	
Regions and Districts						
Region 1	Madhes*	Dhanusha, Bara, Mahottari	Ziguinchor	Ziguinchor, Oussouye, Diouloulou	Lusaka	Lusaka, Rufunsa, Chongwe
Region 2	Bagmati	Makwanpur, Dolakha, Kathmandu	Dakar	Rufisque, Mbao, Keur Massar	Central	Chibombo, Chitambo, Serenje
Region 3	Gandaki Pradesh	Kaski, Myagdi, Nawalparasi	Tambacounda	Tambacoun da, Koumpento um, Goudiry	Luapula	Chipili, Nchelenge, Samfya
Key Informant Interviews Key: Number of KIIs (Participants)	79	9 (79)	62 (63)	6	6 (85)
National level government staff	1	1 (11)	5 (5)	1	1 (12)
Partner organization staff	8	8 (8)	4 (4)	1	1 (15)
Regional health staff		5 (5)	7(7)		6 (8)
District health staff	15 (15)		38 (38)		10 (19)	
Health facility staff	2	3 (23)	6 (6)		7 (10)	
Community leaders	1	5 (15)	2 (2)		10 (10)	
Community health workers **	:	2 (2)	-		1	11 (11)
Focus Group Discussions Key: Number of FGDs (Participants)	30	(191)	19 (1	28)	22	2 (132)
Community health workers **	9	(60)	10 (65)	1	0 (60)
Mothers	9	(60)	9 (6	53)	8	8 (48)
Fathers	ϵ	6 (36)	-			1 (6)
Grandparents	6	5 (35)	-		3 (18)	
Total (per country)	109	9 (270)	81 (191)		88 (217)	
Total (across countries)		-	-	,	27	8 (678)

- 178 *Includes volunteer community health workers, female community health volunteers (FCHV),
- 179 vaccinators, bajenu gox, and neighborhood health committee members.

180 Table 3. Immunization Programming from Key Informants, Organized by

181 **Country and Level of Implementation**

Level of	Policy or intervention	_		
implementation		Nepal	Senegal	Zambia
•				
National level	Introduction of new	v	v	v
programming and	vaccines	Λ	Α	Δ
policies	Strategic reallocation of	x		x
	targeted funds			
Includes partners and	Health post expansion	Х	Х	Х
national government	Codifying health as a	x	x	
	human right		Λ	
	Forums for decision	x	x	x
	making (e.g., ICC)			
	Media engagement	Х	Х	Х
Sub-national	RED/REC implementation	Х	Х	Х
programming	Training CHWs and	x	x	x
	volunteers	**		
Regional and district	Meetings for evidence-	v	v	v
level	based decisions	Λ	Λ	Δ
	Data software, health posts		Х	
	Data software, district-level	X	Х	Х
	Cold chain expansion	Х	Х	Х

Community programming	Community health worker program	X	X	Х
Health post and	Community-level health committees	X	X	X
village/community	Outreach services	Х	Х	Х
level	Micro-planning in health facilities*	X	X	X
	Defaulter tracing**	Х	Х	Х
	Engagement of community leaders	Х	х	х
	Support groups for caregivers	Х	X	Х
	School outreach	Х	Х	Х
	Promotion through media (e.g., radio)	Х	х	х

182 *Micro-planning- integrated set of components prepared to support health system.

**Defaulter tracing- visiting, calling, messaging mothers to remind them about vaccinating their
children.

185 Facilitators to implementation of vaccination programs

Several CFIR constructs were integral to the implementation of immunization programs across the three countries. Of the 14 CFIR Inner Setting constructs, 3 were found to be facilitators to vaccine delivery: 1) networks and communication, 2) implementation climate, which includes: relative priority, incentives and rewards, goals and feedback, and

190 3) readiness for implementation, including available resources and access to knowledge.

191 Two CFIR domains - implementation climate and readiness for implementation - were

192 central to the facilitating factors. However, incentives and rewards, available resources,

193 access to knowledge and information, and patients needs and resources constructs were

- also barriers to implementation of immunization at the local and community level. Table
- 195 4 summarizes findings categorized by CFIR constructs and country.

196 Table 4. Salient CFIR Inner and Outer Setting Constructs: Facilitators and

197 Barriers to Implementation of Routine Vaccination Programming

Construct	Nepal	Senegal	Zambia
Inner Setting (Ministry of Health)		•	
Networks and communication			
The nature and quality of webs of social networks and	+	+	+
the nature and quality of formal and informal			
communications within an organization.			
Implementation climate			
Relative priority			
Individuals' shared perception of the importance of the	+	+	+
implementation within the organization.			
Incentives and Rewards			
Extrinsic incentives such as goal-sharing awards,	+/-	+/-	+/-
performance reviews, promotions, and raises in salary,			

and less tangible incentives such as increased stature or			
respect.			
Goals and Feedback			
The degree to which goals are clearly communicated,			
acted upon, and fed back to staff, and alignment of that	+	+	+
feedback with goals.			
Readiness for implementation	<u> </u>	<u> </u>	<u> </u>
Available resources			
The level of resources dedicated for implementation and			. /
on-going operations, including money, training,	+/-	+/-	+/-
education, physical space, and time.			
Access to knowledge and information			
Ease of access to digestible information and knowledge	+/-	+/-	+/-
about the intervention and how to incorporate it into		+/ -	+/ -
work tasks.			
Outer Setting	1	<u> </u>	<u> </u>
Patient Needs and Resources			
The extent to which patients' /community members'			
needs, as well as facilitators and barriers to meet those	+/-	+/-	+/-
needs, are accurately known and prioritized by the			
organization.			
Cosmopolitanism	+	+	+

The degree to which an organization is networked with			
other external organizations (external networks,			
partners)			
External policies and mandates			
External strategies to spread interventions, including			
policy and regulations, external mandates,	+	+	+
recommendations and guidelines, pay-for-performance,			
collaboratives, and public or benchmark reporting.			

198 + = Facilitator to implementation. - = Barrier to implementation. +/- = Both facilitator and barrier

199 Inner Setting

We found 3 inner setting constructs- 1) network and communications, 2) implementation climate (relative priority, incentives and rewards, and goals and feedback), and 3) readiness for implementation (available resources and access to knowledge and information) as salient facilitators to implementation of vaccination programs in Nepal, Senegal, and Zambia (Table 4). A few representative quotes related to these themes can be found below with additional quotes located in Open Science Framework [19].

206 1. Network and communications

Key informants across all countries viewed communication and coordination between staff at the national and subnational levels as facilitators to successful implementation of childhood vaccinations. Interventions to foster communication and coordination in all three countries included 1) frequent meetings to discuss vaccine data, review and identify improvements, and improve data quality; 2) communication channels between levels of government that support shared ownership of immunization activities; 3) micro-planning at district and community levels to align priorities and tailor strategies; and 4) a clearchain of command that facilitated the flow of information.

"On the relationship side, it is always the sharing of information, we receive 215 instructions or directives. We try to apply them as much as possible, and in return 216 we make our reports, across the medical region, and we always report the 217 difficulties we find at the peripheral level, on the ground and elsewhere. 218 Whenever we need it, we feedback [from the national level] on everything we had 219 expressed as a point for improvement, recommendation. And from time to time, 220 we still feel that the central level is very close to the peripheral level." (National 221 Chief Medical Officer, Senegal) 222

223 2. Implementation climate

Three sub constructs of implementation climate were facilitators of implementation of the immunization programs. These included relative priority, access to knowledge, and incentives and rewards.

Relative Priority. Relative priority is the shared perception of importance of the public 227 health topic and/or program, and it can lead to adoption and/or implementation of public 228 229 health interventions by program leaders and staff. Public health professionals perceived 230 that there was strong political will and commitment by the government to conduct childhood immunization programming in all three countries. Informants from Senegal 231 and Nepal reported codification of health as a human right in their constitution, and in 232 Zambia, equitable access to quality healthcare as a national priority as included in the 233 current Vision 2030 Zambia strategic document. In Nepal, an informant mentioned that 234

immunization programming was prioritized by the government through implementation
of the existing policies that support immunization. In Zambia, there were discussions of
having specific budget line for immunization at the national level, showing the level of
commitment by the Ministry of Health to improve immunization.

"I think I may have mentioned this at one point that the government has been
procuring the vaccines. In fact, we started own procurement [of vaccines] in
2004. We assumed the total budget for vaccines procurement at that time. A
dedicated budget line for vaccine procurement for vaccines was even established
within the yellow book. That helped to make sure we had commodities readily
available within the facilities. (Former EPI manager, Zambia)

"It is the commitment of government to conduct the program and commitment
of policy makers and decision makers, after that commitment of all the health
workers working in implementation level. Commitment and involvement of the
social organizations and bodies in vaccination program, along with that
commitment in the management of vaccines and equipment where it is
necessary." (Former Director General, Nepal)

Incentives and Rewards. All three countries reported use of incentives to motivate personnel to improve their performance and increase their vaccination coverage. These included motivating both health workers to improve their staff performance and caregivers to increase their demand for vaccination. In Zambia, supervisors provided trophies to high-performing facilities, districts, and provinces and partners offered nonmonetary incentives to communities (e.g., certificates). Health professionals were primarily motivated to vaccinate children to reduce disease among children and ensure

healthy communities. Nepal offered incentives at multiple levels: a reward system existed 258 at the district level, and local community health workers were sometimes given a stipend 259 during vaccination campaigns, and they attended training and conferences. Villages, 260 districts and provinces were declared fully immunized and celebrated by the national 261 262 government through the Full Immunization Declaration (FID) Program in a bottom-up approach, and this motivated local government bodies, service providers, community 263 workers, volunteers and parents and facilitated community buy to immunization. 264 Physicians offered postnatal gifts like mirrors, mosquito nets and other incentives to 265 mothers who got their child vaccinated. In Senegal, recognition was common at different 266 levels, including certificates, upward mobility for high performing facility health workers, 267 and awards and recognition for revenue of health posts. 268

"Staff performance is assessed and there's even an incentive to reward staff if
they are performing well at the end of a time. But initially, I think there is every
once in a month where we do the Labor Day celebrations where they award the
staff that are deserving. So, they awarded five staff that are doing well and they
give award just to motivate those staff..." (National Chief Cold Chain Officer,
Zambia)

We organized for Labor Day; we organized a small gathering. Everyone was
there, the service providers, the ASC (community health worker), the bajenu gox,
the community actors in general, and there were certificates of recognition that
were decided beforehand. We identified the recipients, and these names were put
on the certificates. During these gatherings, the doctor gives the certificates along

280 with an envelope of (25,000 francs)." (Sexual and Reproductive Health 281 Coordinator, Senegal)

However, a lack of tangible incentives (money and awards to staff), and inconsistent motivation of community health workers were highlighted as barriers in Nepal and Senegal. Tangible incentives, especially those that were planned in advance (monthly Labor Day celebrations, field allowance or training, for instance) were facilitators.

"They are not at all motivated, it's just when there is training that's the source of
motivation or if there is campaign and others but apart from that they have no
specific motivation." (MoH Staff member, Senegal)

289 "The health workers are the same who worked previously in health institutions 290 so there are not much differences. However, they don't get field allowance now 291 that is why motivation level is quite low." (District Health Officer, Nepal)

Goals and Feedback. All three countries had mechanisms for communicating their 292 immunization goals to ensure feedback was shared across all levels. The national level 293 provided instructions, directions, and support for logistics and capacity building. Public 294 health professionals at the subnational level often applied learning from training and 295 logistical support. In all countries, feedback is given from the national to provincial and 296 to the district level at regular intervals (i.e., monthly, annually). In Zambia, Provincial 297 Medical Officers present their program data to the central level on a quarterly basis -298 including the epidemiology of vaccine preventable diseases - and the national level 299 provides feedback to the provinces. Informants in Nepal stated that feedback is provided 300 to low coverage areas or low performing health facilities, and that their public health staff 301 employ micro-planning and categorization of the health facilities based on that 302

performance. In Senegal, informants indicated that performance was assessed at the start
 of the month with objectives and targets set for each health post. Feedback on indicators
 and achievements were provided from the central and provincial levels during monthly
 meetings.

- 307 "The central level calls the Provincial Medical Officers, on a quarterly basis, 308 sometimes even more often to come and present their data looking at the 309 programs, including EPI, and feedback is given..." (Director of Public Health, 310 Zambia)
- "Recently we have done review at provincial level, which is done yearly. In June 311 each district level data was observed. What is the status level of every district, is 312 everything scheduled? All these were discussed and at the upcoming years what 313 decisions should be made, to make vaccination programs more fruitful what 314 should be done in entire districts. We worked along with family welfare 315 departments and discussion are made. They [family welfare departments] are 316 our responsibilities, and they belong to this group and based on that review is 317 done." (Immunization supervisor, Nepal) 318
- 319 *3. Readiness for implementation*

Three sub-constructs of readiness for implementation were perceived as being both facilitators and barriers to the implementation of the immunization programs. These included available resources, access to knowledge, and incentives and rewards.

Available Resources. Available resources was a strong theme, and consistent across the
 countries; this included budgeting for the national immunization programs, cold chain

infrastructure, and a vaccination workforce. In Zambia, key informants at the sub-325 national and local implementation level mentioned availability of staff and volunteers as 326 an enabler for increases in vaccine coverage and sustainability. In addition, cold chain 327 improvements were essential for increases in vaccine coverage and was prioritized by the 328 329 MOH, who advocated for funding from the Ministry of Finance and from donors to support the infrastructure. In Nepal and Senegal, the government provides the main 330 startup money for the immunization program and there is substantial support by 331 international donor organizations. At the subnational and community levels in Nepal, 332 the local government allocated funds to train and incentivize health workers, maintain 333 health facilities, and hire cold chain technicians. In Senegal, external partners such as 334 GAVI and USAID are mentioned as working at all levels, and some ministry staff noted 335 336 that they fill in the gap where the district or health development committee cannot fund activities. 337

"Basically, the Ministry of Finance decides how much money goes to all the other
sectors but when it comes to the EPI, at least we have some specific budget lines
and we actually do contribute towards coming up with those plans and budgets
but ultimately is the Ministry of Finance that, who determine how much goes to
what." (MoH staff member, Zambia)

³⁴³ "Now ... I cannot say it for sure ... but since last two decades, it [vaccine rate] ³⁴⁴ started to show improvement. We can also say after external support started to ³⁴⁵ come like GAVI is our partner ... they have played great role, even they help ³⁴⁶ community financially to strengthen system they have played major role in it. ³⁴⁷ They are globally big donor in field of vaccine. They have contributed much in

Nepal. Talking about partners in Nepal, UN agencies is UNICEF main mandatory, it works mainly for women and children, they also provide technical support to us and WHO are also providing technical support as well." (National level stakeholder, Nepal)

However, participants at the sub national level reported a lack of available resources which resulted in consistent barriers across the three countries. These constraints included financing for the immunization program, procurement of vaccines, transportation for vaccine provision, human resources, lack of vaccination facilities, and inadequate vaccine supplies.

"At the district level, there are health posts that are suffering in terms of
equipment, and health posts that are suffering from the building. Others are
suffering from the resources of transport for the nurses in charge of vaccination."
(MoH staff member, Senegal)

Access to Knowledge and Information. Across all countries, training, and quality 361 assurance methods were critical. In Zambia, multiple trainings around the Reaching 362 Every District (RED) were mentioned. RED is a WHO strategy to increase vaccination 363 rates through building capacity to increase vaccine delivery by establishing outreach 364 services, planning and management of resources, monitoring and use of data, and linkage 365 of services with the community [23]. Likewise, Senegal stakeholders discussed that 366 knowledge is passed from the national level to lower levels through training sessions. 367 368 modules, and manuals. Some district level staff reported organizing and conducting training for community-level workers. 369

"The first thing you do is train the districts in RED (Reaching Every District and 370 Every Child). Then they do all the micro plans, everything in the district and we'll 371 also go and train them. Let me start with, you have to conduct the TOT, the 372 training of trainers. So, we identify the trainers within the districts and 373 provinces. So, eventually we went to train the districts and the districts are going 374 to train the facilities. So, we go and monitor the facilities assuming they are 375 implementing RED and developing all the micro plans and everything. And that 376 is a continuation in other interventions that are coming on board, RED is also a 377 priority. RED is always a training and RED is always a priority." (Chief Cold 378 Chain Officer, Zambia) 379

Nevertheless, this construct of access to knowledge was discussed across all countries as a common barrier at the levels closer to the community. There was limited, or infrequent, vaccination training and educational community outreach to all staff, organizations, or partners. Education was sometimes provided to only selected staff, or training information was not disseminated to others in the organization or partners.

³⁸⁵ "The training was for one week only. ...It was just presentations by a member of ³⁸⁶ staff from the clinic, ... The selected group then came here at the clinic to be ³⁸⁷ trained on what to do and say to the people in the community. Yes madam, that ³⁸⁸ is all that they did to train us. (Neighborhood Health Committee member, ³⁸⁹ Zambia)

"Training is not given much nowadays. The municipality organizes training only
once a year." (Vaccinator, Nepal)

"It may not be very frequent, but we are given training every 6 months."
(Community health worker, Senegal)

394 Outer Setting

According to key informants, cosmopolitanism, and external policies and mandates were the two outer setting factors recognized as facilitators for implementation of the vaccine program. Other representative quotes related to these themes are found at Open Science Framework web page [19].

399 *Cosmopolitanism*

Cosmopolitanism, the extent to which an organization (e.g., the Ministry of Health) is 400 connected to external partnership or networks, was instrumental in implementation. At 401 each country's national level, there were similar organizations that aided in the 402 implementation of the immunization program. These partners included WHO, UNICEF, 403 and GAVI which offered technical support and financial resources across the three 404 405 countries. In Nepal, WHO and UNICEF fostered a high trust, constant communication, 406 and collaboration and agreed upon division of labor. At the local level, MoH workers collaborated with religious leaders, schools, NGOs, the media, and community health 407 workers (CHWs) to increase vaccination in all the countries. In Nepal and Zambia, school 408 directors and teachers shared information about immunization with children in school, 409 while Nepal focused on strong media partnerships. In Senegal, community-based 410 organizations and neighborhood delegates supported implementation. 411

412 413 "In major vaccination, of Polio campaign to support government technically, financially there was WHO and UNICEF. UNICEF majorly supported

government in capacity development, logistic management, and in few types of
equipment; and after that the global organization such as GAVI, Lions started to
support us and inside Nepal Rotary Club, Lions Club started to support us in Polio
campaign as a sensitizer and mobilize but in government program there was
major support of WHO and UNICEF, where GAVI has constantly supporting us."
(Former Director General, Nepal)

"Like here at our clinic, letters are written...to different zones to make people 420 aware that there will be a vaccination program, again there are letters written 421 to churches that are close by and also to all those that come to this clinic. So, most 422 women or grandmothers are aware of when to bring children here for 423 vaccination. Posters are also stuck for all to know when the vehicle has past. 424 maybe some are at work, but when they return and look on the trees, walls, and 425 all over the markets, they stick them all over even at the shops...even in schools 426 they announce, school children take the message to parents." (Grandmothers, 427 Zambia) 428

429 External Policies and Mandates

External policies and mandates may include political directives, regulations, and external
mandates. Many of the immunization efforts are performed within a WHO, GAVI, or
UNICEF global policy framework. External agencies gave guidance to country leaders
who needed their support and approval to implement policies or adopt new vaccine
technologies.

435 "Nepal hasn't become capable of making policies on its own but as it has made an 436 international commitment, WHO guides us to make necessary changes in the

437 policies. Hence, the decisions are made on that basis. Of that, the regulations are
438 formulated, and according to this." (Ministry of Finance personnel, Nepal)
439 "The most important is the ACD plan, which has become the ACE plan (aka -

- 440 *RED/REC*). Through this plan of Reaching Every District, it is the activities of the
- 441 districts and of the health posts that are funded to enable the position to perform."
- 442 *(EPI and PHC focal point, Senegal)*
- 443 Patient Needs and Resources

Patient needs and resources was reported as a facilitator and barrier to program delivery. 444 This construct identifies whether program implementers are aware of community needs 445 and how they respond to those needs. In all the countries, participants mentioned that 446 they worked together with the healthcare workers to address their needs, and this 447 included the selection of community health workers. These CHWs came from close by 448 communities and understood the needs of communities better. Participants reported that 449 parents in all countries were aware of diseases that could be prevented through childhood 450 vaccination. Participants discussed the changes in infrastructure including better 451 transportation and roads which facilitated access to healthcare. In Senegal, participants 452 reported a mobile strategy where healthcare providers traveled to rural patients to 453 provide vaccination services. Healthcare workers and community actors raised 454 awareness and creating demand for vaccination. Outreach and transportation initiatives 455 were identified as barriers to community needs that the Ministries of Health. 456

"By the way, I think there are a lot of people who are aware of different diseases,
so they have voluntarily gone to vaccinate their children, they know the interest
of vaccination. Certainly, there are people who until now do not believe in

vaccination, but there are also us, community actors, who go every day to the
population to make the population aware of the interests of vaccination. But also,
there are the opinion leaders, who are at the level of neighborhood delegates and
the bajenu gox and others who are involved in relation to vaccination." (Head
nurse for Health Post, Senegal)

465 "The main reason for the existing good immunization coverage is the awareness 466 among people regarding vaccination, which is given to prevent different 467 diseases. Other reasons are the provision of good quality vaccine, the availability 468 of health workers, accessibility of health services, availability of vaccine, road 469 accessibility. In the past, people had to walk for a day to reach the immunization 470 center, but now due to the transportation facility, people can reach there in an 471 hour or even less." (MoH staff member, Nepal)

Interviewees identified unmet needs as a barrier to implementation of immunization
programming. Community barriers included families facing economic hardship, a lack of
knowledge about childhood vaccination, reliance on traditional healers or medicine,
vaccine hesitancy, low literacy, and language barriers.

476 "And the next thing is about the language. In the Hilly area they have different
477 languages and in the Terai area they have different languages. We make the
478 programs in one language, so it does not match with other areas and people do
479 not understand the language." (Former Director General, Nepal)

"It would be nice to explain to mom the benefits of the vaccine her child just took
because most moms are illiterate. You have to take the time to talk to them about
the vaccine and its benefits because otherwise the mother will be discouraged, she

483 will never come back." (Mother, Senegal)

484 **Discussion**

Our findings describe how Nepal, Senegal, and Zambia achieved high routine 485 immunization coverage through investigating facilitators and barriers to program 486 implementation and how they relate to CFIR constructs. CFIR constructs were critical to 487 understanding the success of these countries in increasing childhood immunization. 488 Although there is no "silver bullet" intervention to enhance early childhood vaccination, 489 490 our findings point to critical facilitators that exist at national, sub-national, and local levels of vaccine delivery. The application of the CFIR model enabled a comprehensive 491 492 assessment of implementation context; the use of this framework to assess vaccination programming could be expanded to support adaptation of successful implementation 493 strategies and factors for other countries. 494

Many facilitators to implementation of the national immunization program were shared
by key informants from these exemplar countries. Our findings complement existing
literature that supported the use of CFIR to explore factors related to implementation.

Support from international partners- GAVI, WHO, USAID, UNICEF and others, and local community partners- religious leaders, traditional leaders, local media stations, community based organizations and others, in service provision and delivery have led to successful rollout of childhood vaccines in low-income countries [25, 26]. Adoption and implementation of policy initiatives or mandates like Reaching Every District is instrumental in enhancing vaccination coverage. Within these countries, districts are required to submit plans annually to reach every child to receive funding from the

505 ministries for their vaccination activities through assessment of data, identification of 506 problems and creation of a work plan. This strategy is endorsed by WHO and has assisted 507 LMIC in achieving high coverage rates of childhood vaccinations [27]. Although all 3 508 countries did report that external policies were facilitators, they also reported that access 509 to knowledge, patient needs, and available resources were mixed, and usually negative at 510 the community levels, thus the need for consideration of local context, priorities, and 511 needs alongside external policies.

We identified some common barriers across the countries related to immunization 512 delivery. Some respondents, mostly at regional or local levels, believed that training was 513 missing, not frequent enough, or reached various levels of implementation at national to 514 less local regions. Several studies have demonstrated the impact of training on delivery of 515 immunization programs [15, 28]. Some challenges to vaccination at the community level 516 included language, geography, education or poverty status. In analyzing implementation 517 barriers and strategies in polio eradication initiative in the Democratic Republic of Congo 518 519 and Ethiopia, Deressa et al. found the common barriers include accessibility issues 520 (population movement, geography); gaps in human resources; supply chain; finance and governance and community hesitancy [29]. The CFIR constructs of available resources, 521 training of healthcare and other staff, and patient needs and resources are commonly 522 identified barriers in vaccination studies [16, 30, 31]. An understanding of the contextual 523 factors may support unearthing the systemic issues around immunization. Further 524 research on optimal interventions to increase vaccine confidence and reduce community 525 level barriers can inform further efforts to sustain public demand for childhood vaccines 526 [32, 33].527

The findings from this review have various practical implications for how to enhance the 528 implementation of childhood immunization programs. First, prioritization of health or 529 vaccines by ministries or governments, provision of resources for vaccination programs, 530 supplies, and workforce at all levels and facilities are essential to vaccine program 531 532 implementation [34]. Other qualitative analyses have found insufficient funds and resources and staffing issues as barriers to immunization[35, 36]. Secondly, coordination 533 and engagement of different levels of the country (e.g., ministry, districts/subnational, 534 and local community) and stakeholders was crucial to public education to increase 535 community demand and rollout of the programs [4-6, 37]. Related to this, communication 536 and feedback loops and incentives provided motivation for reaching immunization goals; 537 this has been identified in another study focused on immunization delivery [38]. Third, 538 539 incentives were helpful in motivating staff and provinces or regions to improve vaccination rates. Finally, partnership engagement at all public health levels and 540 community stakeholder engagement is critical to the success of these programs. 541

542 Strengths and limitations

There were several strengths to this study. CFIR informed the qualitative instruments 543 and identified shared factors across countries to fill a gap in the literature related to 544 optimal implementation of vaccine programs globally. It helped examine factors internal 545 to the ministries and public health programs and also external agents that facilitated 546 successful vaccine delivery. Employing CFIR was a strength because it assists in assessing 547 successful determinants of immunization program implementation and barriers to 548 implementation. The stakeholder groups included in this qualitative study were diverse 549 and included the community level to understand their assessment of critical factors that 550

lead to childhood vaccinations. Often, implementation research focuses on program implementers (e.g., public health leaders, implementation staff) and does not include levels of implementation (regional to local) and community members (intervention recipients). Future research could assess other CFIR domains (i.e., process, characteristics of individuals) and their association with implementation success.

This study has several limitations. The data collection instruments focused on the factors 556 that drove catalytic change and did not elicit policies or interventions that were 557 unsuccessful. We focused on the CFIR constructs and sub-constructs of inner and outer 558 setting post-implementation for the majority of this analysis; recall of activities may be 559 more valid with data collection occurring during vaccine program implementation. The 560 CFIR framework has been updated; future research could explore how other domains of 561 CFIR (e.g., individual, process) and new constructs (e.g., critical incidents, local 562 conditions, etc.) contributed to successful immunization program and outcomes such as 563 sustainment of effective intervention strategies employed by these countries [39]. Using 564 565 qualitative methods to understand historical events was challenging; interviewees often spoke about current experiences rather than discussing historical factors. However, 566 research assistants probed respondents to reflect on longitudinal changes in the 567 568 immunization program.

569 **Conclusion**

570 This multiple case study analysis presents the opportunity to explore implementation 571 science determinants that were critical to the successful implementation of childhood 572 vaccination programming in three countries with high immunization coverage. CFIR's

comprehensive and multifaceted domains help capture the complexities of multilevel 573 interventions to increase childhood vaccination. The use of CFIR helps inform why these 574 countries have high vaccination coverage by describing facilitators and barriers to 575 programming implementing immunization at the national. regional. 576 and 577 local/community level. We identified facilitators such as communication of goals and feedback, offering of incentives at multiple levels, training staff central to vaccination 578 education, providing resources to support the program, and maintaining key 579 partnerships. Public health and governmental staff may bolster direct resources to 580 support the immunization infrastructure, communications, collaborations and incentives 581 to improve vaccination. This study identifies a wide range of facilitators and barriers to 582 implementation of vaccine programs across Nepal, Senegal, and Zambia which 583 584 contributes to the limited literature on implementation research in global vaccination.

585 Acknowledgements

We thank Center for Molecular Dynamics, Nepal (CMDN), the Institut de Recherche en 586 Santé de Surveillance Epidemiologique et de Formation (IRESSEF) in Dakar, Senegal and 587 the Center for Family Health Research in Zambia for their partnership in this study. We 588 gratefully acknowledge the participants who gave their time and insights to help us better 589 understand Nepal's, Senegal's, and Zambia's vaccine delivery system, along with 590 facilitators from the Ministry of Health and Population, Department of Health Services, 591 and the Family Welfare Division of Nepal, Ministry of Health and Social Action in 592 Senegal, and the Ministry of Health in Zambia for supporting this research. In addition, 593 we thank Sarah Chesemore, Anna Rapp, Tove Ryman, and Ethan Wong from the Bill and 594 Melinda Gates Foundation; Kate Buellesbach, Nancy Fullman, Nathaniel Gerthe, Gloria 595

- 596 Ikilezi, Caitlyn Mason, David Phillips, and Oliver Rothschild, Jordan-Tate Thomas, and
- 597 Angela Wang from Gates Ventures for technical support; and the Vaccine Exemplars
- 598 Research Advisory Group for their insights, specifically Agnes Binagwaho, Laura Craw,
- 599 Carolina Danovaro, Anuradha Gupta, Heidi Larson, Penelope Masumbu, Kate O'Brien,
- 600 Helen Rees, Lora Shimp, and Aaron Wallace.

601 Data Availability Statement

All relevant data are within the paper and its supporting information files.

603 Funding

This work was supported by the Bill & Melinda Gates Foundation, Seattle, WA
(OPP1195041) with a planning grant from Gates Ventures, LLC, Kirkland, WA. Bill and
Melinda Gates Foundation and Gates Ventures were engaged in the project process,
provided context, when necessary, but all research activities and publications are solely
those of the authors.

609

610 **Competing Interests**

611 The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

613 **References**

614 WHO. Immunization 2019 [Available from: https://www.who.int/news-room/facts-in-1. 615 pictures/detail/immunization. 616 2. WHO. Immunization Agenda 2030 A global strategy to leave no one behind. 2020. 617 WHO. WHO-UNICEF estimates of DTP3 coverage 2020 [updated 15-Jul-2020; cited 2021 23 3. 618 November]. Available from: 619 https://apps.who.int/immunization monitoring/globalsummary/timeseries/tswucoveragedtp3.html. 620 Micek K, Hester KA, Chanda C, Darwar R, Dounebaine B, Ellis AS, et al. Critical success factors for 4. 621 routine immunization performance: A case study of Zambia 2000 to 2018. Vaccine X. 2022;11:100166. 622 5. Hester KA, Sakas Z, Ellis AS, Bose AS, Darwar R, Gautam J, et al. Critical success factors for high 623 routine immunization performance: A case study of Nepal. Vaccine: X. 2022;12:100214-. 624 6. Sakas Z, Hester KA, Rodriguez K, Diatta SA, Ellis AS, Gueye DM, et al. Critical success factors for 625 high routine immunization performance: A case study of Senegal. medRxiv. 2022:2022.01.25.22269847. 626 7. Eccles MP, Mittman BS. Welcome to Implementation Science. Implementation Science. 627 2006;1(1):1. 628 8. Peters DH, Adam T, Alonge O, Agyepong IA, Tran N. Implementation research: what it is and 629 how to do it. BMJ : British Medical Journal. 2013;347:f6753. 630 Damschroder LJ, Aron DC, Keith RE, Kirsh SR, Alexander JA, Lowery JC. Fostering implementation 9. 631 of health services research findings into practice: a consolidated framework for advancing 632 implementation science. Implementation Science. 2009;4(1):50. 633 10. Means AR, Kemp CG, Gwayi-Chore M-C, Gimbel S, Soi C, Sherr K, et al. Evaluating and optimizing 634 the consolidated framework for implementation research (CFIR) for use in low-and middle-income 635 countries: a systematic review. Implementation Science. 2020;15(1):1-19. 636 Nilsen P. Making sense of implementation theories, models, and frameworks. Implementation 11. 637 Science 30: Springer; 2020. p. 53-79. 638 12. Adamu AA, Uthman OA, Gadanya MA, Wiysonge CS. Using the consolidated framework for 639 implementation research (CFIR) to assess the implementation context of a quality improvement 640 program to reduce missed opportunities for vaccination in Kano, Nigeria: a mixed methods study. Hum 641 Vaccin Immunother. 2020;16(2):465-75. 642 Boisson A, Thompson P, Fried B, Shea CM, Ngimbi P, Lumande F, et al. Determinants of the 13. 643 uptake of childhood immunization in Kinshasa Province, the DRC: ordered logit regression analyses to 644 assess timely infant vaccines administered at birth and six-weeks of age. 2022. 645 14. Akinyemi OO, Adebayo A, Bassey C, Nwaiwu C, Kalbarczyk A, Fatiregun AA, et al. Assessing 646 community engagement in Nigeria polio eradication initiative: application of the Consolidated 647 Framework for Implementation Research. BMJ open. 2021;11(8):e048694. 648 Boisson A, Morgan CE, Fried B, Shea CM, Yotebieng M, Ngimbi P, et al. Barriers and facilitators 15. 649 to timely birth-dose vaccines in Kinshasa Province, the DRC: a qualitative study. Journal of Global Health 650 Reports. 2022;6:e2022028. 651 16. Selove R, Foster M, Mack R, Sanderson M, Hull PC. Using an Implementation Research 652 Framework to Identify Potential Facilitators and Barriers of an Intervention to Increase HPV Vaccine 653 Uptake. J Public Health Manag Pract. 2017;23(3):e1-e9. 654 17. Bednarczyk RA, Hester KA, Dixit SM, Ellis AS, Escoffery C, Kilembe W, et al. Protocol: 655 Identification and evaluation of critical factors in achieving high and sustained childhood immunization 656 coverage in selected low- and lower-middle income countries. medRxiv. 2021:2021.12.01.21267018. 657 18. Health EiG. Making Better Decisions in Global Health: Understand Positive Outliers to Inform 658 Policy and Practice. 2023 [Available from: https://www.exemplars.health/. 659 19. OSF. Exemplars in Vaccine Delivery2022.

660 20. Hennink MM. Focus group discussions: Oxford University Press; 2013.

Miles MB, Huberman AM, Saldana J. Qualitative Data Analysis: A Methods Sourcebook: SAGE
Publications; 2019.

66322.Damschroder LJ, Lowery JC. Evaluation of a large-scale weight management program using the664consolidated framework for implementation research (CFIR). Implementation Science. 2013;8(1):51.

665 23. WHO. Essential Programme on Immunization: Reaching Every District (RED) 2023 [Available

from: <u>https://www.who.int/teams/immunization-vaccines-and-biologicals/essential-programme-on-</u>
 immunization/implementation/reaching-every-district-(red).

Sakas Z, Hester KA, Ellis AS, Ogutu EA, Rodriguez K, Bednarczyk RA, et al. Critical success factors
for high routine immunization performance: A multiple case study analysis of Nepal, Senegal, and
Zambia. medRxiv. 2022:2022.11.08.22282076.

Gatera M, Bhatt S, Ngabo F, Utamuliza M, Sibomana H, Karema C, et al. Successive introduction
of four new vaccines in Rwanda: High coverage and rapid scale up of Rwanda's expanded immunization
program from 2009 to 2013. Vaccine. 2016;34(29):3420-6.

Ahonkhai AA, Odusanya OO, Meurice FP, Pierce LJ, Durojaiye TO, Alufohai EF, et al. Lessons for
strengthening childhood immunization in low-and middle-income countries from a successful publicprivate partnership in rural Nigeria. International Health. 2022;14(6):632-8.

677 27. WHO. Microplanning for immunization service delivery using the Reaching Every District (RED)
678 strategy. World Health Organization; 2009.

679 28. Mpabalwani E, Menon J, Phiri G, Malambo A, Mbozi E, Kalesha P, et al. Assessing the delivery
680 and effectiveness of a new immunisation training initiative at district level in Zambia. Medical Journal of
681 Zambia. 2011;38(1):8-12.

29. Deressa W, Kayembe P, Neel AH, Mafuta E, Seme A, Alonge O. Lessons learned from the polio
eradication initiative in the Democratic Republic of Congo and Ethiopia: analysis of implementation
barriers and strategies. BMC Public Health. 2020;20(4):1807.

68530.Escoffery C, Riehman K, Watson L, Priess AS, Borne MF, Halpin SN, et al. Facilitators and Barriers686to the Implementation of the HPV VACs (Vaccinate Adolescents Against Cancers) Program: A

687 Consolidated Framework for Implementation Research Analysis. Prev Chronic Dis. 2019;16:E85.

Soi C, Gimbel S, Chilundo B, Muchanga V, Matsinhe L, Sherr K. Human papillomavirus vaccine
 delivery in Mozambique: identification of implementation performance drivers using the Consolidated
 Framework for Implementation Research (CFIR). Implementation Science. 2018;13(1):151.

Lane S, MacDonald NE, Marti M, Dumolard L. Vaccine hesitancy around the globe: Analysis of
 three years of WHO/UNICEF Joint Reporting Form data-2015-2017. Vaccine. 2018;36(26):3861-7.

693 33. Cooper S, Betsch C, Sambala EZ, McHiza N, Wiysonge CS. Vaccine hesitancy – a potential threat
694 to the achievements of vaccination programmes in Africa. null. 2018;14(10):2355-7.

4. LaFond A, Kanagat N, Steinglass R, Fields R, Sequeira J, Mookherji S. Drivers of routine
immunization coverage improvement in Africa: findings from district-level case studies. Health Policy
Plan. 2015;30(3):298-308.

698 35. Ezezika O, Mengistu M, Opoku E, Farheen A, Chauhan A, Barrett K. What are the barriers and
699 facilitators to polio vaccination and eradication programs? A systematic review. PLOS Global Public
700 Health. 2022;2(11):e0001283.

701 36. Nkwenkeu SF, Jalloh MF, Walldorf JA, Zoma RL, Tarbangdo F, Fall S, et al. Health workers'

perceptions and challenges in implementing meningococcal serogroup a conjugate vaccine in the

routine childhood immunization schedule in Burkina Faso. BMC public health [Internet]. 2020

704 2020/02//; 20(1):[254 p.]. Available from: <u>http://europepmc.org/abstract/MED/32075630</u>

705 <u>https://doi.org/10.1186/s12889-020-8347-z</u>

706 <u>https://europepmc.org/articles/PMC7031928</u>

707 https://europepmc.org/articles/PMC7031928?pdf=render.

70837.Sakas Z, Hester K, Ellis A, Ogutu EA, Rodriguez K, Bednarczyk R, et al. Critical success factors for709high routine immunization performance: A multiple case study analysis of Nepal, Senegal, and Zambia.

710 medRxiv; 2022.

711 38. Guan TH, Htut HN, Davison CM, Sebastian S, Bartels SA, Aung SM, et al. Implementation of a

neonatal hepatitis B immunization program in rural Karenni State, Myanmar: A mixed-methods study.

713 PLoS One. 2021;16(12):e0261470.

39. Damschroder LJ, Reardon CM, Opra Widerquist MA, Lowery J. Conceptualizing outcomes for use

with the Consolidated Framework for Implementation Research (CFIR): the CFIR Outcomes Addendum.

716 Implementation Science. 2022;17(1):7.