| 1 | Word count: 4897 excl. ref. | | | | | | |
|---------|--|--|--|--|--|--|--|
| 2 | Fig. 7 | | | | | | |
| 3 | Tables 3 | | | | | | |
| 4 | | | | | | | |
| 5 | Blood biomarkers in Takotsubo syndrome point to an emerging role for inflammaging in | | | | | | |
| 6 | disease pathophysiology | | | | | | |
| 7 | Nagai, M. ¹ ⁺ ⁺ , Shityakov, S. ² ⁺ , Smetak M ³ , Hunkler, H.J. ⁴ , Bär ^{4,5,6} , C., Schlegel, N. ⁷ , | | | | | | |
| 8 | Thum, T. ^{4,5,6} , Förster C. Y. ³ * | | | | | | |
| 9 10 | ¹ Department of Cardiology, 2-1-1, Kabeminami, Aaskita-ku, Hiroshima City Asa Hiroshima 731-0293, JAPAN | | | | | | |
| 11 | ² Infochemistry Scientific Center, Laboratory of Chemoinformatics, Lomonosova str. 9, Saint- | | | | | | |
| 12 | Petersburg, 191002 Russian federation. | | | | | | |
| 13 | ³ University of Würzburg, Department of Anaesthesiology, Intensive Care, Emergency and Pain | | | | | | |
| 14 | Medicine, Würzburg, Germany | | | | | | |
| 15 | ⁴ Institute of Molecular and Translational Therapeutic Strategies (IMTTS), Hannover Medical School, | | | | | | |
| 16 | Hannover, Germany | | | | | | |
| 17 | ⁵ REBIRTH-Centre for Translational Regenerative Medicine, Hannover Medical School, Hannover, | | | | | | |
| 18 | Germany | | | | | | |
| 19 | ⁶ Fraunhofer Institute for Toxicology and Experimental Medicine (ITEM), Hannover, Germany | | | | | | |
| 20 | ⁷ University of Würzburg, Department of General, Visceral, Transplant, Vascular and Pediatric | | | | | | |
| 21 | Surgery, Würzburg, Germany | | | | | | |
| 22 | [†] These authors contributed equally to this work. | | | | | | |
| 23 | * Correspondence: MN nagai10m@r6.dion.ne.jp | | | | | | |
| 24 | CF foerster_c@ukw.de | | | | | | |
| 25 | | | | | | | |
| 26 | Running title: molecular catecholamine effects on endothelial and myocardial cells | | | | | | |
| 27 | | | | | | | |
| 28 | Abstract: Takotsubo syndrome (TTS), an acute cardiac condition characterized by transient wall | | | | | | |
| 29 | motion abnormalities mostly of the left ventricle, results in difficulties in diagnosing patients. We set | | | | | | |
| 30 | out to present a detailed blood analysis of TTS patients analyzing novel markers to understand the | | | | | | |
| 31 | development of TTS. Significant differences in pro-inflammatory cytokine expression patterns, sex | | | | | | |
| 32 | steroid and glucocorticoid receptor (GR) expression levels were observed in the TTS patient | | | | | | |

33 collective. Remarkably, the measured catecholamine serum concentrations determined from TTS patient blood could be shown to be two orders of magnitude lower than the levels determined from 34 35 experimentally induced TTS in laboratory animals. Consequently, exposure of endothelial cells and 36 cardiomyocytes in vitro to such catecholamine concentrations did not damage the cellular integrity or 37 function of neither endothelial cells forming the blood brain barrier, endothelial cells derived from 38 myocardium nor cardiomyocytes in vitro. Computational analysis was able to link the identified blood 39 markers, specifically, the pro-inflammatory cytokines and glucocorticoid receptor GR to microRNA 40 (miR) relevant in the ontogeny of TTS (miR-15), and inflammation (miR-21, miR-146a), respectively. 41 Amongst the well-described risk factors of TTS (older age, female gender), inflammaging-related 42 pathways were identified to add to these relevant risk factors or prediagnostic markers of TTS.

43

Keywords: Takotsubo syndrome, catecholamines, inflammation, aging, inflammaging, glucocorticoid
receptor, cytokine, epinephrine, norepinephrine, dopamine

46

47 1. Introduction

48 Takotsubo syndrome (TTS), also known as broken heart syndrome, stress induced cardiomyopathy, 49 and Takotsubo cardiomyopathy, is an acute cardiac syndrome with rapid onset of chest pain and dyspnea. ^{1,2} TTS is often triggered by physical and/or emotional stress and is characterized by a 50 51 transient and reversible severe left ventricular dysfunction which typically recovers spontaneously within hours to weeks.³ TTS is a remarkably like acute coronary syndrome (ACS) with almost the 52 53 same clinical presentations and ST elevations; therefore, differential diagnosis is critically important 54 in the emergency department. The prevalence of TTS has been reported to be approximately 2-3% of all patients with clinical appearance of ACS.^{3,4} 55

At present TTS-related heart failure is believed to be chiefly induced by a so-called catecholamine 56 (CAT) storm, e.g. increased circulating and myocardial CAT levels with myocardial toxicity.⁵⁻⁸ 57 58 Central to this unifying hypothesis, the abnormal left ventricular contraction pattern is attributed to 59 this CAT surge which triggers left ventricular wall motion abnormality, e.g. widespread dyskinesia in the apical segments and hyperkinesia in the basal segment of left ventricle (LV) with apical 60 ballooning ^{3,4}; available information is however controversial ⁹⁻¹¹: As a pathophysiological 61 mechanism, it is suggested that local differences in left ventricular (LV) adrenergic receptors may 62 underly this LV wall motion abnormarity.^{12,13} In fact, experimental studies showed that the LV in 63 64 canine has β 2-adrenoceptors (β 2-AR) that are expressed much more in the apical than in the basal

segments,¹⁴ the signaling pathways leading to this aetiopathogenesis at any rate remain unclear. Feola
et al. further supported this hypothesis with a myocardial PET study showing decreased coronary flow
reserve and impaired metabolism in the apical segments during the acute phase of TTS.¹⁵

68 Additional possible pathophysiological mechanisms have been suggested such as induced switch in 69 signal trafficking, and autonomic nervous system dysfunction with cardiac sympathetic activation

70 including the mentioned overstimulation of beta receptors.^{13, 16-19} Female gender and age-related lack

71 of female sex hormones are further well known characteristics of TTS patients.²⁰

72 Given that TTS is remarkably like ACS with almost the same clinical presentation and ST elevations;

valuation valuation of the valuation of

74 of onset. No ECG criteria have been identified that reliably discriminate between TTS and ST-

rsegment elevation myocardial infarction.²¹ The ECG changes are transient and resolve within months
in most cases.

Recent studies have identified microRNAs (miRNAs/miRs) as potentially promising sensitive
and specific biomarkers for cardiovascular diseases such as TTS, including a unique signature of miR1, miR-16, miR-26a and miR-133a, which differentiates TTS patients from STEMI patients.^{22,23}

Using blood samples from real-life patient cohorts, we set out to comprehensively quantify and 80 81 characterize CAT levels and other blood-borne biomarkers (markers of inflammation, physical/ 82 emotional stress hormone and endogenous sex hormone levels and their receptors) to identify a set of 83 novel TTS-associated biomarkers, pointing to inflammaging-associated triggers for TTS rather than 84 direct CAT effects on cardiomyocytes and endothelial cells of the cardiovascular and cerebrovascular 85 segments. By the help of systems biology we delineate new targets and specific molecular pathways 86 which will allow the development of precision medicine in the prevention and therapy of TTS in the 87 future.

88

89 2. Materials and Methods

90 Study population and human patient serum

Patient serum preparation from human blood was performed as described previously.^{24,25} Briefly, human blood was collected using S-Monovette tubes (Sarstedt). We sampled blood from 25 pseudonymized patients and subsequently divided into 5 (6 for cytokine measurements) different groups: postmenopausal healthy patients, n=3, premenopausal healthy patients, n=3, ACS acute (<6 weeks), n=4, TTS acute (first 2 weeks after onset), n= 4, TTS subacute (2-6 weeks after onset), n=8, TTS chronic (>6 weeks), n=3. All patients were diagnosed according to the InterTAK Diagnostic Criteria.²⁶ As healthy controls subjects were selected that did not present with altered coronary arteries

98 in the coronary angiography diagnostics, while minor cardiovascular or endocrinological diseases
99 were accepted (e.g.: arterial hypertension, diabetes mellitus).

For serum preparation, all the blood was drawn from subjects in using S-Monovette collection tubes (Sarstedt) and incubated at room temperature for 60 min. Then, the blood samples were centrifuged at $102 \quad 1500 \times g$ for 15 min, the serum was isolated, immediately frozen and stored at -80°C until the extraction. All the methods used in this study were performed in accordance with the relevant guidelines and regulations. Serum samples were quantified and analyzed individually.

- 105
- 106 Ethics approval and consent to participate

107 The study protocol was approved by the Hiroshima City Asa Hospital Research Committee (01-3-3)
108 as previously described.²⁴

109

110 Study design - Human patient serum

- 111 Patient blood serum preparation was performed as described previously.^{24,25}
- 112

113 Biochemical analyses

Blood samples were drawn after 10-min rest in the supine position in the morning. Adrenaline, noradrenaline, and dopamine concentrations in the plasma were assessed by high-performance liquid chromatography (SRL Inc., Tokyo, Japan). Serum cortisol concentrations were determined by electrochemiluminescence immunoassay (Elecsys®, Roche Diagnostics, Basel, Switzerland). While testosterone was measured using radioimmunoassay kits, 17β -estradiol was measured using an ultrasensitive radioimmunoassay kit (Elecsys®, Roche Diagnostics, Basel, Switzerland).

120

121 Endothelial cell cultures

The mouse brain capillary endothelial cell line cEND and MyEND were immortalized and isolated as described previously²⁷⁻²⁹ and cultured in Dulbecco's modified Eagle's medium (DMEM) with high glucose (Sigma) supplemented with 1-glutamine, MEM-vitamin solution, non-essential amino acids (NEA), Na pyruvate, penicillin/streptomycin (P/S) (all from Sigma) 10% heat-inactivated fetal calf serum (FCS) as described previously.³⁰

- 127
- 128 Human iPSC maintenance and cardiac differentiation

Human induced pluripotent stem cells (iPSC)³³ were cultivated in StemMACS medium (StemMACS 129 iPS-Brew XF with supplement, Miltenyi Biotec, #130-104-368) on Geltrex (Gibco, A1413302) and 130 differentiated into cardiomyocytes via Wnt modulation as described before.³⁴⁻³⁶ Briefly, iPSCs were 131 grown to confluency and passaged with Versene (Gibco, 15040066) in StemMACS medium with 2 132 133 µM Thiazovivin (Selleckchem, S1459). Directed cardiomyocyte differentiation was initiated at 70-134 80% confluency with 5 µM GSK-3 inhibitor XVI (Merck, 361559) in cardio differentiation medium 135 (250 mg human recombinant albumin (Sigma-Aldrich, A9731), 100 mg L-AA (L-ascorbic acid 2-136 phosphate sesquimagnesium salt hydrate, Sigma-Aldrich, A8960) in 500 ml RPMI medium (RPMI 137 1640 + GlutaMAX, Gibco, 72400047) for 48 h. Subsequently, medium was changed to cardio 138 differentiation medium supplemented with 5 mM IWP-2 (Selleckchem, S7085), a Wnt signaling 139 inhibitor for 48 h followed by medium changes with cardio differentiation medium every 48 h. At 140 differentiation day 8, when cells start to contract spontaneously, cardiomyocytes were cultured in 1x 141 B-27 (Gibco, 17504001) in RPMI 1640 + GlutaMAX medium, with medium changes every 2 to 3 142 days.

143 Cardiomyocytes were purified by metabolic selection³⁷ with 4 mM DL-lactate (Merck, L4263, in 1 M 144 HEPES, Carl Roth, HN77.3) in no glucose RPMI medium (RPMI 1640, no glucose, Gibco, 145 11879020) with 250 mg human recombinant albumin and 100 mg L-AA. Experiments were 146 performed in iPSC-derived cardiomyocytes between day 55-80 of differentiation.

147

148 Catecholamine treatment in vitro

149 cEND and MyEND were treated with a physiological and supraphysiological CAT mix – 150 corresponding concentration c1-c3 as determined from TTS patient serum, comp. **table 1**, epinephrine 151 (Sigma-Aldrich, E4250), norepinephrine (Sigma-Aldrich, A0937) and dopamine (Sigma-Aldrich, 152 H8502) in the respective cell culture media for 24 h. Epinephrine and norepinephrine were dissolved 153 in 0.5 M HCl, dopamine in medium, for the dilutions medium was used.

For CAT treatment of iPSC-derived cardiomyocytes, human iPSC-derived cardiomyocytes were treated with a physiological mix of these CATs, epinephrine (Sigma-Aldrich, E4250), norepinephrine (Sigma-Aldrich, A0937) and dopamine (Sigma-Aldrich, H8502)), – corresponding concentration c3/ TTS acute levels as determined from TTS patient serum, comp. **table 1** in RPMI medium supplemented with 1x B-27 (Gibco, 17504001) for 24 h. Additionally to the CAT mix designated c3, cells were treated with a supraphysiological concentration of 5 mM isoprenaline (Sigma-Aldrich, 16504).³⁸

161

- 162 Gene expression analysis in cEND and MyEND cells
- 163 For RT-PCR, RNA isolation and Real-time RT PCR was performed as described.³⁰
- 164
- 165 Western blot analysis in cEND and MyEND cells
- 166 he images were taken with FluorChem FC2 Multi-imager II (Alpha Innotech). The ImageJ software
- 167 was used to determine the density of the protein bands.
- 168
- 169 Extraction of exosomes from human serum and western blot assay
- For extraction of the exosomes, 400 µl of serum was centrifuged (30 minutes, 4°C, 2000 rcf). 300µl of the supernatant was then mixed with 60µl of total exosome isolation reagent (Thermo Fisher Scientific, #4478360) and incubated for 30 minutes at 4°C. After centrifugation (10 minutes, RT, 10000 rcf), the pellet was mixed with 200µl RIPA Buffer containing a protease inhibitor cocktail (ROCHE).
- 175
- 176 antibodies
- 177 primary: GR H-300 (1:200, Santa Cruz Biotechnology, #sc-8992), mouse anti-claudin-5 (1:1000,
- 178 Thermo Fisher Scientific, #35-2500), guinea pig anti-occludin (1:100, Acris, #358-504), rabbit anti-
- 179 ZO-1 (1:1000, Thermo Fisher Scientific, #61-7300), anti-β-actin-HRP (1:25000, Sigma-Aldrich,
- 180 #A3854), anti-goat VE-cadherin (1:200, Santa Cruz Biotechnology, #sc-6458).
- 181 Secondary: goat anti-rabbit HRP (1:15000, System Biosciences, #EXOAB-CD63A-1) was used.
- and rabbit anti-human CD63 (1:1000, System Biosciences, #EXOAB-CD63A-1).
- 183
- 184 Gene expression analysis in cardiomycytes
- 185 RNA were isolated with QIAzol Lysis Reagent (Qiagen, 79306) and reverse transcribed with the 186 Biozym cDNA synthesis kit (Biozym, 3314710X) according to manufacturer's guidelines. Gene 187 expression levels were measured by RT-qPCR with iQ SYBR Green Supermix (Bio-Rad, 1708880). 188 The expression levels of NR4A1 were evaluated with the $\Delta\Delta$ CT method and normalized to the 189 expression of the housekeeper GUSB (β -glucuronidase).
- 190
- 191 Primer sequences (5' to 3'):
- 192 forward reverse
- 193 NR4A1 GTGTGTGGGGACAACGCTTC CCACAGGGCAGTCCTTGTT
- 194 GUSB GACACCCACCACCTACATCG CTTAAGTTGGCCCTGGGTCC

195

196 Cytokine Immunoassay

197 Cytokines INF γ , IL-2, IL-4, IL-6, IL-10, TNF α , EGF, IL-17, IL-17A, MIP3A, CCL20/MIP3A, 198 RANTES were measured in duplicates in the different patients serum by using a multi-analyte 199 immunoassay (Millipore, HCYTOMAG-60K) with Luminex® bead technology according to 200 manufacturer's instruction.

- 201
- 202 Computational analysis

203 The Cytoscape v.3.7.1 software was implemented to generate the human gene-miR interaction 204 pathways linked to the searched miRs (miR-15, miR 26a, miR-21, and miR-146a) and 205 proinflammatory cytokines (INFy, IL-2, IL-4, IL-6, IL-10, TNFa, EGF, IL-17, IL-17A, MIP3A, 206 CCL20/MIP3A, and RANTES). The miRTarBase and TargetScan databases were used for the miR-207 target gene interaction types together with a list of the searched miRs by using the CyTargetLinker app v4.10.39 For the gene-gene interaction network containing proinflammatory cytokines, the 208 GeneMANIA algorithm was applied comprising 911 networks and 24280 genes.^{40,41} Finally, the 209 210 merge tool as a part of the Cytoscape pipeline was implemented to merge previously generated 211 networks combining analyzed miRs and cytokines.

- 212
- 213 Analysis and statistics

214 Data are shown as mean and standard deviation (SD) of biological independent experiments. One-way 215 ANOVA with Dunnett's multiple comparison test or Tukey's t-Test as indicated in the figure legends 216 was performed to evaluated the statistical significance between untreated und treated cells with 217 GraphPad Prism 8. P < 0.05 was considered as significant, * P < 0.05, ** P < 0.01, *** P < 0.001.

218

219 **3. Results**

- 220 Quantification of catecholamines in serum of patients with different heart diseases
- The CAT noradrenaline, adrenaline and dopamine were measured in serum of postmenopausal healthy patients and patients admitted with TTS (acute and subacute phase). Significantly elevated noradrenaline $(5.23 \pm 5.69 \text{ pmol/l})$ vs. $(4.24 \pm 1.52 \text{ pmol/l})$ and dopamine $(0.80 \pm 1.06 \text{ pmol/l})$ vs. $(0.18 \pm 0.08 \text{ pmol/l})$ levels were detected in the TTS subacute group exclusively, all other levels did not differ significantly from the control group, postmenopausal healthy females (Table 1). Among the three CAT, noradrenaline showed highest absolute serum levels among the groups, while dopamine
- and adrenaline varied.

In the following, endothelial cells and cardiomyocytes were treated in vitro with the different CAT concentrations as determined from TTS acute and TTS subacute patient serum as opposed to pharmacological concentrations (concentration c1 pharmacological concentrations⁴²: 150 μ M dopamine, 1 μ M epinephrine, 1 μ M norepinephrine; c2 postmenopausal healthy control; c3 TTS acute; c4 TTS subacute, comp. Table 1).

233

234 The concentration-dependent effects of CAT on cEND brain endothelial cells in vitro

To investigate the concentration-dependent change in protein expression of barrier-forming proteins claudin-5, occludin, VE-cadherin and ZO-1 following CAT exposure in vitro, cEND cells were incubated with the different concentrations of the CAT dopamine, epinephrine and norepinephrine as determined from patient serum comp. Table 1 compared to an exposure to supraphysiological/ pharmacological concentrations ⁴² for 24h.

Changes in protein levels were determined by Western blot analysis and compared to the untreated control group (Figure 1). Only the supraphysiological/ parmacological c1 concentration showed significant effects on the protein expressions: ZO-1, VE-cadherin and occludin expression were significantly reduced (0.451 ± 0.130 -fold) and (0.186 ± 0.069 -fold). Surprisingly, claudin-5 expression was significantly increased (1.761 ± 0.312 -fold). cEND monolayer exposure to CATs in concentration ranges as as determined from TTS patient serum (c2, c3, c4) showed no significant change in the protein expression.

247

The concentration-dependent changes in mRNA expression were determined by qRT-PCR analysis and compared to the untreated control group (Figure 2). Only the supraphysiological/ parmacological c1 concentration showed significant effects on the mRNA expression. Claudin-5 expression was significantly reduced (0.610 \pm 0.059-fold). Also, VE-cadherin and ZO-1 showed a significant reduction (0.582 \pm 0.139-fold) and (0.478 \pm 0.085-fold). Exposure of cEND monolayers with CATs in concentrations as determined from TTS patient serum (c2, c3, c4) and occludin showed no significant change on the mRNA expression (Figure 2).

255

256 The concentration-dependent effects of catecholamines on MyEND cells

To investigate the concentration-dependent change in protein expression, MyEND cells were incubated with the different concentrations of the CATs dopamine, epinephrine and norepinephrine as described for Fig. 2 for 24h. Changes in protein levels were determined by Western blot analysis and compared to the untreated control group (Figure 3). Only the supraphysiological/ parmacological c1

concentration showed significant effects on the protein expressions: Claudin-5 expression levels were significantly decreased in this application (0.275 \pm 0.067-fold), and occludin expression was significantly reduced (0.686 \pm 0.122-fold). The other CAT concentrations applied as determined from TTS patient serum (c2, c3, c4) showed no significant change in the protein expression of claudin-5 or occluding (data not shown).

266

The concentration-dependent changes in mRNA expression were equally determined by RT-qPCR analysis and compared to the untreated control group (Figure 4). Claudin-5 (0.312 ± 0.09 -fold) and occludin (0.473 ± 0.055 -fold) expression were significantly reduced after incubation with CAT in concentration c1 (pharmacological). Occludin expression was significantly reduced as well after incubation with c1. VE-cadherin and ZO-1 showed a significant reduction after c1 incubation (0.438 ± 0.0259 -fold), respectively.

273

274 The concentration-dependent effects of CATs on iPSC-derived cardiomycytess

275 To investigate the effects of physiological concentrations of CATs (CAT) in TTS patients on the 276 myocardium, we treated human iPSC-derived cardiomycytes with CATs for 24 h. To determine 277 changes in cardiomyocytes upon CAT treatment, we analyzed the expression level of nuclear receptor 278 subfamily 4 group A member 1. This hormone receptor is well expressed in the heart and upregulated after β -adrenergic stimulation.¹² Additionally to the CATs, cardiomyocytes were treated with a 279 280 supraphysiological concentration of isoprenaline (Iso). As reported by Borchert et al. the treatment 281 with Iso lead to a significant increase of the cardiac stress marker, whereas physiological CAT 282 concentrations showed no alterations in cardiomyocyte stress levels (Fig. 1a). Multiples of the physiological CAT concentration $(10^3 \text{ and } 10^6 \text{ of the physiological concentration})$ lead to cardiac 283 stress indicated by increased expression of NR4A1 (Fig. 1b), 10⁹-fold of the physiological 284 285 concentration were lethal for the cardiomyocytes.

286

287 Quantification of cytokines and chemokines in patient serum

A previous study demonstrated that treatment with inflammatory cytokines (TNF α , Interleukin-6) leads to compromised endothelial barrier function and reduced or altered junctional protein expression established in vitro model of the BBB, cEND cells ⁴³ and in in vivo studies.⁴⁴ Based on these findings, we hypothesized that inflammation is central to the pathophysiology of TTS and explored the time course and persistence of detectable potentially involved pro-inflammatory cytokines. we

293 hypothesized that inflammation is central to the pathophysiology and natural history of Takotsubo 294 syndrome.

295

We compared the concentration of different cytokines and chemokines in the sera of premenopausal healthy patients, postmenopausal healthy patients, TTS acute, TTS subacute, TTS chronic and ACS (Table 2). Although no statistical significance was found, the concentration of TNF α , IL-2, IL-4 and IL-6 show a tendency to be elevated in the TTS acute group compared to all other groups. In comparison, in the TTS subacute group the concentration of MIP3A, RANTES, EGF, and IL-17 show a tendency to be elevated compared to all other groups. INF γ and IL-10 show higher levels in the postmenopausal healthy group compared to the other groups.

303

304 *Quantification of testosterone and estradiol in serum of patients with various heart diseases and* 305 *correlation between testosterone-to-estradiol ratio*

The sex hormones testosterone and estradiol were measured in serum of healthy postmenopausal patients and patients with TTS (acute, subacute and chronic). The different hormone levels were compared to healthy postmenopausal patients. There was no significant difference between the groups detectable (Table 3).

310

The following ratio between testosterone and estradiol (T:E2) as measured in the serum of patients with various heart diseases was calculated: mean T:E postmeno [5.77+0,2]; premeno diestrous $[0.5\pm$ 0.1]; postmeno estrous [8.1±3.3]; TTS acute [32±0.4]; TTS subacute [6.7±0.1]; TTS chronic [16.11±0.3]. As reference values for healthy premenopausal females the E2-values estrous / diestrous (400 pg/ml /50 pg/ml/) were used, respectively (University of Rochester, medicine labs).

316 The calculated T:E2 ratio was the highest in acute TTS serum (32-fold), followed by chronic TTS (26-

317 fold) and subacute TTS (8-fold). Lowest T:E2 ratios were measured in the healthy groups.

318

319 Protein expression of glucocorticoid receptor α in exosomes of serum from patients with various heart
320 diseases

321 Glucocorticoids are primary stress hormones necessary for life that regulate numerous physiologic 322 processes in an effort to maintain homeostasis.⁴⁵ Glucocorticoids are part of the feedback mechanism 323 in the immune system, which reduces certain aspects of immune function, such as inflammation. The 324 glucocorticoid receptor (GR) is a member of the nuclear hormone receptor superfamily that is

325 activated as a stress-responsive transcription factor upon binding glucocorticoids.⁴⁵ We set out to 326 determine the levels of GR α in patient blood.

For this, after extraction of the exosomes of serum from patients with various heart diseases, Western blot analysis of glucocorticoid receptor α was performed. Changes in protein levels were compared to the healthy postmenopausal group showing a clear trend towards elevation in all TTS group, remarkably the TTS acute and TT chronic groups. Remarkably, even though not significant, values determined from female TTS patients approximate the much higher serum GR levels from male (1,59 of healthy female premenopausal) than as determined from healthy females (postmenopausal healthy set =1; premenopausal healthy = 0.7 ± 0.55).

- 334
- 335 Bioinformatic Analysis

Next, we set out to relate the identified inflammatory cytokines present in TTS patient blood to published miRNA in TTS and aging and age-related disease and inflammaging and the CATs described to play a role in the etiology of TTS.^{5-8, 22, 23, 46, 47}

- A potential miRNA-cytokine-steroid hormone-receptor regulatory network contributing to TTS wasdeveloped:
- 341

342 The Cytoscape algorithm was able to reconstitute a detailed GMI network comprising 2989 nodes and 343 6000 edges associated with the corresponding miRs by utilizing the miRTarBase and TargetScan databases (Figure 5).³⁹ Next, the GGI network was built using the GeneMANIA algorithm for the 344 corresponding blood markers and proinflammatory cytokines comprising 211 nodes and 626 edges 345 (Figure 6).⁴¹ Finally, the merged network was compiled from the GMI and GGI networks containing 346 3152 nodes and 5735 edges to connect the blood markers, pro-inflammatory cytokines, and 347 glucocorticoid receptor gene (NR3C1) to miRs (Figure 7 [A]).⁴⁰ This network was processed by 348 349 subtracting the analyzed nodes to identify only direct gene-miR interactions (Figure 7 [B]). As a 350 result, IL-6, CCL5, CCL20, and INFy were predicted to be interacted with analyzed miRs directly, 351 while the rest of the genes were involved in indirect interaction with the investigated miRs.

352

353 4. Discussion

In the clinical practice, a few blood biomarkers are being used to diagnose and monitor TTS relative to the ACS which is characterised by almost the same clinical presentations and ST elevations. These biomarkers serve to predict response to treatment as well as long-term prognosis; blood troponin T,

357 BNP or creatine kinase-MB (CK-MB) are the most frequently surveyed markers.^{48,49} These markers

- 358 together with clinicopathological findings ^{50,51} do convey only insufficient diagnostic accuracy.
- 359
- 360 TTS patient serum catecholamine levels

361 Takotsubo patients present with supraphysiologic plasma CAT levels (epinephrine, norepinephrine, dopamine), which is generally believed to be a cardiotoxic trigger^{10,11} causal for the abnormal LV 362 contraction pattern in TTS.^{10, 11, 52} Available information remains however controversial^{9, 53} and 363 heavily relies on disease modeling in a dish³⁰ or using the isoprenaline rodent model of TTS³⁸ creating 364 365 a supraphysiological CAT exposure for the cell systems or animals investigated while CAT effects on 366 the cardiovascular system in narrower physiological ranges remain elusive. Supporting the CAT 367 hypothesis, some cross-field references can be found in the literature, e.g. pheochromocytoma-related conditions is a CAT-mediated myocarditis and focal/diffuse myocardial fibrosis condition.⁵⁴ Contrast 368 to this hypothesis, our in vitro assessments did not point to structural endothelial cell (from BBB, 369 370 myocard) or cardiomyocyte damage when exposed to CAT in concentrations as determined from TTS 371 patient blood.

372 Indirect CAT effects on the cardiovascular system in the pathogenesis of TTS should therefore be 373 considered: meanwhile it has been acknowledged that immune and central nervous systems mutually 374 influence on one another in a bidirectional way, eg in the physiological stress response.⁵⁵ Specifically, 375 the contribution of the locus coeruleus, the principal site for brain synthesis of norepinephrine in the 376 genesis of defeat-stress induced inflammatory signaling should be mentioned.⁵⁶

To conclude, CAT serum levels a reliable TTS biomarker, further research is needed to examine the link between CATs and TTS disease ontogeny. In the following we thus concentrated on an elucidation of disparate novel blood-borne biomarkers of stress and sex differences and inflammation in this stress-related disease.

381

382 TTS patient GR serum levels

We did observe a consistent trend towards elevated GR levels in TTS patient serum. Measured GR serum levels in TTS patient blood clearly exceeded the average postmenopausal healthy female levels 1.6-fold and came close to the values determined from male blood donors.

386 Generally, stress responses are inhibited by negative feedback mechanisms, whereby glucocorticoids 387 act to diminish drive (brainstem) and promote transsynaptic inhibition by limbic structures (e.g. 388 hippocampus) including feedback downregulation of GR.⁵⁷ In contrast, elevated GR ^{58,59} expression 389 has been described as a suitable surrogate marker and to positively correlate with glucocorticoid

resistance in chronic stress.⁶⁰ Chronic stress-induced activation of the hypothalamus-pituitary-adrenal axis takes many forms (chronic basal hypersecretion, sensitized stress responses, and even adrenal exhaustion), with manifestation dependent upon factors such as stressor chronicity, intensity, frequency, and modality and involves GR insensitivity and upregulation.⁵⁷ Neural mechanisms driving acute and chronic stress responses can be distinct by the serum GR expression, taking into account age, gender and reproductive cycle biases.⁶¹

Activation of the HPA-axis and sympathetic nervous system overdrive, which are established mechanisms critically involved in the TTS ontogeny are widely accepted to link specifically chronic stress with elevated levels of peripheral pro-inflammatory markers in blood.⁶² Yet, empirical evidence showing that peripheral *levels* of glucocorticoids and/or CATs mediate this effect is equivocal. Recent attention has turned to the possibility that cellular *sensitivity* to these ligands may enhance the effects of inflammatory mediators released under chronic stress.

402 The glucocorticoid resistance in conjunction with β 2-adrenergic receptor signaling pathways in 403 neuronal cells was presented to promote peripheral pro-inflammatory conditions that were postulated 404 to be associated with chronic psychological stress.⁶³

405 However, given the limited size of our study and patient cohort, further research is warranted to 406 examine the link between serum stress, $GR\alpha$ levels and TTS disease progression in female 407 postmenopausal patients.

408

409 Inflammation: cytokines and chemokines in patient serum

The existence of an immune-endocrine relation has been described, pointing out the immunomodulatory effects of the CATs epinephrine, norepinephrine and dopamine on a wide variety of immune functions.⁶⁴ The authors described that the immunomodulatory effects are mediated by specific adrenergic and dopaminergic receptors detectable on the surface of their immunological target cells and that CATs released within the CNS may chiefly influence the activity of the immune system in the periphery.⁶⁴

416 We explored expression levels and time course of expression of various pro- and antiinflammatory 417 cytokines while comparing their levels to healthy pre- and postmenopausal subjects plus ACS 418 patients.

419 Patients with acute TTS had higher serum concentrations of IL-6, CXCL1 (growth-regulated protein)

420 chemokine, TNF α , INF γ , IL-2 and IL-4 compared with control subjects and acute ACS patients, 421 respectively, while EGF-levels were decreased. This observation was in agreement with prior

422 observations published by Scally et al. ⁶⁵ and Santoro,⁶⁶ which also compared inflammatory patterns 423 in TTS and ACS.

424 Based on the data presented, monitoring the elevation of the cytokines IL-6, TNFα, and INFγ serum

425 levels might be suitable to distinguish TTS acute from ACS, again highlighting the importance of

426 inflammation in the etiology of TTS.

427 Novel acute phase markers identified include the elevation of both, IL-2 and IL-4. The exploitability

428 of IL-2 (increase) and IL-10 (decrease) as both disease phase markers and distinction markers

429 ACS:TTS needs further exploitation in bigger patient cohorts and warrants future potential as relapse430 indicators.

The evaluation of IL-2 could represent a novel, TTS acute phase marker, facilitating the discrimination from ACS. Specifically, as IL-2 has been characterized as an anti-inflammatory cytokine, the determination of its levels from patient serum might be a tool to potentially predict propensity to relapse.⁶⁷ However, taking into account the limited number of samples available, a follow-up study including higher sample numbers would be mandatory before final conclusion could be drawn about the suitability of IL-2 as TTS acute phase marker or to differentiate TTS from ACS, or its potential suitability as a relapse marker.

- 438 The subacute cytokine profile is most distinctively characterized by elevations in MIP3A, a so far not 439 yet addressed cytokine in frame of TTS, in EGF, $TNF\alpha$, IL-17 and, interestingly pronounced 440 decreases in IL-10.
- Amongst those, IL-17 might represent a future therapeutic target in TTS: Interleukin-17 (IL-17)
 induces the production of granulocyte colony-stimulating factor (G-CSF) and chemokines such as
 CXCL1 and CXCL2 and is a cytokine that acts as an inflammation mediator.⁶⁸
- 444 Taken together, while the elevation of IL-6, IL-2, IL-4, TNF α , INF γ and CXCL1 can be regarded as 445 TTS acute phase serum markers, a sharp increase in MIP3A and a pronounced decrease in IL-10 was 446 detectable in the TTS subacute, phase which culminated in the chronic phase but also in ACS. In 447 contrast, IL-10 levels were the highest in the postmenopausal healthy group.
- 448

449 Sex differences

The epidemiological link between female sex and the incidence of TTS cannot be fully explained by stress induced CAT or corticosteroid levels, only. Additional factors, such as female sex or hormone receptor regulation and dysregulated immune response could be an important factor for this specific form of heart failure in postmenopausal patients. Consistent with prior studies, we found strong

454 associations between reduced female sex steroid levels and further extended these studies by 455 calculating an T:E2 ratio which was elevated pronouncedly in TTS patients. Several points could be 456 taken into account:

Women mount stronger immune responses not only against foreign but also against selfantigens, and the prevalence of most autoimmune diseases (AD) is greatly increased in women compared to men. An important role underlying the difference in activity of immune cells in men and women is attributed to sex hormones.^{4,5}

461

462 Sexually dimorphic actions of glucocorticoids have been shown to provide a link to inflammatory 463 diseases with gender differences in prevalence.⁵⁹

464 Our data strongly resonate with findings of Walsh et al. ⁶³ who report the consistent observation of 465 elevated levels of peripheral pro-inflammatory markers accompanying chronic stress in humans, 466 which associates with not only heightened GR expression, an assumpted result of decreased cellular 467 GR sensitivity, and alterations in β-AR signaling, which was interpreted to be based on myeloid 468 progenitor cell related stress-signaling pathways.⁶³ This situation could also be associated with TTS 469 pathophysiology. Lastly, the exact role that the ratio T:E2 plays in the development of TTS remains 470 unclear and warrants further investigation.

471

472 Bioinformatic extrapolation: TTS – an inflammaging disorder?

473 The cardiovascular system, and like many other vital organs, the heart is susceptible to the effects of 474 aging. Inflammation is a hallmark of aging, and other coexisting comorbidities linked to age-related 475 decline, such as heart failure, cardiovascular disease, neurodegeneration, vascular dementia and Alzheimer's disease but also age-related hearing loss, age-related macular degeneration and type 2 476 diabetes.^{24,69,70,71} Inflammation is also associated with TTS. Recently, a novel portemanteau of 477 478 inflammation and aging has been coined, "inflammaging", which is defined as chronic, low-grade 479 inflammation accumulated and worsening over age, presumably contributing to the pathogenesis of various age-related pathologies.^{70,72,73} Inflammation has been described to stem from cellular 480 senescence of the immune system, so-called immunosenescence.⁷⁴ Although inflammatory failure,⁷⁵ 481 482 we set out to lay attention on the role of inflammaging in TTS. In humans and experimental animal models, several reports 65,66 documented a significant association between 4 systemic markers of 483 inflammation, i.e. IL-6, IL-10, TNFa, INFy.^{30,38,65,66} 484

485 Our computational analyses predicted IL-6, CCL5, CCL20, and INFγ to interact with the studied miRs

486 directly, while the remaining genes were shown to be involved in indirectly with these miRs. In

487 particular, possible INFγ-signaling pathways were found to be relevant in the ontogeny of TTS (miR-

488 15a and miR-15b). On the other hand, CCL5, CCL20, and IL-6 were predicted to play a pivotal role in

489 inflammation (miR-21 and miR-146a). Similarly, the IL-6 contribution to the TTS lipidome was

490 previously determined by experimental and computational analyses connecting this cytokine to the

491 phosphatidylinositol molecule.⁷⁶

492 Currently, the role of inflammaging in the ontogeny of TTS is only speculative, and further large-scale

493 studies are needed to verify.

494

To conclude, the contributions of various molecular mechanisms, notably inflammaging, oxidative stress, CUMS, gender, and genetic factors shall be evaluated as possible common culprits that interrelate the pathophysiology in the heart and cardiovascular system as part of TTS. Future research using new technologies like big-data analysis of blood samples from real-life patient cohorts will probably gain better insight into underlying mechanisms of different disease phenotypes. Identification of specific molecular pathways and associated biomarkers will then allow the development of new targets for precision medicine.

502

503 Acknowledgment: The assistance provided by Elisabeth Wilken and Mariola Dragan was greatly 504 appreciated. We wish to extend special thanks to Hiroshima Asa Citizens hospital team for technical 505 support. The authors wish to show their appreciation to Dr. C. Maack, Comprehensive heart failure 506 center Würzburg, Germany for inspiring scientific discussions. The research grant DFG FO 315/5-1 to 507 CF is acknowledged.

508

509 Authors' Contributions: MS and HH performed experiments, conducted data and statistical 510 analyses, and prepared the manuscript. SS performed computational analyses. MN, CF and SS, CB, 511 HH prepared the manuscript. CF, CB and TT conceptualized the project and supervised the 512 experiments. All authors contributed to manuscript writing and proofreading.

513

514 Competing interests:

515 The authors declare that they have no competing interests.

516

517 Abbreviations: ACS: acute coronary syndrome; AD: autoimmune disease; BBB: blood-brain barrier;
518 b-FGF: basic fibroblast growth factor; BNP: brain natriuretic peptide; CAT: cathecholamine; CCL:
519 CC-chemokine ligand; cEND: mouse brain capillary endothelial cell line; CK-MB: creatine kinase

520 MB; CM: cardiomyocyte; CNS: Central Nervous System; CUMS: chronic unpredictable mild stress; 521 DMEM: Dulbecco's modified Eagle's medium; ECG: electrocardiogram; EGF: epidermal growth 522 factor; FCS: fetal calf serum; G-CSF: granulocyte colony-stimulating factor; GGI: gene-gene 523 interaction; GMI: gene-miR interaction; GR: glucocorticoid receptor; GSK-3: Glycogen synthase 524 kinase 3; GUSB: β-glucuronidase; hCMEC/D3: human brain vascular endothelial cell line; HPA: 525 human platelet antigen; IL: interleukin; INFy: Interferon gamma; iPSC: induced pluripotent stem 526 cells; iPSC-CMs: induced pluripotent stem cells differentiated into cardiomyocytes; Iso: isoprenaline; 527 LV: left ventricle; MIP3A: Macrophage Inflammatory Protein 3 alpha; miR/miRNA: microRNA; 528 MyEND: myocardial endothelial cell line; NEA: non-essential amino acids; NR3C1: nuclear receptor 529 subfamily 3 group C member 1; NR4A1: nuclear receptor subfamily 4 group A member 1; P/S: 530 penicillin/streptomycin; PET: positronenemissionstomography; RT-qPCR: real-time quantitative 531 PCR; R3-IGF-1: recombinant analog of insulin-like growth factor; RANTES: CC-Chemokin-Ligand-532 5; SD: standard deviation; STEMI: ST-segment elevation myocardial infarction; TNF α : tumor 533 necrosis factor-α; TTC: takotsubo cardiomyopathy; TTS: Takotsubo syndrome; VEGF: Vascular 534 Endothelial Growth Factor; ZO-1: Zonula Occludens 1; β2-AR: β2-adrenergic receptor

535

536 **References**

- 537 1. Elgendy AY, Mahmoud AN, Elgendy IY. Takotsubo syndrome: Still a benign entity? Int J Cardiol.
 538 2017;247:41.
- 539 2. Ghadri JR, Wittstein IS, Prasad A, Sharkey S, Dote K, Akashi YJ, Cammann VL, Crea F, Galiuto
- L, Desmet W, et al. International expert consensus document on Takotsubo syndrome (Part II):
 Diagnostic workup, outcome, and management. Eur Heart J. 2018;39:2047-2062.
- 542 3. Elgendy AY, Elgendy IY, Mansoor H, Mahmoud AN. Clinical presentations and outcomes of
- Takotsubo syndrome in the setting of subarachnoid hemorrhage: A systematic review and metaanalysis. Eur Heart J Acute Cardiovasc Care. 2018;7:236-245.
- 545 4. Lyon AR, Bossone E, Schneider B, Sechtem U, Citro R, Underwood SR, Sheppard MN, Figtree
- 546 GA, Parodi G, Akashi YJ, et al. Current state of knowledge on Takotsubo syndrome: a Position
- statement from the taskforce on Takotsubo syndrome of the Heart Failure Association of the
 European Society of Cardiology. Eur J Heart Fail. 2016;18:8-27.
- 549 5. Angelini P, Tobis JM. Is high-dose catecholamine administration in small animals an appropriate
- model for takotsubo syndrome? Circ J. 2015;79:897.

- 6. Messas N, Willemain R, Bilger A, Rondeau-Lutz M, Kuhnert C, Mertes PM, Morel O, Collange
- 552 O. Takotsubo syndrome triggered by acute intermittent porphyria attack: An unusual stressor for 553 catecholamine-induced cardiomyopathy. Int J Cardiol. 2016;207:28-30.
- 7. Y-Hassan S. Catecholamine levels and cardiac sympathetic hyperactivation-disruption in
 Takotsubo syndrome. JACC Cardiovasc Imaging. 2017;10:95-96.
- 556 8. Y-Hassan S, Sörensson P, Ekenbäck C, Lundin M, Agewall S, Brolin EB, Caidahl K, Cederlund
- 557 K, Collste O, Daniel M, et al. Plasma catecholamine levels in the acute and subacute stages of
- takotsubo syndrome: Results from the Stockholm myocardial infarction with normal coronaries 2
 study. Clin Cardiol. 2021;44:1567-1574.
- 9. Y-Hassan S. Divergence in the results of plasma catecholamine levels in different studies on
 patients with takotsubo syndrome: Why? J Cardiol. 2016;68:89.
- 562 10. Andò G, Boretti I, Tripodi R. Stress cardiomyopathies beyond Takotsubo: does a common
 563 catecholaminergic pathophysiology fit all? Expert Rev Cardiovasc Ther. 2014;12:643-645.
- 564 11. El-Battrawy I, Borggrefe M, Akin I. Catecholamine in takotsubo syndrome. Int J Cardiol.
 565 2017;233:97.
- 566 12. Borchert T, Hübscher D, Guessoum CI, Lam TD, Ghadri JR, Schellinger IN, Tiburcy M, Liaw
- 567 NY, Li Y, Haas J, et al. Catecholamine-dependent β-adrenergic signaling in a pluripotent stem cell
- 568 model of Takotsubo cardiomyopathy. J Am Coll Cardiol. 2017;70:975-991.
- 569 13. Madias JE. Pathophysiology of Takotsubo syndrome: an adrenergic cardiac "chemical
 570 neuritis/myocarditis"? Cardiovasc Revasc Med. 2014;15:50.
- 14. Martins JB, Kerber RE, Marcus ML, Laughlin DL, Levy DM. Inhibition of adrenergic
 neurotransmission in ischaemic regions of the canine left ventricle. Cardiovasc Res. 1980;14:116124.
- 574 15. Feola M, Chauvie S, Rosso GL, Biggi A, Ribichini F, Bobbio M. Reversible impairment of
 575 coronary flow reserve in takotsubo cardiomyopathy: a myocardial PET study. J Nucl Cardiol.
 576 2008;15:811-817.
- 577 16. Palla AR, Dande AS, Petrini J, Wasserman HS, Warshofsky MK. Pretreatment with low-dose β578 adrenergic antagonist therapy does not affect severity of Takotsubo cardiomyopathy. Clin Cardiol.
 579 2012;35:478-481.
- 580 17. Paur H, Wright PT, Sikkel MB, Tranter MH, Mansfield C, O'Gara P, Stuckey DJ, Nikolaev VO,
- 581 Diakonov I, Pannell L, et al. High levels of circulating epinephrine trigger apical cardiodepression
- 582 in a β2-adrenergic receptor/Gi-dependent manner: a new model of Takotsubo cardiomyopathy.
- 583 Circulation. 2012;126:697-706.

- 584 18. Salemi VM, Atik E, Kairalla RA, Queiroz EL, Rosa LV, Kalil Filho R. Takotsubo 585 cardiomyopathy triggered by $\beta(2)$ adrenergic agonist. J Bras Pneumol. 2011;37:560-562.
- 586 19. Tutgun Onrat S, Dural İE, Yalım Z, Onrat E. Investigating changes in β-adrenergic gene
- 587 expression (ADRB1 and ADRB2) in Takotsubo (stress) cardiomyopathy syndrome; a pilot study.
- 588 Mol Biol Rep. 2021;48:7893-7900.
- 589 20. Yoneda H, Nakamura T, Shirao S, Tanaka N, Ishihara H, Suehiro E, Koizumi H, Isotani E,
- 590 Suzuki M, S.A.H.P.S. Group. Multicenter prospective cohort study on volume management after 591 subarachnoid hemorrhage: hemodynamic changes according to severity of subarachnoid 592 hemorrhage and cerebral vasospasm. Stroke. 2013;44:2155-2161.
- 593 21. Bybee KA, Motiei A, Syed IS, Kara T, Prasad A, Lennon RJ, Murphy JG, Hammill SC, Rihal
- 594 CS, Wright RS. Electrocardiography cannot reliably differentiate transient left ventricular apical
- ballooning syndrome from anterior ST-segment elevation myocardial infarction. J Electrocardiol.2007;40:38.e1-6.
- 597 22. Couch LS, Fiedler J, Chick G, Clayton R, Dries E, Wienecke LM, Fu L, Fourre J, Pandey P,
 598 Derda AA, et al. Circulating microRNAs predispose to takotsubo syndrome following high-dose
 599 adrenaline exposure. Cardiovasc Res. 2022;118:1758-1770.
- 600 23. Jaguszewski M, Osipova J, Ghadri JR, Napp LC, Widera C, Franke J, Fijalkowski M, Nowak R,
- Fijalkowska M, Volkmann I, et al. A signature of circulating microRNAs differentiates takotsubo
 cardiomyopathy from acute myocardial infarction. Eur Heart J. 2014;35:999-1006.
- 603 24. Karnati S, Guntas G, Rajendran R, Shityakov S, Höring M, Liebisch G, Kosanovic D, Ergun S,
- Nagai M, Förster CY. Quantitative lipidomic analysis of Takotsubo syndrome patients' serum. Front
 Cardiovasc Med. 2022;9:797154.
- 606 25. Thal SC, Smetak M, Hayashi K, Förster CY. Hemorrhagic cerebral insults and secondary
 607 Takotsubo syndrome: Findings in a novel in vitro model using human blood samples. Int J Mol Sci.
 608 2022;23:11557.
- 609 26. Ghadri JR, Wittstein IS, Prasad A, Sharkey S, Dote K, Akashi YJ, Cammann VL, Crea F, Galiuto
- 610 L, Desmet W, et al. International expert consensus document on Takotsubo syndrome (Part I):
- 611 Clinical characteristics, diagnostic criteria, and pathophysiology. Eur Heart J. 2018;39:2032-2046.
- 612 27. Förster C, Silwedel C, Golenhofen N, Burek M, Kietz S, Mankertz J, Drenckhahn D. Occludin as
- 613 direct target for glucocorticoid-induced improvement of blood-brain barrier properties in a murine
- 614 in vitro system. J Physiol. 2005;565(Pt 2):475-486.
- 615 28. Burek M, Salvador E, Förster CY. Generation of an immortalized murine brain microvascular
- 616 endothelial cell line as an in vitro blood brain barrier model. J Vis Exp. 2012;66:e4022.

- 617 29. Golenhofen N, Ness W, Wawrousek EF, Drenckhahn D. Expression and induction of the stress
- 618 protein alpha-B-crystallin in vascular endothelial cells. Histochem Cell Biol. 2002;117:203-209.
- 619 30. Ittner C, Burek M, Störk S, Nagai M, Förster CY. Increased catecholamine levels and
- 620 inflammatory mediators alter barrier properties of brain microvascular endothelial cells in vitro.
- 621 Front Cardiovasc Med. 2020;7:73.
- 622 31. Weksler BB, Subileau EA, Perriere N, Charneau P, Holloway K, Leveque M, Tricoire-Leignel H,
- 623 Nicotra A, Bourdoulous S, Turowski P, et al. Blood-brain barrier-specific properties of a human 624 adult brain endothelial cell line. FASEB J 2005;19:1872-1874.
- 625 32. Weksler B, IRomero LA, Couraud PO. The hCMEC/D3 cell line as a model of the human blood 626 brain barrier. Fluids Barriers CNS 2013;10:16.
- 627 33. Haase A, Göhring G, Martin U. Generation of non-transgenic iPS cells from human cord blood
- 628 CD34+ cells under animal component-free conditions. Stem Cell Res. 2017;21:71-73.
- 629 34. Lian X, Zhang J, Azarin SM, Zhu K, Hazeltine LB, Bao X, Hsiao C, Kamp TJ, Palecek SP.
- 630 Directed cardiomyocyte differentiation from human pluripotent stem cells by modulating Wnt/β-631 catenin signaling under fully defined conditions. Nat Protoc. 2013;8:162-175.
- 632 35. Foinquinos A, Batkai S, Genschel C, Viereck J, Rump S, Gyöngyösi M, Traxler D, Riesenhuber 633 M, Spannbauer A, Lukovic D. Preclinical development of a miR-132 inhibitor for heart failure
- 634 treatment. Nat Commun. 2020;11:633.
- 635 36. Chatterjee S, Hofer T, Costa A, Lu D, Batkai S, Gupta SK, Bolesani E, Zweigerdt R, Megias D,
- 636 Streckfuss-Bomeke K. Telomerase therapy attenuates cardiotoxic effects of doxorubicin. Mol Ther.
- 637 2021;29:1395-1410.
- 638 37. Tohyama S, Hattori F, Sano M, Hishiki T, Nagahata Y, Matsuura T, Hashimoto H, Suzuki T, 639 Yamashita H, Satoh Y. Distinct metabolic flow enables large-scale purification of mouse and 640 human pluripotent stem cell-derived cardiomyocytes. Cell Stem Cell. 2013;12:127-137.
- 641
- 38. Wilson HM, Cheyne L, Brown PAJ, Kerr K, Hannah A, Srinivasan J, Duniak N, Horgan G,
- 642 Dawson DK. Characterization of the myocardial inflammatory response in acute stress-induced
- 643 (Takotsubo) cardiomyopathy. JACC Basic Transl Sci. 2018;3:766-778.
- 644 39. Kutmon M, Kelder T, Mandaviya P, Evelo CT, Coort SL. CyTargetLinker: a cytoscape app to 645 integrate regulatory interactions in network analysis. PLoS One. 2013;8:e82160.
- 646 40. Fareed MM, Ullah S, Aziz S, Johnsen TA, Shityakov S. In-silico analysis of non-synonymous
- 647 single nucleotide polymorphisms in human β -defensin type 1 gene reveals their impact on protein-
- 648 ligand binding sites. Comput Biol Chem. 2022;98:107669.

- 649 41. Franz M, Rodriguez H, Lopes C, Zuberi K, Montojo J, Bader GD, Morris Q. GeneMANIA
- 650 update 2018. Nucleic Acids Res. 2018;46(W1):W60-W64.
- 42. Grouzmann E, Cavadas C, Grand D, Moratel M, Aubert JF, Brunner HR, Mazzolai L. Blood
- sampling methodology is crucial for precise measurement of plasma catecholamines concentrations
- 653 in mice. Pflugers Arch. 2003;447:254-258.
- 43. Förster C, Kahles T, Kietz S, Drenckhahn D. Dexamethasone induces the expression of
- 655 metalloproteinase inhibitor TIMP-1 in the murine cerebral vascular endothelial cell line cEND. J
- 656 Physiol. 2007;580(Pt.3):937-949.
- 657 44. Yi X, Xu C, Huang P, Zhang L, Qing T, Li J, Wang C, Zeng T, Lu J, Han Z. 1-
- 658 Trifluoromethoxyphenyl-3-(1-Propionylpiperidin-4-yl) urea protects the blood-brain barrier against
- 659 ischemic injury by upregulating tight junction protein expression, mitigating apoptosis and
- 660 inflammation In Vivo and In Vitro Model. Front Pharmacol. 2020;11:1197.
- 661 45. Davies L, Paraskevopoulou E, Sadeq M, Symeou C, Pantelidou C, Demonacos C, Krstic-
- 662 Demonacos M. Regulation of glucocorticoid receptor activity by a stress responsive transcriptional
 663 cofactor. Mol Endocrinol. 2011;25:58-71.
- 46. Barcena ML, Aslam M, Pozdniakova S, Norman K, Ladilov Y. Cardiovascular inflammaging:
 mechanisms and translational aspects. Cells. 2022;11:1010.
- 47. Bleve A, Motta F, Durante B, Pandolfo C, Selmi C, Sica A. Immunosenescence, inflammaging,
 and frailty: Role of myeloid cells in age-related diseases. Clin Rev Allergy Immunol. 2023;64:123144.
- 669 48. Nagai M, Kobayashi Y, Kobatake H, Dote K, Kato M, Oda N, Kunita E, Kagawa E, Yamane A,
- Osawa A, et al. Happy heart syndrome: a case of Takotsubo syndrome with left internal carotid
 artery occlusion. Clin Auton Res. 2020;30:347-350.
- 49. Osawa A, Nagai M, Dote K, Kato M, Oda N, Kunita E, Kagawa E, Yamane A, Kobatake H,
- 673 Shiota H. A mid-ventricular variant of Takotsubo syndrome: was it triggered by insular cortex
- 674 damage? ESC Heart Fail. 2021;8:3408-3412.
- 50. Bybee KA, Prasad A. Stress-related cardiomyopathy syndromes. Circulation. 2008;118:397-409.
- 676 51. Nagai M, Dote K, Kato M, Sasaki S, Oda N, Kagawa E, Nakano Y, Yamane A, Higashihara T,
- 677 Miyauchi S, et al. The insular cortex and Takotsubo cardiomyopathy. Curr Pharm Des.
 678 2017;23:879-888.
- 679 52. Singh T, Khan H, Gamble DT, Scally C, Newby DE, Dawson D. Takotsubo syndrome:
- Pathophysiology, emerging concepts, and clinical implications. Circulation. 2022;145:1002-1019.

- 53. Nazir S, Lohani S, Tachamo N, Ghimire S, Poudel DR, Donato A. Takotsubo cardiomyopathy
- associated with epinephrine use: A systematic review and meta-analysis. Int J Cardiol. 2017;229:6770.
- 684 54. Ferreira VM, Marcelino M, Piechnik SK, Marini C, Karamitsos TD, Ntusi NAB, Francis JM,
- Robson MD, Arnold JR, Mihai R, et al. Pheochromocytoma is characterized by catecholamine-
- 686 mediated myocarditis, focal and diffuse myocardial fibrosis, and myocardial dysfunction. J Am Coll
- 687 Cardiol. 2016;67:2364-2374.
- 688 55. Grippo AJ, Scotti MA. Stress and neuroinflammation. Mod Trends Pharmacopsychiatry.
 689 2013;28:20-32.
- 690 56. Finnell JE, Moffitt CM, Hesser LA, Harrington E, Melson MN, Wood CS, Wood SK. The
- 691 contribution of the locus coeruleus-norepinephrine system in the emergence of defeat-induced
- 692 inflammatory priming. Brain Behav Immun. 2019;79:102-113.
- 693 57. Herman JP, McKlveen JM, Ghosal S, Kopp B, Wulsin A, Makinson R, Scheimann J, Myers B.
- Regulation of the hypothalamic-pituitary-adrenocortical stress response. Compr Physiol.2016;6:603-621.
- 58. Bardo MT, Chandler CM, Denehy ED, Carper BA, Prendergast MA, Nolen TL. Effect of the
 glucocorticoid receptor antagonist PT150 on acquisition and escalation of fentanyl selfadministration following early-life stress. Exp Clin Psychopharmacol. 2023;31:362-369.
- 59. Duma D, Collins JB, Chou JW, Cidlowski JA, Sexually dimorphic actions of glucocorticoids
 provide a link to inflammatory diseases with gender differences in prevalence. Science signaling
 2010;3:ra74.
- 60. González Ramírez C, Villavicencio Queijeiro A, Jiménez Morales S, Bárcenas López D, Hidalgo
- 703 Miranda A, Ruiz Chow A, Tellez Cardenas L, Guardado Estrada M, et al. The NR3C1 gene
- 704 expression is a potential surrogate biomarker for risk and diagnosis of posttraumatic stress disorder.
- 705 Psychiatry Res. 2020;284:112797.
- 61. Sapolsky RM, Krey LC, McEwen BS. The neuroendocrinology of stress and aging: the
 glucocorticoid cascade hypothesis. Endocr Rev. 1986;7:284-301.
- 62. Nagai M, Förster CY, Dote K. Sex hormone-specific neuroanatomy of Takotsubo syndrome: Is
 the insular cortex a moderator? Biomolecules. 2022;12:110.
- 710 63. Walsh CP, Bovbjerg DH, Marsland AL. Glucocorticoid resistance and β2-adrenergic receptor
- 711 signaling pathways promote peripheral pro-inflammatory conditions associated with chronic
- 712 psychological stress: A systematic review across species. Neurosci Biobehav Rev. 2021;128:117-
- 713 135.

- 714 64. Oberbeck R. Catecholamines: physiological immunomodulators during health and illness. Curr
- 715 Med Chem. 2006;13:1979-1989.
- 716 65. Scally C, Abbas H, Ahearn T, Srinivasan J, Mezincescu A, Rudd A, Spath N, Yucel-Finn A,
- 717 Yuecel R, Oldroyd K, et al. Myocardial and systemic inflammation in acute stress-induced
- 718 (Takotsubo) cardiomyopathy. Circulation. 2019;139:1581-1592.
- 719 66. Santoro F, Costantino MD, Guastafierro F, Triggiani G, Ferraretti A, Tarantino N, Saguner A, Di
- Biase M, Brunetti ND. Inflammatory patterns in Takotsubo cardiomyopathy and acute coronary
- syndrome: A propensity score matched analysis. Atherosclerosis. 2018;274:157-161.
- 722 67. Šerstņova K, Pilmane M, Vitenberga-Verza Z, Melderis I, Gontar Ł, Kochański M, Drutowska A,
- 723 Maroti G, Prieto-Simon B. Expression of anti-inflammatory markers IL-2, IL-10, TGF-β1, βDEF-2,
- 724 βDEF-3 and Cathelicidin LL37 in dairy cattle milk with different health status of the udder. Pol J
- 725 Vet Sci. 2022;25:237-248.
- 68. Kuwabara T, Ishikawa F, Kondo M, Kakiuchi T. The Role of IL-17 and related cytokines in
 inflammatory autoimmune diseases. Mediators Inflamm. 2017;2017:3908061.
- 728 69. Blasko I, Stampfer-Kountchev M, Robatscher P, Veerhuis R, Eikelenboom P, Grubeck-
- 729 Loebenstein B. How chronic inflammation can affect the brain and support the development of
- Alzheimer's disease in old age: the role of microglia and astrocytes. Aging Cell. 2004;3:169-176.
- 731 70. Ferrucci L, Fabbri E. Inflammageing: chronic inflammation in ageing, cardiovascular disease,
 732 and frailty. Nat Rev Cardiol. 2018;15:505-522.
- 733 71. Shityakov S, Nagai M, Ergün S, Braunger BM, Förster CY. The Protective effects of
- neurotrophins and microRNA in diabetic retinopathy, nephropathy and heart failure via regulating
 endothelial function. Biomolecules. 2022;12:1113.
- 736 72. Baylis D, Bartlett DB, Patel HP, Roberts HC. Understanding how we age: insights into
 737 inflammaging. Longev Healthspan. 2013;2:8.
- 738 73. Franceschi C, Garagnani P, Parini P, Giuliani C, Santoro A. Inflammaging: a new immune739 metabolic viewpoint for age-related diseases. Nat Rev Endocrinol. 2018;14:576-590.
- 740 74. Müller L, Di Benedetto S, Pawelec G. The immune system and its dysregulation with aging.
 741 Subcell Biochem. 2019;91:21-43.
- 742 75. Adamo L, Rocha-Resende C, Prabhu SD, Mann DL. Reappraising the role of inflammation in
- heart failure. Nature reviews. Cardiology 2020;17:269-285.
- 744 76. Karnati S, Garikapati V, Liebisch G, Van Veldhoven PP, Spengler B, Schmitz G, Baumgart-Vogt
- E. Quantitative lipidomic analysis of mouse lung during postnatal development by electrospray
- ionization tandem mass spectrometry. PLoS One. 2018;13:e0203464.

747

748 Figure legends

749 Figure 1. Changes of protein expression in cEND cells after incubation with the CATs 750 norepinephrine, epinephrine and dopamine in different concentration (c1 pharmacological concentrations) ⁴²: 150µM dopamine, 1µM epinephrine, 1µM norepinephrine; c2 postmenopausal 751 healthy control; c3 TTS acute; c4 TTS subacute, comp. table 1). a) Western blots displaying xpression 752 753 of the different proteins after 24h of incubation. b) densitometric evaluation. Changes of protein 754 expression was normalized to β -actin. Data are the means (±SEM) of 3 independent experiments. 755 Statistical significance was evaluated using Dunnett's multiple comparisons test. *: P<0.05; ***: P < 756 0.001: ****: P < 0.00001

757

Figure 2. Changes of mRNA expression in cEND cells after incubation with the CATs norepinephrine, epinephrine and dopamine in different concentrations (c1 pharmacological concentrations ⁴²: 150µM dopamine, 1µM epinephrine, 1µM norepinephrine; c2 postmenopausal healthy control; c3 TTS acute; c4 TTS subacute, comp. Table 1). Changes of protein expression was normalized to β-actin. Data are the means (±SEM) of 3 independent experiments. Statistical significance was evaluated using Dunnett's multiple comparisons test. ***: P < 0.001; ****: P < 0.00001

765

Figure 3. Physiological CAT concentrations did not alter cardiac stress marker NR4A1 in 766 767 cardiomyocytes. A Human iPSC-derived cardiomyocytes were treated with physiological 768 concentrations of a CAT mix as determined from acute TTS (c3 comp. Table 1) and 769 supraphysiological concentrations of isoprenaline (Iso) for 24 h. The expression level of the cardiac 770 stress marker nuclear receptor subfamily 4 group A member 1 (NR4A1) was analyzed by RT-qPCR, 771 mean \pm SD of three independent experiments is shown. B Human iPSC-derived cardiomyocytes were 772 treated with physiological concentrations of CAT as determined from acute TTS (c3) and multiples 773 $(10^3, 10^6)$ of it. Expression of NR4A1 was determined by RT-qPCR, mean \pm SD of triplicates is 774 shown. Statistical significance was analyzed with one-way ANOVA with Dunnett's correction in 775 comparison to basal, ** P < 0.01, *** P < 0.001, ns P > 0.05.

776

Figure 4. Changes in the protein expression of glucocorticoid receptor α in exosomes from serum from patients with various heart diseases.

a) Expression of glucocorticoid receptor α in the exosomes. (densitometric evaluation)

b) Membranes of the Western blots used in this experiment.

781 Glucocorticoid receptor α protein expression was normalized to CD63. Data are the means (±SD) of n

782 independent patients (postmenopausal healthy: n=3, premenopausal healthy: n=3, ACS acute: n=4,

783 TTS acute: n=4, TTS subacute: n=8, TTS chronic: n=3). Statistical significance was evaluated using

784 Tukey's multiple comparisons test.

785

Figure 5. Human GMI (gene-miR interaction) network generated by the CyTargetLinker algorithm human using the miRTarBase (blue) and TargetScan (red) databases. The marked nodes are depicted in yellow. The network is displayed using the yFiles organic layout.

789

Figure 6. Human GGI (gene-gene interaction) network generated by the GeneMANIA plugin for the
corresponding blood markers and proinflammatory cytokines (CCL20, CCL5, EGF, IL2, IL4, IL6,
IL10, IL17A, INFγ, NR3C1, and TNF). The marked nodes are depicted in yellow. The network is

793 displayed using the vFiles organic layout.

794

Figure 7. The merged (A) and subtracted (B) networks produced by merging GMI and GGI networks or creating a new network (8 nodes and 5 edges) from selected nodes and all of their edges. The marked nodes are depicted in yellow. The merged network is displayed using the prefuse forcedirected layout. **Table 1.** Measured concentration of different CATs (CAT) in serum of postmenopausal healthy patients and patients with TTS in pmol/l \pm SD.*: P < 0.05; [TTS acute] = c2, [TTS subacute] = c3 in the following sections</td>

| | postmenopausal healthy n= 3 | TTS acute n=4 | TTS subacute n=8 |
|---------------|-----------------------------|-----------------|------------------|
| Noradrenaline | 4.24 ± 1.52 | 3.65 ± 2.37 | 5.23 ± 5.69 * |
| Adrenaline | 0.22 ± 0.11 | 0.13 ± 0.1 | 0.19 ± 0.2 |
| Dopamine | 0.18 ± 0.08 | 0.21 ± 0.12 | 0.80 ± 1.06 * |

| | premenopausal healthy n = 3 | postmenopausal healthy n= 3 | TTS acute n = 4 | TTS subacute n = 8 | TTS chronic n = 3 | ACS $n = 4$ |
|--------|-----------------------------|-----------------------------|-----------------------|---------------------|-------------------|-------------------|
| | | | | | | |
| INFγ | 2,41 ± 1,48 | $18,7\pm7,18$ | 65,6 ± 71,99 | $26,\!79\pm32,\!46$ | 19,81 ± 30,50 | $18,46 \pm 32,03$ |
| IL-2 | $0,32 \pm 0,33$ | 1,49 ± 1,23 | $4,87 \pm 8,012$ | 2,2 ± 3,263 | 1,53 ± 2,243 | $0,37 \pm 0,3641$ |
| IL-4 | $2,34 \pm 3,38$ | $1,59 \pm 0,88$ | 50,77 ± 8,68 | $2,75 \pm 4,29$ | 0,80 ± 1,10 | $0,70 \pm 0,92$ |
| IL-6 | $0,34 \pm 0,34$ | 6,64 ± 15,41 | $38,01 \pm 45,06$ | 8,98 ± 15,13 | $1,45 \pm 1,45$ | $0,97 \pm 0,77$ |
| IL-10 | $12,13 \pm 19,54$ | 30,21 ± 27,33 | $24,\!48 \pm 17,\!52$ | $10,55 \pm 10,74$ | 3,88 ± 6,68 | 3,41 ± 5,79 |
| TNFα | 15 ± 5,67 | $18,53 \pm 18,51$ | $41,04 \pm 47,72$ | 25,8 ± 16,62 | 17,02 ± 5,262 | $15,6 \pm 12,43$ |
| | | postmenopausal healthy n= 3 | TTS acute n = 4 | TTS subacute n = 8 | TTS chronic n = 3 | ACS $n = 4$ |
| EGF | | 57,05 | 22,19 | 85,14 | 26,14 | 47,85 |
| IL-17 | | $3,40 \pm 1,70$ | $1{,}91\pm0{,}84$ | $25,67 \pm 52,31$ | 3,13 | 3,52 |
| MIP3A | | $147,3\pm104,4$ | $111,3 \pm 68,08$ | $249,6\pm168,5$ | 105,3 | 72,69 |
| | | postmenopausal healthy n= 3 | TTS acute n = 4 | TTS subacute n = 8 | | |
| RANTES | | 161542 ± 190762 | 69217 ± 81541 | 264518 ± 381721 | | |

Table 2. Measured concentration of different cytokines and chemokines in patients sera in pg/ml \pm SD.

| | postmenopausal healthy n= 3 | TTS acute n = 4 | TTS subacute n = 8 | TTS chronic n = 3 |
|--------------|-----------------------------|-----------------|--------------------|-------------------|
| Testosterone | 202.5 ± 120.4 | 160 | 287.5 ± 257.5 | 145 ± 63.6 |
| Estradiol | 35.25 ± 54.04 | 5 | 43.13 ± 66.69 | 9.5 ± 6.364 |

Table 3: Measured concentrations of testosterone and 17 β -estradiol in pg/ml in the serum of patients with various heart disease. Data are the means (\pm SD) of n independent measures.

Figure 1.



Figure 2.



Figure 3.



Figure 4.



Glucocorticoid receptor α in exosomes

Figure 5.



Figure 6.



Figure 7.

