

Assessment of Geriatric Depression among Elderly Peoples Residing at Rural Area of Kheda District, Gujarat.

Ms. Vaibhavi Parmar¹ Mr. Kailash Nagar² Prof. Virendra Jain³

1. Final year M.Sc. Nursing Students, Dinsha Patel College of Nursing, Nadiad.
2. Assistant Professor & HOD of Community Health Nursing Department, Dinsha Patel College of Nursing, Nadiad
3. Principal of Dinsha Patel College of Nursing, Nadiad

Corresponding Author: - Mr. Kailash Nagar (Email Id: dpcnkailash@gmail.com)

<https://orcid.org/0000-0002-0564-2761>,

Researcher: Ms. Vaibhavi parmar (Email Id: vaibhavip0907gmail.com)

ABSTRACT

Background: Depression is a major mental health problem, which is yet to be recognized as an important public health challenge. About 322 million people affected with depression worldwide.

Depression among the elderly population in India has been recognized as one of the major public health problems with a prevalence of 8 to 22%. It causes significant suffering and accounts for 5.7% of the years lived with disability, to make matters worse, it has been reported that in many cultures and societies, deteriorating mental status, say it dementia or depression, has been perceived as a part of normal ageing.

Aims and Objectives: The aim of this study is to explore the magnitude and risk factors of depression in elderly people residing in rural area of Gujarat. **Objective:** 1. To assess the level of depression among elderly people residing at rural area. 2. To find out Significant Association between the level of depression selected demographic variables among elderly people at rural area.

Methodology

Design and Setting: A cross sectional descriptive study design was used. Non probability convenient sampling technique was used to drawn samples. 100 elderly persons were screened. Geriatric depression scale (GDS) was used to assess depressive level. A self-rating questionnaire tool was used to collect data and some socio-demographics and clinical variables. Prior to data collection written setting permission obtain from authority of the village and written consent from was obtained from the participants. Data for study was collected by door to door visit in selected community. Chi square was employed to determine association of collected data. **Tool consists of following Section A:** Socio-demographic variables. **Section B:** Geriatric Depression Scale (GDS).

The GDS questions are answered 'yes' or 'no', instead of a five-category response set. This simplicity enables the scale to be used with ill or moderately cognitively impaired individuals. The scale is commonly used as a routine part of a comprehensive geriatric assessment. One point is assigned to each answer and the cumulative score is rated on a scoring grid. The grid sets a range of 0–9 as 'normal', 10–19 as 'mildly depressed', and 20–30 as 'severely depressed'.

Results: Results showed that 61% was Male and 39% was Female. The majority of had 71% primary education. Regarding work majority 60% are labor, regarding financial dependency most 62% was independent, 35(35%) dependent. This study reveals that level of depression among geriatric peoples, 51 (51%) peoples had no depression, 33 (33%) having mild depression and 16 (16%) having severe depression level.

Conclusion: It is concluded that nearly 50% of geriatric peoples having no depression and mild depression is more prevalent among elderly in selected rural area of Gujarat. So majority of the elderly peoples were found normal.

KEYWORDS: (GDS) Geriatric Depression scale, Elderly, Rural area, world health organization.

INTRODUCTION: Geriatric depression is a mental and emotional disorder affecting older adults. Geriatric depression scale is used to quantify the severity of depression. Empty nest syndrome (loneliness in elderly is common causative factor for depression) is a factor should be rectified by social skills training.¹ Late onset bipolar disorder is rare. Depression is common among elderly.² It is a mental health condition that can affect people of all ages. While it is normal to feel down sometimes, if you feel this way for 2 weeks or more, or your mood is affecting your ability to cope with everyday life, you may be experiencing depression.³ In elderly people, depression mainly affects those with chronic medical illnesses and cognitive impairment, causes suffering, family disruption, and disability, worsens the outcomes of many medical illnesses, and increases mortality.⁴

According to WHO, factors increasing depression risk in older adults include genetic, chronic disease and disability, pain, frustration with limitations in activities of daily living, personality traits (dependent, anxious or avoidant), adverse life events (separation, divorce, bereavement, poverty, social isolation) and lack of adequate social support. Many studies have demonstrated a relationship between depression and various socioeconomic variables such as advanced age, low education, poverty and manual occupation. Thus, an older adult patient suffering from depression often has a combination of psychological, physical and social needs.⁵

More than 1 in 10 older people, and more than 3 in 10 people living in residential aged-care, experience depression. It's important to remember that not all older people become depressed, and just because you are older, you don't need to accept that you will become depressed, or that your depression can't be treated. Technical advances have facilitated the exploration of factors related to geriatric depression and have helped generate novel biological and psychosocial treatment approaches.⁶

Based on UNDESA's 2019 report, the population aged over 65 years old would reach 703million and was predicted to double by 2050, that is, 1.5 billion in which the proportion of older adult is 16%. It means that 1 of 6 people in the world's population is elderly.⁷ Depression is one of the most common illnesses worldwide, with more than 264 million people

affected. Various studies in the elderly population have estimated the prevalence of depression across India, with results ranging from 6% to 62%. The objectives of this study were to estimate the prevalence of depression among the elderly population using a Geriatric Depression Scale (GDS) and to find out the association between various sociodemographic parameters and depression among elderly people.⁸

Depression or the occurrence of depressive symptomatology is a prominent condition amongst older people, with a significant impact on the well-being and quality of life. Many studies have demonstrated that the prevalence of depressive symptoms increases with age (Kennedy, 1996). Depressive symptoms not only have an important place as indicators of psychological well-being but are also recognized as significant predictors of functional health and longevity. Longitudinal studies demonstrate that increased depressive symptoms are significantly associated with increased difficulties with activities of daily living (Penninx *et al.*, 1998). Community-based data indicate that older persons with major depressive disorders are at increased risk of mortality (Bruce, 1994).⁹

Attention guides thought and behaviour. Information that is attended becomes available to higher-order cognitive processes such as working memory and decision-making. In order to promote wellbeing, it is important that attention selects stimuli associated with rewarding outcomes (Anderson, 2013).¹⁰ Because of its devastating consequences, late life depression is an important public health problem. It is associated with increased risk of morbidity, increased risk of suicide, decreased physical, cognitive and social functioning, and greater self-neglect, all of which are in turn associated with increased mortality.¹⁰

OBJECTIVE:

1. To assess the level of depression among elderly people residing at rural area of Kheda district
2. To find out Significant Association between the level of depression selected demographic variables among elderly people at rural area of Kheda, district.

MATERIALS AND METHODS

Research Approach: -Quantitative Research approach was used for the current study.

Research Design: - The design of the study will be used Non-Experimental Cross-Sectional Survey Research Design.

Research Variables: There are two types of variables considered under the study as follows

1. **Research variables:** In present study research variable is the level of depression among old age people.
2. **Demographic variables:** Age, gender, religion, education, marital status, type of family, locality of house, no. of children, frequency of visit of children or relative, hobbies, financial support, Previous occupation, old age pension

Sampling method: -

The sample of the study will be selected by using **Non probability convenient sampling technique.** According to inclusive criteria as well as availability of samples.

Study population: - In the present study, the target populations are old age people residing at old age residing at rural area of selected areas of Kheda district”.

Study Sample: - Geriatric peoples residing at rural area of selected areas of Kheda district”.

Study Setting: PHC Pethai has been selected for the main study setting and data collect from the elderly people.

Sample Size: - 100 geriatric people

SAMPLE CRITERIA

Inclusion criteria

1. Old age people who are aged 60year and above.
2. Old age people who are willing to participate in the study.
3. Old age people who are available at the time of study.

Exclusion criteria

1. Those who are present at data collection.
2. Those who are seriously ill
3. Those who are not able to interact (hearing loss and language problems)

Tool for Data Collection:

Section A: Sociodemographic variables.

Section B: Geriatric Depression Scale.

RESULTS

TABLE–1: Socio-demographic variables of geriatric peoples residing at rural area of Kheda district N=100


Demographic Variables	Frequency	Percentage
Age In Years		
a) 60- 65 Years	24	24%
b) 66-70 Years	40	40%
c) 71-75 Years	24	24%
d) >75 Years	12	12%
Gender		
a) Male	61	61%
b) Female	39	39%
c) Transgender		
Religion Status		
a) Hindu	73	73%
b) Christian	0	0%
c) Muslim	27	27%
d) Others	0	0%
Educational Status		
a) Primary	71	71%
b) Secondary	20	20%
c) Higher Secondary	8	8%
d) Illiterate	1	1%
Marital Status		
a) Married	50	50%
b) Unmarried	1	1%
c) Divorced	1	1%
d) Widow/Widower	48	48%
Number Of Child		
a) One	11	11%
b) Two	28	28%
c) More Than Two	56	56%
d) No Children	5	5%
Previous occupation		
a) Private	6	6%
b) Government	6	6%
c) Labor	60	60%
d) Farmer	28	28%
Employment status		
a) Working	48	48%
b) Non- working	52	52%
Currently staying with		
a) Son	73	73%
b) Daughter	12	12%
c) Separated	8	8%
d) Any Other relative	7	7%
Healthy Habits		
a) Reading books,	39	39%
b) Exercise, Yoga	2	2%
c) Listening music	10	10%
d) Chatting in a group	49	49%
Any Bad Habits		
a) Tobacco chewing	17	17%
b) Smoking	9	9%
c) Alcohol	6	6%
d) No bad habits	68	68%

Do you get Retirement pension	14	14%
a) Yes	86	86%
b) No		
Do you get 1000 Rs. pension Yojana	38	38%
a) Yes	62	62%
b) No		
Type of family		
a) Joint family	75	75%
b) Nuclear family	16	16%
c) Extended family	9	9%
Disease status		
a) Acute illness	25	25%
b) Chronic illness	23	23%
c) No illness	52	52%
Financial dependency		
a) Independent	62	62%
b) Dependent	35	35%
c) Partially depended	3	3%

TABLE-2.1: Distribution of the geriatric Peoples according to Level the Depression

Level of Depression	Frequency (%)
No Depression	51 (51%)
Mild Depression	33 (33%)
Severe Depression	16 (16%)
Total	100 (100%)

The above table 2.1 shows level of depression among geriatric peoples, 51 (51%) peoples had no depression, 33 (33%) having mild depression and 16 (16%) having severe depression level.



Graph no.1 showing distribution according to level of depression among geriatric peoples

Table-2.2: Range, Mean, and Standard Deviation of depression among the geriatric peoples residing at rural area of Kheda district

Level of knowledge	Max. score	Range	Mean	SD
Overall	24	23	10.97	6.014

The above table 2.2 shows depression score range was 1-24, mean score was 10.97, and standard deviation was 6.014

TABLE-3.1: Association between the level of depression with selected demographic variables of geriatric peoples.

Demographic Variables	Level of Depression			χ^2 -Value	P-Value
	Normal	Mild	Severe		
Age In Years					
60- 65 Years	14	8	2	5.049 df=6 SN	P>0.05 Sig 0.538
66-70 Years	22	12	6		
71-75 Years	10	10	4		
>75 Years	5	3	4		
Gender				3.156 df=2 SN	P>0.05 Sig 0.206
Male	27	22	12		
Female Transgender	24	11	4		
Religion Status				4.163 df=2 SN	P>0.05 Sig 0.125
Hindu	33	25	12		
Christian	0	0	0		
Muslim	18	5	4		
Others					
Educational Status				4.837 df=6 SN	P>0.05 Sig 0.565
Primary	36	24	11		
Secondary	8	7	5		
Higher	6	2	0		
Secondary Illiterate	1	0	0		
Marital Status				3.479 df=6 SN	P>0.05 Sig 0.747
Married	27	17	6		
Unmarried	1	0	0		
Divorced Widow/Widowe	1 22	0 16	0 10		
No. of Child				3.516 df=6 SN	P>0.05 Sig 0.742
One	7	3	1		
Two	12	9	7		
More Than Two No Children	29 3	20 1	7 1		
Previous occupation				12.94 df=6 S	P<0.05 Sig 0.040
Private	2	1	3		
Government	4	2	0		
Labor	27	21	12		
Farmer	18	9	1		

Employment status				0.618	P>0.05
Working	26	14	8	df=2	Sig
Non- working	25	19	8	SN	0.734
staying with					
Son	39	25	9		P>0.05
Daughter	4	3	5	7.375	Sig
Separated	5	2	1	df=6	0.288
Any relative	3	3	1	SN	
Healthy Habits					P<0.05
Reading books,.	26	9	4	13.54	Sig
Exercise, yoga,	2	0	0	df=6	0.037
Listening music	3	6	1	S	
Chatting group	20	18	11		
Any Bad Habits					P>0.05
Tobacco chewing	7	5	5	6.012	Sig
Smoking	5	2	2	df=6	0.422
Alcohol	2	2	2	SN	
No bad habits	37	7	7		
Retirement pension					P>0.05
Yes	9	5	0	3.204	Sig
No	42	28	16	df=2	0.018
				SN	
1000 Rs. pension Yojana					P>0.05
Yes	20	15	3	3.317	Sig
No	31	18	13	df=2	0.189
				SN	
Type of family					P>0.05
Joint family	40	24	11	3.382	Sig
Nuclear family	8	4	4	df=4	0.496
Extended family	3	5	1	SN	
Disease status					P<0.05
Acute illness	10	13	2	13.06	Sig
Chronic illness	12	3	8	2	0.011
No illness	29	17	6	df=4	
				S	
Financial dependency					P<0.05
Independent	34	24	4	15.31	Sig
Dependent	16	7	12	df=4	0.004
Partially depended	1	2	0	S	

(S)= Significant At 0.05 Level

(Ns)= Not Significant

(S) = Significant At 0.05 Level

The table no.3.1 envisages the outcome of chi-square analysis being carried out to bring out the association between the mean difference level of depression among geriatric peoples living at rural area of Kheda district with their selected demographic variables, Age, Religion Status, Educational Status, Marital status,

Number of Child, Occupation, habits, Pension etc. were accounted for determining the association with level of depression.

Out of which Previous Occupation, Any Good Habit, Diseases status and Financial dependency of the demographic variable found significant associated at P value <0.05 with Depression level.

Hence null hypothesis was rejected and research hypothesis was accepted as above mentioned selected demographic variables **its evidence that there is significant difference between level of depression and geriatric people's demographic variables.**

Conclusion: On the basis of analysis of this study the following conclusion were drawn:

The purpose of the present study is to assess the geriatric depression among rural area of Kheda District.

The study consisted of 100 samples that were selected on the basis of Non probability convenient sampling techniques. Based on the objective the data analysis was done. After the assessing the geriatric depression among old age people score was majority had no depression 51(51%), Mild Depression 33 (33%), Severe Depression 16 (16%). This study is in line with previous studies showing the high prevalence of depression in elderly. Results suggest a proper screening for depression among elderly. The high prevalence of depression observed among older adults emphasize on the need of increased community support and availability of health care services for better care of the elderly. There is also an urgent need for greater awareness of depression among family members and community at large. At the same time, it is important to increase community support and create networks for better geriatric care, in accordance with WHO findings.

Conflict of Interest: There is not any conflict of interest between the all authors

Source of Funding: Self-funding

Ethical Clearance: The study was approved by the institutional ethical committee of Dinsha Patel college of nursing, research committee, there are total 15 members in the committee from various field. **The ethical approval reference number is DPCN-IEC/4062100003** and a formal written permission was gathered from the authority of or Principal of Institute prior to data collection

Statement of Informed consent: Yes, informed consent form was taken from the participants prior to data collection.

Acknowledgement

Special thanks to all the participants of the study and Medical officer of the selected PHC for provide us permission for data collection.

REFERENCES

1. P. Prakash 'textbook of mental health and psychiatric nursing', first edition 2020, CBS publisher and distributors, Ansari road daryaganj new Delhi- page no 210-208.
2. R. Shravani, "A guide to mental health and psychiatric nursing ", 4th edition 2016, jaypee brothers medical publisher, Kathmandu Nepal, page no – 208, 212-213.
3. <https://www.healthdirect.gov.au/depression-in-older-people#:~:text=The%20%20main%20causes%20of,del ay%20%E2%80%94%20speak%20to%20your%20doct or.>
4. Alexopoulos, George S. "Depression in the elderly." *Lancet (London, England)* vol. 365,9475 (2005): 1961-70. doi:10.1016/S0140-6736(05)66665-2.
5. Goyal, A., & Kajal, K. S. (2014). Prevalence Of Depression In Elderly Population In The Southern Part Of Punjab. *Journal Of Family Medicine And Primary Care*, 3(4), 359–361. <https://doi.org/10.4103/2249-4863.148109>
6. Barua, Ankur, et al. "Socio-demographic Factors of Geriatric Depression". *Indian Journal of Psychological Medicine*, vol. 32, no. 2, July 2010, pp. 87-92. doi: 10.4103/0253-7176.78503.
7. UNDESA (United Nation for Department of Economic and Social Affair). World Population Prospect 2019. UNDESA; 2019
8. Kumar, Abhishek et al. "Screening of Depression in Elderly Population Using a Geriatric Depression Scale in the Field Practice Area of Urban Health Training Centre Attached to SMS Medical College, Jaipur." *Cureus* vol. 13,6 e15859. 23 Jun. 2021, doi:10.7759/cureus.15859
9. Blazer DG. Depression in late life: Review and commentary. *J. Gerontol. A Biol. Sci. Med. Sci.* 2003;58:249–65. [PubMed] [Google Scholar]
10. Published in final edited form as: Vis cogn. 2014; 22(8): 1014–1017. doi:10.1080/13506285.2014.960
11. B.T Basvanthappa 'psychiatric mental health nursing', first edition 2007 , jaypee brothers , medical publisher new Delhi, page no-500-510.
12. Gail w.stuart, 'principal and practice of psychiatric nursing', 9th edition Elsevier karnal road new Delhi, page no – 305.
13. Christian A, Nagar K. Understanding Patients Experiences Living with Diabetes Mellitus: A Qualitative Study, Gujarat, India. *J Pharm Res Int.* 2021;33(58A):464-471. doi: 10.9734/jpri/2021/v33i58A34139 . Epub 2021 Dec 15. PMID: 35024509
14. Suthar DB, Nagar K. A Study To Assess The Effectiveness Of Planned Teaching Programme On Prevention Of Selected Life Style Diseases In Terms Of Knowledge And Attitude Among Male Adults At Selected PHC Of Kheda District. *Indian J Forensic Med Toxicol.* 2021 May 17;15(3):15732. doi: 10.37506/ijfmt.v15i3.15732. PMID: 34434007.
15. Jinal Patel, Rahul Parmar, Heena Solanki, Bhumi Pando, Fiza Vohra, Prachi Patel, Kailash Nagar, & Virendra Jain. (2022). Occupational Health Problems Among Tobacco Processing Factory Workers, at Kheda District Gujarat: A Cross Sectional Study. *Journal of Pharmaceutical Negative Results*, 1378–1387. <https://doi.org/10.47750/pnr.2022.13.S09.165>.
16. Prachi Patel, & Kailash Nagar. (2022). Effectiveness of Educational Package Regarding Knowledge, Attitude and Utilization of Selected Contraceptive Methods among Postnatal Mother at Selected PHC of Kheda District Gujarat. *Journal of Pharmaceutical Negative Results*, 1039–1050. <https://doi.org/10.47750/pnr.2022.13.S04.121>.
17. Ms. Shweta Prajapati, & Mr. Kailash Nagar. (2022). Safety Education on Knowledge, Attitude and Practice Towards Road Traffic Signs and Regulations Among College Students At, Gujarat. *Journal of Pharmaceutical Negative Results*, 948–958. <https://doi.org/10.47750/pnr.2022.13.S04.112>