

1 **Profiling the effect of micronutrient levels on vital cardiac markers**

2 Hari Krishnan Krishnamurthy^{1*}, Swarnkumar Reddy², Vasanth Jayaraman¹, Karthik
3 Krishna¹, Qi Song², Tianhao Wang¹, Kang Bei¹, John J. Rajasekaran¹

4 1 Vibrant Sciences LLC., San Carlos, CA, United States of America, 2 Vibrant America
5 LLC., San Carlos, CA, United States of America

6 Short title: Micronutrients and cardio markers

7 Corres: * hari@vibrantsci.com (HKK)

8

9 Corresponding Author:

10 Hari Krishnan Krishnamurthy

11 Director – Biomedical Engineering

12 Vibrant Sciences LLC

13 1021 Howard Avenue

14 San Carlos, CA 94070

15 Ph: 5094325707

16

17

18

19

20

21

22 Number of Tables: 03

23 Number of Figures: 02

24 Word count: 3815

25

26

27

28

29

30 NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

31 **Abstract**

32 Cardiovascular diseases (CVD) are among the most preventable chronic disorders accounting
33 for about one-third of general mortality around the globe. Micronutrients have been shown to
34 have a significant impact on cardiovascular health. Micronutrients have been looked at as the
35 most adoptable lifestyle choice which could reduce the burden of disease around the world. In
36 this context, it is important to study the levels of micronutrients and see their correlation to
37 cardiac disease biomarkers. The present study, has attempted to investigate the relationship
38 between the diverse class of micronutrients and serum levels of the key lipids and lipoproteins.
39 A retrospective analysis was carried out between the serum levels of micronutrients and vital
40 cardiovascular markers. The study was carried out in a group of 358 individuals tested for the
41 Cardio Health and Micronutrients Panel at Vibrant America Clinical Laboratory. The study
42 population was categorized based on the serum concentration of lipids and lipoproteins into 3
43 groups ‘Low’ ‘Normal’ and ‘High’ and the levels of micronutrients were compared among
44 these groups. The results revealed a significant association of several cardiovascular markers
45 with vitamins including Vit D, Vit E, Vit K, and minerals including zinc, iron, calcium,
46 magnesium, and amino acids including leucine, isoleucine, and valine. Quantitative analysis
47 by Pearson’s correlation exhibited a negative correlation of asparagine with serum levels of
48 cholesterol and LDL. Amino acids such as cysteine, isoleucine, and valine were found to have
49 a significant negative correlation with HDL. A positive correlation was observed between
50 valine and serum levels of LDL and Apo B. Vitamins such as Vit A, Vit D3, Vit E, and Vit K1
51 were found to have a strong positive correlation with levels of total cholesterol and
52 triglycerides. The study summarizes micronutrients and modulation of several lipid markers
53 which are critical for the management of cardiovascular diseases. Micronutrients such as
54 vitamins B1, B3, asparagine, and glutamine have a strong positive association, and fat-soluble
55 vitamins, and BCAA has a strong negative association with cardiovascular health.

56

57 Keywords: micronutrients, cardiovascular diseases, amino acids, lipids, vitamins, branched-
58 chain amino acids

59

60

61

62 **Introduction**

63 Micronutrients are the most essential nutrients for a whole range of physiological functioning
64 of a living system. Dietary deficiency of micronutrients leads to pathogenesis, progression, and
65 morbidity in various critical clinical disorders including cardiovascular diseases (CVD) [1].
66 The pathobiology of CVD remains complex and it is generally regarded to be a result of various
67 genetic predispositions interacting with environmental factors. More than 40% of the CVD are
68 related to nutritional factors and more than 90% of such cases were attributed to preventable
69 factors with nutrition as a major determinant [2]. Atherosclerosis and hypertension are the
70 predominant chronic cardiovascular diseases that are profoundly preventable with a multi-
71 component strategy including nutrient, diet, and lifestyle interventions [1]. Atherosclerosis,
72 characterized by the chronic inflammation of arteries affecting the function of the heart results
73 from the deposition of fats, cholesterol, and other substances on the inner lining of the artery
74 eventually resulting in cardiovascular disease or heart failure [3]. It is estimated that $\geq 2\%$ of
75 the adult population has atherosclerosis which tends to increase up to 10% or higher in people
76 above 70 years and continues to increase with the age of the population [4]. Hypertension is
77 another predominant CVD, which is clinically referred to as a chronic sustained increase in
78 arterial blood pressure. Hypertension is the most widespread CVD with more than 35% of the
79 population affected globally, A report from the World health organization stated this might
80 increase beyond 50% in 2025 [3]. Cardiac disease, being a complex multifunctional etiology,
81 the precise mechanism of pathophysiology remains unclear in both atherosclerosis and
82 hypertension. Atherosclerosis is characterized by the impaired flow of blood and hypertension
83 is characterized by a chronic increase in blood pressure, both these conditions result in
84 abnormal cardiac functioning and vascular damage [5].

85 Cardiovascular diseases include a wide spectrum of risk factors such as genetic, behavioral,
86 and environmental factors. In most cases CVDs commonly result from the interplay of multiple
87 risk factors, common classical risk factors include age, any pre-existing medical illness such
88 as diabetes, hypertension, predisposition genetics, or behavioral factors like smoking. Among
89 these risk factors, environmental factors are the most easily modifiable and play a pivotal role
90 in cardiovascular health. The environmental factors are the least investigated risk factors and
91 are noted as the most preventable risk factors of CVDs. Environmental factors include the
92 intake of micronutrients such as vitamins, minerals, and amino acids. The micronutrients
93 include organic compounds such as Vit A, Vit C, Vit D, Vit E, and Vit K, dietary minerals
94 including calcium, sodium, potassium, magnesium, etc, and trace elements like iron,

95 chromium, copper, selenium, and zinc [6, 7, 8, 9]. Several clinical reports have shown a
96 significant correlation between the deficiency of one or more micronutrients and cardiovascular
97 diseases. However, those reports were limited by the diversity of the micronutrients studied.
98 For instance, Rai et al. [10] evaluated the effect of two micronutrients copper and zinc on lipid
99 profiles in healthy adults, and another study by Islam et al. [6] reported that no significant
100 impact was observed by consumption of vitamin D with calcium on serum lipids. The current
101 study aimed to broadly investigate the correlation between cardiovascular markers and the
102 levels of 32 micronutrients. We also attempted to hypothesize the pathophysiology of
103 micronutrient deficiency through which the cardio biomarkers could be affected.

104 **Material and methods**

105 Study population

106 A total of 358 candidates with a mean age of 48 ± 15 years who were tested for cardiovascular
107 panel and micronutrient panel between December 2020 and July 2021 at Vibrant America
108 Clinical Laboratory were included in the study. The study was categorized as a retrospective
109 analysis of observed clinical data and hence was exempted from formal ethical review by
110 Western IRB (work order #1-1098539-1) (Washington USA). The study was conducted in a
111 free-living general population with no clinical indications of abnormalities. The subjects were
112 classified based on the serum levels of lipids and lipoproteins as detailed in Table 1.

113 Cardiovascular Panel

114 Blood samples were processed for the separation of serum and further analyzed for a
115 cardiovascular panel comprised of lipids (total cholesterol, LDL, HDL, and triglycerides),
116 apolipoproteins (Apo A1, Apo B), and a lipoprotein marker lipoprotein (a). Total cholesterol
117 was measured by the cholesterol dehydrogenase method via the Beckman Coulter AU680
118 analyzer. Serum levels of LDL, HDL, and triglycerides were measured by an enzymatic-
119 colorimetric method using the Beckman Coulter AU680 analyzer. Other cardiovascular
120 markers such as Apo A1, Apo B, and Lp (a) were also measured by a particle-enhanced
121 immunoturbidimetric assay via Roche Cobas 6000 c 501 analyzers

122 Micronutrient Panel

123 Serum levels of micronutrients were determined using Waters TQ-XS Tandem mass
124 spectrometer coupled with LCMS, Waters GC-MS, and Perkin Elmer NexION ICP-MS using
125 standard protocols.

126 Statistical Analysis

127 Clinical data were subjected to retrospective analysis from de-identified subjects using Java for
 128 windows version 1.8.161. Non-parametric Mann-Whitney U test was used to compare the
 129 micronutrients with normal and altered lipid and lipoprotein concentrations. Pearson's
 130 correlation was carried out to analyze the univariant relationship between serum lipids,
 131 lipoproteins, and micronutrients with significance set at $p < 0.05$. All statistical analysis was
 132 performed using GraphPad Prism Version 7.00 and a descriptive statistic was used to define
 133 the continuous variables (mean \pm SD, median, minimum and maximum).

134 **Results**

135 The study group comprised 165 males and 193 females with a mean age group of 47.9 ± 15.4 ,
 136 table 1 shows the descriptive statistics of the lipid and lipoprotein profile of 358 subjects. The
 137 study aimed to evaluate the significant relationship of the wide range of micronutrients
 138 including vitamins, dietary minerals, trace elements, and organic compounds with cholesterol
 139 (total, LDL, HDL), triglycerides as well as apo A, apo B, and Lp (a) levels.

140 Table 1: Serum Lipid, and Lipoprotein levels of candidates

Cardio marker	n=358	Frequency (n)	Mean \pm SD
	Male	165	48.1 ± 16.6
	Female	193	47.8 ± 14.4
Cholesterol	Low	-	-
≤ 199 mg/dL	Normal	296	183.3 ± 29.9
	High	62	265.5 ± 26.6
Low-density	Low	-	-
lipoprotein	Normal	212	98.3 ± 21.1
≤ 99 mg/dL	High	146	163.8 ± 27.7
High-density	Low	36	35.1 ± 5.4
lipoprotein	Normal	301	59.5 ± 15.6
≥ 56 mg/dL	High	2	-
Triglyceride	Low	-	-
≤ 149 mg/dL	Normal	323	87.6 ± 36.0
	High	35	314.2 ± 168.4

Apolipoprotein A1	Low	47	116.8 ± 17.0
	Normal	311	169.0 ± 30.4
≥120 mg/dL	High	2	-
Apolipoprotein B	Low	-	-
	Normal	270	88.9 ± 18.0
≤89 mg/dL	High	88	144.0 ± 30.8
Lipoprotein (a)	Low	-	-
≥30 mg/dL	Normal	157	14.1 ± 6.4
	High	84	69.4 ± 34.9

141

142 The subjects were classified based on the levels of serum lipids and lipoproteins. In the case of
 143 total cholesterol (TC), LDL, triglycerides, and Apo B the subjects were categorized based on
 144 lipid concentrations higher than the reference range and subjects within the reference range
 145 while in the case of HDL and apo A the subjects were divided into lipid concentrations lower
 146 than the reference range and subjects within the reference range. A significant statistical
 147 relationship was observed between various micronutrients and serum lipid and lipoprotein
 148 concentrations as shown in table 2. An increase in the serum levels of Vit E, Vit D3, and Mg
 149 were found to be statistically significant with increased levels of serum cholesterol and LDL.
 150 A decrease in serum levels of HDL was significantly affected by the serum levels of Vit D
 151 25(OH), asparagine, glutamine, and serine. A high level of serum triglycerides is significantly
 152 associated with increased levels of various micronutrients including Vit A, Vit D 25(OH), Vit
 153 E, Vit K1, and amino acids such as glutamine, serine, isoleucine, valine, and leucine.

154

155 Table 2: Significant association of micronutrients with serum lipids and lipoprotein

CHOLESTEROL					
	Greater than Reference		Within range (n=296)		P
	Range (n=62)				(<i>p</i> <0.05)
Vitamin E	16±5.7	14.4 (7.9-29.4)	12.9±4.2	11.9 (5.3-30.7)	0.0001
Vitamin D3	1.1±0.2	1.1 (0.4-1.7)	0.8±0.2	0.8 (0.3-1.6)	0.0001
Vitamin K1	2.1±1.5	1.5 (0.1-7.3)	1.4±1.3	1.2 (0.05-13.9)	0.0003
Calcium	9.8±0.3	9.8(9-10.6)	9.6±0.4	9.6 (8.5-11.2)	0.0001

Magnesium	2.2±0.1	2.2 (1.9-2.6)	2.1±0.1	2.1 (1.0-2.6)	0.0001
LOW-DENSITY LIPOPROTEIN					
	Greater than Reference Range (n=146)		Within range (n=212)		P (<i>p</i> <0.05)
Vitamin E	14.8±4.8	13.6(7.3-29.4)	12.3±4.2	11.2(5.3-30.7)	0.0001
Vitamin D3	1.04±0.26	1.02(0.44-1.78)	0.84±0.81	0.81(0.32-1.63)	0.0001
Iron	108±33.3	108.1(35.5-201.8)	99.1±41.4	91.4(25.3-238.9)	0.0024
Magnesium	2.23±0.16	2.2(1.5-2.66)	2.16±0.19	2.1(1.0-2.6)	0.0002
HIGH-DENSITY LIPOPROTEIN					
	Less than Reference Range (n=36)		Within range (n=301)		P (<i>p</i> <0.05)
Vitamin D 25 OH	40.7±24.9	33.6(12.4-136)	48.3±22.8	41.7(8.8-133)	0.0091
Asparagine	105.6±42.6	110.1(34.1-220.6)	53.9±11.6	52.9(28.1-108.0)	0.0001
Glutamine	212.5±164.3	124.6(61.0-671.4)	508.6±80.7	510.9(202.2- 752.4)	0.0001
Serine	128.0±36.2	124.7(61.0-282.9)	149.4±34.2	145.6(59.6-274.5)	0.0001
TRIGLYCERIDES					
	Greater than Reference Range (n=36)		Within range (n=323)		P (<i>p</i> <0.05)
Vitamin E	20.1±17	16.7(8.5-118.7)	12.9±4.3	11.9(5.3-29.4)	0.0001
Vitamin A	88.4±27.3	82(52.8-151.9)	75.8±23.3	72.1(34.1-151.5)	0.0061
Vitamin K1	3.14±2.7	2.3(0.22-13.9)	1.42±0.99	1.22(0.05-7.32)	0.0001
Vitamin D25 OH	33.9±13.6	30.7(15.4-71.2)	48.6±23.2	41.8(8.8-136)	0.0001
Glutamine	463.7±107.6	472.9(118.7-608.3)	508.4±78.7	507.6(202.2-752)	0.0290
Serine	136.5±32.8	130.6(76.0-198.5)	149.6±34.7	145.6(59.6-282.9)	0.0368
Carnitine	32.6±16.5	31.5(11.6-118.7)	26.5±7.2	26.6(4.1-50.5)	0.0014
Isoleucine	84.0±27.5	84.2(35.7-142.7)	67.1±19.1	64.6(24.0-139.7)	0.0002
Valine	275.0±63.3	281.1(175.3-442.4)	241.7±52.8	233.6(134.9- 442.7)	0.0020
Leucine	179.9±36.8	182.7(113.7-233.5)	161.7±34.1	157.4(79.7-249.2)	0.0033
APOLIPOPROTEIN A					
	Less than Reference Range (n=47)		Within range (n=323)		P (<i>p</i> <0.05)

Vitamin E	12.2±4.8	10.9(4.0-23.4)	13.5±4.7	12.4(5.6-30.7)	0.0369
Vitamin D3	0.84±0.24	0.81(0.41-1.5)	0.94±0.28	0.91(0.32-2.19)	0.0198
Asparagine	51.0±10.6	49.4(34.1-79.1)	54.5±12.1	53.3(28.1-113.0)	0.0444
Calcium	9.4±0.3	9.5(8.7-10.4)	9.6±0.39	9.6(8.5-11.1)	0.0022
Zinc	0.69±0.19	0.64(0.45-1.50)	0.71±0.15	0.69(0.39-2.23)	0.0315
Iron	85.6±36.6	78.5(25.3-182.4)	105.5±38.4	101.6(33.2-238.9)	0.0015

APOLIPOPROTEIN B

	Greater than Reference Range (n=87)		Within range (n=270)		P (<i>p</i> <0.05)
Vitamin E	16.0±5.5	15.3(7.3-29.0)	12.5±4.1	11.7(4.0-30.7)	0.0001
Vitamin A	82.4±23.7	77.7(34.1-145.8)	75.8±24.2	72.1(37.1-151.9)	0.0096
Vitamin D3	0.87±0.24	0.83(0.32-1.6)	1.09±0.30	1.08(0.44-2.19)	0.0001
Vitamin K1	1.5±1.3	1.2(0.05-13.9)	1.8±1.4	1.4(0.10-7.3)	0.0416
Selenium	146.4±24.0	144.8(93.4-210.8)	142.3±30.3	139(95.7-360.4)	0.0451
Potassium	4.5±0.35	4.5(3.6-5.6)	4.5±0.40	4.49(3.1-6.22)	0.0377
Calcium	9.7±0.3	9.7(8.7-10.6)	9.6±0.3	9.6(8.5-11.1)	0.0055
Zinc	0.73±0.11	0.72(0.39-1.0)	0.70±0.17	0.68(0.44-2.23)	0.0058
Iron	109.3±34.2	107.7(45.6-201.3)	100.8±39.9	97.7(25.3-238.9)	0.0239
Magnesium	2.2±0.1	2.2(1.9-2.9)	2.1±0.19	2.1(1.0-2.6)	0.0002
Isoleucine	72.7±19.9	70.7(34.0-121.4)	67.6±20.7	64.2(24.0-142.7)	0.0138
Valine	261.9±55.2	262.7(146.5-404.0)	240.2±54.2	232.1(134.9)	0.0004
Leucine	172±34.8	168.8(113.7-248.2)	160.5±34.2	157.4(79.7-249.2)	0.0138
Serine	140.6±33.0	135.4(75.0-209.7)	151.1±34.7	147.2(59.6-282.9)	0.0189

156

157 Decreased levels of serum Apo A are significantly associated with decreased serum levels of
 158 Vit E, Vit D3, asparagine, and minerals such as calcium, zinc, and iron. The elevated levels of
 159 Apo B associated with increasing the serum LDL and non-HDL fats are found to be statistically
 160 significant with all classes of micronutrients including vitamins, minerals, amino acids, and
 161 trace elements.

162 Quantitative analysis by Pearson's correlation showed serum concentration of cholesterol had
 163 a strong positive correlation with serum levels of Vit E ($r=0.4086$, $p<0.0001$), Vit D3
 164 ($r=0.5125$, $p<0.001$), and Vit K1 ($r=0.2334$, $p<0.0001$) The cholesterol levels are also

165 considerably correlated with Vit A and iron (<0.05). The serum levels of cholesterol were
 166 found to have a significantly negative correlation with folate and asparagine. LDL was found
 167 to have a significant positive correlation with serum levels of Vit E ($r=0.3398$, $p<0.0001$) and
 168 Vit D3 ($r=0.4774$, $p<0.0001$).

169 Pearson’s correlation was carried out to analyze the univariant relationship between serum
 170 lipids, lipoproteins, and micronutrients with significance set at $p<0.05$. Significant results are
 171 shown in Table 3. The directionality of the variation of micronutrients is represented in Fig 2.
 172 It was interesting to note the significant association of branched-chain amino acids with several
 173 markers in the cardiovascular panel. Increased levels of BCAA (branched-chain amino acids)
 174 Leucine, Isoleucine, and Valine showed significant association with decreased levels of HDL
 175 and increased levels of triglycerides Fig 1. Increased levels of Valine were shown to have a
 176 significant association with increased levels of LDL and Apo B. These associations show the
 177 significant association between branched-chain amino acids and lipid dysregulation.
 178 Additionally, amino acids asparagine and glutamine were found to be inversely correlated with
 179 serum LDL. A significant inverse correlation was observed between glutamine and serine with
 180 serum levels of Apo B.

181 Figure 1. Correlation between serum triglycerides with serum BCAA

182 a. Leucine, b. Isoleucine, c. Valine

183 Table 3: Pearson correlation between micronutrients with serum lipids and lipoproteins

	r	p
CHOLESTEROL		
Vitamin E	0.4086	<i><0.0001</i>
Vitamin A	0.1471	<i>0.0053</i>
Vitamin D3	0.5125	<i><0.0001</i>
Vitamin K1	0.2334	<i><0.0001</i>
Vitamin B9	-0.104	<i>0.0493</i>
Iron	0.1067	<i>0.0437</i>
Asparagine	-0.125	<i>0.018</i>
LOW-DENSITY LIPOPROTEIN		
Vitamin D3	0.4774	<i><0.0001</i>
Iron	0.1199	<i>0.0233</i>

Asparagine	-0.1217	<i>0.0213</i>
Glutamine	-0.1074	<i>0.0423</i>
Carnitine	0.1446	<i>0.0061</i>
Valine	0.1471	<i>0.0053</i>
HIGH-DENSITY LIPOPROTEIN		
Cysteine	-0.2289	< <i>0.0001</i>
Isoleucine	-0.2298	< <i>0.0001</i>
Valine	-0.2317	< <i>0.0001</i>
Leucine	-0.2405	< <i>0.0001</i>
TRIGLYCERIDES		
Vitamin E	0.3832	< <i>0.0001</i>
Vitamin A	0.2278	< <i>0.0001</i>
Vitamin B6	-0.1057	<i>0.0457</i>
Vitamin D3	0.1342	<i>0.011</i>
Vitamin K1	0.5732	< <i>0.0001</i>
Cysteine	0.1626	<i>0.002</i>
Isoleucine	0.2838	< <i>0.0001</i>
Valine	0.2234	< <i>0.0001</i>
Leucine	0.2139	< <i>0.0001</i>
Arginine	0.1363	<i>0.0098</i>
APOLIPOPROTEIN B		
Glutamine	-0.1388	<i>0.0086</i>
Serine	-0.1269	<i>0.0165</i>
Valine	0.1259	<i>0.0173</i>
LIPOPROTEIN (a)		
Vitamin B1	-0.1818	<i>0.0046</i>

184

185

Figure 2. Role of micronutrients in regulating serum lipids

186

Micronutrients in green font reduce the risk by modulating the particular component of the lipid

187

panel, Micronutrients in red font increase the risk by modulating the particular component of the lipid

188

panel

189

190 **Discussion**

191 Micronutrient deficiency is a major cause of the pathogenesis, progression, mortality, and
192 morbidity in developing various chronic health issues including CVD. It is regarded as a vital
193 component of the socio-economic development of a society and hence effective strategies have
194 been framed to address deficiencies at the population level by adding emphasis on dietary
195 improvement, supplementation, food fortification, and global public health control.
196 Understanding the role of individual micronutrients and proper intake of the essential
197 micronutrients plays a vital role in preventing micronutrient-related diseases. Micronutrient
198 deficiency due to malnutrition can be prevented by the above-mentioned strategies. But in the
199 context of the current study, the knowledge on the role of individual micronutrients and timely
200 dietary supplementation of micronutrients plays a crucial role in improving physiological
201 functions [11]. Micronutrients such as vitamins, minerals, and amino acids are known to
202 improve cardiovascular health by regulating the serum levels of lipids and lipoproteins.
203 However, the association of micronutrients with cardiovascular markers remains unclear [12].
204 The present study details the vital micronutrients with significant association with primary
205 cardiovascular markers such as cholesterol, LDL, HDL, triglycerides, Apo A, Apo B, and
206 Lp(a). The study specifies a wide range of micronutrients which includes 13 vitamins, 9
207 minerals, and 8 amino acids.

208 The results of the present study signify a strong positive correlation with high significance
209 between fat-soluble vitamins Vit A, Vit D3, Vit E, and Vit K1 with cholesterol and
210 triglycerides. Few studies have discussed the association of fat-soluble vitamins with lipid
211 profile, Piran et al. [13] attempted to study the association of fat-soluble vitamins A, E, and D.
212 They reported a correlation only between vitamin E and lipid profile, they also suggested the
213 supplementation of vitamin E for overweight subjects to decrease LDL-C levels. Several
214 substantial shreds of evidence have proved that significant changes in mean serum retinol levels
215 resulting from dietary intake or when supplemented to reduce the risk of cancer alter lipid
216 metabolism by increasing circulating levels of triglycerides and cholesterol. Pastorino et al.
217 [14] have stated retinol-induced liver damage by abnormal elevation of alkaline phosphatase
218 with overconsumption of vitamin A. Several reports have demonstrated the association of
219 vitamin D with serum lipid profiles, a study by Dibada [15] reported an inverse correlation
220 where the supplementation of vitamin D resulted in reducing serum cholesterol, LDL, and
221 triglycerides, but they reported no association between vitamin D and HDL. A randomized trial
222 by Barbarawi et al. [16] observed no association between vitamin D supplementation and
223 reduced risk of major adverse cardiovascular events, myocardial infarction, stroke, etc. In the

224 present result, vitamin D was found to have a positive correlation with cholesterol and
225 triglycerides. As Barbarawi et al. [16] stated the present result also showed no association with
226 serum HDL. At optimal levels of serum concentrations, vitamin D might be beneficial by
227 reducing cholesterol, LDL, and triglycerides. High doses of Vit D supplements result in
228 irregular heartbeat and can also raise blood calcium levels which lead to heart failure (Galior
229 et al., 2018). An increase in Vit E might interfere with blood clotting by inhibiting platelet
230 aggregation and Vitamin K-mediated clotting factors which might lead to hemorrhagic stroke
231 (Natural Medicines Comprehensive Database). Vitamin E below the reference range results in
232 acute heart muscle damage. Vitamins B9 (folate) were found to have a negative correlation
233 with cholesterol and Vit D3 was found to have a negative correlation with serum LDL.
234 Interestingly no vitamins exhibited any significant correlation with HDL and Apo A. The
235 current results signify that an overdose of fat-soluble vitamins could result in adverse health
236 effects by modulating serological biomarkers. Water-soluble vitamins tend to accumulate less
237 in tissues as they are readily soluble in water and excreted [18]. Among all the analyzed
238 minerals, iron was found to be positively correlated with serum levels of cholesterol ($r=0.1067$,
239 $p<0.0437$) and LDL ($r=0.1199$, $p<0.0233$).

240 Numerous studies have demonstrated the significance of micronutrients on cardiovascular
241 health. For instance, Ma et al. [19] suggested that the deficiency in levels of trace minerals such
242 as zinc, copper, iron, and selenium are directly associated with cardiovascular diseases, and
243 supplementation of these trace elements can prevent CVD. A review by Panchal et al. [20] has
244 demonstrated the importance of selenium, vanadium, and chromium in improving metabolic
245 syndrome. Li et al. [21] in their study on plasma metabolomics stated that plasma metabolites
246 and serum micronutrient levels can be used to predict the risk of CVD. Extensive research has
247 increased the understanding of the association between micronutrients and cardiac health.
248 Micronutrients such as vitamins and minerals are widely investigated, whereas amino acids are
249 hardly investigated. A recent report by Tharrey et al. [22] has detailed the association between
250 amino acid intake and cardiovascular mortality and they also reported the negative effect of a
251 few amino acids on cardiac health. The intake of protein and peptide-based health supplements
252 has increased in recent years which could result in increased levels of non-essential amino
253 acids, the effect of this on lipid and lipoprotein levels was analyzed in our study.

254 Levels of a few vital minerals such as magnesium, potassium, calcium, and zinc have also been
255 known to influence cardiac health. For instance, deficiency in levels of magnesium and
256 potassium results in elevated blood pressure. Calcium levels need to be at an optimum since

257 the deficiency causes pulmonary hypertension, while the excess results in plaque build-up in
258 arteries.

259 To our knowledge, the present study is the first to report on the association of various amino
260 acids with serum lipid and lipoprotein components. The study observed a statistical
261 significance between various amino acids and serum levels of lipids and lipoproteins as
262 represented in Fig 2. Increased levels of asparagine and glutamine were shown to decrease the
263 levels of LDL. Deprivation of asparagine and glutamine results in lethal effects on endothelial
264 cells (EC). The supplementation of these non-essential amino acids can assist in the recovery
265 of EC, can also fight against nitrification stress, and aid in angiogenesis. A previous report by
266 Luo et al. [23] on the interactive effects of asparagine and aspartate stated that the subjects with
267 type 2 diabetes were observed to have low levels of HDL-C and reported a negative correlation
268 between serum levels of asparagine and HDL-C. Serum levels of glutamine and serine were
269 found to have no influence on plasma lipids and also regular dietary supplementation of these
270 amino acids has various clinical advantages [24]. A study by Mansour et al. [25] demonstrated
271 the advantage of glutamine supplementation as it remarkably reduced the total cholesterol and
272 blood pressure in patients with type 2 diabetes. They also proposed that supplementation of
273 glutamine can be an effective pharmaconutrient in controlling diabetes, obesity, and other
274 chronic metabolic disorders. Similarly, adequate supplementation of serine could help in
275 increased antioxidant and anti-fatty streak (lesion in the development of atherosclerosis)
276 activity. The increase in the serum levels of triglycerides is significantly associated with low
277 serum levels of amino acids such as glutamine, serine, leucine, isoleucine, and valine. An
278 increase in the serum levels of triglycerides results in atherosclerosis which leads to high
279 chances of stroke and other related heart diseases [26]. A significant positive association was
280 observed between elevated levels of Apo B and low serum levels of amino acids such as serine,
281 and isoleucine. Serum levels of Apo B above adequate levels are associated with the
282 accumulation of LDL-C and non-HDL-C which are associated with various CVD.

283 The present study brings out an interesting correlation between cardiovascular markers and
284 BCAA's (branched-chain amino acids). BCAA's, leucine, isoleucine, and valine are nonpolar,
285 and hydrophobic, and are vital nitrogen sources for the synthesis of glutamine and alanine [27].
286 Several epidemiological studies have proved the association between BCAAs and increased
287 risk of type 2 diabetes [28]. The increase in the levels of BCAA's increases the oxidation of
288 BCAA's in muscles which inhibits the fatty acid oxidation and results in blunted insulin
289 signaling. Diabetes and CVD are the major causes of morbidity and mortality. A report by

290 Magnusson et al., [28] identified BCAA as a strong predictor of CVD development and also as
291 an early marker for the association between diabetes and CVD. Another study by Tobias et al.,
292 [29] proved, that circulating BCAA's as strong predictors of type 2 diabetes. The quantitative
293 analysis of BCAA's - leucine, isoleucine, and valine with the cardio markers showed a clear
294 association. A strong negative correlation between HDL and BCAAs and a strong positive
295 correlation between triglycerides and BCAAs (Table 3). This signifies the increase in the serum
296 level of BCAA decreases the HDL and increases the triglycerides. While in the case of LDL
297 and Apo B a positive correlation was observed with serum level of valine which signifies the
298 increase in the serum levels of valine also increases the levels of LDL and Apo B which could
299 result in CVD.

300 The present results identify a few micronutrients with the possibility of being pathogenic, for
301 instance, the positive correlation between vitamin D and serum cholesterol and LDL. The same
302 association has been previously reported by numerous studies including a report by Giri et al.
303 [30]. Grimes [31] has reported that in the absence of sunlight the inactive vitamin D is diverted
304 to the synthesis of cholesterol. Even so, intestinal absorption is the only source of fat-soluble
305 vitamins and their mechanism of intestinal absorption remains unclear. Recent studies revealed
306 the participation of several membrane proteins in cholesterol absorption evidenced by
307 advancements in genome-editing, genome-wide association, and gene mutation analysis on
308 cholesterol and intense studies in cholesterol absorption inhibitors. Interestingly, these analyses
309 also revealed that cholesterol transporters can also transport fat-soluble vitamins [32]. In
310 association with these studies, the present study identifies the serum levels of fat-soluble
311 vitamins as effective markers of elevated cholesterol transportation and can be used as an early
312 predictor of CVD. It's a well-established fact that increased serum BCAA was associated with
313 elevated triglycerides and reduced HDL [33]. The present results also identify a similar
314 pathology between serum HDL, Triglycerides and BCAA. The BCAA in the present study
315 exhibited a significant negative correlation with HDL and a strong positive correlation with
316 triglycerides. Consumption of energy-dense and high-palatable foods has increased in recent
317 years, and consumption of BCAAs is one such major behavioral change received considerable
318 attention in recent times. The most recent study by Latimer et al. (2021) reported that an
319 increase in BCAA consumption resulted in the dramatic growth of the heart and also increased
320 the progression of cardiac diseases.

321 One of the main limitations of our study is that it includes data from free-living people with
322 limited information on diet and lifestyle choices. The current study signifies the association of

323 vital cardio markers with a diverse class of micronutrients. Further research is required for a
324 better understanding of the clinical significance of these micronutrients and also to understand
325 the biochemical interference of vitamins, minerals, and amino acids in lipid metabolism.

326 **Conclusions**

327 The current study examines the explicit correlation of micronutrients with various cardio
328 markers. Given the broad set of micronutrients evaluated their significance in regulating vital
329 cardio markers was better understood. The study highlights the negative correlation of various
330 vitamins and amino acids on cardiovascular health which was a significant observation. The
331 deficiency of various micronutrients leads to a significant increase in cardiovascular disease
332 risk due to dysregulation of lipid and lipoprotein markers. Similarly increased levels of certain
333 micronutrients especially branched-chain amino acids lead to increased cardiovascular risk due
334 to lipid and lipoprotein dysregulation. It is important to look at both deficiencies and
335 overconsumption of micronutrients to optimize nutrient intake. The study suggests the proper
336 monitoring of body vitals and optimal intake of micronutrients are to be considered for their
337 effective roles in metabolic pathways to implement a risk reduction strategy for cardiovascular
338 health.

339

340

341

342

343

344

345

346

347

348

349

350

351

352 **Acknowledgments**

353 We thank Qingqing Yue, Junior Graphic Designer for assisting with designing figures and
354 Vibrant America LLC for supporting this research.

355 **Statement of Ethics**

356 An ethics statement was not required for this study type

357 **Author Contributions**

358 Hari Krishnamurthy, Karthik Krishna, and Tianhao Wang performed the research. Hari
359 Krishnamurthy, John J. Rajasekaran, Karenah Rajasekaran, and Vasanth Jayaraman designed
360 the study. Qi Song, Kang Bei, and Swarnkumar Reddy analyzed the data. Hari Krishnamurthy
361 and Swarnkumar Reddy wrote the article.

362 **Conflicts of Interest**

363 Krishnamurthy, Jayaraman, Krishna, Wang, Bei, and Rajasekaran are employees of Vibrant
364 Sciences LLC. Reddy, Song, and Rajasekaran are employees of Vibrant America LLC. Vibrant
365 America is a commercial diagnostic lab that could benefit from increased testing of
366 micronutrients and cardiovascular biomarkers.

367 **Institutional Review Board Statement**

368 IRB exemption (work order #1-1098539-1) was determined by the Western Institutional
369 Review Board (WIRB) for Vibrant America Biorepository to use de-linked and deidentified
370 remnant human specimen and medical data for research purposes.

371 **Data Availability**

372 All relevant data are within the manuscript and its Supporting Information files. Any
373 additional data will be available upon request from Vibrant America LLC by sending an
374 email to our bioinformatics team at bioinformatics@vibrant-america.com or
375 hari@vibrantsci.com

376 **Supplementary Material**

377 Table S1: Association of serum cholesterol with micronutrients
378 Table S2: Association of serum LDL with micronutrients
379 Table S3: Association of serum HDL with micronutrients
380 Table S4: Association of serum triglycerides with micronutrients
381 Table S5: Association of serum Apo A with micronutrients
382 Table S6: Association of serum Apo B with micronutrients
383 Table S7: Association of serum Lipoprotein A with micronutrients
384 Table S8: Pearson correlation between serum cholesterol and micronutrients
385 Table S9: Pearson correlation between serum LDL and micronutrients
386 Table S10: Pearson correlation between serum HDL and micronutrients
387 Table S11: Pearson correlation between serum triglycerides and micronutrients
388 Table S12: Pearson correlation between serum Apo A and micronutrients
389 Table S13: Pearson correlation between serum Apo B and micronutrients
390 Table S14: Pearson correlation between serum lipoprotein A and micronutrients

391 **References**

- 392 1. Getz, G.S. and Reardon, C.A., 2007. Nutrition and cardiovascular disease. *Arterioscler*
393 *Thromb Vasc Biol.* **2007**, **27**, 2499.
- 394 2. Tappia, P.S. and Blewett, H. Nutrition and cardiovascular health. *Int. J. Mol. Sci.* **2020**,
395 *21(7)*, 2284.
- 396 3. Chiu, H.F., Venkatakrishnan, K., Golovinskaia, O. and Wang, C.K. Impact of
397 Micronutrients on Hypertension: Evidence from Clinical Trials with a Special Focus
398 on Meta-Analysis. *Nutrients* **2021**, *13*, 588.
- 399 4. Kkeveetil, C.V., Thomas, G. and Chander, S.J.U. Role of micronutrients in congestive
400 heart failure: A systematic review of randomized controlled trials. *Tzu Chi Medical*
401 *Journal.* **2016** *28 (4)*, 143-150.
- 402 5. Ekpenyong CE. Micronutrient vitamin deficiencies and cardiovascular disease risk:
403 advancing current understanding. *Eur J Prev Med.* **2017**, *5*, 1.

- 404 6. Islam MZ, Shamim AA, Ahmed A, Akhtaruzzaman M, Kärkkäinen M, Lamberg-
405 Allardt C. Effect of vitamin D, calcium and multiple micronutrients supplementation
406 on lipid profile in pre-menopausal Bangladeshi garment factory workers with
407 hypovitaminosis D. *J Health Popul Nutr.* **2014**, *32*, 687.
- 408 7. Lima, L.W., Checchio, M.V., dos Reis, A.R., de Cássia Alves, R., Tezzoto, T. and
409 Gratão, P.L. Selenium restricts cadmium uptake and improve micronutrients and
410 proline concentration in tomato fruits. *Biocatal. Agric. Biotechnol.* **2019**, *18*, 101057.
- 411 8. Liu, L., Yin, X., Ikeda, K., Sullivan, D.H. and Eisen, H.J. Micronutrients, inflammation
412 and congestive heart failure among the elderly: nutritional perspectives on primary
413 prevention and clinical treatment. *Clin. Exp. Pharmacol. Physiol.* **2007**, *34*, S14-S16.
- 414 9. Cascino TM, Hummel SL. Nutrient deficiencies in heart failure: a micro problem with
415 macro effects? *J. Am. Heart Assoc.* **2018**, *7(17)*, e010447.
- 416 10. Rai K, N., Kumari, N. S., Gowda Km, D., & Kr, S. The Evaluation of Micronutrients
417 and Oxidative Stress and their Relationship with the Lipid Profile in Healthy adults. *J.*
418 *Clin. Diagnostic Res.* **2013**, *7(7)*, 1314–1318.
- 419 11. Bruins MJ, Kupka R, Zimmermann MB, Lietz G, Engle-Stone R, Kraemer K.
420 Maximizing the benefits and minimizing the risks of intervention programs to address
421 micronutrient malnutrition: symposium report. *Matern Child Nutr.* **2016**, *12*, 940.
- 422 12. Ma, X., Jiang, S., Yan, S., Li, M., Wang, C., Pan, Y., Sun, C., Jin, L., Yao, Y. and Li,
423 B. Association between copper, zinc, Iron, and selenium intakes and TC/HDL-C ratio
424 in US adults. *Biol. Trace Elem. Res.* **2020**, *197(1)*, 43-51.
- 425 13. Piran, S., Sarmasti, S., Shabani, M., Kakavandi, N., Hosseni, B., Khosravi, M., Resaee,
426 S., Soltanmohammadi, E., Naseri, F., Mohammadi, A. and Najafi, M. Association
427 Between Fat-soluble Vitamins and Lipid Profile in the Overweight Population. *Recent*
428 *Pat. food, Nutr. Agric.* **2020**, *11(1)*, 56-62.
- 429 14. Pastorino, U., Chiesa, G., Infante, M., Soresi, E., Clerici, M., Valente, M., Belloni, P.A.
430 and Ravasi, G., 1991. Safety of high-dose vitamin A. *Oncology*, **1991**, *48(2)*, 131-137.
- 431 15. Dibaba DT. Effect of vitamin D supplementation on serum lipid profiles: a systematic
432 review and meta-analysis. *Nutr Rev.* **2019**, *77(12)*, 890-902.
- 433 16. Barbarawi M, Kheiri B, Zayed Y, Barbarawi O, Dhillon H, Swaid B, Yelangi A, Sundus
434 S, Bachuwa G, Alkotob ML, Manson JE. Vitamin D Supplementation and
435 Cardiovascular Disease Risks in More Than 83 000 Individuals in 21 Randomized
436 Clinical Trials: A Meta-analysis. *JAMA Cardiol.* **2019**, *4(8)*, 765-776.

- 437 17. Galior K, Grebe S, Singh R. Development of Vitamin D Toxicity from Overcorrection
438 of Vitamin D Deficiency: A Review of Case Reports. *Nutrients*, **2018**, *10(8)*, 953.
- 439 18. Lykstad J, Sharma S. *Biochemistry, Water Soluble Vitamins*. StatPearls Publishing,
440 Treasure Island (FL); 2021. PMID: 30860745.
- 441 19. Ma, X., Jiang, S., Yan, S., Li, M., Wang, C., Pan, Y., Sun, C., Jin, L., Yao, Y. and Li,
442 B. Association between copper, zinc, iron, and selenium intakes and TC/HDL-C ratio
443 in US adults. *Biol. Trace Elem. Res.* **2020**, *197(1)*, 43-51.
- 444 20. Panchal, S.K., Wanyonyi, S. & Brown, L. Selenium, Vanadium, and Chromium as
445 Micronutrients to Improve Metabolic Syndrome. *Curr Hypertens Rep.* **2017**, *19(3)*, 1-
446 11.
- 447 21. Li, K.J., Jenkins, N., Luckasen, G., Rao, S., and Ryan, E.P. Plasma metabolomics of
448 children with aberrant serum lipids and inadequate micronutrient intake. *PLoS one.*
449 **2018**, *13(10)*, p.e0205899.
- 450 22. Tharrey, M., Mariotti, F., Mashchak, A., Barbillon, P., Delattre, M., Huneau, J.F. and
451 Fraser, G.E. Patterns of amino acid intake are strongly associated with cardiovascular
452 mortality, independently of the sources of protein. *Int J Epidemiol*, **2020**, *49(1)*, 312-
453 321
- 454 23. Luo, H.H., Feng, X.F., Yang, X.L., Hou, R.Q. and Fang, Z.Z., 2020. Interactive effects
455 of asparagine and aspartate homeostasis with sex and age for the risk of type 2 diabetes
456 risk. *Biol. Sex Differ.* **2020**, *11(1)*, 1-10.
- 457 24. Xiao, F., Du, Y., Lv, Z., Chen, S., Zhu, J., Sheng, H. and Guo, F. Effects of essential
458 amino acids on lipid metabolism in mice and humans. *J. Mol. Endocrinol.* **2016**, *57(4)*,
459 223-231.
- 460 25. Mansour, A., Mohajeri-Tehrani, M.R., Qorbani, M., Heshmat, R., Larijani, B. and
461 Hosseini, S. Effect of glutamine supplementation on cardiovascular risk factors in
462 patients with type 2 diabetes. *Nutrition*, **2015**, *31(1)*, 119-126.
- 463 26. Park, M.S.C. and Liepa, G.U. Effects of dietary protein and amino acids on the
464 metabolism of cholesterol-carrying lipoproteins in rats. *J Nutr.* **1982**, *112(10)*, 1892-
465 1898.
- 466 27. Szpetnar, M., Luchowska-Kocot, D., Boguszewska-Czubara, A. et al. The Influence of
467 Manganese and Glutamine Intake on Antioxidants and Neurotransmitter Amino Acids
468 Levels in Rats' Brain. *Neurochem Res.* **2016**, *41*, 2129–2139.

- 469 28. Martin Magnusson, Olle Melander, Bo Israelsson, Anders Grubb, Leif Groop, Stefan
470 Jovinge; Elevated Plasma Levels of Nt-proBNP in Patients with Type 2 Diabetes
471 Without Overt Cardiovascular Disease. *Diabetes Care*, **2004**, 27 (8), 1929–1935.
- 472 29. Tobias, D. K., Mora, S., Verma, S., & Lawler, P. R. Altered branched chain amino acid
473 metabolism: toward a unifying cardiometabolic hypothesis. *Curr. Opin. Cardiol.* **2018**,
474 33(5), 558–564.
- 475 30. Giri, R., Rai, R., Verma, R.K. and Verma, S. Correlation between Vitamin D and lipid
476 profile in patients with ischemic stroke. *Int J Res Med Sci.* **2016**, 4(6), 2309-12.
- 477 31. Grimes, D.S., Hindle, E. and Dyer, T. Sunlight, cholesterol and coronary heart disease.
478 *QJM: Int. J. Med.* **1996**, 89(8), 579-590.
- 479 32. Yamanashi, Y., Takada, T., Kurauchi, R., Tanaka, Y., Komine, T. and Suzuki, H.
480 Transporters for the intestinal absorption of cholesterol, vitamin E, and vitamin K. *J.*
481 *Atheroscler. Thromb.* **2017**, 24, 16007.
- 482 33. Yang, P., Hu, W., Fu, Z., Sun, L., Zhou, Y., Gong, Y., Yang, T. and Zhou, H., 2016.
483 The positive association of branched-chain amino acids and metabolic dyslipidemia in
484 Chinese Han population. *Lipids Health Dis.* **2016**, 15(1), 1-8.
- 485 34. Latimer, M.N., Sonkar, R., Mia, S., Frayne, I.R., Carter, K.J., Johnson, C.A., Rana, S.,
486 Xie, M., Rowe, G.C., Wende, A.R. and Prabhu, S.D., 2021. Branched chain amino acids
487 selectively promote cardiac growth at the end of the awake period. *J. Mol. Cell. Cardiol.*
488 **2021**, 157, 31-44.

489

490

491

492

493

494

495

496

497

498

499
500
501
502
503
504

Table S1: Association of serum cholesterol with micronutrients

	Greater than Reference Range (n=62)		Within range (n=278)		P (P<0.05)
ASPARAGINE	53.8±12.7	51.4(34.1-93.1)	54.1±12.7	52.9(28-113.0)	0.8067
GLUTAMINE	505.7±82.7	512.8(210-752)	504.2±80.6	503.1(202.2-741)	0.8270
SERINE	144.9±34.3	138(75-213.4)	149.3±35.3	146.1(59.6-282.9)	0.2869
CYSTEINE	17.3±8.7	17.0(2.2-46.2)	17.7±8.9	16.2 (1.3-45.2)	0.8426
SELENIUM	139±32	140.1(-47-207.5)	143.3±30.5	139 (93.4-360.4)	0.8771
VITAMIN E	16±5.7	14.4 (7.9-29.4)	12.9±4.2	11.9 (5.3-30.7)	0.0001
CHOLINE	14.4±5.6	13.3(6.1-35.1)	14.1±5.7	12.5(1.0-39.6)	0.5605
CARNITINE	28.8±7.7	29.1 (12.6-44.3)	26.5±7.1	26.6(4.1-50.5)	0.0317
SODIUM	141.9±2.9	142 (137-148)	140.1±22.0	141 (-222-155)	0.2091
POTASSIUM	4.5±0.4	4.6 (3.1-5.6)	4.5±0.47	4.4 (3.72-6.22)	0.1658
CALCIUM	9.8±0.3	9.8(9-10.6)	9.6±0.4	9.6 (8.5-11.2)	0.0001
MANGANESE	0.6±0.2	0.6 (0.3-1.7)	0.7±0.4	0.6 (0.2-6.8)	0.6281
ZINC	0.7±0.1	0.7 (0.4-.9)	0.7±0.1	0.6 (0.3-2.2)	0.0160
COPPER	1.0±0.2	0.9 (0.6-2.1)	1.05±0.3	1.0 (0.3-3.4)	0.7309
CHROMIUM	0.2±0.1	0.2 (0.05-0.6)	0.2±0.1	0.22 (0.01-1.7)	0.6393
IRON	108.7±33.3	102.0 (34.8-206.5)	101.9±38.8	99.9 (25.3-238.9)	0.1217
MAGNESIUM	2.2±0.1	2.2 (1.9-2.6)	2.1±0.1	2.1 (1.0-2.6)	0.0001
VITAMIN A	80.7±25.1	77.8(34.1-142.9)	76.0±23.6	71.5 (37.4-151.9)	0.0942
VITAMIN_B1	21.3±13.5	19.4 (1.0-70.3)	21.6±15.0	18.3 (0.8-136.8)	0.8489
VITAMIN_B2	26.8±25.3	19.4 (6.2-174.1)	28.1±21.9	21.9 (4.8-140.8)	0.4462
VITAMIN_B3	17.4±7.2	18.1 (3.4-34.6)	19.7±8.8	19.1 (2.7-110.1)	0.0559
VITAMIN_B6	15.3±13.0	11.3 (1.1-70.8)	17.5±19.6	12.3 (1.6-236.7)	0.4704
VITAMIN_B12	787.9±358.1	758 (245.1-1958)	719±319	650 (196-1923)	0.1395
VITAMIN_B5	99.6±74.4	70.8 (23.5-315.2)	107.5±89.2	71.0 (13.8-482.1)	0.6705
VITAMIN_C	0.4±0.2	0.4 (.05-1.2)	0.4±0.3	0.3 (.05-3.74)	0.5634
VITAMIN_D3	1.1±0.2	1.1 (0.4-1.7)	0.8±0.2	0.8 (0.3-1.6)	0.0001
VITAMIN_K1	2.1±1.5	1.5 (0.1-7.3)	1.4±1.3	1.2 (0.05-13.9)	0.0003
VITAMIN_K2	0.8±1.7	0.2 (0.11-11.2)	0.6±1.0	0.2 (0.1-11.0)	0.1909
FOLATE	13.6±4.4	13.6(2.9-19.9)	13.6±3.9	13.9 (2.8-19.9)	0.8426
VITAMIN D 25 OH	50.2±26.6	41.2 (11.4-133)	46.5±22.5	40.9 (8.8-136)	0.5307

ISOLEUCINE	68.0±17.2	66.9 (35.7-115.7)	68.4±21.1	64.6 (24.0-142.7)	0.6996
VALINE	255.7±54.7	248.3 (146.5-400)	242.3±55.4	233.2 (134.9-442)	0.0484
LEUCINE	166.5±37.4	159.0 (79.7-247.6)	162.6±34.2	159.1 (80.3-249.2)	0.5826
CITRULLINE	31.4±7.6	31.3 (17.3-52.1)	29.6±8.1	28.8 (10.1-69.3)	0.0774
ARGININE	121.8±32	117.5 (65.1-220.6)	123.2±33.8	119.1 (49.6-299.2)	0.7459

505

506

507

508

Table S2: Association of serum LDL with micronutrients

	Greater than Reference Range (n=146)		Within range (n=212)		P (P<0.05)
ASPARAGINE	52.9±11.2	51.7(31.1-108.0)	54.8±12.7	53.1(28.1-113.0)	0.2238
GLUTAMINE	496.9±80.5	502.3(204.1-752.4)	510.6±80.3	507.6(202.2-741.1)	0.1594
SERINE	148.4±33.9	144.1(75-227.9)	148.4±35.4	146.4(59.6-282.9)	0.9280
CYSTEINE	18.5±8.7	17.9(2.8-46.2)	17.3±9.1	15.6(1.3-45.2)	0.1121
SELENIUM	142.2±23.6	139.4(93.4-210.8)	143.6±32.1	141.4(95.7-360.4)	0.8131
VITAMIN E	14.8±4.8	13.6(7.3-29.4)	12.3±4.2	11.2(5.3-30.7)	0.0001
CHOLINE	14.2±5.6	13.0(5.8-39.6)	14.0±5.7	12.6(1.0-36.6)	0.7838
CARNITINE	28.0±7.3	28.3(10.2-50.5)	26.1±7.1	26.3(4.1-43.6)	0.0258
SODIUM	141.7±3.4	142 (133-151)	139.7±25.1	141(129-155)	0.3443
POTASSIUM	4.5±0.3	4.5(3.1-5.6)	4.5±0.4	4.5(3.6-6.2)	0.7340
CALCIUM	9.6±0.3	9.6 (8.5-10.6)	89.6±0.4	9.6(8.6-11.2)	0.5248
MANGANESE	0.69±0.26	0.63(0.2-1.85)	0.74±0.54	0.63(0.3-6.82)	0.9838
ZINC	0.72±0.12	0.71(0.39-1.39)	0.70±0.18	0.67(0.44-2.23)	0.0373
COPPER	1.05±0.29	1.02(0.37-2.53)	1.05±0.31	1.0 (0.4-3.4)	0.8362
CHROMIUM	0.2±0.1	0.2(0.02-1.0)	0.26±0.17	0.2(0.011-1.79)	0.4714
IRON	108±33.3	108.1(35.5-201.8)	99.1±41.4	91.4(25.3-238.9)	0.0024
MAGNESIUM	2.23±0.16	2.2(1.5-2.66)	2.16±0.19	2.1(1.0-2.6)	0.0002
VITAMIN A	78.3±22.7	74.6(34.1-151.5)	76.2±24.8	72.2(37.1-151.9)	0.2012
VITAMIN_B1	20.8±13.5	18.0(1.0-75.9)	22.4±15.9	18.8(0.83-136.8)	0.4553
VITAMIN_B2	27.2±22.8	21.1(5.7-174.1)	29.2±25.8	21.6(4.8-222.5)	0.7505
VITAMIN_B3	19.4±10.5	18.6(3.4-110.1)	19.0±6.7	18.2(2.7-40.6)	0.7766
VITAMIN_B6	16.0±13.2	12.2(1.1-84.2)	18.2±21.3	12.3(1.4-236.7)	0.6179
VITAMIN_B12	744.3±328.8	657.2(245.1-1958)	726.1±323.4	663.9(196-1923)	0.6856
VITAMIN_B5	103.6±87.5	70.6(13.8-482.1)	111.1±9.2	71.5(19.2-407.4)	0.5429
VITAMIN_C	0.45±0.29	0.41(0.05-2.62)	0.45±0.31	0.39(0.08-3.74)	0.7418
VITAMIN_D3	1.04±0.26	1.02(0.44-1.78)	0.84±0.81	0.81(0.32-1.63)	0.0001
VITAMIN_K1	1.65±1.2	1.29(0.05-7.32)	1.56±1.4	1.25(0.15-13.9)	0.3875
VITAMIN_K2	0.68±1.2	0.28(0.10-11.2)	0.72±1.17	0.29(0.082-11.0)	0.5063
FOLATE	13.8±4.0	14.1(2.9-19.9)	13.6±3.9	13.9(2.8-19.9)	0.5133
VITAMIN D 25 OH	47.8±22.6	41.5(10.4-133)	46.3±23.5	40.5(8.8-136)	0.3757

ISOLEUCINE	69.0±20.1	67.0(31.0-139.7)	68.6±21.0	64.4(24.0-142.7)	0.6712
VALINE	250.5±55.1	242.0(146.5-442.7)	241.1±54.2	232.5(134.9-442.4)	0.0903
LEUCINE	166.4±35.1	159.3(79.7-248.2)	161.5±34.4	158.2(87.7-249.2)	0.2160
CITRULLINE	30.1±7.6	29.4(10.3-51.8)	29.9±8.3	28.8(10.1-69.3)	0.7509
ARGININE	122.2±33.2	118.9(61.0-220.6)	123.3±33.8	118.4(49.6-299.2)	0.7600

509

510

511

512

Table S3: Association of serum HDL with micronutrients

	Less than Reference Range (n=36)		Within range (n=301)		P (P<0.05)
ASPARAGINE	105.6±42.6	110.1(34.1-220.6)	53.9±11.6	52.9(28.1-108.0)	0.0001
GLUTAMINE	212.5±164.3	124.6(61.0-671.4)	508.6±80.7	510.9(202.2-752.4)	0.0001
SERINE	128.0±36.2	124.7(61.0-282.9)	149.4±34.2	145.6(59.6-274.5)	0.0001
CYSTEINE	20.4±7.8	20.1(6.7-42.7)	17.8±9.5	16.5(2.5-46.2)	0.0364
SELENIUM	141.1±24.8	142.2(96.8-210.8)	143.0±31.5	140.6(93.4-360.4)	0.7682
VITAMIN E	15.8±5.3	16.2(6-30.7)	13.2±4.5	12.1(5.3-29.4)	0.0011
CHOLINE	13.3±4.8	11.7(6.1-27.3)	14.2±5.8	12.9(1.0-39.6)	0.3783
CARNITINE	27.3±6.6	26.9(12.8-40.3)	26.9±7.3	27.2(4.1-50.5)	0.7748
SODIUM	141.5±3.5	142(134-150)	141.4±3.2	141(129-155)	0.9044
POTASSIUM	4.4±0.43	4.3(3.1-5.3)	4.5±0.39	4.5(3.6-6.2)	0.2055
CALCIUM	9.6±0.5	9.5(8.8-11.2)	9.6±0.3	9.6(8-11.1)	0.3304
MANGANESE	0.70±0.2	0.68(0.32-1.2)	0.72±0.4	0.62(0.2-6.8)	0.3071
ZINC	0.68±0.1	0.6(0.4-0.9)	0.7±0.1	0.6(0.3-2.2)	0.2098
COPPER	1.1±0.3	1.0(0.4-2.5)	1.04±0.30	0.9(0.3-3.4)	0.3136
CHROMIUM	0.23±0.10	0.21(0.06-0.52)	0.2±0.16	0.2(0.01-1.7)	0.6318
IRON	92.2±33.6	86.5(25.3-192.1)	104.1±38.2	100(29.8-238.9)	0.0983
MAGNESIUM	2.1±0.2	2.1(1.2-2.5)	2.1±0.18	2.2(1.0-2.6)	0.8566
VITAMIN A	84.8±27.0	80.0(43.3-151.9)	76.7±23.9	72.3(34.1-151.5)	0.0661
VITAMIN_B1	21.1±14.0	17.7(1.0-56.8)	21.4±15.0	18.4(0.8-136.8)	0.8646
VITAMIN_B2	21.8±14.3	17.2(6.4-64.2)	28.1±21.8	22.0(4.8-140.8)	0.0848
VITAMIN_B3	19.4±5.5	19.4(5.4-30.3)	19.4±8.8	18.9(2.7-110.1)	0.7625
VITAMIN_B6	13.5±13.6	9.1(2.9-80.9)	17.6±19.0	12.4(1.1-236.7)	0.1048
VITAMIN_B12	642±304.0	532.2(245.1-1593)	742.3±331.1	674.3(150-1958)	0.0509
VITAMIN_B5	104.3±84.9	71.8(25.8-330.0)	106.0±87.9	68.3(13.8-482.1)	0.8070
VITAMIN_C	0.43±0.23	0.39(0.08-0.88)	0.44±0.29	0.39(0.05-3.74)	0.9910
VITAMIN_D3	0.93±0.24	0.93(0.41-1.46)	0.92±0.27	0.88(0.32-1.78)	0.5533
VITAMIN_K1	2.3±2.7	1.6(0.2-13.9)	1.5±1.0	1.2(0.05-7.3)	0.0290
VITAMIN_K2	0.74±0.87	0.32(0.10-3.8)	0.71±1.2	0.28(0.082-11.2)	0.2940
FOLATE	12.0±4.1	11.0(2.9-19.6)	13.7±3.9	13.9(2.8-19.9)	0.0146
VITAMIN D 25 OH	40.7±24.9	33.6(12.4-136)	48.3±22.8	41.7(8.8-133)	0.0091

ISOLEUCINE	76.8±24.3	73.2(35.7-142.7)	67.9±20.2	65.1(24.0-139.7)	0.0193
VALINE	256.6±57.0	247.8(175.3-363.9)	244.1±54.8	234.8(134.9-442.1)	0.2253
LEUCINE	169.3±35.8	168.7(113.7-241.0)	163.4±34	158.3(80.3-249.2)	0.3703
CITRULLINE	28.3±8.3	28.7(10.3-44.9)	30.2±8.0	29.1(10.1-69.3)	0.3014
ARGININE	129.6±33.0	119.5(78.3-205.3)	122.5±34.1	119.2(49.6-299.2)	0.3083

513

514

515

516

Table S4: Association of serum triglycerides with micronutrients

	Greater than Reference Range (n=36)		Within range (n=323)		P (P<0.05)
ASPARAGINE	53.4±15.5	49.9(31.1-118.7)	54.3±12.2	53.1(28.1-113.0)	0.3363
GLUTAMINE	463.7±107.6	472.9(118.7-608.3)	508.4±78.7	507.6(202.2-752)	0.0290
SERINE	136.5±32.8	130.6(76.0-198.5)	149.6±34.7	145.6(59.6-282.9)	0.0368
CYSTEINE	22.1±18.1	19.4(4.9-118.7)	17.6±9.1	15.7(1.3-46.2)	0.0573
SELENIUM	142.6±22.8	141(96.8-207.5)	143.0±29.5	140.2(93.4-360.4)	0.7712
VITAMIN E	20.1±17	16.7(8.5-118.7)	12.9±4.3	11.9(5.3-29.4)	0.0001
CHOLINE	17.8±18.1	14.1(7.0-118.7)	14.0±5.7	12.6(1.0-39.6)	0.1661
CARNITINE	32.6±16.5	31.5(11.6-118.7)	26.5±7.2	26.6(4.1-50.5)	0.0014
SODIUM	141.5±4.8	142.5(118.7-150)	141.5±3.4	141(129-155)	0.3449
POTASSIUM	7.7±19.0	4.5(4.1-118.7)	4.5±0.3	4.5 (3.1-6.2)	0.6792
CALCIUM	12.8±18.1	9.7(9.0-118.7)	9.6±0.4	9.6(8.2-11.1)	0.0518
MANGANESE	0.7±0.2	0.67(0.32-1.5)	0.72±0.46	0.63(0.2-6.8)	0.1321
ZINC	0.68±0.13	0.702(0.4-1.0)	0.71±0.16	0.69(0.39-2.23)	0.5508
COPPER	1.01±0.23	1.0(0.48-1.5)	1.0±0.3	1.0(0.3-3.4)	0.6959
CHROMIUM	0.23±0.11	0.22(0.05-0.51)	0.25±0.16	0.22(0.01-1.7)	0.6972
IRON	101.2±29.1	97.5(50.7-160)	102.8±39.5	100(25.3-238.9)	0.9361
MAGNESIUM	2.2±0.15	2.2(1.9-2.6)	2.1±0.19	2.2(1.0-2.6)	0.7556
VITAMIN A	88.4±27.3	82(52.8-151.9)	75.8±23.3	72.1(34.1-151.5)	0.0061
VITAMIN_B1	19.8±15.7	16.9(1.0-66.4)	21.9±14.9	18.6(0.8-136.8)	0.2701
VITAMIN_B2	26.8±15.5	24.5(6.2-64.7)	27.9±22.9	20.8(4.8-174.1)	0.4594
VITAMIN_B3	17.5±5.6	16.9(8.7-27.8)	19.4±8.7	18.9(2.7-110.1)	0.1979
VITAMIN_B6	13.2±10.8	9.8(2.9-49.1)	17.7±19.0	12.4(1.1-236.7)	0.1156
VITAMIN_B12	688.9±321.3	552.1(281.5-1593)	738.2±325.9	667.8(196-1958)	0.2503
VITAMIN_B5	112.6±97.3	70.5(27.4-437.4)	107.5±88.3	71.2(13.8-482.1)	0.6561
VITAMIN_C	0.51±0.42	0.41(0.14-2.6)	0.44±0.28	0.40(0.05-3.74)	0.6053
VITAMIN_D3	0.96±0.29	0.93(0.40-1.42)	0.91±0.26	0.88(0.32-1.78)	0.3611
VITAMIN_K1	3.14±2.7	2.3(0.22-13.9)	1.42±0.99	1.22(0.05-7.32)	0.0001
VITAMIN_K2	1.07±1.9	0.32(0.10-11.2)	0.66±1.09	0.28(0.082-11.0)	0.1134
FOLATE	12.4±4.5	12.6(2.8-19.6)	13.8±3.9	14.1(2.9-19.9)	0.0839
VITAMIN D 25 OH	33.9±13.6	30.7(15.4-71.2)	48.6±23.2	41.8(8.8-136)	0.0001

ISOLEUCINE	84.0±27.5	84.2(35.7-142.7)	67.1±19.1	64.6(24.0-139.7)	0.0002
VALINE	275.0±63.3	281.1(175.3-442.4)	241.7±52.8	233.6(134.9-442.1)	0.0020
LEUCINE	179.9±36.8	182.7(113.7-233.5)	161.7±34.1	157.4(79.7-249.2)	0.0033
CITRULLINE	29.3±7.5	30.8(10.3-41.7)	30.0±8.1	28.8(10.1-69.3)	0.8478
ARGININE	134.4±38.1	125.0(76.9-211.3)	121.6±32.8	117.7(49.6-299.2)	0.0539

517

518

519

520

Table S5: Association of serum Apo A1 with micronutrients

	Less than Reference Range (n=47)		Within range (n=323)		P (P<0.05)
ASPARAGINE	51.0±10.6	49.4(34.1-79.1)	54.5±12.1	53.3(28.1-113.0)	0.0444
GLUTAMINE	498.7±78.4	490.2(320.3-719.9)	506.4±81.7	509(202.2-752.4)	0.2889
SERINE	147.0±34.8	150.5(76.0-225.4)	148.8±34.6	144.1(59.6-282.9)	0.8934
CYSTEINE	20.7±8.7	20.2(6.7-42.7)	17.7±9.2	16.4(2.2-60.3)	0.0210
SELENIUM	140.6±34.4	131.5(96.8-298.7)	143.7±28.4	141.7(93.4-360.4)	0.1749
VITAMIN E	12.2±4.8	10.9(4.0-23.4)	13.5±4.7	12.4(5.6-30.7)	0.0369
CHOLINE	13.7±5.8	13.2(6.0-36.6)	14.1±5.6	12.7(1.0-39.6)	0.5099
CARNITINE	26.3±6.5	25.2(12.8-40.3)	27.1±7.3	27.2(4.1-50.5)	0.4189
SODIUM	140.8±2.9	141(134-148)	141.6±3.5	141(129-155.0)	0.1868
POTASSIUM	4.4±0.4	4.4(3.1-5.3)	4.5±0.3	4.5(3.6-6.22)	0.0501
CALCIUM	9.4±0.3	9.5(8.7-10.4)	9.6±0.39	9.6(8.5-11.1)	0.0022
MANGANESE	0.70±0.39	0.64(0.2-2.6)	0.7±0.4	0.64(0.24-6.8)	0.2491
ZINC	0.69±0.19	0.64(0.45-1.50)	0.71±0.15	0.69(0.39-2.23)	0.0315
COPPER	1.0±0.29	1.10(0.48-1.59)	1.05±0.30	0.99(0.37-3.41)	0.5663
CHROMIUM	0.23±0.11	0.22(0.04-0.56)	0.2±0.16	0.22(0.01-1.79)	0.4849
IRON	85.6±36.6	78.5(25.3-182.4)	105.5±38.4	101.6(33.2-238.9)	0.0015
MAGNESIUM	2.1±0.22	2.16(1.2-2.5)	2.2±0.18	2.2(1.03-2.95)	0.0638
VITAMIN A	72.0±23.2	65.0(38.2-145.8)	78.3±24.4	73.5(34.1-151.9)	0.0573
VITAMIN_B1	20.5±14.3	16.8(.89-66.4)	21.7±15.1	18.5(0.83-136.8)	0.4571
VITAMIN_B2	24.0±18.8	17.3(5.3-84.3)	28.0±22.8	21.9(4.8-174.1)	0.1599
VITAMIN_B3	19.3±5.4	19.8(5.4-30.3)	19.1±8.8	18.3(2.7-110.4)	0.5138
VITAMIN_B6	20.2±34.8	12.3(2.9-236.7)	16.6±13.9	12.4(1.1-84.2)	0.9154
VITAMIN_B12	637.5±271.7	559.4(245.1-1600)	750.3±330.1	691.3(196-1958)	0.0198
VITAMIN_B5	104.0±81.5	73.7(20.2-330.0)	107.2±87.9	70.0(13.8-482.1)	0.9560
VITAMIN_C	0.41±0.24	0.32(0.086-1.04)	0.45±0.31	0.41(0.05-3.74)	0.2341
VITAMIN_D3	0.84±0.24	0.81(0.41-1.5)	0.94±0.28	0.91(0.32-2.19)	0.0198
VITAMIN_K1	1.3±1.1	0.99(0.15-5.8)	1.6±1.3	1.2(0.05-13.9)	0.1330
VITAMIN_K2	0.69±0.72	0.34(0.10-3.24)	0.72±1.2	0.28(0.082-11.2)	0.1358
FOLATE	12.1±4.0	12.2(2.2-19.6)	13.8±4.0	14.2(2.8-19.9)	0.0050
VITAMIN D 25 OH	44.1±29.5	31.4(8.8-136)	48.1±21.9	41.8(10.4-133)	0.0177

ISOLEUCINE	73.7±23.9	72.2(35.7-142.7)	68.1±20.0	65.3(24.0-139.7)	0.1804
VALINE	251.1±63.6	238.2(149.8-442.4)	244.6±53.8	236.9(134.9-442.1)	0.6595
LEUCINE	162.7±32.0	161.9(102-233.4)	163.4±35.1	158.8(79.7-249.2)	0.9295
CITRULLINE	28.2±8.1	27.8(10.3-47.9)	30.4±8.4	29.4(10.1-79.4)	0.1294
ARGININE	122.2±26.7	115.8(78.3-205.3)	123.5±34.3	120(49.6-299.2)	0.8107

521

522

523

524

Table S6: Association of serum Apo B with micronutrients

	Greater than Reference Range (n=87)		Within range (n=270)		P (P<0.05)
ASPARAGINE	53.4±11.1	53.5(34.1-79.1)	54.2±12.3	52.8(28.1-113.0)	0.7013
GLUTAMINE	489.6±84.4	492.2(204.1-752.4)	510.5±79.7	507.8(202.2-741.0)	0.0643
SERINE	140.6±33.0	135.4(75.0-209.7)	151.1±34.7	147.2(59.6-282.9)	0.0189
CYSTEINE	18.4±9.5	17.9(2.8-60.3)	18.0±9.1	16.5(2.2-45.2)	0.6201
SELENIUM	146.4±24.0	144.8(93.4-210.8)	142.3±30.3	139(95.7-360.4)	0.0451
VITAMIN E	16.0±5.5	15.3(7.3-29.0)	12.5±4.1	11.7(4.0-30.7)	0.0001
CHOLINE	14.4±5.3	13.2(6.6-39.6)	13.9±5.7	12.4(1.0-36.6)	0.3090
CARNITINE	27.8±7.7	28.2(9.8-44.3)	26.7±7.1	26.6(4.1-50.5)	0.1555
SODIUM	141.5±3.4	141(133-151)	141.5±3.4	141(129-155)	0.8920
POTASSIUM	4.5±0.35	4.5(3.6-5.6)	4.5±0.40	4.49(3.1-6.22)	0.0377
CALCIUM	9.7±0.3	9.7(8.7-10.6)	9.6±0.3	9.6(8.5-11.1)	0.0055
MANGANESE	0.73±0.29	0.67(0.34-2.1)	0.73±0.49	0.63(0.24-6.82)	0.1635
ZINC	0.73±0.11	0.72(0.39-1.0)	0.70±0.17	0.68(0.44-2.23)	0.0058
COPPER	1.0±0.32	1.0(0.6-2.5)	1.0±0.3	1.0(0.3-3.4)	0.4660
CHROMIUM	0.24±0.11	0.22(0.05-0.59)	0.25±0.17	0.22(0.011-1.7)	0.9988
IRON	109.3±34.2	107.7(45.6-201.3)	100.8±39.9	97.7(25.3-238.9)	0.0239
MAGNESIUM	2.2±0.1	2.2(1.9-2.9)	2.1±0.19	2.1(1.0-2.6)	0.0002
VITAMIN A	82.4±23.7	77.7(34.1-145.8)	75.8±24.2	72.1(37.1-151.9)	0.0096
VITAMIN_B1	19.3±14.1	16.8(1.17-70.3)	22.2±15.2	18.6(0.83-136.8)	0.0516
VITAMIN_B2	27.1±23.8	20.9(6.2-174.1)	28.5±24.8	21.2(4.8-222.5)	0.7427
VITAMIN_B3	18.7±11.9	17.8(3.4-110.1)	19.3±7.0	19.1(2.7-40.6)	0.0954
VITAMIN_B6	15.5±14.3	11.7(2.9-84.2)	17.6±19.1	12.5(1.1-236.7)	0.2511
VITAMIN_B12	750.2±345.3	659.5(290.2-1958)	730.6±318.6	666.8(196-1923)	0.8363
VITAMIN_B5	110.1±89.9	72.0(13.8-437.4)	96.5±76.8	68.6(23.5-482.1)	0.4485
VITAMIN_C	0.45±0.29	0.40(0.05-3.74)	0.45±0.32	0.39(0.05-2.6)	0.7818
VITAMIN_D3	0.87±0.24	0.83(0.32-1.6)	1.09±0.30	1.08(0.44-2.19)	0.0001
VITAMIN_K1	1.5±1.3	1.2(0.05-13.9)	1.8±1.4	1.4(0.10-7.3)	0.0416
VITAMIN_K2	0.6±1.0	0.29(0.082-11.0)	0.86±1.6	0.28(0.10-11.2)	0.9964
FOLATE	13.0±4.3	13.3(2.8-19.8)	13.8±3.9	14.2(2.2-19.9)	0.1956
VITAMIN D 25 OH	47.6±22.3	41.7(8.8-136)	47.5±25.3	40.1(15.4-133)	0.4508

ISOLEUCINE	72.7±19.9	70.7(34.0-121.4)	67.6±20.7	64.2(24.0-142.7)	0.0138
VALINE	261.9±55.2	262.7(146.5-404.0)	240.2±54.2	232.1(134.9)	0.0004
LEUCINE	172±34.8	168.8(113.7-248.2)	160.5±34.2	157.4(79.7-249.2)	0.0138
CITRULLINE	30.1±8.0	28.8(10.1-69.3)	30.2±9.6	29.7(10.3-79.4)	0.9509
ARGININE	123.7±33.8	119.4(49.6-299.2)	122.2±32.0	117.5(61.1-211.3)	0.7427

525

526

527

528

Table S7: Association of serum Lipoprotein (a) with micronutrients

	Greater than Reference Range (n=87)		Within range (n=270)		P (P<0.05)
ASPARAGINE	54.0±10.0	54.2(31.1-80.6)	53.5±12.3	52.7(28.1-108.0)	0.4975
GLUTAMINE	508.3±81.7	514.8(320.3-710.3)	505.3±80.5	506.9(202.2-741.0)	0.8216
SERINE	151.4±35.3	150.7(59.6-274.5)	148.0±37.8	142.5(70.5-241.1)	0.4179
CYSTEINE	17.9±8.6	7.02(2.8-45.1)	19.0±9.9	17.4(3.3-45.2)	0.6336
SELENIUM	139.2±21.9	138.3(96.6-210.8)	141.5±25.9	138.0(93.4-274.7)	0.7976
VITAMIN E	13.5±4.8	12.0(5.6-29.4)	13.1±4.2	12.4(5.3-28.2)	0.9456
CHOLINE	13.6±5.6	11.9(5.8-36.6)	14.6±6.1	13.2(1.0-39.6)	0.1288
CARNITINE	26.5±7.1	26.7(12.5-50.5)	26.8±7.7	27.0(4.1-44.3)	0.7048
SODIUM	141.5±3.5	142(129-150)	141.6±3.6	141(133-155)	0.7881
POTASSIUM	4.5±0.4	4.5(3.1-5.6)	4.5±0.39	4.4(3.7-6.2)	0.4339
CALCIUM	9.6±0.4	9.5(8.7-10.6)	9.6±0.4	9.6(8.5-10.8)	0.6777
MANGANESE	0.70±0.29	0.64(0.24-1.85)	0.74±0.32	0.69(0.3-2.6)	0.2023
ZINC	0.72±0.15	0.69(0.45-1.5)	0.7±0.14	0.69(0.3-1.3)	0.4235
COPPER	1.0±0.29	1.0(0.10-2.1)	1.0±0.29	1.0(0.5-2.5)	0.4000
CHROMIUM	0.24±0.12	0.23(0.02-0.68)	0.25±0.14	0.22(0.02-1.0)	0.8911
IRON	100.8±38.4	95(33.7-238.9)	105.7±38.4	102.0(25.3-201.3)	0.3270
MAGNESIUM	2.1±0.9	2.1(1.2-2.6)	2.1±0.2	2.1(1.0-2.6)	0.4372
VITAMIN A	76.8±25.4	71.2(34.3-151.2)	77.9±25.9	71.5(36.9-151.5)	0.7935
VITAMIN_B1	16.8±11.1	14.9(1.0-49.2)	22.7±13.7	20.7(0.8-78.8)	0.0008
VITAMIN_B2	26.7±20.4	20.3(5.7-135.7)	30.1±30.4	22.2(5.7-222.5)	0.4808
VITAMIN_B3	19.9±6.9	19.8(3.4-35.4)	19.7±10.2	18.9(2.7-110.1)	0.3823
VITAMIN_B6	17.2±22	11.9(1.6-236.7)	17.1±16.9	11.0(1.4-84.2)	0.6969
VITAMIN_B12	684.3±264.5	599.1(196-1381)	776.9±364.0	713.1(200-1923)	0.1376
VITAMIN_B5	103.8±81.9	68.6(13.8-360.9)	101.8±86.0	70.0(21.5-482.1)	0.8976
VITAMIN_C	0.43±0.24	0.38(0.08-1.2)	0.44±0.33	0.4(0.05-3.7)	0.8100
VITAMIN_D3	0.93±0.28	0.91(0.3-1.7)	0.94±0.26	0.94(0.32-1.6)	0.4137
VITAMIN_K1	1.5±1.1	1.2(0.05-5.8)	1.5±1.1	1.32(0.1-7.3)	0.5896
VITAMIN_K2	0.69±1.2	0.29(0.08-11.2)	0.61±0.78	0.28(0.10-4.8)	0.9556
FOLATE	13.5±4.2	13.7(2.9-19.9)	13.8±3.7	13.9(3.4-19.8)	0.7113
VITAMIN_D_25_OH	46.7±21.1	40.9(10.4-128)	47.7±23.4	41.2(12.4-133)	0.9703

ISOLEUCINE	68.8±19.7	66.3(33.5-129.7)	68.6±21.6	65.6(24.0-142.7)	0.8235
VALINE	243.9±56.8	236.2(149.8-442.7)	243.1±53.4	240.0(146.5-409.1)	0.9007
LEUCINE	164.4±33.4	164.5(90.3-235.1)	161.1±33.9	157.2(80.3-248.2)	0.3296
CITRULLINE	29.6±7.8	28.6(10.1-51.2)	30.1±9.2	29.6(10.3-69.3)	0.5236
ARGININE	124.7±39.2	118.3(56.1-299.2)	125.4±32.8	122.4(49.6-220.6)	0.5700

529

530

531

532

Table S8: Pearson correlation between serum cholesterol and micronutrients

	r	p
ASPARAGINE	-0.125	0.018
GLUTAMINE	-0.1127	0.0331
SERINE	-0.02601	0.6238
CYSTEINE	-0.01392	0.793
SELENIUM	0.02659	0.616
VITAMIN E	0.4086	<0.0001
CHOLINE	0.03297	0.5341
CARNITINE	0.09912	0.061
SODIUM	0.01349	0.7992
POTASSIUM	0.009154	0.863
CALCIUM	0.01211	0.8193
MANGANESE	-0.08336	0.1154
ZINC	-0.06639	0.2101
COPPER	-0.05695	0.2826
CHROMIUM	-0.07196	0.1743
IRON	0.1067	0.0437
MAGNESIUM	0.01114	0.8337
VITAMIN A	0.1471	0.0053
VITAMIN B1	-0.08282	0.1178
VITAMIN B2	-0.02302	0.6643
VITAMIN B3	-0.03355	0.5268
VITAMIN B6	-0.08714	0.0997
VITAMIN B12	0.03583	0.4992
VITAMIN B5	-0.0739	0.1629
VITAMIN C	-0.02218	0.6757
VITAMIN D3	0.5125	<0.0001
VITAMIN K1	0.2334	<0.0001
VITAMIN K2	0.07045	0.1835
FOLATE	-0.104	0.0493
VITAMIN D 25 OH	0.06205	0.2416
ISOLEUCINE	0.007779	0.8834
VALINE	0.09774	0.0647
LEUCINE	0.04598	0.3857
CITRULLINE	0.02438	0.6458
ARGININE	0.02322	0.6614

533
534
535
536
537
538
539
540

Table S9: Pearson correlation between serum LDL and micronutrients

	r	p
ASPARAGINE	-0.1217	0.0213
GLUTAMINE	-0.1074	0.0423
SERINE	-0.04485	0.3975
CYSTEINE	0.0241	0.6495
SELENIUM	-0.00553	0.917
VITAMIN E	0.3398	<0.0001
CHOLINE	-0.00725	0.8912
CARNITINE	0.1446	0.0061
SODIUM	0.02695	0.6113
POTASSIUM	0.02737	0.6058
CALCIUM	0.03125	0.5556
MANGANESE	-0.09346	0.0774
ZINC	-0.07538	0.1546
COPPER	-0.07262	0.1704
CHROMIUM	-0.08254	0.119
IRON	0.1199	0.0233
MAGNESIUM	0.03085	0.5607
VITAMIN A	0.09463	0.0737
VITAMIN_B1	-0.07151	0.177
VITAMIN_B2	-0.02853	0.5906
VITAMIN_B3	-0.00581	0.9127
VITAMIN_B6	-0.05988	0.2585
VITAMIN_B12	0.03573	0.5003
VITAMIN_B5	-0.04077	0.4419
VITAMIN_C	0.00019	0.9971
VITAMIN_D3	0.4774	<0.0001
VITAMIN_K1	0.06084	0.2509
VITAMIN_K2	0.0651	0.2192
FOLATE	-0.07946	0.1335
VITAMIN_D_25_OH	0.07967	0.1324
ISOLEUCINE	0.01793	0.7354
VALINE	0.1471	0.0053
LEUCINE	0.09979	0.0593
CITRULLINE	0.001029	0.9845

ARGININE	-0.03571	0.5007
----------	----------	--------

541

542

543

544

545

546

547

548

Table S10: Pearson correlation between serum HDL and micronutrients

	r	p
ASPARAGINE	0.007988	0.8803
GLUTAMINE	0.01765	0.7392
SERINE	0.06766	0.2016
CYSTEINE	-0.2289	<0.0001
SELENIUM	0.08967	0.0902
VITAMIN E	-0.099	0.0613
CHOLINE	0.01779	0.7373
CARNITINE	-0.1653	0.0017
SODIUM	-0.08206	0.1212
POTASSIUM	-0.0799	0.1313
CALCIUM	-0.08061	0.1279
MANGANESE	0.03657	0.4903
ZINC	0.03521	0.5067
COPPER	0.04878	0.3574
CHROMIUM	0.0416	0.4326
IRON	0.03515	0.5074
MAGNESIUM	-0.08087	0.1267
VITAMIN A	-0.03156	0.5517
VITAMIN_B1	0.02135	0.6873
VITAMIN_B2	0.02877	0.5874
VITAMIN_B3	0.01155	0.8276
VITAMIN_B6	-0.01147	0.8288
VITAMIN_B12	0.07736	0.1441
VITAMIN_B5	-0.01955	0.7123
VITAMIN_C	-0.01446	0.7852
VITAMIN_D3	0.02496	0.6378
VITAMIN_K1	-0.04944	0.3509
VITAMIN_K2	-0.0078	0.8831
FOLATE	-0.02846	0.5915
VITAMIN_D_25_OH	0.0655	0.2163
ISOLEUCINE	-0.2298	<0.0001
VALINE	-0.2317	<0.0001
LEUCINE	-0.2405	<0.0001

CITRULLINE	0.0138	0.7948
ARGININE	-0.03223	0.5432

549

550

551

552

553

554

555

556

Table S11: Pearson correlation between serum triglycerides and micronutrients

	r	p
ASPARAGINE	0.001283	0.9807
GLUTAMINE	-0.1002	0.0582
SERINE	-0.09025	0.0882
CYSTEINE	0.1626	0.002
SELENIUM	0.001458	0.9781
VITAMIN E	0.3832	<0.0001
CHOLINE	0.02606	0.6232
CARNITINE	0.1316	0.0127
SODIUM	0.03108	0.5578
POTASSIUM	0.0221	0.6769
CALCIUM	0.02466	0.6419
MANGANESE	-0.05805	0.2733
ZINC	-0.0603	0.2552
COPPER	-0.05772	0.2761
CHROMIUM	-0.06203	0.2417
IRON	-0.03861	0.4665
MAGNESIUM	0.02212	0.6766
VITAMIN A	0.2278	<0.0001
VITAMIN_B1	-0.04035	0.4466
VITAMIN_B2	-0.05275	0.3196
VITAMIN_B3	-0.02601	0.6238
VITAMIN_B6	-0.1057	0.0457
VITAMIN_B12	-0.1209	0.0221
VITAMIN_B5	-0.0363	0.4936
VITAMIN_C	-0.02185	0.6803
VITAMIN_D3	0.1342	0.011
VITAMIN_K1	0.5732	<0.0001
VITAMIN_K2	0.0625	0.2382
FOLATE	-0.01247	0.8141
VITAMIN_D_25_OH	-0.1507	0.0043
ISOLEUCINE	0.2838	<0.0001
VALINE	0.2234	<0.0001

LEUCINE	0.2139	<0.0001
CITRULLINE	-0.0145	0.7845
ARGININE	0.1363	0.0098

557

558

559

560

561

562

563

564

Table S12: Pearson correlation between serum Apo A and micronutrients

	r	p
ASPARAGINE	-0.04291	0.419
GLUTAMINE	0.003761	0.9435
SERINE	0.0321	0.5455
CYSTEINE	-0.05867	0.2689
SELENIUM	0.0395	0.4569
VITAMIN E	-0.05202	0.327
CHOLINE	-0.03578	0.5004
CARNITINE	-0.0026	0.9609
SODIUM	-0.06147	0.2467
POTASSIUM	-0.0526	0.3216
CALCIUM	-0.05579	0.2931
MANGANESE	0.05431	0.3062
ZINC	0.04085	0.4417
COPPER	0.05533	0.2971
CHROMIUM	0.05078	0.3387
IRON	0.01219	0.8185
MAGNESIUM	-0.05607	0.2907
VITAMIN A	0.03416	0.52
VITAMIN_B1	0.009489	0.8582
VITAMIN_B2	-0.05013	0.345
VITAMIN_B3	-0.04226	0.426
VITAMIN_B6	-0.05052	0.3412
VITAMIN_B12	0.03913	0.4611
VITAMIN_B5	-0.05052	0.3412
VITAMIN_C	-0.00363	0.9455
VITAMIN_D3	-0.05824	0.2724
VITAMIN_K1	-0.04603	0.3859
VITAMIN_K2	-0.04088	0.4413
FOLATE	0.07275	0.1702
VITAMIN_D_25_OH	-0.07913	0.1356
ISOLEUCINE	-0.04946	0.3514

VALINE	-0.02571	0.6283
LEUCINE	-0.06696	0.2069
CITRULLINE	0.002819	0.9577
ARGININE	0.003076	0.9538

565

566

567

568

569

570

571

572

Table S13: Pearson correlation between serum Apo B and micronutrients

	r	p
ASPARAGINE	-0.03841	0.4694
GLUTAMINE	-0.1388	0.0086
SERINE	-0.1269	0.0165
CYSTEINE	-0.02353	0.6577
SELENIUM	-0.01502	0.7774
VITAMIN E	0.06915	0.1924
CHOLINE	0.032	0.5467
CARNITINE	0.03729	0.4825
SODIUM	0.04303	0.4176
POTASSIUM	0.03938	0.4582
CALCIUM	0.03969	0.4547
MANGANESE	-0.00535	0.9198
ZINC	-0.00317	0.9525
COPPER	0.01326	0.8028
CHROMIUM	-0.00639	0.9042
IRON	0.08265	0.119
MAGNESIUM	0.04038	0.4469
VITAMIN A	0.1008	0.057
VITAMIN_B1	-0.02215	0.6767
VITAMIN_B2	-0.0354	0.505
VITAMIN_B3	-0.09294	0.0795
VITAMIN_B6	-0.02192	0.6798
VITAMIN_B12	0.003217	0.9517
VITAMIN_B5	-0.00175	0.9737
VITAMIN_C	0.01495	0.7784
VITAMIN_D3	0.04051	0.4455
VITAMIN_K1	-0.00555	0.9167
VITAMIN_K2	-0.07411	0.1623
FOLATE	0.1001	0.0589
VITAMIN D 25 OH	-0.01943	0.7144

ISOLEUCINE	0.09702	0.0671
VALINE	0.1259	0.0173
LEUCINE	0.03799	0.4743
CITRULLINE	-0.06951	0.1901
ARGININE	-0.1126	0.0334

573

574

575

576

577

578

579

580

Table S14: Pearson correlation between serum lipoprotein A and micronutrients

	r	p
ASPARAGINE	-0.01713	0.7914
GLUTAMINE	0.06125	0.3437
SERINE	-0.07783	0.2287
CYSTEINE	0.03708	0.5668
SELENIUM	0.04334	0.5031
VITAMIN E	0.008081	0.9007
CHOLINE	-0.06643	0.3044
CARNITINE	-0.05322	0.4108
SODIUM	-0.03759	0.5614
POTASSIUM	0.09305	0.1498
CALCIUM	0.000307	0.9962
MANGANESE	-0.01672	0.7963
ZINC	0.03884	0.5485
COPPER	0.04358	0.5007
CHROMIUM	-0.01997	0.7577
IRON	0.07018	0.2778
MAGNESIUM	0.01067	0.8691
VITAMIN A	0.0261	0.6869
VITAMIN_B1	-0.1818	0.0046
VITAMIN_B2	0.02598	0.6882
VITAMIN_B3	-0.01185	0.8548
VITAMIN_B6	-0.00366	0.955
VITAMIN_B12	-0.08534	0.1867
VITAMIN_B5	0.01869	0.7729
VITAMIN_C	0.08509	0.188
VITAMIN_D3	0.05586	0.388
VITAMIN_K1	0.004489	0.9447
VITAMIN_K2	-0.04037	0.5328
FOLATE	-0.03829	0.5541

VITAMIN D 25 OH	0.07921	0.2205
ISOLEUCINE	-0.02975	0.6458
VALINE	-0.0444	0.4927
LEUCINE	0.02835	0.6614
CITRULLINE	-0.02425	0.708
ARGININE	-0.0665	0.3039

581

582




583

584

585

586

587



Figure

