1 Characterisation of placental, fetal brain and maternal cardiac structure and function in

2 pre-eclampsia using MRI

3 Short title: Maternal-fetal changes in pre-eclampsia

4	Megan Hall MRCOG, BSc (Hons) ^{1,2} , Antonio de Marvao MRCP, PhD ^{1,3,4} , Ronny Schweitzer
5	MD ^{3,4} , Daniel Cromb MRCPCH, BSc (Hons) ^{2,} , Kathleen Colford, BSc (Hons), PGDip ² , Priya
6	Jandu, MSc ⁵ , Declan P O'Regan, FRCP, FRCR, PhD ⁴ , Alison Ho, MRCOG, PhD ^{1,2} , Anthony
7	Price, PhD ^{2,6} , Lucy C. Chappell, FRCOG, PhD, FMedSci ¹ , Mary A. Rutherford, MRCPCH, FRCR,
8	MD ² , Lisa Story MRCOG, PhD ^{1,2} , Pablo Lamata, PhD ⁶ , Jana Hutter, PhD ^{2,6}
9	1. Department of Women and Children's Health, King's College London, UK
10	2. Centre for the Developing Brain, King's College London, UK
11	3. School of Cardiovascular Medicine, King's College London, UK
12	4. MRC London Institute of Medical Sciences, Imperial College London, UK
13	5. GKT School of Medical Education, King's College London, UK
14	6. Centre for Medical Engineering, King's College London, UK
15	
16	Corresponding author: Megan Hall
17	First Floor, South Wing

18 St Thomas' Hospital

19 London

20 SE1 7UH

21 <u>megan.hall@kcl.ac.uk</u>

- 22 +447906542307
- 23

24 Total word count: 3754

25 Abstract

26 Background

- 27 Pre-eclampsia is a multiorgan disease of pregnancy that has short- and long-term
- implications for the woman and fetus, whose immediate impact is poorly understood. We
- 29 present a novel multi-system approach to MRI investigation of pre-eclampsia, with
- 30 acquisition of maternal cardiac, placental, and fetal brain anatomical and functional

31 imaging.

32 Methods

33 A prospective study was carried out recruiting pregnant women with pre-eclampsia, chronic

34 hypertension, or no medical complications, and a non-pregnant female cohort. All women

35 underwent a cardiac MRI, and pregnant women underwent a fetal-placental MRI. Cardiac

- 36 analysis for structural, morphological and flow data was undertaken; placenta and fetal
- brain volumetric and T2* data were obtained. All results were corrected for gestational age.

38 Results

Seventy-eight MRIs were obtained during pregnancy. Pregnancies affected by pre-eclampsia demonstrated lower placental and fetal brain T2*. Within the pre-eclampsia group, three placental T2* results were within the normal range, these were the only cases with normal placental histopathology. Similarly, three fetal brain T2* results were within the normal range; these cases had no evidence of cerebral redistribution on fetal Dopplers. Cardiac MRI analysis demonstrated higher left ventricular mass in pre-eclampsia with 3D modelling revealing additional specific characteristics of eccentricity and outflow track remodelling.

46 Conclusions

- 47 We present the first holistic assessment of the immediate implications of pre-eclampsia on
- 48 the placenta, maternal heart, and fetal brain. As well as having potential clinical implications
- 49 for the risk-stratification and management of women with pre-eclampsia, this gives an
- 50 insight into disease mechanism.
- 51

52 What is new?

53	•	We propose a comprehensive MRI protocol for maternal cardiac and fetal-placental
54		imaging that is safe in pregnancy and acceptable to women, and report on the
55		largest set of functional placental MRI data in pre-eclamptic pregnancies
56	•	The findings suggest altered oxygenation and microstructure in the placenta, which
57		is associated with similar changes in the fetal brain
58	•	The study identifies a specific pattern of remodelling in the maternal left ventricle
59		that is associated with pre-eclampsia
60		
61	What	are the clinical implications?
62	•	We demonstrate that MRI provides maternal and fetal insights into pre-eclamptic
62 63	•	We demonstrate that MRI provides maternal and fetal insights into pre-eclamptic pregnancies that cannot be obtained by other means during pregnancy
62 63 64	•	We demonstrate that MRI provides maternal and fetal insights into pre-eclamptic pregnancies that cannot be obtained by other means during pregnancy Functional placental MRI may be able to differentiate between preeclampsia disease
62 63 64 65	•	We demonstrate that MRI provides maternal and fetal insights into pre-eclamptic pregnancies that cannot be obtained by other means during pregnancy Functional placental MRI may be able to differentiate between preeclampsia disease severities, while fetal brain imaging may help refine neurodevelopmental prognostic
62 63 64 65 66	•	We demonstrate that MRI provides maternal and fetal insights into pre-eclamptic pregnancies that cannot be obtained by other means during pregnancy Functional placental MRI may be able to differentiate between preeclampsia disease severities, while fetal brain imaging may help refine neurodevelopmental prognostic assessment of children exposed to pre-eclampsia in utero
62 63 64 65 66 67	•	We demonstrate that MRI provides maternal and fetal insights into pre-eclamptic pregnancies that cannot be obtained by other means during pregnancy Functional placental MRI may be able to differentiate between preeclampsia disease severities, while fetal brain imaging may help refine neurodevelopmental prognostic assessment of children exposed to pre-eclampsia in utero A specific 3D pattern of left ventricular remodelling and thickening was found to be
62 63 64 65 66 67 68	•	We demonstrate that MRI provides maternal and fetal insights into pre-eclamptic pregnancies that cannot be obtained by other means during pregnancy Functional placental MRI may be able to differentiate between preeclampsia disease severities, while fetal brain imaging may help refine neurodevelopmental prognostic assessment of children exposed to pre-eclampsia in utero A specific 3D pattern of left ventricular remodelling and thickening was found to be discriminant of the presence of pre-eclampsia and may provide a basis for improved
62 63 64 65 66 67 68 69	•	We demonstrate that MRI provides maternal and fetal insights into pre-eclamptic pregnancies that cannot be obtained by other means during pregnancy Functional placental MRI may be able to differentiate between preeclampsia disease severities, while fetal brain imaging may help refine neurodevelopmental prognostic assessment of children exposed to pre-eclampsia in utero A specific 3D pattern of left ventricular remodelling and thickening was found to be discriminant of the presence of pre-eclampsia and may provide a basis for improved clinical risk stratification

71 Non-standard abbreviations and acronyms

- 72 ADC = apparent diffusion coefficient
- 73 bpm = beats per minute
- 74 CO = cardiac output
- 75 EDV = end diastolic volume
- 76 ESV = end systolic volume
- 77 FGR = fetal growth restriction
- 78 FOV = field of view
- 79 GCS = global circumferential strain
- 80 GLS = global longitudinal strain
- 81 GRS = global radial strain
- 82 IQR = interquartile range
- 83 LV = left ventricular
- 84 LVM = left ventricular mass
- 85 TE = echo time
- 86 TR = repetition time

87 Introduction

88	Pre-eclampsia occurs when placental malperfusion, and a resultant release of soluble
89	factors into the circulation, causes maternal vascular endothelial injury with subsequent
90	hypertension and multi-organ damage. ¹ The Incidence is between 2 and 8%, with
91	geographical variation. ² Globally, 14% of maternal deaths are associated with hypertensive
92	diseases of pregnancy, ³ which can occur secondary to multiple complications including renal
93	and hepatic failure, stroke, eclampsia, pulmonary oedema, and disseminated intravascular
94	coagulopathy. Cardiac manifestations include reduced cardiac output, ⁴ increased vascular
95	resistance, left ventricular hypertrophy, ⁵ and diastolic dysfunction. ⁶ After the postnatal
96	period, women who have had pre-eclampsia remain at an increased risk of vascular disease
97	and ischaemic heart disease. ^{7,8} While it is well recognised that maternal biochemical
98	abnormalities can be detected prior to clinical onset of pre-eclampsia, there is also some
99	evidence that cardiovascular maladaptation precedes the diagnosis of pre-eclampsia:
100	echocardiography has demonstrated abnormalities in left ventricular mass in the late third
101	trimester prior to the development of term pre-eclampsia. ^{9,10}
102	Cardiac magnetic resonance imaging (MRI) combines excellent spatial and temporal
103	resolution, and is the gold-standard non-invasive method of assessing cardiac morphology
104	and function. In contrast with echocardiography, cardiac MRI is not limited by geometric
105	assumptions in assessing ventricular volumes and image acquisition is less operator-
106	dependent. This allows highly reproducible and accurate assessment of cardiac structure
107	and function, ¹¹ making it the ideal modality for assessing differences between small
108	groups. ¹² We have provided pilot data demonstrating feasibility of cardiac MRI in pregnancy,
109	but large prospective studies are still required. ¹³

110	For the fetus, pre-eclampsia is associated with fetal growth restriction (FGR) and
111	complications such as placental abruption; excluding congenital anomalies, 5% of stillbirths
112	occur in women with pre-eclampsia (Cantwell et al, 2011). In infants born to women with
113	pre-eclampsia, increased rates of hypertension and cardiac dysfunction, stroke, cognitive
114	dysfunction and psychiatric morbidity have been demonstrated. ¹⁴
115	Postnatally obtained placental histopathological correlates with pre-eclampsia include both
116	villous and vascular lesions, ¹⁵ but sufficient <i>in vivo</i> and <i>in vitro</i> modelling of placental
117	development in disease is lacking. MRI offers an opportunity for in vivo study of the
118	placenta in pre-eclampsia, as well as evaluation of maternal and fetal structures. Recent
119	studies applying functional MRI techniques to the placenta have proposed reduced
120	oxygenation in the pre-eclamptic placenta via demonstration of increased heterogeneity
121	and decreased mean T2*. ^{16,17} There is evidence of decreased placental diffusivity in small
122	for gestational age fetuses. ^{18,19} Furthermore, advances in acquisition and reconstruction
123	techniques have allowed more detailed investigation of the fetal brain.
124	Here we describe a multi-system approach to MRI investigation of pre-eclampsia, with
125	acquisition of maternal cardiac, placental, and fetal brain anatomical and functional

126 imaging.

127 Methods

128	A prospective observational study was performed between 2019 and 2022 at Guy's and St
129	Thomas' NHS Foundation Trust (London Dulwich Ethics Committee 08/LO/1958). Additional
130	high-risk placental and fetal datasets were obtained from a previous study (London Dulwich
131	Ethics Committee 16/LO/1573). Additional non-pregnant control datasets were obtained
132	from a previous study (London - West London & GTAC Research Ethics Committee
133	17/LO/0034).
134	Women were recruited into four categories: non-pregnant controls, pregnant controls,
135	women with pre-eclampsia, and pregnant women with chronic hypertension; a fifth
136	category 'complicated pregnancies' was created retrospectively for women who had been
137	recruited as controls but had developed other pregnancy complications and this group was
138	excluded from further analysis. Women with diabetes and a hypertensive disorder were
139	analysed within the appropriate hypertensive cohort. For all groups inclusion criteria
140	included a body mass index (BMI) <40kg/m ² (with the MRI bore size the limiting factor),
141	aged 16-50 years, and no contraindication to MRI such as non-compatible metal implants or
142	claustrophobia. All women could undergo up to three scans.
143	Inclusion criteria for the control group was women with no previous cardiovascular or
144	inflammatory cardiac history. Pregnant controls were excluded if they had a multi-fetal
145	pregnancy or fetal congenital anomalies. Pregnant controls were moved into the
146	complicated pregnancy cohort if they developed any other complications of pregnancy, had
147	an infant born preterm, below the 3 rd centile, above the 97 th centile, or that required
148	neonatal intensive care for any reason. Additionally, age and ethnicity matched non-
149	pregnant controls, having undergone equivalent imaging, were recruited from the UK Digital

150	Heart Project (UKDHP) at Imperial College London. Enrolment into this study excluded
151	participants who had known cardiovascular or metabolic disease, as previously described. ²⁰
152	Case inclusion criteria was a diagnosis of pre-eclampsia (including superimposed on chronic
153	hypertension) defined by the International Society for the Study of Hypertension in
154	Pregnancy as hypertension with one of the following new-onset conditions at 20 weeks'
155	gestation or beyond: proteinuria, acute kidney injury, hepatic dysfunction, neurological
156	features, haemolysis or thrombocytopaenia, or fetal growth restriction (Brown et al, 2018).
157	Women with chronic hypertension were recruited as a separate cohort.
158	Demographic and clinical information, including on previous obstetric history, was collected
159	on all participants. Neonatal outcomes including gestation at birth, mode of delivery,
160	condition at birth, need for neonatal support, and birthweight were collected.
161	MRI
162	After informed consent was obtained, participants had an MRI on a clinical 1.5T Philips
163	Ingenia scanner using a combined 24-channel posterior and torso (dStream) coil. Maternal
164	comfort in the supine position was achieved by careful positioning of the head and legs on
165	elevated cushions, back padding, and additional cushions as requested. The MRI was
166	performed in two sessions (maternal cardiac, and fetal-placental) each of which lasted
167	around 30 minutes. A break was given in the middle of the session. In order to reduce any
168	bias that could arise from maternal anxiety at the start of the scan, women with an odd case
169	study ID underwent fetal-placental imaging first, and those with an even ID underwent
170	cardiac imaging first.

- 171 An obstetrician or midwife was present for all scans. Maternal temperature was recorded
- 172 before and after the scan. Continuous oxygen saturations were measured, and blood
- 173 pressure was taken every 10 minutes.
- 174 The protocol is illustrated in Figure 1.
- 175 Fetal-placental MRI
- 176 Fetal brain and whole uterus structural imaging was performed in five planes (transverse
- 177 centred on fetal brain; two opposing sagittal oblique planes; maternal coronal plane and
- 178 maternal sagittal plane) in order to obtain whole uterus and dedicated placental images.
- 179 MRI was undertaken using 2D-single-shot Turbo Spin Echo (resolution 1.5x1.5x2.5mm, field
- of view (FOV) 320x320x110mm, echo time (TE) 180ms). Fetal body imaging was performed
- 181 for clinical reporting only.
- 182 Placental T2* mapping was obtained using a multi-echo gradient-echo sequence (2.5mm
- isotropic, FOV=300x300x110mm, TE=11, 58, 117, 176ms). Placental diffusion-weighted
- imaging used a single-shot echo-planar-imaging sequence with parameters adapted to meet
- the expected diffusivity (1 b=0, 6 b=375, 6 b=750; transverse relaxation (TR)=6.6ms,
- 186 TE=78ms, matrix = 512x512x56, resolution 2x2x4mm). Both sequences were repeated twice
- 187 to improve data robustness.

188 Cardiac MRI

Anatomical and functional cardiac MRI sequences were acquired using commonly employed sequences adapted to reduce acoustic noise and heating, and so improve suitability in pregnancy. Views taken included: 4 chamber, 2 chamber, 3 chamber, short axis and outflow (acquired using dynamic balanced stead state free precession sequences with 30 cardiac phases, 11 second breath-hold, FA=60, TE/TR~2/4ms, resolution=1.7x1.7x8mm); and a short

medRxiv preprint doi: https://doi.org/10.1101/2023.04.24.23289069; this version posted April 25, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

194	axis stack (14 slices, sense=2, resolution 2x2x10mm). Phase contrast flow sequences of the
195	ascending and descending aorta, and the superior vena cava were acquired with individually
196	adapted encoding velocities (30 phases, 14 second breath-holds, sense=2,
197	resolution=2.5x2.5x8mm). A pragmatic decision was made to obtain inferior vena cava
198	images during the 'fetal-placental' portion of the MRI as the coil was appropriately placed.
199	The scanning parameters are summarised in Table 1. Images acquired are demonstrated in
200	Figure 2.
201	Non-pregnant controls from the healthy control group underwent an equivalent cardiac MRI
202	protocol on a 1.5-T Philips Achieva system (Best, the Netherlands) as previously described. ²⁰
203	Analysis
204	The placenta was visually inspected on a T2 image. Manual outlining was performed on a
205	functional scan mapped to subsequent scans. Monoexponential fitting was performed to
206	obtain T2* maps. Mean placenta T2*, histogram skewness and kurtosis, and mean
207	diffusivity (as the apparent diffusion coefficient – ADC) were calculated.
208	Brain volume (supratentorial tissues excluding brainstem and cerebrospinal fluid) was
209	manually segmented on functional maps. Mean T2* values were obtained by averaging the
210	T2* maps from manually outlined brains.
211	The cardiac MRIs were analysed using cvi42 post-processing software (Version 5.1.4, Circle
212	Cardiovascular Imaging Inc., Calgary, Canada), using standard clinical methodology, by
213	operators masked to the clinical diagnosis. ²¹
214	The 3D left ventricular morphology was studied by the construction of a statistical shape
215	model from the segmentations of the short axis stack at the end diastolic frame as reported

216	in previous studies. ²²⁻²⁴ Briefly, 3D meshes were built using a computational anatomy tool
217	kit, ²⁵ anatomical modes of variation were found by Principal Component Analysis, and the
218	impact of pre-eclampsia was assessed by an optimised linear discriminant analysis of the
219	first 12 modes.
220	Flow data was processed using CVI42 Flow version 5.10.3. The magnitude image with the
221	sharpest contrast was used to determine vessel contours. Contours were then propagated
222	to phase contrast images (with manual correction as required) in all temporal phases.
223	All parameters were first assessed against gestational age, and then as cases against
224	controls. Matching was undertaken to nearest gestational age, with further analysis using
225	the Mann-Whitney U Test.
226	Statistical Analysis
226 227	Statistical Analysis Statistical analysis was performed with R version 3.6.0 (R Foundation for Statistical
226 227 228	Statistical Analysis Statistical analysis was performed with R version 3.6.0 (R Foundation for Statistical Computing) and RStudio Server version 1.043, unless otherwise stated. Variables are
226 227 228 229	Statistical Analysis Statistical analysis was performed with R version 3.6.0 (R Foundation for Statistical Computing) and RStudio Server version 1.043, unless otherwise stated. Variables are expressed as percentages if categorical, mean ± SD if continuous and normal, and median
226 227 228 229 230	Statistical Analysis Statistical analysis was performed with R version 3.6.0 (R Foundation for Statistical Computing) and RStudio Server version 1.043, unless otherwise stated. Variables are expressed as percentages if categorical, mean ± SD if continuous and normal, and median (interquartile range) if continuous and non-normal. Baseline anthropometric data were
226 227 228 229 230 231	Statistical Analysis Statistical analysis was performed with R version 3.6.0 (R Foundation for Statistical Computing) and RStudio Server version 1.043, unless otherwise stated. Variables are expressed as percentages if categorical, mean ± SD if continuous and normal, and median (interquartile range) if continuous and non-normal. Baseline anthropometric data were compared by using Kruskal-Wallis tests and, if differences were identified, a Wilcoxon test
226 227 228 229 230 231 232	Statistical Analysis Statistical analysis was performed with R version 3.6.0 (R Foundation for Statistical Computing) and RStudio Server version 1.043, unless otherwise stated. Variables are expressed as percentages if categorical, mean ± SD if continuous and normal, and median (interquartile range) if continuous and non-normal. Baseline anthropometric data were compared by using Kruskal-Wallis tests and, if differences were identified, a Wilcoxon test was used for pairwise comparisons with Benjamini-Hochberg adjustment for multiple
226 227 228 229 230 231 232 233	Statistical Analysis Statistical analysis was performed with R version 3.6.0 (R Foundation for Statistical Computing) and RStudio Server version 1.043, unless otherwise stated. Variables are expressed as percentages if categorical, mean ± SD if continuous and normal, and median (interquartile range) if continuous and non-normal. Baseline anthropometric data were compared by using Kruskal-Wallis tests and, if differences were identified, a Wilcoxon test was used for pairwise comparisons with Benjamini-Hochberg adjustment for multiple testing. Imaging parameters in 2 or more groups were compared by using analysis of
226 227 228 229 230 231 232 233 233 234	Statistical Analysis Statistical analysis was performed with R version 3.6.0 (R Foundation for Statistical Computing) and RStudio Server version 1.043, unless otherwise stated. Variables are expressed as percentages if categorical, mean ± SD if continuous and normal, and median (interquartile range) if continuous and non-normal. Baseline anthropometric data were compared by using Kruskal-Wallis tests and, if differences were identified, a Wilcoxon test was used for pairwise comparisons with Benjamini-Hochberg adjustment for multiple testing. Imaging parameters in 2 or more groups were compared by using analysis of covariance, adjusted for relevant clinical covariates. When differences were significant, a
2226 2227 2228 229 230 231 232 233 233 234 235	Statistical Analysis Statistical analysis was performed with R version 3.6.0 (R Foundation for Statistical Computing) and RStudio Server version 1.043, unless otherwise stated. Variables are expressed as percentages if categorical, mean ± SD if continuous and normal, and median (interquartile range) if continuous and non-normal. Baseline anthropometric data were compared by using Kruskal-Wallis tests and, if differences were identified, a Wilcoxon test was used for pairwise comparisons with Benjamini-Hochberg adjustment for multiple testing. Imaging parameters in 2 or more groups were compared by using analysis of covariance, adjusted for relevant clinical covariates. When differences were significant, a Tukey post hoc test was applied for pairwise multiple comparisons.

236

237 Results

238 Study participants

239	Seventy-eight scans were obtained from 65 pregnant women (with nine women having two scans,
240	and two having three scans). An additional 38 scans were obtained from healthy non-pregnant
241	controls. Cardiac assessment was not completed in two pregnant participants: one owing to
242	claustrophobia, and one to worsening hypertension. For a further 16 women, no cardiac
243	data were obtained. Nine women developed pregnancy complications after being scanned
244	as a control and so were moved to the complicated pregnancy cohort and excluded.
245	Thirteen women with pre-eclampsia were recruited, one of whom had three MRIs. Of the
246	13, six had pre-eclampsia superimposed on chronic hypertension. Table 2 summarises the
247	demographics of all cohorts. Supplementary Table 1 details relevant obstetric history and
248	outcomes of the pre-eclamptic cohort.
249	Placenta
250	In the cross-sectional control group, there was a decrease in mean T2* values across
251	gestation (p<0.01). This was also seen as a trend (not reaching significance) in the chronic
252	hypertensive group. The pre-eclamptic cohort had significantly lower T2* values throughout
253	gestation, and a gradual decline was not seen. Both skewness and kurtosis increased
254	throughout gestation in the control group, with the chronic hypertensive group overlying
255	this, although not reaching significance. The pre-eclamptic group had higher skewness and
256	kurtosis than both other groups, and placental volume was reduced (p=0.02) (Figure 3).
257	Three women with pre-eclampsia had placental T2* values similar to controls; of note these
258	were the only three women without any vascular abnormalities on placental histopathology
259	(Table 3, Supplementary Table S1).

260 Visual inspection of control versus pre-eclamptic placentas revealed a more variable lobule

- size and less consistent signal intensity in women with pre-eclampsia. There was also an
- 262 increase in low signal areas as compared to controls.

263 Fetal brain

- 264 Fetal brain T2* values were demonstrated to decline with advancing gestation among
- 265 controls. This relationship was preserved in the chronic hypertensive group. Among women
- with pre-eclampsia, the fetal brain T2* values were significantly reduced through all
- 267 gestations (p<0.01) and did not have a linear decline with gestation (Figure 4). There were
- three cases in which brain T2* was in line with controls; in all these cases fetal Doppler
- studies were normal (Table 3, Supplementary Table S1). One case demonstrated an
- abnormal T2* values but with preserved fetal Doppler studies (Table 3, Supplementary
- Table S1). Brain volume was reduced in the presence of pre-eclampsia (p=0.03) (Figure 4).
- 272 Z scores were generated for mean placental T2* values and mean fetal brain T2* values in
- 273 order to allow for direct comparison: there was a direct correlation between normal and
- low scores in both domains (Figure 5).

275 Major differences in the placental and fetal findings of the control and pre-eclampsia groups

are summarised in Supplementary Figure 1.

277 Cardiac MR

Systolic (p<0.001) and diastolic (p=0.03) blood pressure were lower in the pregnant controls
(107 / 66 mmHg) than in pre-eclamptic participants (121 / 77 mmHg). There were no other
baseline anthropometric differences between groups.

281	After adjusting for body surface area (BSA), non-pregnant controls had lower heart rate (68
282	bpm vs 74.8 bpm; p = 0.02), cardiac output (5.5 L / min vs 6.2 L/ min; p = 0.02), LV mass
283	(77.4 g vs 87 g; p = 0.01) and concentricity (LV mass / LV end-diastolic volume; 0.57 g / ml vs
284	0.64 g / L; $p = 0.01$) than pregnant participants. The LV mass of the pre-eclamptic group
285	(98.05 \pm 24.5 g) was higher than both the pregnant controls (83.3 \pm 12.08 g; p = 0.04) and
286	the non-pregnant group (77.4 \pm 13.8 g; p=0.04). The study of the 3D anatomy revealed a
287	specific thickening pattern (existence of regional increase in wall thickness wall locations of
288	mid antero-lateral and postero-septal) together with changes in eccentricity (the ventricular
289	cross-section displayed a dilation in the axis oriented in the direction of the outflow tract)
290	and the onset of a bulge below the outflow track, see Figure 3. A detailed inspection of the
291	modes of anatomical variation further reveals that the thickening pattern associated to pre-
292	eclampsia was linked to a localised basal concentric remodelling and not to the
293	complementary basal eccentric remodelling (see modes 11 and 13 in Supplementary Figure
294	S2).

295 Discussion

296 Summary of main findings

We have demonstrated a comprehensive anatomical and functional MRI protocol for
cardiac and fetal-placental imaging that is safe in pregnancy and acceptable to women, with
97% of scans completed. As far as we are aware, this is the largest set of functional placental
MRI data in pre-eclamptic pregnancies, and confirms previous findings suggestive of altered
oxygenation and microstructure. We have demonstrated altered T2* in the fetal brain, in
proportion to that seen in the placenta. Finally, we have characterised the remodelling
pattern of the maternal left ventricle associated with the presence of pre-eclampsia.

304	Conventional cardiac MRI analysis demonstrated higher LVM in pre-eclampsia than in
305	uncomplicated pregnancies, while 3D statistical modelling revealed specific characteristics
306	of eccentricity and outflow track remodelling beyond the thickening of the walls.
307	Strengths and limitations
308	This is the largest study to date investigating maternal, fetal and placental implications of
309	pre-eclampsia in vivo. We utilise an optimised protocol ²⁶ and includes novel findings related
310	to the T2* values. To our knowledge this is the first published study making a holistic
311	assessment of the fetal, placental and maternal implications of pre-eclampsia. The MRI
312	sequence described here is safe in pregnancy and acceptable to women, and gives insight
313	into pre-eclamptic pregnancies that cannot be obtained by other means during pregnancy.
314	The use of consistent maternal positioning and scanning protocols means that the data are
315	robust and comparable between subjects.
316	Functional placental data build on that previously presented to demonstrate abnormalities
317	in microstructure and oxygenation in the pre-eclamptic placenta. ^{17,18} The inclusion of a non-
318	pregnant cardiac control group delineates changes expected in pregnancy from those that
319	are pathological. Motion correction and reconstruction of fetal images has allowed for the
320	first analysis of T2* brain maps in fetuses affected by pre-eclampsia. The inclusion of a
321	chronic hypertensive group demonstrates where phenotypes are driven by pre-eclampsia,
322	rather than the gestational effects of hypertension alone.
323	There are several limitations to this study: firstly, recruitment to the pre-eclampsia group
324	was challenging, largely due to the often short intervals from diagnosis to delivery and so
325	the study is relatively small, perhaps obscuring other significant differences between cases
326	and controls. Secondly, the lack of long-term follow up limits understanding of how much

327	the observed maternal cardiac and fetal brain changes directly impact chronic health
328	outcomes. Thirdly, the small sample size makes it difficult to comment on confounders
329	within the pre-eclampsia group, in particular the presence of diabetes, and early versus late
330	onset disease.
331	Comparison to other work
332	The placental phenotype in control pregnancies is in line with previous work: the reduction
333	in T2* across gestation in normal pregnancies has been demonstrated using both our
334	imaging protocols ²⁶ and those that of others. ^{27,28} As previously demonstrated, there is
335	overlap between chronic hypertensive pregnancies and normal pregnancies in terms of
336	mean T2*, skewedness and kurtosis. ¹⁶ Pregnancies affected by pre-eclampsia are typically
337	outliers in all domains. This is to be expected given the higher rate of hypoxic vascular
338	anomalies seen in the pre-eclamptic placentae compared to that affected by chronic
339	hypertension alone. ²⁹
340	In all cases where the placental T2 st in pre-eclamptic pregnancies lay within the normal
341	range, there was normal placental histopathology and delivery at later gestations,
342	suggesting a less severe clinical phenotype. Pre-existing work on placental perfusion MRI in
343	women with early and late pre-eclampsia, where women with later onset disease has
344	demonstrated preserved placental perfusion. ³⁰ The decreased placental perfusion seen in

the early pre-eclampsia group, as well as histopathological evidence for greater maternal

malperfusion in early onset pre-eclampsia as compared to late,³¹ provide basis for the

variation seen in our cohort. Our findings go further to support the ability of functional

348 placental MRI to differentiate between disease severities.

medRxiv preprint doi: https://doi.org/10.1101/2023.04.24.23289069; this version posted April 25, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

349	Novel to this study is the association between placental T2* and fetal brain T2*. The
350	negative correlation between decreasing brain T2* and increasing gestation in the normal
351	pregnancy is already established, and thought to be secondary to a number of structural
352	changes including decreasing water density and increasing myelination. ³² Preservation of
353	brain volume with a decrease in T2 st in the pre-eclampsia group may reflect a reduction in
354	oxygenation, with or without reduction in angiogenesis. This seems particularly likely given
355	that this does not seem to occur in the absence of reduced placental T2*, and that in all but
356	one case there is corresponding Doppler evidence of cerebral redistribution. Long-term
357	neurodevelopmental and behavioural outcomes of children born to mothers with pre-
358	eclampsia are recognised, and a small study has shown differences in neuroanatomy
359	persisting at 7-10yrs; ³³ although this is difficult to disentangle from the effects of
360	prematurity, our findings may suggest an antenatal antecedent.
361	Looking into the mother's health, novel to this study is the 3D thickening pattern of the left
362	ventricle that is found discriminant of the presence of pre-eclampsia. The extra workload in
363	pre-eclampsia is indeed increasing the cardiac mass and mass to volume ratio, as previously
364	reported in much larger echocardiography studies. ⁵ This demonstrates the higher power of
365	cardiac MRI to identify differences between small groups, and its role in cardiovascular
366	research in pregnancy. Our preliminary data further reveal an intriguing pattern of
367	thickening, eccentricity and bulging as illustrated in Figure 6. The fact that the other groups
368	of non-pregnant controls and chronic hypertension did not show differences with control
369	pregnancy along this axis suggests that this is an acute maladaptive remodelling process
370	specific to pre-eclampsia. In relationship with this finding, the chronic remodelling of the left
371	ventricle 5 to 10 years after a hypertensive pregnancy manifest as a global concentric (and
372	not eccentric) thickening of the left ventricle. ³⁴ Intriguingly in the response of pre-eclampsia

373	during pregnancy it is only the base of the heart where this concentric (and not eccentric)
374	thickening pattern is seen (see Supplementary Figure S1). This collection of observations
375	leads us to hypothesize that it is the base of the heart the region that is first recruited and
376	most intensively working in response to pre-eclampsia, and that the rapid onset of
377	hypertension causes a level of uneven thickening pattern and septal bulging.
378	Implications
379	While clinical fetal MRI is largely confined to assessment of structural anomalies, there is
380	evidence here of a potential role for other high risk obstetric conditions. We have
381	demonstrated feasibility of a comprehensive fetal, placental and maternal cardiac MRI in
382	terms of safety, data acquisition and acceptability to women.
383	Future work into determining risk of developing, likely clinical severity and implications of
384	pre-eclampsia should combine MRI, obstetric ultrasound and biomarkers, as it is likely that a
385	combination of investigations could yield greatest information.
386	While we have demonstrated a correlation of findings associated with pre-eclampsia,
387	further work should be done in defining this phenotype. As well as increasing the cohort
388	size, attention should be paid to early and late onset disease, as these are likely to be
389	mechanistically different and have different clinical end point. ³⁵ While fetal growth
390	restriction has previously been associated with reduced placental T2*, ^{19,36} e we have been
391	unable to delineate the implications of pre-eclampsia on this finding. Creation of a cohort
392	where the interplay between these three findings could be investigated would be of value.
393	MRI could have implications for earlier diagnosis of pre-eclampsia; high-risk women (for
394	example those with early onset growth restriction) who are not yet diagnosed with pre-
395	eclampsia should be included in future work, as this could have significant implications for

396	their management. Moreover, the 3D specific remodelling pattern linked to pre-eclampsia
397	could have implications for prognosis and risk models for future cardiovascular events. Any
398	future work on cardiac imaging should include longitudinal and postnatal data in order to
399	better delineate the mechanisms of continued cardiac risk, and for stratification of risk in
400	individuals.
401	In terms of perinatal outcomes, we have demonstrated a relationship between pre-
402	eclampsia and reduced fetal brain T2 * , and also between reduced fetal brain T2 * and fetal
403	cerebral Doppler redistribution. While there is some evidence of abnormal neurocognitive
404	outcomes in children who have been affected by redistribution (Beukers et al, 2017), further
405	neurocognitive follow up of children with paired MRI and ultrasound data could improve
406	understanding of the longer-term clinical implications of pre-eclampsia on the child.
407	In terms of techniques used, while the relationship between T2* and deoxyhaemoglobin is
408	well established, it is subject to influence from other factors. Therefore, additional
409	functional imaging, such as diffusion techniques, linked with recent advanced analysis
410	techniques ³⁷ may lead to greater insight of the mechanisms of pre-eclampsia.
411	Conclusions
412	MRI of pregnancies affected by pre-eclampsia reveals a reduction in mean placental and
413	fetal brain T2*, with preservation of the volume of both organs. There is also an increase in
414	maternal cardiac work and specific thickening patterns of the left ventricle. This all
415	demonstrates that MRI is clinically valuable in risk stratification, investigation, and
416	management planning of women at high risk of pre-eclampsia, or who have been diagnosed
417	with the condition.

418 Acknowledgments

- 419 We are grateful to the women who participated in this study.
- 420 Disclosures
- 421 The authors declare no conflicts of interest.

422 Funding

- 423 This work was supported by core funding from the Wellcome/EPSRC Centre for Medical
- 424 Engineering [WT203148/Z/16/Z], by the NIH Human Placenta Project grant 1U01HD087202-
- 425 01 (Placenta Imaging Project (PIP)), by the Wellcome Trust, Sir Henry Wellcome Fellowship
- to JH [201374/Z/16/Z] and Senior Research Fellowship to PL [209450/Z/17/Z]; National
- 427 Institute for Health Research (NIHR) Imperial College Biomedical Research Centre; Medical
- 428 Research Council (MC_UP_1605/13); British Heart Foundation (RG/19/6/34387,
- 429 RE/18/4/34215); Academy of Medical Sciences (SGL015/1006) and Fetal Medicine
- 430 Foundation (495237). For the purpose of open access, the authors have applied a creative
- 431 commons attribution (CC BY) licence to any author accepted manuscript version arising.

432 Table 1: Scanning parameters

Placental and Fetal MRI	
Anatomical	T2 weighted Turbo Spin Echo
	TR 30s; TE 180ms; Matrix size 384x384x150;
	Resolution 1.25x1.25x2.5mm
T2* static	Multi-Echo Gradient Echo
	TR 23s; TE
	12.599ms/65.34ms/118.08ms/170.821ms/223.562ms;
	Dynamics – 2; Matrix size 224x224x850; Resolution
	2.5x2.5x2.5mm
Diffusion	Pulsed Gradient Echo Spin Echo, Echo Planar Imaging
	TR 6.6s; TE 78ms; Matrix 512x512x56; Resolution
	2x2x4mm; b 375 (6 directions); b 750 (6 directions)
T2* dynamic	Multi-Echo Gradient Echo
	TR 6.4s; TE 7.84ms/60.574ms/113.308ms/116.041ms;
	Dynamics – 30; Matrix size 256x256360; Resolution
	2.5x2.5x2.5mm
Cardiac MRI	
Cine bSSFP	bSSFP
	30 phases; FA 60; TR 4ms; TE 2ms; Resolution
	1.7.1.7.8mm; ~11 second breath-hold
Short axis stack	bSSFP
	TR 4.1s; TE 2ms; 14 slices; Sense 2; Resolution
	2x2x10mm
Phase contrast (breath-hold	30 phases; Sense 2; Resolution 2.5x2.5x8mm; 14
option)	second breath-hold
Phase contrast (free breathing	30 phases; Sense 2; Resolution 2.5x2.5x5mm; NSA 4
option)	

433 TR: repetition time; TE echo time; bSSFP balanced Steady State Free Precession

434 Table 2: Demographics of participants

Cohort (number of scans)	Pregnant controls (n=32)	Non- pregnant controls (n=38)	Pre- eclampsia (n=15)	Chronic hypertension (n=22)	Complicated pregnancy (n=9)
Age (years), median (IQR)	34.4	32.0	33.5	36.8	33.1
	(31.8-37.3)	(27.0-39.0)	(30.8-37.8)	(33.8-41.5)	(31.9-35.3)
BMI (kg/m ²), median (IOB)	24.7	25.3	28.1	31.9	23.1
incular (locity	(22.9-27.2)	(21.2-25.4)	(24.1-33.2)	(27.4-33.8)	(21.9-24.4)
	, , ,		, ,	, , , , , , , , , , , , , , , , , , ,	. ,
Gestational age	28.1		32.7	27.8	27.3
scan (weeks), median (IQR)	(23-5-32.1)		(30.0-34.1)	(25.0-32.0)	(24.4-33.3)
Gestational age at birth, mean	39.6		33.4	37.8	36.3
(SD)	(1.07)		(2.77)	(0.7)	(2.87)
Birth weight	3394		1781	2973	2370
(SD)	(402)		(552)	(636)	(978)
Birth weight (grams), range	2710-4175		695-3070	1560-4840	1010-3220
Previous pre- eclampsia (%)	0	0	3 (25)	2 (8)	0

435 BMI: body mass index; IQR: interquartile range

436 Table 3: Relationship between fetal brain and placental T2* and clinical parameters

437

Participant ID	Placental	Placental	Fetal	Fetal Dopplers	Fetal growth
	T2*	histopathology	brain		restriction
			Т2*		
1	Low	Abnormal	Low	Redistribution	Yes
2	Low	Other vascular	Low	Normal	Yes
		abnormality			
3	Low	Abnormal	Low	Redistribution	Yes
4	Low	Abnormal	Low	Redistribution	No
5	Normal	Normal	Normal	Normal	No
6	Low	Abnormal	Low	Redistribution	Yes
7	Low	Abnormal	Low	Redistribution	Yes
8	Normal	No pre-eclampsia	Normal	Normal	Yes
		related changes			
9	Low	Not available	Low	Redistribution	No
10	Low	Abnormal	Low	Redistribution	Yes
11	Low	Abnormal	Low	Redistribution	Yes
12	Low	Abnormal	Low	Redistribution	Yes
13	Normal	Normal	Normal	Normal	No

438

440 Figure 1: Schematic of study protocol



- 442 US: ultrasound; UtAD: uterine artery Doppler; AFI: amniotic fluid volume; UmbAD: umbilical
- 443 artery Doppler; MCA: middle cerebral artery Doppler; DV: ductus venosus Doppler; BP blood
- 444 pressure; CW cardiac work; CI: cardiac index; DW-SS EPI: diffusion weighted single shot echo
- 445 planar imaging; 4C: 4 chamber; 2C 2 chamber; 3C 3 chamber; SA: short axis; aAo ascending
- 446 aorta; dAo descending aorta; SVC superior vena cava.

447 Figure 2: Summary of images acquired on protocol



449 Figure 3: Placental anatomical and T2* by group



451 Figure 1: A) mean placental T2* by gestation; B) placental mass by gestation; C) skewness

- 452 and D) kurtosis by gestation. Black dot = healthy pregnancy; blue = chronic hypertension;
- 453 *orange = pre-eclampsia.*

454 Figure 4: Fetal brain anatomical and T2* by group



456 Figure 2: A) Fetal brain mean T2* by gestation; B) fetal brain volume by gestation. Black dot

457 = healthy pregnancy; blue = chronic hypertension; orange = pre-eclampsia.

458 Figure 5: Comparison of fetal brain mean T2* z-score and placental mean T2* z-score





460 Black dot = healthy pregnancy; blue = chronic hypertension; orange = pre-eclampsia.



461 Figure 6: Left ventricular remodelling associated with pre-eclampsia.

462 Figure 3: Left ventricular remodelling associated with pre-eclampsia (PE), illustrating

463 thickened walls and eccentricity & bulging located at the outflow track. **Top panel:** extremes

464 of the anatomical mode that best discriminate pregnant controls to pregnant pre-eclamptic

subjects - The 3D and bullseye views are colour-coded by thickness (in mm) and have black

466 and white spheres at the sides of the outflow track for easy mapping between both views –

- 467 A: Anterior; P: Posterior; L: Lateral; S: Septal. Bottom panel: box-plot of distributions of
- 468 pregnant controls (black), pre-eclampsia (blue), chronic hypertension (orange) and non-
- 469 pregnant controls (green), together with metrics of the discriminant performance of the
- 470 linear discriminant analysis between pregnant controls and pre-eclampsia (AUC: Area Under
- 471 the Curve; RS: Resubstitution; L1: Leave-1 out cross-validation).

473 Supplementary material

474 Supplementary Table S1: Medical history and obstetric outcomes of the pre-eclamptic cohort

Ethnicity (n)	
White	2
Black	5
South Asian	
Other	2
Parity (n)	
Nullinarous	9
Parous	4
Gestational age at PFT diagnosis (weeks) (range)	27.0 (20.7 – 35.0)
Pre-eclampsia superimposed on chronic hypertension (n)	6
PIGE (pg/ml) (n)	-
>12	2
<12	2
Proteinuria (n)	13
Acute kidney injury (n)*	5
Henatic dysfunction (n)*	8
Haemolysis (n)*	1
Neurological involvement (n)*	0
Thrombocytonaenia (n)*	2
Fetal growth restriction (n)	10
Gestational age at MBI (weeks) (range)	320(263 - 353)
Antenatal aspirin use (n)	7
Antibupartensives at time of MPI (n)	/
Single agent	Q
	8
Three agents	1
Maximum antihypertensives (n)	±
Single agent	5
	4
Three agents	3
Four agents	1
Evidence of redistribution on ultrasound at time of MRI (n)	-
Yes	9
No	4
Mode of delivery (n)	
Caesarean section in labour	2
Caesarean section prior to labour	10
Spontaneous vaginal	1
Indication for delivery (n)	
Maternal	4
Fetal	9
Neonatal complications (n)	

NICU admission	12
Respiratory support	4
Necrotising enterocolitis	2
Hypoglycaemia	5
Jaundice	6
Placental histopathology (n)	
Abnormalities consistent with pre-eclampsia	8
Other abnormalities	2
Normal	2
Not available	1

476 PIGF: placental growth factor.

477 *Diagnosed at any stage in pregnancy or postnatal period

478 Supplementary Figure S1: Summary of major findings in the control and pre-eclamptic cohorts



- 479 Placental mean T2* p<0.001; placental volume p=0.02; placental T2* skewness p=0.03; placental T2*
 480 kurtosis p=0.01.
- 481 Fetal brain mean $T2^* p < 0.001$; fetal brain volume p = 0.03.
- 482 Cardiac output p=0.22; cardiac work p=0.905; mean systolic blood pressure p<0.001; mean diastolic
- 483 blood pressure p=0.01; mean heart rate p=0.159
- 484 BP: blood pressure



485 Supplementary Figure S2: Anatomical models of variation that most correlate with

496 References

498 499	1.	Chappell LC, Cluver CA, Kingdom J, Tong S. Pre-eclampsia. <i>Lancet</i> . 2021;398:341-354. doi: 10.1016/S0140-6736(20)32335-7
500	2.	Steegers EA, von Dadelszen P. Duvekot JJ. Pijnenborg R. Pre-eclampsia. Lancet.
501		2010;376:631-644. doi: 10.1016/S0140-6736(10)60279-6
502	3.	Say L, Chou D, Gemmill A, Tunçalp Ö, Moller AB, Daniels J, Gülmezoglu AM, Temmerman M,
503		Alkema L. Global causes of maternal death: a WHO systematic analysis. <i>Lancet Glob Health</i> .
504		2014;2:e323-333. doi: 10.1016/S2214-109X(14)70227-X
505	4.	Valensise H, Vasapollo B, Gagliardi G, Novelli GP. Early and late preeclampsia: two different
506		maternal hemodynamic states in the latent phase of the disease. <i>Hypertension</i> .
507		2008;52:873-880. doi: 10.1161/HYPERTENSIONAHA.108.117358
508	5.	Castleman JS, Ganapathy R, Taki F, Lip GY, Steeds RP, Kotecha D. Echocardiographic
509		Structure and Function in Hypertensive Disorders of Pregnancy: A Systematic Review. Circ
510		Cardiovasc Imaging. 2016;9. doi: 10.1161/CIRCIMAGING.116.004888
511	6.	Melchiorre K, Sutherland G, Sharma R, Nanni M, Thilaganathan B. Mid-gestational maternal
512		cardiovascular profile in preterm and term pre-eclampsia: a prospective study. BJOG.
513		2013;120:496-504. doi: 10.1111/1471-0528.12068
514	7.	Bellamy L, Casas JP, Hingorani AD, Williams DJ. Pre-eclampsia and risk of cardiovascular
515		disease and cancer in later life: systematic review and meta-analysis. BMJ. 2007;335:974.
516		doi: 10.1136/bmj.39335.385301.BE
517	8.	Leon LJ, McCarthy FP, Direk K, Gonzalez-Izquierdo A, Prieto-Merino D, Casas JP, Chappell L.
518		Preeclampsia and Cardiovascular Disease in a Large UK Pregnancy Cohort of Linked
519		Electronic Health Records: A CALIBER Study. Circulation. 2019;140:1050-1060. doi:
520		10.1161/CIRCULATIONAHA.118.038080
521	9.	Thilaganathan B. Maternal Cardiac Dysfunction Precedes Development of Preeclampsia.
522		Hypertension. 2020;76:321-322. doi: 10.1161/HYPERTENSIONAHA.120.15281
523	10.	Garcia-Gonzalez C, Georgiopoulos G, Azim SA, Macaya F, Kametas N, Nihoyannopoulos P,
524		Nicolaides KH, Charakida M. Maternal Cardiac Assessment at 35 to 37 Weeks Improves
525		Prediction of Development of Preeclampsia. Hypertension. 2020;76:514-522. doi:
526		10.1161/HYPERTENSIONAHA.120.14643
527	11.	Grothues F, Smith GC, Moon JC, Bellenger NG, Collins P, Klein HU, Pennell DJ. Comparison of
528		interstudy reproducibility of cardiovascular magnetic resonance with two-dimensional
529		echocardiography in normal subjects and in patients with heart failure or left ventricular
530		hypertrophy. <i>Am J Cardiol</i> . 2002;90:29-34. doi: 10.1016/s0002-9149(02)02381-0
531	12.	Bellenger NG, Davies LC, Francis JM, Coats AJ, Pennell DJ. Reduction in sample size for
532		studies of remodeling in heart failure by the use of cardiovascular magnetic resonance. J
533		Cardiovasc Magn Reson. 2000;2:271-278. doi: 10.3109/10976640009148691
534	13.	Colford K, Price AN, Sigurdardottir J, Fotaki A, Steinweg J, Story L, Ho A, Chappell LC, Hajnal
535		JV, Rutherford M, et al. Cardiac and placental imaging (CARP) in pregnancy to assess
536		aetiology of preeclampsia. <i>Placenta</i> . 2022;122:46-55. doi: 10.1016/j.placenta.2022.03.012
537	14.	Hakim J, Senterman MK, Hakim AM. Preeclampsia is a biomarker for vascular disease in both
538		mother and child: the need for a medical alert system. <i>Int J Pediatr</i> . 2013;2013:953150. doi:
539		10.1155/2013/953150
540	15.	Falco ML, Sivanathan J, Laoreti A, Thilaganathan B, Khalil A. Placental histopathology
541		associated with pre-eclampsia: systematic review and meta-analysis. Ultrasound Obstet
542	4.6	<i>Gynecol.</i> 2017;50:295-301. doi: 10.1002/uog.17494
543	16.	Ho A, Hutter J, Slator P, Jackson L, Seed PT, Mccabe L, Al-Adnani M, Marnerides A, George S,
544		Story L, et al. Placental magnetic resonance imaging in chronic hypertension: A case-control
545		study. <i>Placenta</i> . 2021;104:138-145. doi: 10.1016/j.placenta.2020.12.006

546 547 548 549	17.	Ho AEP, Hutter J, Jackson LH, Seed PT, Mccabe L, Al-Adnani M, Marnerides A, George S, Story L, Hajnal JV, et al. T2* Placental Magnetic Resonance Imaging in Preterm Preeclampsia: An Observational Cohort Study. <i>Hypertension</i> . 2020;75:1523-1531. doi: 10.1161/HYPERTENSIONAHA.120.14701
550 551 552	18.	Slator PJ, Hutter J, Palombo M, Jackson LH, Ho A, Panagiotaki E, Chappell LC, Rutherford MA, Hajnal JV, Alexander DC. Combined diffusion-relaxometry MRI to identify dysfunction in the human placenta. <i>Magn Reson Med</i> . 2019:82:95-106. doi: 10.1002/mrm.27733
553 554 555	19.	Sinding M, Peters DA, Frøkjaer JB, Christiansen OB, Petersen A, Uldbjerg N, Sørensen A. Placental magnetic resonance imaging T2* measurements in normal pregnancies and in those complicated by fetal growth restriction. <i>Ultrasound Obstet Gynecol</i> . 2016;47:748-754.
556		doi: 10.1002/uog.14917
557	20.	de Marvao A, Dawes TJ, Shi W, Durighel G, Rueckert D, Cook SA, O'Regan DP. Precursors of
558		Hypertensive Heart Phenotype Develop in Healthy Adults: A High-Resolution 3D MRI Study.
559		JACC Cardiovasc Imaging. 2015;8:1260-1269. doi: 10.1016/j.jcmg.2015.08.007
560	21.	Petersen SE, Aung N, Sanghvi MM, Zemrak F, Fung K, Paiva JM, Francis JM, Khanji MY,
561		Lukaschuk E, Lee AM, et al. Reference ranges for cardiac structure and function using
562		cardiovascular magnetic resonance (CMR) in Caucasians from the UK Biobank population
563		cohort. J Cardiovasc Magn Reson. 2017;19:18. doi: 10.1186/s12968-017-0327-9
564	22.	Marciniak M, van Deutekom AW, Toemen L, Lewandowski AJ, Gaillard R, Young AA, Jaddoe
565		VWV, Lamata P. A three-dimensional atlas of child's cardiac anatomy and the unique
566		morphological alterations associated with obesity. <i>Eur Heart J Cardiovasc Imaging</i> .
567		2022;23:1645-1653. doi: 10.1093/ehjci/jeab271
568	23.	Hermida U, Stojanovski D, Raman B, Ariga R, Young AA, Carapella V, Carr-White G, Lukaschuk
569		E, Piechnik SK, Krämer CM, et al. Left ventricular anatomy in obstructive hypertrophic
570		cardiomyopatny: beyond basal septal hypertrophy. Eur Heart J Cardiovasc Imaging. 2022.
5/1	24	dol: 10.1093/enjci/jeac233
572	24.	Balabali G, Halliudy BP, Hallillersley D, Killalul CA, Plasau SK, Bishop Wij, Lalilald P. Leil
575		$2022 \cdot 24 \cdot 1127$ 1147, doi: 10.1002/ouropaco/ouab206
575	25	2022,24.1137-1147. doi: 10.1035/ediopace/edab300
576	25.	and robust method to generate nationt-specific cubic Hermite meshes. Med Image And
570		2011:15:801-813, doi: 10.1016/i media 2011.06.010
578	26	Hutter Slator PL Jackson Gomes ADS Ho A Story O'Muircheartaigh Teixeira RPAG
579	20.	Channell I.C. Alexander D.C. et al. Multi-modal functional MRI to explore placental function
580		over gestation. Magn Reson Med. 2019:81:1191-1204. doi: 10.1002/mrm.27447
581	27.	Sinding M. Sørensen A. Hansen DN. Peters DA. Frøkiær JB. Petersen AC. T2* weighted
582		placental MRI in relation to placental histology and birth weight. <i>Placenta</i> . 2021:114:52-55.
583		doi: 10.1016/j.placenta.2021.07.304
584	28.	Schabel MC, Roberts VHJ, Gibbins KJ, Rincon M, Gaffney JE, Streblow AD, Wright AM, Lo JO,
585		Park B, Kroenke CD, et al. Quantitative longitudinal T2* mapping for assessing placental
586		function and association with adverse pregnancy outcomes across gestation. PLoS One.
587		2022;17:e0270360. doi: 10.1371/journal.pone.0270360
588	29.	Stanek J. Placental pathology varies in hypertensive conditions of pregnancy. <i>Virchows Arch</i> .
589		2018;472:415-423. doi: 10.1007/s00428-017-2239-3
590	30.	Sohlberg S, Mulic-Lutvica A, Lindgren P, Ortiz-Nieto F, Wikström AK, Wikström J. Placental
591		perfusion in normal pregnancy and early and late preeclampsia: a magnetic resonance
592		imaging study. Placenta. 2014;35:202-206. doi: 10.1016/j.placenta.2014.01.008
593	31.	Ogge G, Chaiworapongsa T, Romero R, Hussein Y, Kusanovic JP, Yeo L, Kim CJ, Hassan SS.
594		Placental lesions associated with maternal underperfusion are more frequent in early-onset
595		than in late-onset preeclampsia. J Perinat Med. 2011;39:641-652. doi:
596		10.1515/jpm.2011.098

597	32.	Vasylechko S, Malamateniou C, Nunes RG, Fox M, Allsop J, Rutherford M, Rueckert D, Hajnal
598		JV. T2* relaxometry of fetal brain at 1.5 Tesla using a motion tolerant method. Magn Reson
599		Med. 2015;73:1795-1802. doi: 10.1002/mrm.25299
600	33.	Figueiró-Filho EA, Mak LE, Reynolds JN, Stroman PW, Smith GN, Forkert ND, Paolozza A,
601		Rätsep MT, Croy BA. Neurological function in children born to preeclamptic and
602		hypertensive mothers - A systematic review. Pregnancy Hypertens. 2017;10:1-6. doi:
603		10.1016/j.preghy.2017.07.144
604	34.	Boardman H, Lamata P, Lazdam M, Verburg A, Siepmann T, Upton R, Bilderbeck A, Dore R,
605		Smedley C, Kenworthy Y, et al. Variations in Cardiovascular Structure, Function, and
606		Geometry in Midlife Associated With a History of Hypertensive Pregnancy. Hypertension.
607		2020;75:1542-1550. doi: 10.1161/HYPERTENSIONAHA.119.14530
608	35.	Lisonkova S, Joseph KS. Incidence of preeclampsia: risk factors and outcomes associated with
609		early- versus late-onset disease. Am J Obstet Gynecol. 2013;209:544.e541-544.e512. doi:
610		10.1016/j.ajog.2013.08.019
611	36.	He J, Chen Z, Chen C, Liu P. Comparative study of placental T2* and intravoxel incoherent
612		motion in the prediction of fetal growth restriction. <i>Placenta</i> . 2021;111:47-53. doi:
613		10.1016/j.placenta.2021.06.005
614	37.	Slator PJ, Hutter J, Marinescu RV, Palombo M, Jackson LH, Ho A, Chappell LC, Rutherford M,
615		Hajnal JV, Alexander DC. Data-Driven multi-Contrast spectral microstructure imaging with
616		InSpect: INtegrated SPECTral component estimation and mapping. Med Image Anal.
617		2021;71:102045. doi: 10.1016/j.media.2021.102045
618		