

1 **Emerging viruses are an underestimated cause of undiagnosed febrile illness in Uganda**


2

3 Authors

4 Shirin Ashraf*¹, Hanna Jerome*¹, Daniel Lule Bugembe³, Deogratius Ssemwanga^{2,3}, Timothy
5 Byaruhanga², John Timothy Kayiwa², Robert Downing², Jesus F. Salazar-Gonzalez³, Maria G.
6 Salazar³, James G. Shepherd¹, Chris Davis¹, Nicola Logan¹, Vattipally B. Sreenu¹, Gavin S. Wilkie¹,
7 Ana da Silva Filipe¹, Alfred Ssekagiri^{1, 2}, Prossy Namuwulya², Henry Bukenya², Brian K. Kigozi²,
8 Weronika Witkowska McConnell¹, Brian J. Willett¹, Stephen Balinandi², Julius Lutwama²,
9 Pontiano Kaleebu^{2,3}, Josephine Bwogi^{1,2}, Emma C. Thomson^{1,4}

10

11 *Contributed equally

12  Contributed equally

13 Institutions

14 1 MRC-University of Glasgow Centre for Virus Research

15 2 Uganda Virus Research Institute

16 3 MRC/UVRI & LSHTM Uganda Research Unit

17 4 LSHTM

18

19

20 **SUMMARY**

21 **Background:** Viruses that cause acute febrile illness (AFI) in sub-Saharan Africa cause a spectrum
22 of disease from mild to life-threatening. Viral infection is often undiagnosed, as routine
23 diagnostics are insufficient to capture the diversity of circulating pathogens.

24

25 **Methods:** 1281 patients with fever of 2-7 days were prospectively recruited from three sites in
26 Uganda as part of the CDC-UVRI AFI Study and screened with enhanced diagnostics. Plasma from
27 233 undiagnosed patients was analysed using metagenomic next-generation sequencing (mNGS).
28 Confirmatory testing was carried out by PCR and serology.

29

30 **Findings:** Thirty-eight viral pathogens were identified by mNGS in 35/233 (15%) undiagnosed
31 patients including Measles, Hepatitis A/B/E viruses, Human immunodeficiency virus-1,
32 Rhinovirus, Rotavirus-like virus, Human herpesvirus 6B, Human parainfluenza virus 3 and
33 Enteroviruses. Four high-consequence arboviruses were found in six patients; Crimean-Congo
34 haemorrhagic fever virus, Rift Valley fever virus, dengue virus and yellow fever virus. Le Dantec
35 virus, last reported in 1969, was detected and confirmed by serology in one patient (and a
36 contact of that patient). The majority of patients (23/30; 76%) diagnosed with an acute viral
37 infection were treated with antibiotics and/or (12/30; 40%) antimalarials.

38

39 **Interpretation:** AFI in Uganda is commonly associated with undiagnosed viral infection, including
40 high-consequence and rarely reported viruses. This highlights an ongoing risk to public health
41 and the need for improved vigilance. MNGS alongside diagnostic serology is a powerful method
42 for clinical surveillance to investigate circulating viral pathogens. Cost-effective diagnostic assays
43 should be adapted according to regional needs for testing.

44

45 **Funding:** Medical Research Council and Wellcome Trust

46

47 **Keywords:** genomics, febrile illness, sub-Saharan Africa, haemorrhagic fever viruses

48 INTRODUCTION

49 East Africa is a biodiverse region and a hotspot for viral zoonoses. In the last decade, Uganda has
50 reported outbreaks of haemorrhagic fever caused by Ebola (EBOV), Sudan (SUDV), Marburg
51 (MARV), Rift Valley fever (RVFV), yellow fever (YFV) and Crimean-Congo haemorrhagic fever
52 (CCHFV) viruses¹. Many vector-borne viral infections are prevalent in Uganda, and several were
53 described here for the first time, including Zika virus (ZIKV), Semliki Forest virus (SFV),
54 chikungunya virus (CHIKV), O’Nyong Nyong virus (ONNV) and West Nile virus (WNV)².
55 Anthropogenic change, as a result of frequent travel, increasing commerce, and shifting global
56 climate patterns raises the risk of spread within and outside zoonotic hotspots³. Many infections
57 are likely missed due to the scarcity of diagnostic tests, increasing the risk of spread of high
58 consequence viruses. Further, they may be misattributed to malaria or bacterial infection,
59 increasing the risk of antimicrobial resistance. In this study, we investigated the occurrence of
60 undiagnosed viral infections as a cause of acute febrile illness (AFI).

61

62 AFI is a common cause of presentation to healthcare centres (HC) across sub-Saharan Africa
63 (SSA), despite a significant reduction in malaria prevalence following successful intervention
64 campaigns⁴. Communicable diseases are estimated to account for up to 36% productivity losses
65 in SSA⁵. The contribution of viral infection to AFI in Africa is at least 8-60%, likely an under-
66 estimate as population studies have relied on traditional diagnostic tests which are limited to
67 selected pathogens⁶⁻⁹. A thorough understanding of the burden of viral disease is essential for
68 implementing prevention strategies by limiting vector-borne transmission, introducing
69 vaccination where needed and available, rapid cost-effective diagnostics and treatments.
70 Metagenomic next-generation sequencing (mNGS) allows for unbiased identification of viral
71 genomes in clinical samples, including known and novel pathogens. In this study, we used mNGS
72 to identify viral infections in patients in Uganda presenting with AFI that remained undiagnosed
73 after routine serological and PCR-based assays.

74 **METHODS**

75 **Patients and sampling**

76 1281 patients were recruited prospectively with informed consent into the AFI study from three
77 study sites in Uganda and assigned an anonymized study number, not known to anyone outside the
78 recruitment team at each healthcare provision site; Ndejje HC IV (Wakiso), St. Paul's HC IV (Kasese),
79 and Adumi HC IV (Arua) between April 2011 and January 2013 (**Figure 1a**). Inclusion criteria were >2
80 years age with (a) a reported fever lasting 2-7 days or $\geq 38^{\circ}\text{C}$ temperature on admission, or (b)
81 symptoms consistent with brucellosis or typhoid fever. Cases with clinical evidence of an alternative
82 diagnoses such as otitis media were excluded. Samples were obtained at presentation (acute sample)
83 and 14-21 days later (convalescent sample). Diagnostic assays including serology, blood culture and
84 blood films were carried out to identify and exclude active or recent infection with malaria, typhoid,
85 leptospirosis, rickettsiae, CHIKV, dengue (DENV), WNV, YFV and ONNV. 210 patients remained
86 undiagnosed and acute plasma were retrospectively tested using mNGS. In addition, 23 samples
87 were obtained from two outbreaks of febrile illness that occurred within the time period of the
88 study.

88

89 **Ethics statement**

90 Ethical approval for the AFI study was granted by the Uganda Virus Research Institute (UVRI)
91 Research Ethics Committee (GC/127/10/02/19) and the Uganda National Council for Science and
92 Technology (HS767).

93

94 **Sample processing and mNGS**

95 RNA was extracted from 200 μl of plasma using the Agencourt RNAdvance Blood Kit (Beckman
96 Coulter), according to manufacturer's instructions, including DNase treatment at 37°C for 15min.
97 RNA was reverse transcribed using Superscript III (Invitrogen) followed by dsDNA synthesis with
98 NEBNext Enzyme (New England Biolabs). DNA libraries were prepared using LTP low-input Library
99 preparation kit (KAPA Biosystems) as previously described¹⁰. Resulting libraries were quantified
100 with the Qubit 3.0 fluorometer (Invitrogen) and their size determined using a 4200 TapeStation
101 (Agilent). Libraries were pooled in an equimolar ratio and sequenced on the Illumina MiSeq and
102 NextSeq platforms.

103

104 **Bioinformatic analysis**

105 Raw Fastq files were searched directly for evidence of viral sequences using Diamond BLASTX
106 **(Figure S1a)**¹¹. *De novo* assembly was carried out using dipspades and contigs were identified
107 using Diamond BLASTX against the nr database¹². Viral hits detected with blastx were confirmed
108 with blastn and mapped to the closest reference genome using *Tanoti*, a mapper developed for
109 use with highly diverse viral genomes (github.com/vbsreenu/Tanoti). A diagnosis threshold of 28
110 nucleotides was estimated as a reliable cut-off following an analysis of all common human viral
111 pathogens and removal of genomic repeat regions, well under the minimum sequence length in
112 our samples **(Figure S1b)**. Maximum-likelihood phylogenetic analysis was carried out using
113 IQTREE and 1000 ultra-fast bootstrap replicates using relevant reference sequences¹³.
114 Uncorrected pairwise-distances were estimated using MEGA 10.0¹⁴.

115

116 **Capture ELISA**

117 Rhabdovirus (Le Dantec; LDV and Adumi virus) glycoprotein genes (synthesized by Eurofins
118 Genomics) excluding the transmembrane regions (identified using TMHMM and TMPred) were
119 amplified and cloned into the secretory mammalian expression vector pHLSec containing a C-
120 terminal 6xHistidine tag^{15,16}. Human embryonic kidney cells (HEK 293T) were maintained in
121 Dulbecco's modified Eagle's medium (DMEM) supplemented with 100IU/ml penicillin, 100µg/ml
122 streptomycin, 2mM glutamine and 10% foetal bovine serum. Cells were transfected (Fugene 6,
123 Promega) and cell supernatant containing secreted recombinant glycoproteins was harvested at
124 48-hours and expression confirmed by Western Blotting against 6xHis. ELISA plates (ImmulonHB)
125 were coated with rabbit anti-His antibody overnight and blocked with 2.5% BSA/PBS at 37°C for 1
126 hour. Cell supernatant containing glycoprotein was added to the wells followed by patient
127 serum, both for 1 hour at 37°C. This was followed by HRP-conjugated goat anti-human IgG at
128 room temperature for 1 hour. Wells were washed after each step with 0.1% Tween-20/PBS.
129 Reactions were developed using TMB substrate, stopped with 0.16M sulphuric acid and read on
130 a spectrophotometer (Pherastar).

131

132 **Pseudotype neutralization assay**

133 The use of Vesicular Stomatitis Virus (VSV) in which the glycoprotein sequence has been replaced
134 (VSVΔG*luc*) to generate pseudotyped viruses has been described previously¹⁷. Glycoprotein
135 genes were cloned into the eukaryotic expression vector and transfected into HEK-293T cells
136 using polyethylenimine followed by infection with VSVΔG*luc*-VSV-G at a multiplicity of infection

137 of 0.02. Cells were incubated for one hour at 37°C, washed with PBS and re-incubated in DMEM.
138 Supernatants were harvested 72-hours post infection. Pseudoparticle activity was assessed on
139 HEK-293T cells by measuring luciferase activity at 72-hours on a Microbeta 1450 Jet luminometer
140 (Perkin Elmer).
141 For neutralisation assays, 1×10^4 HEK-293T cells were plated in a white 96-well plate and
142 incubated at 37°C for one hour. Four-fold serum dilutions were prepared in DMEM ranging from
143 1:8 to 1:131072. Serum was mixed with either VSV Δ Gluc-VSV-G or VSV Δ Gluc-LDV-G at a
144 concentration expected to generate a luciferase reading of 1×10^6 counts per second and
145 incubated for 1 hour at 37°C. 50 μ l per well of the pseudotype/serum mixture was added to the
146 cells and incubated for 24 hours. Luciferase substrate was added to each well (Steadylite plus™,
147 Perkin Elmer) and luminescence measured on a Chameleon V plate scintillation counter (Hidex
148 Oy).

149 **RESULTS**

150 **Demographics and clinical analysis**

151 15% of patients with undiagnosed AFI (35/233) had evidence of acute viral infection by mNGS
152 (**Figure 1b**). Demographic and clinical data was available for 30/35 patients with AFI. The median
153 age at clinical presentation was 7.5 years (range 2-40). 80% of cases occurred in children under
154 the age of 16. Males were more commonly represented than females (20/30; 67% versus 10/30;
155 33%). The majority had been recently bitten by mosquitoes (22/30; 73%) and/or other
156 arthropod vectors, including biting flies (3/30; 10%), ticks (3/30; 10%), fleas (2/30; 7%) and lice
157 (1/30; 3%). Ten percent reported haemorrhagic symptoms but none of these had evidence of a
158 viral haemorrhagic fever (VHF) virus by mNGS. At presentation, the median temperature was
159 38.2°C. Other clinical symptoms are described in **Table 1**. Common presumptive diagnoses
160 included malaria, respiratory tract infection, typhoid fever and unknown. The majority 23/30
161 (77%) of patients were presumptively treated with antibiotics, 12 (40%) were treated with
162 antimalarials and 5 (17%) with antihelminth therapy.

163 Of the 35 patients diagnosed by mNGS, 6 (17%) had a virus associated with VHF (CCHFV, RVFV,
164 YFV and DENV), 9 (25%) with viruses associated with respiratory infection (human parainfluenza
165 3, rhinovirus C and measles virus), 8 (22%) with hepatitis causing viruses (HAV, HBV, HEV), and 5
166 (14%) with gastroenteritis associated viruses (**Table 2, Supplementary Table S1**). Six patients
167 (17%) were infected with human immunodeficiency virus-1 (HIV-1). One patient was infected
168 with the rhabdovirus LDV. Two of these cases, one each of yellow fever and RVF were correctly
169 identified in the context of a known outbreak.

170

171 **Viruses associated with VHF**

172 Genomes from four VHF-associated viruses were detected in six febrile patients; CCHFV (patient
173 138-2), RVFV (patients 117-3, RVF58), DENV (patient ARB248) and YFV (patients YFV685,
174 ARB176).

175

176 The CCHF patient (138-2) was a male in his 30s from Kasese district. He presented with fever
177 (37.9°C), headache, nausea and joint pain. He had been bitten by mosquitoes but had not been
178 aware of tick bites and had not recently slaughtered animals. He was treated with vitamins and
179 paracetamol and recovered. Partial (10-25% coverage) CCHFV sequences across all three
180 segments; large (L), medium (M) and small (S) were detected by mNGS (**Figure 2a**). The L and S

181 segments had >90% nucleotide identity to previously identified genomes from Uganda and
182 Democratic Republic of the Congo, respectively (**Table 2, S1**). The M segment had 81%
183 nucleotide identity to a genome from Uganda, suggestive of higher divergence or possible
184 reassortment (**Figure 2d**). Convalescent serology was not available from the patient, who
185 recovered and had moved abroad.

186

187 Two cases of RVF (117-3, RVF58) were detected. RVF58 was suspected due to occupational risk
188 (contact with slaughtered animals) and suggestive symptoms during an outbreak, and was
189 confirmed by serology. 117-3 was a woman in her 30s from a rural part of Arua district who
190 gave a 3-day history of fever but whose temperature was recorded as 36.3°C on admission. She
191 described chills, headache, eye, neck and joint pain, nausea, diarrhoea and jaundice. She had
192 been bitten by mosquitoes and fleas. She was given paracetamol and attended a traditional
193 healer for further treatment. Convalescent serology was not available for this patient. Viral reads
194 most closely matched the L segment of a Ugandan strain in case 1 (RVF58) and the M segment of
195 a Kenyan strain in case 2 (117-3).

196

197 Two cases of YFV infection and one DENV infection (genotype 3) were detected during suspected
198 YFV outbreaks and confirmed by convalescent serology. All genomes identified by mNGS were
199 related to East/Central Africa lineages (**Table 2, S1; Figure 2 b,c,e,f**).

200

201 While haemorrhagic symptoms were recorded for 3/30 (10%) AFI study cohort patients, none of
202 the these had a known VHF virus detected. The study patients described above with CCHFV and
203 RVFV did not report classical haemorrhagic symptoms.

204

205 **Le Dantec virus**

206 Patient 220-2 was a male aged between 5 and 10 years from Kasese district. He presented to hospital
207 with a 38.5°C fever of 4 days, with headache, abdominal pain, vomiting and joint pains. He reported having
208 been bitten by mosquitoes. He was treated with ciprofloxacin and paracetamol for suspected
209 typhoid fever and made a full recovery. By mNGS, we detected the full genomic sequence of a
210 previously rarely reported rhabdovirus; Le Dantec virus (LDV; **Figure 3a**). This was confirmed by
211 direct RT-PCR from patient plasma and Sanger sequencing. The LDV genome had 94% nucleotide
212 identity to the RefSeq LDV genome (NC_034443.1), isolated for the first time in 1965 in Senegal

213 and sequenced from an archival sample in 2015¹⁸ (**Figure 3d**). In keeping with an active infection,
214 we demonstrated immunological response to LDV glycoprotein by ELISA and pseudotype
215 neutralisation assay, in both the patient and in a contact of the patient who had been unwell at
216 the same time, using convalescent serum (**Figure 3b,c**).

217

218 **Viruses associated with acute hepatitis**

219 Hepatitis A (HAV) and hepatitis E (HEV) viruses were detected in 5 (patients ARB169, 218-1, 169-
220 1, 184-2, 145-3) and 2 (patients 20-2, 60-1) patients respectively. Clinical data was available for
221 four of the HAV cases. The age of infection ranged from 2 to 15 years. All were febrile at
222 presentation. Three of four patients presented with vomiting, one had diarrhoea and one had
223 jaundice. All HAV genomes were genotype IB and related to previously reported sequences from
224 Uganda (**Figure 4a, S3e**). The HEV patients were aged between 0 and 5 years, and febrile at
225 presentation. One had diarrhoea and vomiting and the other had undifferentiated fever.
226 The two HEV isolates were genotype 1e and closely related to a Nigerian sequence (**Figure 4b, S3d**).

227

228 **Viruses associated with gastroenteritis**

229 Gastrointestinal viruses were detected in plasma from 5 patients, including 4 enteroviruses (the
230 echoviruses E3, E4 and E9 and Coxsackie B3 virus from patients 212-1, 316-3, 165-1, 195-1
231 respectively) and a fragment of a highly divergent rotavirus H-like virus in one patient (96-1)
232 (**Figure 4c, S3f**).

233

234 **Viruses associated with respiratory infection**

235 Respiratory viruses were detected in plasma from 9 patients and included six cases of measles
236 morbillivirus (patients 215-1, 217-1, 242-1, 223-1, 113-1, 203-1), two cases of human
237 parainfluenza virus 3 (patients 88-1, 114-1) and one case of rhinovirus C (patient 40-2) (**Figure**
238 **4c,d and S3b,c**). The measles virus sequences closely matched recently reported Ugandan B3
239 genomes. The rhinovirus C clustered with subtypes RV-C12 and RV-C44.

240

241 **Bloodborne viruses**

242 The bloodborne viruses HBV (patient 192-1) and HIV-1 (patients RVF58, ARB176, 143-2, 390-1,
243 199-2, 214-1) may have occurred as acute or chronic infections, detected in a population with a
244 high seroprevalence of both viruses. HBV was detected in a teenage patient (192-1)

245 presenting with headache, anorexia and abdominal pain. It was most closely related to a
246 sequence from Rwanda. HIV was identified as a co-infection in two patients, one with RVFV
247 (RVF58) and another with YFV (ARB176) (**Figure 4e, S3a**); suggesting chronic infection. Patient
248 214-1 was infected with HIV-1 subtype A1, clustering with another Ugandan sequence. All other
249 HIV-1 sequences were subtype D.

250

251 **Herpesviruses**

252 Three patient samples (082-2, 185-2 and 143-2) contained human herpesvirus-6B (HHV6B)
253 genome, the cause of roseola infantum, related to strains from USA, Uganda and DRC. Seven
254 patients (187-2, 189-2, ARB244, ARB265, ARB454, ARB688 and HEV2) had evidence of
255 cytomegalovirus (HHV5) in the bloodstream; this may have been acute or chronic infection. No
256 varicella zoster virus (VZV) infections were detected.

257

258 **Viruses detected that are not known to be associated with disease**

259 MNGS also detected viruses often found in human populations that have not been definitively
260 associated with disease. These included the annelloviruses: Torque Teno virus (TTV), Torque
261 Teno mini virus (TTMV), Torque Teno midi virus (TTMdV); and human pegiviruses (HPgV) (**Table**
262 **S2**).

263 We also detected a second and novel rhabdovirus sequence in several AFI samples by *de novo*
264 assembly (Adumi virus). This virus was likely to be a contaminant of blood collection tubes, as
265 discussed in Supplementary Information.

266 **DISCUSSION**

267 In this prospective cohort study of AFI in Uganda, we used mNGS to identify viral causes of acute
268 undiagnosed fever in 210/1281 (16.4%) patients who tested negative after extensive screening
269 for well-characterised pathogens, and an additional 23 samples obtained from outbreaks during
270 the same time period. The cohort was prospectively recruited from patients presenting to three
271 HCs in rural and urban Uganda. We detected viral pathogens in 35/233 (15.5%) patients,
272 including 6/233 (2.6%) associated with VHF. Lack of clinical diagnosis of such viruses means that
273 appropriate public health measures may not be taken to prevent onward transmission and
274 appropriate treatment may not be given. The most commonly detected viruses reflected acute
275 gastrointestinal or respiratory infection. The detection of active blood-borne virus infection
276 reflects a chronic burden of disease and a need to widen treatment in the Ugandan population.
277 Importantly, acute viral infection was invariably misdiagnosed as bacterial or malaria and treated
278 with antibiotics or antimalarials. This carries an associated risk of antimicrobial drug resistance.
279 We also noted vaccine-preventable infections in children (HAV, HBV, measles, Rotavirus)
280 indicating the need for a more comprehensive vaccination schedule.

281

282 AFI is widespread in SSA and in the absence of specialised point-of-care diagnostics; clinical
283 symptoms associated with different viruses may be indistinguishable. Even high-consequence
284 VHFs such as Lassa fever and Ebola virus disease present with haemorrhage in less than 20% of
285 cases^{19,20}. We identified CCHFV in a patient with undifferentiated fever who did not report
286 contact with ticks or slaughtered animals, highlighting the limitations of current risk algorithms in
287 identifying such patients in an endemic area. A high CCHFV seroprevalence has recently been
288 reported in farming communities and domesticated animals in Uganda, suggesting that
289 unrecognised infection may be common²¹.

290

291 We also detected the mosquito-borne infections RVFV, DENV and YFV; of which RVF can also be
292 caused by contact with infected livestock. RVF is widespread across Africa and may cause a
293 spectrum of illness from mild to severe. It is associated with haemorrhage and encephalitis in
294 around 1% of human cases²². DENV is transmitted by *Aedes* mosquitoes and is endemic in Asia,
295 South America and sub-Saharan Africa. It is relatively less well reported from the African
296 continent, although studies suggest high seroprevalence^{23,24}. Worldwide, there are an estimated
297 100 million clinical cases per year, of which around 5% progress to dengue shock syndrome or

298 dengue haemorrhagic fever. We detected DENV-3 in this study, in keeping with previous reports
299 of genotypes 1-3 on the African continent. YFV is endemic in Africa and South America and is
300 transmitted by *Aedes* and *Haemagogus* mosquitoes. Severe disease with haemorrhage occurs in
301 up to 12% of those infected²⁵. An effective live vaccine exists for YFV and has recently been used
302 to control outbreaks in Brazil and DRC. It is required for travellers to Uganda and other endemic
303 regions, but the local unvaccinated population may remain vulnerable to infection.

304

305 We detected two rhabdoviruses in this study; LDV, a potential pathogen and Adumi virus, a likely
306 contaminant. Rhabdoviruses are highly divergent and ubiquitous in host range, infecting plants,
307 invertebrates and vertebrates. Human disease is well-described; rabies lyssavirus causes fatal
308 neurological infection and Chandipura virus causes acute neurological disease. Other, less well
309 described members have also been suggested as human pathogens, including Bas Congo virus
310 that was associated with a single outbreak of haemorrhagic disease in the DRC. Additionally,
311 studies have also identified rhabdoviruses in blood from healthy asymptomatic individuals^{26,27}.
312 LDV is an emerging pathogen and has only two historical documentations; in a child with
313 hepatosplenomegaly and fever in Senegal in 1965 (sequenced in 2015) and in a middle-aged
314 shipyard worker with fever and severe neurological symptoms in Wales in 1969, who had
315 offloaded a ship inbound from Nigeria. LDV belongs to the genus *Ledantevirus* and closely related
316 species have been found in rodents and bats. Neurological pathology has been described in
317 rodents infected experimentally with the virus²⁸. In our study, we demonstrated an immune
318 response and the development of neutralising antibodies in the infected individual and a
319 household contact who had similar symptoms at the time.

320

321 Vaccine-preventable diseases remain common in Uganda, even for viruses covered under routine
322 immunisation, indicating insufficient coverage to generate herd immunity. Uganda has an active
323 Expanded Programme of Immunization (EPI) against measles, HBV and rotavirus. YFV vaccination
324 was introduced in October 2022. Vaccine coverage is lower in some neighbouring countries and
325 this may increase the risk of imported infection. Measles vaccine in Uganda is often given as a
326 single dose and booster vaccines during outbreaks. Such an approach may limit severe illness but
327 has lower efficacy than the two-dose regimen used in other countries²⁹.

328

329 We detected a viral pathogen in 15.5% of undiagnosed cases, however, the high detection of
330 viruses associated with respiratory and gastrointestinal infection in plasma highlights the high
331 burden that these diseases carry in Uganda. The inclusion of respiratory and stool samples (not
332 available in this study) in future studies will likely reveal a much higher prevalence of infection.
333 Additionally, the inclusion of serological assays would increase detection of infections outside
334 the viraemic phase. We have likely under-estimated the burden of viral disease in this study for
335 these reasons.

336

337 Viral metagenomics, although a powerful and sensitive method for unbiased genomic discovery
338 is prone to limitations. Caution must be practiced when interpreting NGS data to exclude
339 artefacts, which can arise from cross-contamination from the environment, within sequencing
340 reagents and from other samples processed or sequenced together. In several studies, viral reads
341 have subsequently been related to contamination within kits or from the environment. For
342 example, Nextera XT reagents have been associated with detection of Merkel cell
343 polyomavirus³⁰. In this study, we found a novel rhabdovirus in several patient samples, most
344 likely as a result of contaminated blood tubes with a virus of mosquito origin. This highlights the
345 need for careful interpretation of mNGS data, confirmation by other methods and the
346 development of standards and validation methods to interpret experimental information.

347

348 **CONCLUSIONS**

349 Febrile illness is widespread in Uganda and a significant proportion of undiagnosed disease can
350 be attributed to viral infections, several of which have pandemic potential and can cause high
351 mortality and morbidity. Metagenomic next generation sequencing is a powerful tool to
352 investigate the viral aetiology of AFI in high-risk populations and can be applied to develop low-
353 cost clinical diagnostics. The finding of multiple highly pathogenic and poorly-described viruses in
354 a single multi-site study in Uganda highlights the depth of the unknown with regard to circulating
355 human viruses and the risk for onward transmission in a rapidly changing world.

356 **References :**

- 357 1. Carrington LB, Wills B. Lessons from history: viral surveillance in 1940s East Africa: Epidemiological
358 notes on some viruses isolated in Uganda, GWA Dick, Transactions of the Royal Society of Tropical
359 Medicine and Hygiene, 1953; 47 (1): 13–48. Trans R Soc Trop Med Hyg. 2018;112:413–4.
- 360 2. Mayanja MN, Mwiine FN, Lutwama JJ, Ssekagiri A, Egesa M, Thomson EC, et al. Mosquito-borne
361 arboviruses in Uganda: History, transmission and burden. Journal of General Virology [Internet].
362 2021 [cited 2023 Jan 17];102:001680. Available from:
363 <https://www.microbiologyresearch.org/content/journal/jgv/10.1099/jgv.0.001680>
- 364 3. Mordecai EA, Ryan SJ, Caldwell JM, Shah MM, LaBeaud AD. Climate change could shift disease
365 burden from malaria to arboviruses in Africa. Lancet Planet Health. 2020;4:e416–23.
- 366 4. Stoler J, Awandare GA. Febrile illness diagnostics and the malaria-industrial complex: a socio-
367 environmental perspective. BMC Infect Dis. 2016;16:1–9.
- 368 5. Murray CJL, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. Disability-adjusted life
369 years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the
370 Global Burden of Disease Study 2010. The lancet. 2012;380:2197–223.
- 371 6. Prasad N, Murdoch DR, Reyburn H, Crump JA. Etiology of severe febrile illness in low-and middle-
372 income countries: a systematic review. PLoS One. 2015;10:e0127962.
- 373 7. Crump JA, Morrissey AB, Nicholson WL, Massung RF, Stoddard RA, Galloway RL, et al. Etiology of
374 severe non-malaria febrile illness in Northern Tanzania: a prospective cohort study. PLoS Negl Trop
375 Dis. 2013;7:e2324.
- 376 8. Maze MJ, Bassat Q, Feasey NA, Mandomando I, Musicha P, Crump JA. The epidemiology of febrile
377 illness in sub-Saharan Africa: implications for diagnosis and management. Clinical Microbiology and
378 Infection. 2018;24:808–14.
- 379 9. Baba M, Logue CH, Oderinde B, Abdulmaleek H, Williams J, Lewis J, et al. Evidence of arbovirus co-
380 infection in suspected febrile malaria and typhoid patients in Nigeria. The Journal of Infection in
381 Developing Countries. 2013;7:51–9.
- 382 10. Jerome H, Taylor C, Sreenu VB, Klymenko T, Filipe ADS, Jackson C, et al. Metagenomic next-
383 generation sequencing aids the diagnosis of viral infections in febrile returning travellers. Journal
384 of Infection. 2019;79:383–8.
- 385 11. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. Nat Methods.
386 2015;12:59–60.
- 387 12. Safonova Y, Bankevich A, Pevzner PA. dipSPAdes: assembler for highly polymorphic diploid
388 genomes. Journal of Computational Biology. 2015;22:528–45.
- 389 13. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective Stochastic
390 Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol [Internet]. 2015 [cited
391 2022 Aug 8];32:268–74. Available from:
392 <https://academic.oup.com/mbe/article/32/1/268/2925592>
- 393 14. Kumar S, Tamura K, Nei M. MEGA: molecular evolutionary genetics analysis software for
394 microcomputers. Bioinformatics. 1994;10:189–91.
- 395 15. Aricescu AR, Lu W, Jones EY. A time-and cost-efficient system for high-level protein production in
396 mammalian cells. Acta Crystallogr D Biol Crystallogr. 2006;62:1243–50.

- 397 16. Hofmann K, Stoffel W. TMpred, Prediction of Transmembrane Regions and Orientation. 1993.
- 398 17. Whitt MA. Generation of VSV pseudotypes using recombinant Δ G-VSV for studies on virus entry,
399 identification of entry inhibitors, and immune responses to vaccines. *J Virol Methods*.
400 2010;169:365–74.
- 401 18. Walker PJ, Firth C, Widen SG, Blasdel KR, Guzman H, Wood TG, et al. Evolution of genome size and
402 complexity in the Rhabdoviridae. *PLoS Pathog*. 2015;11:e1004664.
- 403 19. Schieffelin JS, Shaffer JG, Goba A, Gbakie M, Gire SK, Colubri A, et al. Clinical illness and outcomes
404 in patients with Ebola in Sierra Leone. *New England journal of medicine*. 2014;371:2092–100.
- 405 20. (CDC C for DC and P. Imported Lassa fever--New Jersey, 2004. *MMWR Morb Mortal Wkly Rep*.
406 2004;53:894–7.
- 407 21. Atim SA, Ashraf S, Belij-Rammerstorfer S, Ademun AR, Vudriko P, Nakayiki T, et al. Risk factors for
408 Crimean-Congo Haemorrhagic Fever (CCHF) virus exposure in farming communities in Uganda.
409 *Journal of Infection*. 2022;85:693–701.
- 410 22. Paweska JT. Rift Valley fever. In: *Emerging Infectious Diseases*. Elsevier; 2014. p. 73–93.
- 411 23. Simo FBN, Bigna JJ, Kenmoe S, Ndangang MS, Temfack E, Moundipa PF, et al. Dengue virus
412 infection in people residing in Africa: a systematic review and meta-analysis of prevalence studies.
413 *Sci Rep*. 2019;9:1–9.
- 414 24. Dieng I, Ndione MHD, Fall C, Diagne MM, Diop M, Gaye A, et al. Multifoci and multiseroypes
415 circulation of dengue virus in Senegal between 2017 and 2018. *BMC Infect Dis*. 2021;21:1–11.
- 416 25. Johansson MA, Vasconcelos PFC, Staples JE. The whole iceberg: estimating the incidence of yellow
417 fever virus infection from the number of severe cases. *Trans R Soc Trop Med Hyg [Internet]*. 2014
418 [cited 2023 Jan 23];108:482–7. Available from:
419 <https://academic.oup.com/trstmh/article/108/8/482/2765182>
- 420 26. Stremlau MH, Andersen KG, Folarin OA, Grove JN, Odi A, Ehiane PE, et al. Discovery of novel
421 rhabdoviruses in the blood of healthy individuals from West Africa. *PLoS Negl Trop Dis*.
422 2015;9:e0003631.
- 423 27. Grard G, Fair J, Chiu C, Leroy E. Bas-Congo virus: A novel rhabdovirus associated with acute
424 hemorrhagic fever. In: *Emerging infectious diseases*. Elsevier; 2014. p. 13–24.
- 425 28. Woodruff AW, Ansdell VE, Bowen ET. Le Dantec virus infection in a patient who had not been to
426 West Africa. *Br Med J*. 1977;2:1632.
- 427 29. Wichmann O, Hellenbrand W, Sagebiel D, Santibanez S, Ahlemeyer G, Vogt G, et al. Large measles
428 outbreak at a German public school, 2006. *Pediatric Infectious Disease Journal [Internet]*. 2007
429 [cited 2023 Jan 17];26:782–6. Available from:
430 https://journals.lww.com/pidj/Fulltext/2007/09000/Large_Measles_Outbreak_at_a_German_Public_School,.3.aspx
431
- 432 30. Asplund M, Kjartansdóttir KR, Mollerup S, Vinner L, Fridholm H, Herrera JAR, et al. Contaminating
433 viral sequences in high-throughput sequencing viromics: a linkage study of 700 sequencing
434 libraries. *Clinical Microbiology and Infection*. 2019;25:1277–85.

435

1 **Emerging viruses are an underestimated cause of undiagnosed febrile illness in Uganda**


2

3 Authors

4 Shirin Ashraf*¹, Hanna Jerome*¹, Daniel Lule Bugembe³, Deogratius Ssemwanga^{2,3}, Timothy
5 Byaruhanga², John Timothy Kayiwa², Robert Downing², Jesus F. Salazar-Gonzalez³, Maria G.
6 Salazar³, James G. Shepherd¹, Chris Davis¹, Nicola Logan¹, Vattipally B. Sreenu¹, Gavin S. Wilkie¹,
7 Ana da Silva Filipe¹, Alfred Ssekagiri^{1,2}, Prossy Namuwulya², Henry Bukenya², Brian Kigozi²,
8 Weronika Witkowska McConnell¹, Brian J. Willett¹, Stephen Balinandi², Julius Lutwama²,
9 Pontiano Kaleebu^{2,3}, Josephine Bwogi², Emma C. Thomson^{1,4}

10

11 *Contributed equally

12  Contributed equally

13 Institutions

14 1 MRC-University of Glasgow Centre for Virus Research

15 2 Uganda Virus Research Institute


16 3 MRC/UVRI & LSHTM Uganda Research Unit

17 4 LSHTM


18

19


20 **Figure 1. Undiagnosed viruses identified in the AFI study.** (a) AFI study sites and locations.
 21 Viruses identified from undiagnosed samples are listed by site. (b) Viruses detected by mNGS
 22 grouped as zoonoses/arboviruses (green), hepatitis (yellow), blood-borne infections (red),
 23 respiratory viruses (blue), gastrointestinal infections (brown), viruses associated with rash
 24 (purple) and uncharacterized (grey). Icons show the number of patients infected with each virus.
 25



26 **Figure 2. Viruses associated with haemorrhagic fever.** (a-c) Coverage plots for CCHFV, YFV,
 27 DENV identified by mNGS showing number of reads (y-axis) at each nucleotide position across
 28 the genome (x-axis) for each virus. Reference orthonairovirus and flavivirus genomes indicate
 29 regions covered by sequencing reads. (d) Phylogenetic tree of full-length nucleotide sequence of
 30 CCHFV showing placement of patient isolate for L, M, S segments. Tip labels show strain name
 31 and country of isolation. Colours indicate geographically distributed lineages of CCHFV. (e) Full
 32 genome nucleotide tree of DENV genotype 3 sequences showing country and year of isolation. (f)
 33 Full genome nucleotide tree of YFV sequences showing country and year of isolation.
 34 Evolutionary scale is showing below for each tree. Ultrafast bootstrap values >90 are shown for
 35 nodes. Patient samples are highlighted in grey. Substitution models used: GTR+F+I+G4 (CCHF-L,
 36 DENV-3), GTR+F+R3 (CCHF-M), TIM2+F+I+G4 (CCHF-S), GTR+F+R2 (YFV).
 37




38 **Figure 3. Le Dantec virus (LDV).** (a) Coverage plots for LDV identified by mNGS showing number
 39 of reads (y-axis) at each nucleotide position across the genome (x-axis) for each virus.
 40 Sequencing depth for different genes across the genome is shown. (b) ELISA analysis showing IgG
 41 antibody response to recombinant LDV glycoprotein in patient 220-2, contact of patient and a
 42 healthy individual from Uganda. Plasmid containing an Ephrin B2 insert was used as a mock
 43 recombinant protein. The plot shows absorbance OD values at 450nm (y-axis) against increasing
 44 serum dilutions (x-axis). (c) Pseudotype neutralisation assay showing luciferase reporter activity
 45 (y-axis) against increasing serum dilutions (x-axis) for patient 220-2, contact and healthy serum
 46 using LeDantec pseudotype virus compared with VSV control. (d) Full genome maximum
 47 likelihood phylogenetic nucleotide tree using the GTR+F+R4 model for ledantevirus sequences
 48 illustrating the reported host for each virus. Evolutionary scale and ultrafast bootstrap values >90
 49 are shown. The patient sample is highlighted in grey.



51

52

53 **Figure 4. Other viral pathogens identified from patient samples.** Coverage plots showing
 54 number of mapped reads (y-axis) against genome position (x-axis) for each patient, for (a) HAV,
 55 (b) HEV, (c) Enteroviruses including Echoviruses E3, E9/25, E4 and Rhinovirus C, (d) Measles virus;
 56 and (e) HIV-1. Reference genome drawings indicate depth for different regions along the
 57 sequence.



58

59

60

1 **Emerging viruses are an underestimated cause of undiagnosed febrile illness in Uganda**


2

3 Authors

4 Shirin Ashraf*¹, Hanna Jerome*¹, Daniel Lule Bugembe³, Deogratius Ssemwanga^{2,3}, Timothy
5 Byaruhanga², John Timothy Kayiwa², Robert Downing², Jesus F. Salazar-Gonzalez³, Maria G.
6 Salazar³, James G. Shepherd¹, Chris Davis¹, Nicola Logan¹, Vattipally B. Sreenu¹, Gavin S. Wilkie¹,
7 Ana da Silva Filipe¹, Alfred Ssekagiri^{1,2}, Prossy Namuwulya², Henry Bukenya², Brian Kigozi²,
8 Weronika Witkowska McConnell¹, Brian J. Willett¹, Stephen Balinandi², Julius Lutwama²,
9 Pontiano Kaleebu^{2,3}, Josephine Bwogi², Emma C. Thomson^{1,4}

10

11 *Contributed equally

12  Contributed equally

13 Institutions

14 1 MRC-University of Glasgow Centre for Virus Research

15 2 Uganda Virus Research Institute

16 3 MRC/UVRI & LSHTM Uganda Research Unit

17 4 LSHTM

18

19

20 **Table 1. Demographics and clinical characteristics of patients diagnosed with viral infections**
21

n	30
Sex = Male/Female (%)	20/10 (66.67/33.33)
Age (median [IQR])	7.50 [3.75, 13.5]
Age bracket in years (%)	
0 - 5	12 (40)
6 - 15	12 (40)
16 - 25	2 (6.67)
26 - 35	2 (6.67)
36 - 45	2 (6.67)
Study Clinic (%)	
St Paul's HC	9 (30.00)
Ndejje HC	18 (60.00)
Adumi HC	3 (10.00)
Clinical features	
BaselineTemperature (mean (SD))	38.21 (0.85)
Fever_duration (mean (SD))	3.6 (1.55)
Rigors (%)	7 (23.3)
Headache (%)	19 (63.3)
Neck_pain (%)	6 (20)
Jaundice (%)	3 (10)
Nausea (%)	11 (36.7)
Vomiting (%)	12 (40)
Abdominal_pain (%)	13 (43.3)
Diarrhoea (%)	9 (30)
Myalgia (%)	6 (20)
Arthralgia (%)	9 (30)
Back_pain (%)	6 (20)
Haemorrhage_any (%)	3 (10)
Haemoptysis (%)	1 (3.3)
Haemochezia (%)	2 (6.7)
Contact with ectoparasites (%)	23 (76.67)
Mosquitoes (%)	22 (73.33)
Ticks (%)	3 (10.00)
Fleas (%)	2 (6.67)
Lice (%)	1 (3.33)
Biting flies (%)	3 (10.00)
Health Centre Diagnosis (%)	
Gastroenteritis	1 (3.3)
Malaria	3 (10)
Respiratory tract infection	9 (30)
Typhoid	17 (56.6)
Unknown	23 (76.7)
Antibacterial therapy (%)	12 (40)
Antimalarial therapy (%)	5 (16.7)
Anthelmintic therapy (%)	1 (3.3)

22

23 **Table 2. Viruses detected by mNGS in undiagnosed patients in the AFI study**

24

Virus	No. of patients	Nearest reference accession(s)	Country of nearest reference
Viruses associated with haemorrhagic fever			
CCHFV	1	MW464976 (L), MW464977 (M), DQ211650 (S)	Uganda (L and M) and DRC (S)
RVFV	2	MG953422, MG273456	Uganda, Kenya
YFV	2	AY968065, JN620362	Uganda
DENV-3	1	MN964273	China ex Ethiopia
Viruses associated with acute hepatitis			
HAV	5	MH685714	Uganda
HEV	2	MH918640	Nigeria
Blood-borne viruses			
HBV	1	MK512473	Rwanda
HIV-1	6	MW006081, MW006060, AY322184, EU110093	Uganda, Kenya
Respiratory viruses			
Measles B3	6	ON642798, ON642800, ON642797, ON642801	Uganda
Respirovirus 3	2	MN145875, MF973181	China, USA
Rhinovirus C	1	JF317017	China
Viruses associated with gastroenteritis			
Rotavirus H-like virus	1	KX362515	Vietnam
Echovirus E3, E4 and E9/25	3	MF678296, KM024043, MF554740	Australia (E25), Greece (E3), USA (E4)
Coxsackievirus B3	1	KY262575	China
Herpesviruses associated with rash and fever			
HHV6B	3	KY315537, MF511176, MF994829	USA, Uganda, DRC
Rarely described viruses associated with acute febrile illness			
LDV	1	NC_034443	Senegal

25

