

1 **Serendipitous detection of *Anopheles stephensi* in Kisumu, Kenya in June 2022**

2
3 Bryson Alberto Ndenga, Sammy Wambua, Kevin Omondi Owuor, Rodney Omukuti, Salome
4 Chemutai, Daniel Arabu, Irene Miringu, Carren Bosire, Kavinya Mwendwa, Christabel Achieng
5 Winter, Francis Maluki Mutuku, Donal Bisanzio, Angelle Desiree LaBeaud, Keli Nicole Gerken

6
7 Author affiliations: Kenya Medical Research Institute, Kisumu, Kenya (B.A. Ndenga, K.O.
8 Owuor, C.A. Winter); Pwani University, Kilifi, Kenya; Research and Conservation Support
9 Sciences, Kilifi, Kenya; and University of Glasgow, Glasgow, UK (S. Wambua); Pwani
10 University, Kilifi, Kenya, and Research and Conservation Support Sciences, Kilifi, Kenya (R.
11 Omukuti, S. Chemutai, D. Arabu, I. Miringu); Technical University of Mombasa, Mombasa,
12 Kenya (C. Bosire, K. Mwendwa, F.M. Mutuku); RTI International, Washington, DC, USA (D.
13 Bisanzio); Stanford University School of Medicine, Stanford, California, USA (A.D. LaBeaud,
14 K.N. Gerken).

15
16 **Abstract**

17 In June 2022, a pool of five mosquitoes that were morphologically classified as *Anopheles*
18 *gambiae* and caught in Kisumu (Kenya) were tested for blood-meal analysis. Of the 19.6%
19 (11/56) amplicon sequence variants assigned to mosquito species using basic local alignment
20 search tool (BLAST), one had 15 hits matching *Anopheles stephensi*.

21 **Research letter**

22 *Anopheles stephensi* is a native urban malaria vector in south Asia and the Arabian
23 Peninsula. It is capable of transmitting both *Plasmodium falciparum* and *P. vivax* and is

24 projected to expand further as urbanization increases, especially in Africa (1). It was first
25 reported in Africa in Djibouti in 2012 (2) then in Ethiopia in 2019 (3), Sudan in 2018 (4),
26 Somali-land in 2020 (5), Nigeria in 2020 (6) and for the first time in Marsabit County in northern
27 Kenya in December 2022 (7). This demonstrates a clear and rapid expansion of its geographical
28 range within the continent.

29 During a One Health study on livestock and vector dynamics at urban slaughterhouses,
30 we carried out mosquito collections using ovitraps, larval/pupae sampling, Prokopack aspirator
31 and Biogents (BG)-trap baited with CO₂. This study was titled “Urban Rift Valley Fever Virus as
32 a New Ecological Niche: Continuous Introduction from Animal Products” and performed at the
33 Mamboleo Slaughterhouse (0° 3' 25.4" N, 34° 47' 10.9" E, Figure) in Kisumu City, Kenya, and
34 was approved by Stanford University (61386), Kenya Medical Research Institute Scientific and
35 Ethics Review Unit (KEMRI/SERU/CGHR/03-07-390/4293) and obtained research license from
36 the National Commission for Science, Technology & Innovation (NACOSTI/P/21/13557).

37 Adult mosquitoes resting indoors and outdoors were sampled using a Prokopack aspirator
38 from May 16th 2022 to July 1st 2022. Sampling lasted one hour and was conducted between
39 08:45 HRS and 01:30 HRS. Subsequent samplings were completed biweekly. Collected
40 mosquitoes were killed using a pyrethrum aerosol spray before morphological identification in a
41 field station and recorded on a data form. Female mosquitoes were further sorted according to
42 their blood-feeding stages as unfed, blood-fed, half-gravid or gravid. Mosquitoes were preserved
43 in silica gel self-indicating 6-20mesh (Blue) - 500gm (Loba Chemie Pvt. Ltd, Jehangir Villa,
44 107, Wodehouse Road, Colaba, Mumbai 400 005, India) with the intention for further blood
45 meal testing.

46 A total of 2,891 mosquitoes were collected: 66.2% (1,782) indoors and 33.8% (765)
47 outdoors; 50.7% (1,465) males and 49.3% (1,426) females. Species identified morphologically
48 were: 0.3% (9) *Aedes aegypti*; 6.3% (182) *Anopheles coustani*; 0.3% (8) *An. funestus*; 10.9%
49 (315) *An. gambiae*, and 82.2% (2,377) *Culex* spp. Over a half, 61.6% (194/315), of the *An.*
50 *gambiae* were collected outdoors and 38.4% (121) indoors; 46.3% (146) were males; 19.0% (60)
51 un-blood-fed; 25.1% (79) blood-fed; 6.0% (19) half-gravid and 3.5% (11) gravid. A sample of
52 the blood-fed mosquitoes were transported on dry ice to Pwani University Biosciences Research
53 Centre (PUBReC) in Kilifi, Kenya, for determination of blood meal sources.

54 Genomic DNA was extracted from the mosquito abdomens using the TIANamp Genomic
55 DNA Kit (Tiangen, Beijing, China). Quality and quantity of DNA was analyzed by NanoDrop
56 2000C spectrophotometer (Thermo Scientific Inc, USA) and verified on 1% agarose gel
57 electrophoresis. To identify blood meal sources, we PCR-amplified ~300 base pair (bp) of the
58 cytochrome b barcode (8) which was high-throughput-sequenced, at Macrogen Inc., Seoul
59 Korea, using the Illumina 300 × 2 bp platform (Illumina, USA). The amplicon sequence data
60 were analyzed with the DADA2 (version 1.21.0) bioinformatics pipeline (Callahan et al., 2016)
61 implemented in the R programming language (version 4.1.1). After quality preprocessing, high
62 quality reads were denoised and clustered into unique sequences i.e., amplicon sequence variants
63 (ASVs). We assigned taxonomy to the ASVs using two approaches. We assigned taxonomy,
64 first, by exact ASV matching against DADA2-trained MIDORI2 reference database (version
65 GB254) which includes eukaryotic mitochondrial sequences (9). ASVs that were not assigned by
66 this approach were compared against the NCBI database using BLAST (10). Raw sequences are
67 available in SRA database of NCBI under project accession number PRJNA966766.

68 In all, 248 ASVs were inferred, 192 of which were classified to 14 vertebrate species
69 (Mammalia and Aves) by exact matching against MIDORI2 reference database. Other than
70 vertebrates, 19.6% (11/56) of the ASVs assigned using BLAST matched to mosquito species one
71 of which was *An. stephensi* with a total of 15 hits (89-93% identity, 67-85% coverage, E-value <
72 1e-46 (Table). The reads matching *An. stephensi* were tracked to a pool of five (5) mosquitoes
73 that had been morphologically classified as *An. gambiae* in the field. These mosquitoes were
74 collected from Mamboleo Slaughterhouse in Kisumu City, Kenya in June 2022.

75 Our findings report the first detection of *An. stephensi* south of the Equator. Furthermore,
76 this detection represents the presence of an invasive vector in an urban area that is malaria
77 endemic. Our study suggests that the detections of *An. stephensi* within Africa (1-5, 7) could be
78 dramatically underreported. The initial false identification of this vector also indicates that to
79 fully capture this vector's boundary, molecular methods and confirmatory testing may be
80 required. The presence of a malaria vector adapted to breeding in open water containers
81 prevalent in urban areas is particularly concerning. *Anopheles stephensi* can effectively transmit
82 *P. falciparum* and *P. vivax* and entry into malaria endemic zones could shift malaria transmission
83 dynamics considerably.

84

85 **Acknowledgements**

86 We thank Charles Otieno Adipo for sampling the vectors, the management of Mamboleo
87 Slaughterhouse for allowing us to sample mosquitoes in the premises and our colleagues in
88 KEMRI and Stanford University for their support to approve this research study. This study was
89 funded by the Stanford University Center for Innovations in Global Health Seed Grant 2021 and

90 the Robert E Shope ASTMH (American Society of Tropical Medicine and Hygiene) to Dr. Keli
91 Nicole Gerken.

92

93 **About the Author**

94 Bryson Alberto Ndenga, PhD is a Senior Research Scientist in Kenya Medical Research
95 Institute, Centre for Global Health Research. His primary interest is in the biology, ecology and
96 control of mosquitoes that transmit malaria, dengue, chikungunya and Rift Valley fever.

97

98 **Financial interest**

99 The authors declare that they have no financial interests.

100

101 **Conflicts of interest**

102 The authors declare that they have no conflicts of interests.

103


104 **References**

- 105 1. Sinka M, Pironon S, Massey N, Longbottom J, Hemingway J, Moyes C, et al. A new
106 malaria vector in Africa: Predicting the expansion range of *Anopheles stephensi* and
107 identifying the urban populations at risk. PNAS. 2020;117.
- 108 2. Faulde M, Rueda L, Khaireh B. First record of the Asian malaria vector *Anopheles*
109 *stephensi* and its possible role in the resurgence of malaria in Djibouti, Horn of Africa.
110 Acta Trop. 2014;139:39-43.

- 111 3. Tadesse F, Ashine T, Teka H, Esayas E, Messenger L, Chali W, et al. *Anopheles*
112 *stephensi* mosquitoes as vectors of *Plasmodium vivax* and *P. falciparum*, horn of Africa,
113 2019. Emerg Infect Dis. 2021;27.
- 114 4. Ahmed A, Khogali R, Elnour M, Nakao R, Salim B. Emergence of the invasive malaria
115 vector *Anopheles stephensi* in Khartoum State, Central Sudan. Parasit Vectors. 2021;14.
- 116 5. Ali S, Samake J, Spear J, Carter T. Morphological identification and genetic
117 characterization of *Anopheles stephensi* in Somaliland. Parasit Vectors. 2022;15.
- 118 6. World Health Organization initiative to stop the spread of *Anopheles stephensi* in Africa.
119 Geneva: WHO; 2022.
- 120 7. Ochomo E, Milanoi S, Abong'o B, Onyango B, Muchoki M, Omoke D, et al. Molecular
121 surveillance leads to the first detection of *Anopheles stephensi* in Kenya. Res Square. In
122 press 2023.
- 123 8. Hadj-Henni L, De-Meulemeester T, Depaquit J, Noël P, Germain A, Helder R, et al.
124 Comparison of vertebrate cytochrome b and prepronociceptin for blood meal analyses in
125 Culicoides. Front Vet Sci. 2015;2.
- 126 9. Leray M, Knowlton N, Machida R. MIDORI2: A collection of quality controlled,
127 preformatted, and regularly updated reference databases for taxonomic assignment of
128 eukaryotic mitochondrial sequences. Environ DNA. 2022;4:894–907.
- 129 10. Altschul S, Gish W, Miller W, Myers E, Lipman D. Basic local alignment search tool. J
130 Mol Bio. 1990;215:403–410.

131

132 Address for correspondence: Dr. Bryson Alberto Ndenga, Kenya Medical Research Institute,
 133 Centre for Global Health Research, P.O. Box 1578 Code 40100 Kisumu City, Kenya, email:
 134 bndenga@yahoo.com, phone number: +254 722 932 350.



135
 136 **Figure 1.** Map showing the location in Kisumu, Kenya where *Anopheles stephensi* (green
 137 triangle) was collected in June 2022 and other sites in Africa where it has been reported (Source:
 138 *Anopheles stephensi* reports (red points): <https://apps.who.int/malaria/maps/threats/>). Suitability
 139 map created using data obtained from Sinka et al. 2020¹.

140
 141 **Table 1.** Summary of coverage and identity values for each of the top ten (10) hits in the BLAST
 142 search matching *Anopheles stephensi*.

143

Description	Maximum score	Query coverage	E value	Identity	Accession
<i>Anopheles stephensi</i> strain Indian chromosome 2L	326	84%	2.00E-84	92.51%	CP032300.1

<i>Anopheles stephensi</i> strain Indian chromosome 3L	320	85%	8.00E-83	92.07%	CP032302.1
<i>Anopheles stephensi</i> strain Indian chromosome 3R	320	82%	8.00E-83	92.07%	CP032301.1
<i>Anopheles stephensi</i> strain Indian chromosome 2R	315	81%	4.00E-81	91.63%	CP032299.1
<i>Anopheles stephensi</i> strain SDA-500 chromosome 3R	315	83%	4.00E-81	91.63%	CP032234.1
<i>Anopheles stephensi</i> strain SDA-500 chromosome 2R	309	81%	2.00E-79	91.19%	CP032232.1
<i>Anopheles stephensi</i> strain SDA-500 chromosome 2L	303	84%	9.00E-78	90.75%	CP032233.1
<i>Anopheles stephensi</i> strain SDA-500 chromosome 3L	298	88%	4.00E-76	90.31%	CP032235.1
<i>Anopheles ziemanni</i> genome assembly, chromosome: 2	291	83%	7.00E-74	89.52%	OX030919.2

144

145