

Strain measures of the left ventricle and left atrium are composite measures of left heart geometry and function

Fredrika Fröjd¹, Dhnanjay Soundappan², Peder Sörensson¹, Andreas Sigfridsson¹,

Eva Maret¹, Jannike Nickander^{1*}, Martin Ugander^{1,2*}

* equal contribution

1. Department of Clinical Physiology, Karolinska University Hospital, and Karolinska Institutet, Stockholm, Sweden
2. Kolling Institute, Royal North Shore Hospital, and University of Sydney, Sydney, Australia

Abstract

Background: Left ventricular (LV) global longitudinal strain (GLS) and circumferential strain (GCS), and left atrial (LA) strain (LAS) are indicators of poor clinical prognosis. However, it is unclear how they relate to each other and to LV and LA geometry. The aim was to clarify these relationships to inform clinical and research applications. **Methods:** Patients referred for cardiovascular magnetic resonance imaging were retrospectively identified. Univariable and multivariable linear regression models evaluated associations between GLS, GCS, LAS, LV mass, the volumes and dimensions of the LV and LA, and mitral annular plane systolic excursion (MAPSE). **Results:** In patients (n=66, median [interquartile range] age 62 [53–72] years, 82% male, LV ejection fraction 48 [34–56]%, range 6–69%), GLS associated with both GCS ($R^2=0.86$, $p<0.001$) and LAS ($R^2=0.51$, $p<0.001$), and LAS associated with GCS ($R^2=0.42$, $p<0.001$). GLS, GCS, and LAS were all univariably associated with MAPSE, LV mass, and the volumes and dimensions of the LV and LA ($p<0.001$ for all). In multivariable analysis, GLS associated with MAPSE and LV length ($R^2=0.85$, $p<0.001$); GCS with MAPSE, LV end-systolic volume, and LV mass ($R^2=0.80$, $p<0.001$); and LAS with LA end-diastolic volume and MAPSE ($R^2=0.67$, $p<0.001$). **Conclusions:** MAPSE and LV length alone can accurately estimate GLS. GLS and GCS provide similar information. LV and LA strains can be understood as geometrically coupled composite measures of MAPSE, and the size, function, and dimensions of the LV and LA. The composite of these geometrical relationships likely explains the excellent prognostic strength of strain measures.

Key words: strain, MAPSE, left ventricular function, left ventricular size

Introduction

Strain of the left ventricle (LV), measured as global longitudinal strain (GLS) or global circumferential strain (GCS), and strain of the left atrium (LA), measured as left atrial strain (LAS), are increasingly being used for assessment of LV function and for patient prognosis. GLS and GCS have been identified as more sensitive¹ and reproducible² parameters of LV systolic function compared to LV ejection fraction (LVEF), and can provide incremental prognostic value beyond LVEF³. While LVEF is used to characterize heart failure (HF) with reduced LVEF (HFrEF) and HF with preserved LVEF (HFpEF), LV strain has been shown to be a more accurate measure for detecting the presence of LV dysfunction and evaluating disease progression over time^{4,5}.

In patients with HFpEF, strain is often impaired⁶. Mathematical modelling has been used to describe how GLS, GCS, LV wall thickness, and LV end-diastolic volume associate with LVEF⁷. Those authors concluded that changes to LV end-diastolic volume and mass explain how LVEF may be preserved despite a reduced GLS. Moreover, LAS, also referred to as LA reservoir strain, has emerged as a promising measure in the assessment of LV diastolic dysfunction. LAS has been shown to correlate more closely with invasive measurements compared to conventional diastolic measures including e' , E/e' , E/A , and left atrial volume index^{8,9}. Notably, the LV apex and the posterior aspect of the LA remain relatively stationary throughout the cardiac cycle. Consequently, LV and LA strain measures are inherently coupled through the atrioventricular valve plane displacement, often referred to as mitral annular plane systolic excursion (MAPSE). Not surprisingly, GLS and LAS have been shown to be closely correlated, and GLS and LA volume alone have been shown to provide an accurate estimate of LAS¹⁰.

Strain analysis is performed by tracking the myocardial deformation over the course of the cardiac cycle using dedicated state-of-the-art image analysis software. The use of such dedicated software could add perceived complexity to a parameter that is relatively straightforward. Mathematically speaking, strain is no more than a fractional change in length for what is effectively a mid-wall line through the myocardium, which is measured in either the longitudinal (GLS, LAS) or circumferential (GCS) direction. However, an understanding of the geometric relationships associated with these strain measures might provide insight into the clinical interpretation of these measures. Therefore, the aim of this study was to empirically clarify how LV and LA strain measures relate to conventional measures of LV and LA dimensions and function and to each other, in order to inform the understanding and interpretation of LV and LA strain measures in clinical and research applications. The hypothesis was that strain measures are highly correlated with each other and with conventional LV measures, and that these measures can be used to approximate strain measures with high accuracy.

Methods

Population

We retrospectively screened patients that had undergone clinically indicated cardiovascular magnetic resonance imaging (CMR). The study was approved by the local human subject research ethics committee and all participants provided written informed consent. The population was selected to represent a variable spectrum of LVEF, including 20-25 patients from each of the following ranges: LVEF<30%, 30-40%, 40-50%, 50-60%, and >60%, while excluding those with distinct myocardial pathologies such as hypertrophic cardiomyopathy, amyloidosis, hemosiderosis, Anderson-Fabry's disease, congenital heart defects, atrial fibrillation during CMR, or insufficient image quality to perform LV strain

analysis. Further exclusion criteria included inadequate tracking of the myocardium, defined as visually inadequate tracking of >2 segments in any of the long-axis or short-axis views, and visually apparent foreshortening or the presence of ≥ 3 of the atrial appendage and pulmonary veins in both 2- and 4-chamber views.

CMR imaging

Clinical CMR scans were performed at 1.5T or 3T MAGNETOM (Aera or Skyra, Siemens Healthcare, Erlangen, Germany) with an 18-channel and 32-channel phased-array body and spine coil, respectively. The exam included retrospectively ECG-gated balanced steady-state free precession cine imaging in short axis (8 mm slice thickness, 1.6 mm slice gap) and long axis 2-, 3- and 4-chamber views (8 mm slice thickness) with 30 phases per cardiac cycle.

Post-processing analysis

All post-processing analyses were performed using freely available software Segment version 3.0 R9405e (<http://segment.heiberg.se>). LV volumes were assessed by manual delineation of the endo- and epicardium according to guidelines (Schulz-Menger et al., 2020), excluding papillary muscles for LV mass assessment. LV GLS, LV GCS were separately analyzed using a semi-automated feature-tracking module using cine images by a single observer (FF) blinded to clinical data and geometric measures. Endo- and epicardial borders were manually traced in end-diastole and automatically propagated throughout the cardiac cycle. LV GLS was measured as the mean longitudinal strain in two-, three- and four-chamber views, and LV GCS as the mean circumferential strain in an apical, midventricular, and basal short-axis view. Delineations were adjusted manually if ≥ 2 segments did not track myocardial motion adequately as assessed visually.

LA dimensions and volumes were measured using the bi-plane area-length method in the two- and four-chamber views by a blinded single observer (DS), excluding the left atrial appendage and pulmonary veins. LAS was assessed as the strain measured from end diastole to end systole in a two- or four-chamber view or the mean of both. LV and LA measurements were performed by two different observers.

MAPSE was measured as the mean distance in millimeters traveled by the mitral annular insertion points from end-diastole to end-systole as the average of two manual caliper measurements per view in all three long-axis views. LV length (LVL) was measured as the distance from the most apical point of the epicardium to the midpoint between the mitral annular insertion points in end-diastole. The longitudinal contribution to LV stroke volume was calculated as the maximal LV short-axis epicardial area multiplied by MAPSE as previously described and validated (Carlsson M, et al., 2007).

Statistical analysis

Statistical analysis was performed using RStudio Team (2022). Patient characteristics were summarized as the median [interquartile range] for continuous variables and as counts and percentages for categorical variables. Univariable linear regression was used to assess correlations. Multivariable linear regression was used to identify measures associated with GLS, GCS, and LAS using models with variables selected using variable inflation factors (VIF) <5 to reduce the effects of multicollinearity. Agreements between GCS and GLS were assessed using Bland-Altman plots. A p-value less than 0.05 was considered statistically significant.

Results

Patient characteristics

A flowchart of patient selection is shown in Figure 1. The final study population included 66 patients: 14 (18%) were female, age was 62 [53–72] years and LVEF was 48 [34–57]%, full range 9–69%. Characteristics of the study population are shown in Table 1.

Global longitudinal strain

A summary of the relationships between strain values and geometric measures is illustrative in representative patients in Figure 2. GLS was -15.5% [-17.2 – -11.6] in the study population. GLS was highly correlated with GCS ($R^2=0.86$, $p<0.001$). GLS was also highly correlated with conventional LV metrics, including LV mass, EDV, ESV, LV diameter, MAPSE, and LV length, but not with BSA or age. In multivariable regression analysis using GLS as the dependent variable, MAPSE, LV mass, and LV ESV together contributed to the model with the strongest association ($R^2=0.89$, $p<0.001$). MAPSE modeled together with other parameters of LV size yielded models of similar strength of association (MAPSE and LVEDV, $R^2=0.87$, $p<0.001$; MAPSE and LV length, $R^2=0.85$, $p<0.001$; MAPSE and LV maximal diameter, $R^2=0.85$, $p<0.001$). In these multivariable models, the association with LV mass did not remain statistically significant. LVEDV and LVM were correlated ($R^2=0.51$, $p<0.001$). Results from univariable and multivariable linear regression analysis are presented in Table 2. GLS was highly associated with MAPSE indexed to LV length ($R^2=0.83$, $p<0.001$) and with MAPSE indexed to the sum of LV length and LV diameter ($R^2=0.84$, $p<0.001$). GLS could be estimated with high accuracy from MAPSE indexed to LV length. In our series of patients, the linear relationship between MAPSE/LVL could be expressed using the following formula:

$$GLS = \frac{MAPSE \times -100}{LVL} - 3.96$$

Global circumferential strain

GCS was -16.2% [-18.75 – -11.6] in the study population. Similarly to GLS, GCS was highly correlated with LV mass, stroke volume, LVEDV, LVESV, MAPSE, LV length, LVEF, and LV diameter, but not with BSA (Table 3). In multivariable analyses, LVESV and MAPSE together produced the best model for estimating GCS.

Left atrial strain

LAS was 24 [20–29] % in the study population. Results from the uni- and multivariable analysis of LAS, LA, and LV parameters are summarized in Table 4. LAS was correlated with GLS ($R^2=0.51$, $p<0.001$). LAEDV and MAPSE together associated with LAS ($R^2=0.65$, $p<0.001$). MAPSE was correlated with LAS ($R^2=0.46$, $p<0.001$), and this correlation increased when MAPSE was indexed to LA length ($R^2=0.51$, $p<0.001$). Measured LAS and LAS calculated according to a previously published model defined as LA volume/LV volume times GLS associated with LAS ($R^2=0.31$, $p<0.001$).

Longitudinal and circumferential contribution to left ventricular stroke volume

The longitudinal contribution to LV stroke volume was 62 [58 – 66]% in the population. The longitudinal contribution to LV stroke volume did not associate with GLS ($p=0.68$) or MAPSE indexed to LV length ($p=0.12$). Similarly, GCS was not associated with the short-axis contribution to LV stroke volume ($p=0.92$).

Discussion

The main findings of the study are that LV strain is a composite measure of both LV function and geometry that is closely associated with MAPSE and LV dimensions, and LA strain is similarly closely associated with MAPSE and LA volume. All strain measures are closely correlated with each other. Insight into and consideration of these associations between strain measures and conventional geometric measures of LV and LA size and function can inform clinical interpretation.

GLS and MAPSE were closely correlated, and the magnitude of the association increased further when MAPSE was indexed to LV length, to LV maximal diameter, or both, reflecting the dependency of GLS on both the atrioventricular plane displacement and LV size. A simple formula comprised of MAPSE and LV length could be used in our sample to estimate GLS with high accuracy. Since strain measures are subject to sizeable inter-vendor variability¹², estimating GLS using MAPSE indexed to LV length provides a potential vendor-independent and accurate estimate of GLS. In line with these findings, MAPSE indexed to LV length has previously been suggested as a surrogate measure for GLS in the case of limited image quality¹³, and was recently found to provide similar prognostic accuracy to GLS¹⁴.

LV mass was correlated with GLS, but the association remained significant only in a multivariable model using LVESV ($p=0.04$), and not in the other multivariable models. This may be explained by a high correlation between LVM and LV size (EDV and LVL). In our study we excluded patients with HCM.

GCS was also highly correlated with MAPSE and, similarly to GLS, indexing MAPSE to LV length or LV diameter further increased the strength of this correlation, which indicates that

GLS and GCS both provide similar information. Indeed, GLS and GCS were also highly correlated with each other. While the direct correlation between GLS and GCS has not previously been reported, both GLS and GCS have been found to be highly linearly correlated with LVEF, indirectly indicating a high correlation (Onishi et al., 2015). Beyond both being measures of myocardial shortening, GCS and GLS are physically coupled through the endocardial inward movement that results from longitudinal shortening, comparable to the coupling of MAPSE to fractional shortening (Carlsson 2007). GLS is further mechanistically coupled to GCS through the radial inward movement resulting from circumferential shortening.

The median longitudinal contribution to LV stroke volume in our study was 62%, and this was comparable to findings in previous studies showing that longitudinal function is similar across a broad range of values for LVEF^{16 17}. MAPSE or GLS were not associated with the longitudinal contribution to stroke volume and GCS was similarly not correlated with the short-axis contribution to stroke volume. Consequently, while MAPSE and GLS are measures of longitudinal shortening, they are not necessarily appropriate to describe longitudinal function when defined as the longitudinal contribution to stroke volume.

LAS was, similarly to LV strain, highly correlated with both MAPSE and LA dimensions. Consequently, LAS also associated with GLS, in agreement with previous findings^{10,18,19}, as well as with GCS. A previous study found that LAS could be calculated from GLS by correcting for the LA to LV volume ratio. Similarly, our results suggest that LAS is mainly determined by LA size and LV function, although the correlation between LAS and LAS calculated from GLS and the LA/LV volume ratio was not as strong in our study as compared to previously presented models ($R^2=0.31$ vs 0.65)¹⁰.

LAS has been found to identify patients with increased filling pressures with incremental accuracy compared to conventional echocardiographic diastolic parameters^{18,20,21}. LA enlargement is a known consequence of longstanding elevated filling pressures, and MAPSE has been found to be correlated with echocardiographic measures of diastolic dysfunction (Willenheimer et al., 1999). It is also known that diastolic and systolic dysfunction often co-occur. Naturally, as the myocardial shortening during systole must be equal to the myocardial lengthening during diastole, MAPSE must, mathematically speaking, be a composite measure of both. These associations may consequently explain the correlation between LAS and diastolic dysfunction. However, the majority of studies comparing LAS to invasive pressure measurements have omitted LV functional indices from performance analyses. Moreover, one study found that LAS only associated with diastolic dysfunction among patients with reduced GLS ($\leq 18\%$)²³, supporting the notion that GLS is closely associated with LAS. However, one study did include GLS and found LAS provided prognostic value independently of GLS and LAVi in a population with reduced LVEF¹⁹. Notably, that study did not consider that the correlation between GLS and LAS is biased by the relative sizes of the LV and the LA¹⁰.

Limitations

Female patients were underrepresented in our study, and this may be seen as a limitation. However, while females are known to have smaller hearts than males on a group level, there is no indication that female and male hearts of the same size exhibit different proportions or geometric scaling. Hence, the results should be reasonably applicable across both sexes. Also, the majority of patients included in our study had some degree of focal scarring as we included patients referred for CMR and excluded several other common clinical indications for CMR assessment, such as those with hypertrophic cardiomyopathy, valvular, and

congenital heart disease. Focal scarring might bias strain measurements differently depending on the location in the myocardium. However, the exceptionally high correlation between GLS and GCS indicates that this potential effect is unlikely to have a large magnitude of effect on the findings. Moreover, the magnitude of decrease in MAPSE has been found to associate with infarct size but not infarct location²⁴. LA and LV measurements were performed by two different observers, which may have contributed to a lower correlation between LA strain and MAPSE. All images used for strain analyses were optimized for the LV and not the LA. This could potentially affect LA length and volume calculations, and may have, in part, conferred a source of bias as there is a known correlation between an increasing LA-to-LV angle with LA and LV remodeling²⁵. This could potentially cause increasing measurement error with the presence of cardiac disease. However, both volume measurements and strain measurements were performed in the same images, which should have limited the impact of such potential measurement error.

Conclusions

LV and LA strains can be understood as geometrically coupled composite measures of MAPSE, and the size, function, and dimensions of the LV and LA. MAPSE and LV length alone can be used to accurately estimate GLS, which in turn provides similar information to GCS. These highly correlated composite geometrical relationships between strain and geometric measures of known prognostic significance likely explain the excellent prognostic value of strain measures.

Table 1. Baseline characteristics

| | |
|--|-----------------------|
| Females, n/total (%) | 14/66 (18) |
| Age, years | 62 [53 – 72] |
| LVEF, % | 48 [34 – 57] |
| MAPSE, mm | 11.1 [8.1 – 13.3] |
| GLS, % | -15.5 [-17.2 – -11.6] |
| GCS, % | -16.2 [-18.8 – -11.5] |
| BSA, m ² | 2 [1.8 – 2.1] |
| BMI, kg/m ² | 27.5 [24 – 30] |
| LAS, % | 24 [19.9 – 26.6] |
| LVSF, ml | 88 [77 – 112] |
| LV short-axis diameter, mm | 84 [78 – 91] |
| LVM, g | 124 [99 – 150] |
| LVESV, ml | 108 [80 – 170] |
| LVEDV, ml | 216 [166 – 280] |
| LA volume, ml | 88 [71 – 112] |
| LAVi, ml/m ² | 43 [36 – 56] |
| Values are expressed as median [interquartile range]. LVEF, left ventricular ejection fraction; MAPSE, mitral annular plane systolic excursion; GLS, global longitudinal strain; GCS, global circumferential strain; BSA, body surface area; BMI, body mass index; LAS, left atrial strain; LVSF, left ventricular stroke volume; LVM, left ventricular mass; LVESV, left ventricular end-systolic volume; LVEDV, left ventricular end-diastolic volume; LAVi, left atrial volume index. | |

Table 2. Associations between GLS and left heart measures.

| | Univariable | | Multivariable | | |
|--|----------------|------------------|----------------------|------------------|-----------------------|
| | R ² | p | Standardized β | p | Global R ² |
| LVEF, % | 0.82 | <0.001 | | | 0.89 |
| GCS, % | 0.86 | <0.001 | | | |
| MAPSE, mm | 0.76 | <0.001 | -0.59 | <0.001 | |
| Indexed MAPSE, %, | 0.83 | <0.001 | | | |
| LVL, mm | 0.23 | <0.001 | | | |
| SV, ml | 0.19 | <0.001 | | | |
| SA area, cm² | 0.53 | <0.001 | | | |
| SA area cm² \times MAPSE/SV | 0.01 | 0.69 | | | |
| LVM, g | 0.33 | <0.001 | 0.12 | 0.04 | |
| LVEDV, ml | 0.50 | <0.001 | | | |
| LVESV, ml | 0.69 | <0.001 | 0.37 | <0.001 | |
| LAS, % | 0.51 | <0.001 | | | |
| LA, ml | 0.13 | <0.01 | | | |


Table 3. Associations between GCS and left heart measures.

| | Univariable | | Multivariable | | |
|--|----------------|------------------|----------------------|------------------|-----------------------|
| | R ² | p | Standardized β | p | Global R ² |
| EF, % | 0.82 | <0.001 | | | 0.80 |
| GLS, % | 0.86 | <0.001 | | | |
| MAPSE, mm | 0.61 | <0.001 | -0.41 | <0.001 | |
| MAPSE/LVL, % | 0.84 | <0.001 | | | |
| LVL, mm | 0.21 | <0.001 | | | |
| SV, ml | 0.12 | <0.001 | | | |
| SA area, cm² | 0.53 | <0.001 | | | |
| SA area cm² \times MAPSE/SV | 0.01 | 0.69 | | | |
| LVM, g | 0.33 | <0.001 | 0.02 | 0.08 | |
| LVEDV, ml | 0.50 | <0.001 | | | |
| LVESV, ml | 0.69 | <0.001 | 0.56 | <0.001 | |
| LAS, % | 0.43 | <0.001 | | | |
| LAESV, ml | 0.13 | <0.01 | | | |

Table 4. Associations between LAS and left heart measures.

| | Univariable | | Multivariable | | |
|-----------------|----------------|------------------|----------------------|------------------|-----------------------|
| | R ² | p | Standardized β | p | Global R ² |
| LAEDV, mm | 0.46 | <0.001 | -0.49 | <0.001 | 0.66 |
| LAESV, mm | 0.18 | <0.01 | | | |
| LAES width, mm | 0.10 | <0.01 | | | |
| LAED width, mm | 0.37 | <0.001 | | | |
| LAES length, mm | 0.13 | <0.01 | | | |
| LAED length, mm | 0.42 | <0.001 | | | |
| GLS, % | 0.51 | <0.001 | | | |
| MAPSE, mm | 0.46 | <0.001 | 0.49 | <0.001 | |
| LVL, mm | 0.08 | 0.01 | | | |
| LVEDV, ml | 0.25 | <0.001 | | | |
| LVM, g | 0.13 | <0.01 | | | |

Figure 1. Flowchart of patient inclusion.






Figure 2. Representative patients illustrating the association between strain and geometric measures. All images are shown in the same magnification. End-diastolic (ED) and end-systolic (ES) long-axis (top row) and short-axis (lower row) images of three representative patients with normal (left), hypertrophied (middle), and dilated (right) geometry shown together with measures of left ventricular ejection fraction (LVEF), global longitudinal strain (GLS), global circumferential strain (GCS), mitral annular plane systolic excursion (MAPSE), LV mass (LVM) and end-diastolic volume (EDV), stroke volume (SV), left atrial strain (LAS), left atrial end-systolic volume (LAESV) and left atrial length (LAL). The white lines denote the LV and left atrial length (LAL) in ED and the white dotted lines LV and LA length in ES. The white double-headed arrows denote septal and lateral MAPSE. Across all patients and magnitudes of LV function, approximately 60% of the LV stroke volume is generated by the longitudinal movement of the atrioventricular plane, also referred to as MAPSE, that simultaneously empties and fills the LV and LA. Thus, MAPSE mechanically couples the function of the LV and LA to each other. Meanwhile, the LV apex and the posterior aspect of the LA are effectively stationary throughout the cardiac cycle. Note that across these representative patients, 1) Intra-individual GLS and GCS are very similar, 2) GLS is a function of both MAPSE and LV volume or length, and 3) LAS is function of MAPSE and LA volume or length.




Figure 3. Agreement between Global Longitudinal Strain (GLS) and GLS estimated as $\text{MAPSE}^* \cdot 100 / \text{LVL} - 3.96$.

1. Oikonomou EK, Kokkinidis DG, Kampaktsis PN, et al. Assessment of Prognostic Value of Left Ventricular Global Longitudinal Strain for Early Prediction of Chemotherapy-Induced Cardiotoxicity A Systematic Review and Meta-analysis Author Audio Interview Supplemental content. *JAMA Cardiol.* 2019;4(10):1007-1018.
2. Karlsen S, Dahlslett T, Grenne B, et al. Global longitudinal strain is a more reproducible measure of left ventricular function than ejection fraction regardless of echocardiographic training. *Cardiovasc Ultrasound.* 2019;17(1):1-12.
3. Fröjd F, Fridman Y, Bering P, et al. Extracellular Volume and Global Longitudinal Strain Both Associate With Outcomes But Correlate Minimally. *JACC Cardiovasc Imaging.* 2020;13(11).
4. Hochstadt A, Arnold J, Rosen R, et al. Longitudinal diastolic strain slope as an early sign for systolic dysfunction among patients with active cancer. *Clinical Research in Cardiology.* Published online November 21, 2020:1-10.

5. De E, Gripp A, Escudini De Oliveira G, et al. Global Longitudinal Strain Accuracy for Cardiotoxicity Prediction in a Cohort of Breast Cancer Patients During Anthracycline and/or Trastuzumab Treatment. Published online 2017.
6. Kammerlander AA, Kraiger JA, Nitsche C, et al. Global Longitudinal Strain by CMR Feature Tracking Is Associated With Outcome in HFPEF. *JACC Cardiovasc Imaging*. 2019;12(8P1):1585-1587.
7. Stokke TM, Hasselberg NE, Smedsrud MK, et al. Geometry as a Confounder When Assessing Ventricular Systolic Function. *J Am Coll Cardiol*. 2017;70(8):942-954.
8. Inoue K, Khan FH, Remme EW, et al. Determinants of left atrial reservoir and pump strain and use of atrial strain for evaluation of left ventricular filling pressure. *Eur Heart J Cardiovasc Imaging*. 2021;23(1):61-70.
9. Frydas A, Morris DA, Belyavskiy E, et al. Left atrial strain as sensitive marker of left ventricular diastolic dysfunction in heart failure. *ESC Heart Fail*. 2020;7(4):1956-1965.
10. Mălăescu GG, Mirea O, Capotă R, Petrescu AM, Duchenne J, Voigt JU. Left Atrial Strain Determinants During the Cardiac Phases. *Cardiovascular Imaging*. 2022;15(3):381-391.
11. M C, M U, E H, H A. The quantitative relationship between longitudinal and radial function in left, right, and total heart pumping in humans. *Am J Physiol Heart Circ Physiol*. 2007;293(1).
12. Dobrovie M, Barreiro-Pérez M, Curione D, et al. Inter-vendor reproducibility and accuracy of segmental left ventricular strain measurements using CMR feature tracking. *Eur Radiol*. 2019;29:6846-6857.
13. Huang SJ, Ting I, Huang AM, Slama M, McLean AS. Longitudinal wall fractional shortening: an M-mode index based on mitral annular plane systolic excursion

- (MAPSE) that correlates and predicts left ventricular longitudinal strain (LVLS) in intensive care patients. *Crit Care*. 2017;21(1).
14. Xue H, Artico J, Davies RH, et al. Automated In-Line Artificial Intelligence Measured Global Longitudinal Shortening and Mitral Annular Plane Systolic Excursion: Reproducibility and Prognostic Significance. *J Am Heart Assoc*. Published online February 8, 2022.
 15. Onishi T, Saha SK, Delgado-Montero A, et al. Global longitudinal strain and global circumferential strain by speckle-tracking echocardiography and feature-tracking cardiac magnetic resonance imaging: Comparison with left ventricular ejection fraction. *Journal of the American Society of Echocardiography*. 2015;28(5):587-596.
 16. Carlsson M, Ugander M, Mosén H, Buhre T, Arheden H. Atrioventricular plane displacement is the major contributor to left ventricular pumping in healthy adults, athletes, and patients with dilated cardiomyopathy. *Am J Physiol Heart Circ Physiol*. 2007;292(3).
 17. Asgeirsson D, Hedström E, Jögi J, et al. Longitudinal shortening remains the principal component of left ventricular pumping in patients with chronic myocardial infarction even when the absolute atrioventricular plane displacement is decreased. *BMC Cardiovasc Disord*. 2017;17(1).
 18. Inoue K, Khan FH, Remme EW, et al. Determinants of left atrial reservoir and pump strain and use of atrial strain for evaluation of left ventricular filling pressure.
 19. Carluccio E, Biagioli P, Mengoni A, et al. Left Atrial Reservoir Function and Outcome in Heart Failure With Reduced Ejection Fraction. *Circ Cardiovasc Imaging*. 2018;11(11):e007696.

20. Negishi K. Incremental Diagnostic Value of Left Atrial Strain Over Left Atrial Volume: An Analogy of Glucose Level and Glycosylated Hemoglobin? *. *JACC Cardiovasc Imaging*. 2018;11(10):1416-1418.
21. Hewing B, Theres L, Spethmann S, Stangl K, Dreger H, Knebel F. Left atrial strain predicts hemodynamic parameters in cardiovascular patients. *Echocardiography*. 2017;34(8):1170-1178.
22. Willenheimer R, Israelsson B, Cline C, Rydberg E, Broms K, Erhardt L. Left atrioventricular plane displacement is related to both systolic and diastolic left ventricular performance in patients with chronic heart failure. *Eur Heart J*. 1999;20(8):612-618.
23. Inoue K, Khan FH, Remme EW, et al. Determinants of left atrial reservoir and pump strain and use of atrial strain for evaluation of left ventricular filling pressure. *Eur Heart J Cardiovasc Imaging*. Published online January 26, 2021.
24. Pahlm U, Ostfeld E, Seemann F, et al. Evolution of left ventricular function among subjects with ST-elevation myocardial infarction after percutaneous coronary intervention. *BMC Cardiovasc Disord*. 2020;20(1):309.
25. Al-Mohaissen MA, W Chow BJ, Lee T, Chan KL. Left atrial-left ventricular angle, a new measure of left atrial and left ventricular remodeling. 2022;38:435-445.