

1 Concordance of weight status between mothers and children: A secondary
2 analysis of the Pakistan Demographic and Health Survey VII

3

4

5

6 Faiz Alam ^{1,2,3}, Mohammed K. Ali ^{3,4,5}, Shivani A. Patel ^{3,5}, Romaina Iqbal ^{6*}

7

8 ¹ Yale School of Public Health, Yale University, New Haven, Connecticut, United States
9 of America

10 ² School of Medicine, Emory University, Atlanta, Georgia, United States of America

11 ³ Hubert Department of Global Health, Rollins School of Public Health, Emory
12 University, Atlanta, Georgia, United States of America

13 ⁴ Department of Family and Preventive Medicine, School of Medicine, Emory University,
14 Atlanta, Georgia, United States of America

15 ⁵ Global Diabetes Research Center, Woodruff Health Sciences Center, Emory
16 University, Atlanta, Georgia, United States of America

17 ⁶ Department of Community Health Sciences, Aga Khan University, Karachi, Sindh,
18 Pakistan

19

20 * Corresponding Author

21 Email: romaina.iqbal@aku.edu (RI)

22 **Abstract**

23 Familial concordance of nutritional status is an emerging field of study that may guide the
24 development of interventions that operate beyond the individual and within the family
25 context. Little published data exist for concordance of nutritional status within Pakistani
26 households. We assessed the associations between weight status of mothers and their
27 children in a nationally representative sample of households in Pakistan using
28 Demographic and Health Survey data. Our analysis included 3465 mother-child dyads,
29 restricting to children under-five years of age with body mass index (BMI) information on
30 their mothers. We used linear regression models to assess the associations between
31 maternal BMI category (underweight, normal weight, overweight, obese) and child's
32 weight-for-height z-score (WHZ), accounting for socio-demographic characteristics of
33 mothers and children. We assessed these relationships in all children under-five and also
34 stratified by age of children (younger than 2 years and 2 to 5 years). In all children under-
35 five and in children 2 to 5 years, maternal BMI was positively associated with child's WHZ,
36 while there was no association between maternal BMI and child WHZ for children under-
37 two. The findings indicate that the weight status of mother's is positively associated with
38 that of their children. These associations have implications for interventions aimed at
39 healthy weights of families.

40

41

42 Introduction

43 There is a growing body of literature focusing on concordance of cardiometabolic
44 disease risk factors and outcomes within families and households, suggesting that NCD
45 prevention programs may be more effective if the family – rather than the individual – was
46 targeted for intervention. Studies show increased likelihood of adults living with someone
47 with a chronic condition, such as obesity, diabetes, hypertension, hyperlipidemia, and
48 other cardiometabolic conditions, having that same condition. This includes concordance
49 between parent-child [1,2], sibling [3,4], spousal dyads [5], and between any co-residing
50 members of households [6,7]. These studies investigate the genetic predisposition to
51 disease [6], as can be seen in strictly familial relationships, and also the effect of the
52 shared environment [8] and health behavior influences [9,10], such as in co-residing
53 studies.

54 Intergenerational concordance within households – and specifically that between
55 parents and children – may reflect a composite of genetic, epigenetic, and shared
56 environmental factors. The magnitude of intergenerational concordance of NCD risk
57 within households in Pakistan is largely unknown. Studies focusing on parent-child
58 concordance have been primarily done in high-income countries, such as in the United
59 Kingdom and Australia [11–16]. There is some evidence from low- and middle-income
60 countries (LMICs) regarding concordance of biomarkers such as hemoglobin A1c and C-
61 reactive protein [2]. Specifically in South Asia, studies show statistically significant
62 associations between child under-five obesity and maternal weight, but do not further
63 stratify these relationships by child age [17,18].

64 Within Pakistan, there is particular importance of the parent-child dyad. Pakistan
65 has one of the highest average household sizes and percentage of multigenerational
66 households in the world, suggesting extended sharing of environment between
67 generations. Moreover, Pakistan is following a global trend of increasing life expectancy,
68 signaling a large population of individuals at higher risk of developing NCDs that the
69 Pakistani healthcare system will have to manage in the future [19]. Additionally, children
70 in Pakistan are becoming increasingly vulnerable to different forms of malnutrition: in the
71 2018 Pakistan National Nutrition Survey (PNNS), wasting was noted in 17.7% of children,
72 the highest in Pakistan's history, and prevalence of overweight children under-five almost
73 doubled from 5% in 2011 to 9.5% in 2018. Similarly, women ages 15-49 experienced an
74 increase in overweight and obesity increasing from 28% to 37.8% from 2011 to 2018 [20].
75 This parallel increase in excess weight across a wide range of Pakistani age
76 demographics may have important implications for cardiometabolic disease at the
77 population level. Thus, there is a need to study Pakistani household concordance of NCD
78 risk factors to further understand the extent, if present, of household concordance in
79 Pakistan, especially with an additional focus on parent-child concordance to advocate for
80 interventions that involve the full household, and not only prevention for adults.

81 In the present study, we analyzed the concordance of maternal and child weight
82 status, a critical risk factor for NCDs. We further investigate the dynamics of these
83 associations across different age groups of children under 5.

84 **Materials and Methods**

85 **Data sources and participants**

86
87 Pakistan's Demographic and Health Survey, Round VII (PDHS-7) survey was
88 conducted in 2017-18 by the National Institute of Population Studies and funded by the
89 United States Agency for International Development (USAID). PDHS-7 was the 7th
90 round of a nationally representative survey, with information about household,
91 demographic, and maternal and child health indicators. The survey design and sample
92 size calculations were formulated to provide reliable information at the national level and
93 subnational levels, as well as for urban and rural areas separately, with data collected
94 from the four provinces of Punjab, Sindh, Khyber Pakhtunkhwa, and Balochistan, the
95 regions of Azad Jammu and Kashmir and Gilgit Baltistan, the Islamabad Capital
96 Territory, and the Federally Administered Tribal Areas. The sampling design used a
97 two-stage stratified approach and, because of non-proportional sample allocation, the
98 sample weights were generated by the PDHS team. Participants were recruited and
99 data was collected from November 2017 to April 2018

100 The respondents included all ever-married women aged 15-49 in all selected
101 households. In one-third of the selected households, height and weight were directly
102 measured for ever-married women 15-49 and children under-five years of age. For this
103 study, we included all children under-five who had WHZ information and BMI
104 information on their mothers, leading to a sampling-weight adjusted sample size of
105 3465. Fig 1 depicts the sampling scheme and relevant response rates for PDHS-7 [21].

Fig 1. Flow diagram of household and participant selection for the 2017-18 Pakistan Demographic and Health Survey.

106 The study was reviewed and approved by the National Bioethics Committee of
107 the Pakistan Health Research Council and the Inner City Fund/ ICF Institutional Review
108 Board. Adult participants were consented both, for themselves and for their children
109 [21]. Prior to obtaining access to and performing analysis on the PDHS-7 data, the
110 researchers of the present study submitted a concept note, gained data access, and
111 obtained permission from USAID. The data were de-identified and anonymized by the
112 DHS program prior to being shared, therefore not considered human subjects research
113 by an author-affiliated institution (Emory University, Atlanta, GA, USA) and not requiring
114 further ethical clearance.

115 **Study Measures**

116

117 **Child weight status: Weight-for-height z-score (WHZ)**

118 Child weight and height measurements were taken during a single visit. Weight
119 measurements were taken using SECA scales (model no. SECA 878U) and height
120 measurements using a Shorr Board (recumbent if child's age was less than 24 months
121 and standing if older). Anthropometry data in PDHS-7 was collected by two female
122 enumerators in each field team (total of 44 enumerators) who jointly took
123 measurements. All enumerators were trained beforehand to standardize procedures for
124 anthropometry, including hands-on training to measure ten children twice to assess
125 accuracy and precision of measurements and further training for those enumerators
126 who were out of range more three or more times [21].

127 The dependent variable was weight-for-height z-score (WHZ) for children under-
128 five per the 2006 Child Growth Standards released by the World Health Organization,
129 reliably shown to describe both overweight (WHZ > 2) and wasting (WHZ < -2) [22].

130 Every one unit in z-score represents one standard deviation from the mean weight-for-
131 height from a study of more than eight thousand children recruited from Brazil, Ghana,
132 India, Norway, Oman and the United States of America [23,24]. For example, a WHZ of
133 -2 indicates that the child's weight is two standard deviations below the mean for their
134 height and sex [25].

135 **Maternal weight status: Body Mass Index (BMI)**

136 The primary exposure of interest for this study was maternal BMI. BMI was
137 measured in the same way as child WHZ, with SECA 878U scales and Shorr Boards
138 (standing height measurement). Maternal BMI was coded as obese (30 or greater),
139 overweight (25 to less than 30), healthy weight (18.5 to less than 25), and underweight
140 (less than 18.5), which was the reference category ([ref]).

141 **Covariates**

142 Several other covariates were also considered in the analysis. A covariate was
143 included in statistical the models if it had evidence of correlation with child weight
144 outcome and if there was sufficient variation in the covariate for it to add information to
145 the model, e.g. if a factor has 99% 'yes' response rate and 1% 'no' rate, then its use
146 was avoided because it would not meaningfully explain variation in the outcome.
147 Through this process, the following eight covariates were selected for analysis: urban/
148 rural specific wealth index [18,22,26–28], sex of child [22,26], mother's employment
149 status[26,27,29], type of place of residence [28,30–32], mother's youngest child under-
150 five [28,33,34], mother's oldest child under-five [28,33,34], child age in months, and
151 maternal age in years [29,33,34].

152 Household wealth status was measured as quintiles of assets as provided by the
153 PDHS-7 survey. Wealth index quintile categories were labeled as richest, rich, middle,
154 poor, poorest[ref]. Child sex was coded as male or female[ref]. Type of place of
155 residence was coded as urban or rural[ref]. Mother's employment status, youngest child
156 under-five and oldest child under-five were all codes as no or yes[ref].

157 **Analysis**

158
159 A dyadic dataset of mothers and their children was created by merging
160 anthropometric and demographic characteristics of mothers and children per DHS
161 guidance for analysis. Mother's characteristics were treated as exposures for the child
162 [35]. Furthermore, in studying cardiometabolic risk concordance between younger
163 children and their mothers, it is important to consider the impact of children being born
164 small for gestational age (SGA) can have on anthropometric concordance. This is
165 particularly important consideration in Pakistan, where the prevalence of SGA births is
166 >45% [36]. As approximately 85% of children born SGA have experienced sufficient
167 catch up growth by age 2 [37–39], stratifying analysis by this age point in relation to
168 concordance which, to our knowledge, has not been done before, could yield important
169 findings regarding the 'unmasking' of parent-child concordance. Thus, we stratified
170 children by two years of age in our analysis and modeling.

171 All analysis and modeling accounted for the complex survey design [35]. All data
172 restructuring and analysis was done using IBM SPSS Statistics v29.0.0.0 with Complex
173 Samples v29. The code used for restructuring and analysis can be found via the
174 following link: [doi:10.5281/zenodo.7794384](https://doi.org/10.5281/zenodo.7794384).

175 **Descriptive statistics**

176
177 Reported statistics (sampling weight-adjusted) included background
178 characteristics of the participants which include average age of mother (in years) and
179 children (in months), mean BMI of mothers, and mean child WHZ with 95% confidence
180 intervals. Other reported adjusted population descriptive statistics were percentage of
181 dyads that were urban vs rural, percentage of male and female children, percentage of
182 mothers currently employed, percentage of children being the youngest child,
183 percentage of children being the oldest under-five, and the percentage of the sample
184 size in each wealth index quintile. The mean child WHZ with 95% confidence interval
185 and sampling weight-adjusted sample size was also reported for each factor level.

186 **Modeling associations between maternal and child weight status**

187
188 We estimated linear regression models to investigate concordance between
189 maternal BMI and child WHZ, measured continuously. Three linear models were
190 generated: 1) all children under-five years, 2) all children under-two years, 3) all children
191 ages 2 to 5, with the models adjusted for the 8 covariates. We reported the R^2 value and
192 the Wald F for each model and reported the parameter estimates with 95% confidence
193 interval and t-test value for each categorical level. For each model, the statistically
194 significant variables were remodeled including only these variables and an additional
195 term for each two-variable combination to assess for interaction between the statistically
196 significant variables. Cases with missing values for any variables were treated as
197 invalid. Associations for all models were considered statistically significant when $p <$
198 0.05.

199 **Further Considerations for Analysis**

200
201 We assessed multicollinearity between the independent variables using the
202 variance inflation factor (VIF) calculation. We used a conservative cutoff of 3, as an
203 indicator to further investigate relationships [40]. Using this cutoff, no multicollinearity
204 was observed among the independent variables. To account for multiple comparisons
205 and counter the increased risk of Type I error, the Holm-Bonferroni method was used to
206 adjust the P-value of 0.05 accordingly [41,42]. Regarding outliers, the DHS data is
207 screened before dissemination for plausibility of Z-scores. For children who had WHZ
208 that were below -5 standard deviations or above +5 standard deviations were flagged
209 and their WHZ was not reported. Similarly, women who had BMI below 12 or above 60
210 were also flagged and did not have their BMI reported [35]. Further consideration of
211 outliers was not made because of these adjustments by the PDHS data processing
212 team.

213 **Results**

214 For the analysis, out of 4671 sample unweighted cases, 4130 had complete
215 variable information (with 541 missing) and were included, which led to a sampling-weight
216 adjusted sample size of 3465. With cases that had information on maternal BMI and child
217 WHZ, only two did not have complete information on the other covariates.

218 **Summary Statistics**

219
220 Of the overall sampling weight-adjusted sample of 3465 children (Fig. 1), 1406
221 (40.6%) children were less than 2 years old, and 2059 (59.4%) children were ages 2 to
222 5. The mean age of mothers was 29.0 years (95% CI [28.6, 29.4]) and children under-
223 five years was 28.8 months (95% CI [28.0, 29.5]). The mean maternal BMI was 24.6

224 (95% CI [24.2, 25.1]). The mean child WHZ for the entire sample was -0.30 (95% CI [-
225 0.36, -0.24]), mean child WHZ for children under-two was -0.41 (95% CI [-0.50, -0.31]),
226 and mean child WHZ for children ages 2 to 5 was -0.23 (95% CI [-0.29, -0.16]). The
227 greatest mean child WHZ was in children who belonged to the richest quintile of
228 households at 0.04 (95% CI [-0.09, 0.17]), and the category with the lowest mean child
229 WHZ was children with mothers who were underweight at -0.66 (95% CI [-0.80, -0.51]).
230 Table 1 summarizes maternal, child, and household characteristics of the analyzed
231 maternal-child dyads and the mean child WHZ for each factor level is summarized in
232 Table 2.

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247 Table 1: Percentage of sample in each factor level and mean values of continuous variables (n=3465),
 248 with 95% confidence interval from PDHS-7 dataset used in analysis.

Variable	All	Children under two	Children 2 to 5 years
Maternal Characteristics			
		Percent (95% CI)	
Maternal BMI Category			
30+	15.7 (13.1 – 18.8)	13.1 (10.4 – 16.4)	17.5 (14.5 – 20.9)
25 to < 30	27.6 (24.8 – 30.6)	27.9 (24.5 – 31.7)	27.4 (24.3 – 30.7)
18.5 to <25	46.2 (42.9 – 49.6)	46.7 (42.6 – 50.8)	45.9 (42.1 – 49.8)
<18.5	10.4 (8.5 – 12.7)	12.3 (9.9 – 15.1)	9.2 (7.2 – 11.7)
Mother currently employed			
No	86.7 (83.9 – 89.0)	34.6 (30.1 – 39.4)	31.4 (27.6 – 35.5)
Yes	13.3 (11.0 – 16.1)	65.4 (60.6 – 69.9)	68.6 (64.5 – 72.4)
		Mean (95% CI)	
Maternal BMI	24.6 (24.2, 25.1)	24.3 (23.8, 24.8)	24.9 (24.4, 25.4)
Maternal Age (yrs.)	29.0 (28.6, 29.4)	27.4 (26.9, 27.9)	30.3 (29.6, 30.5)
Child Characteristics			
		Percent (95% CI)	
Youngest child			
No	34.3 (32.8 – 35.9)	5.5 (4.2 – 7.2)	54 (51.3 – 56.7)
Yes	65.7 (64.1 – 67.2)	94.5 (92.8 – 95.8)	46 (43.3 – 48.7)
Oldest child under-five			
No	32.5 (30.9 – 34.1)	56.4 (52.9 – 59.8)	16.2 (14.3 – 18.2)
Yes	67.5 (65.9 – 69.1)	43.6 (40.2 – 47.1)	83.8 (81.8 – 85.7)
Sex of child			
Male	51.1 (49.1 – 53.1)	51.2 (47.3 – 55.0)	51.0 (48.7 – 53.3)
Female	48.9 (46.9 – 50.9)	48.8 (45.0 – 52.7)	49.0 (46.7 – 51.3)
		Mean (95% CI)	
Child WHZ	-0.30 (-0.36, -0.24)	-0.41 (-0.50, -0.31)	-0.23 (-0.29, -0.16)
Child Age (mos.)	28.8 (28.0, 29.5)	11.0 (10.5, 11.5)	40.9 (40.4, 41.5)
Household Characteristics			
		Percent (95% CI)	
Area			
Urban	32.7 (29.1 – 36.6)	34.6 (30.1 – 39.4)	31.4 (27.6 – 35.5)
Rural	67.3 (63.4 – 70.9)	65.4 (60.6 – 69.9)	68.6 (65.4 – 72.4)
Wealth index (urban/ rural specific)			
Richest	18.3 (14.7 – 22.5)	19.1 (14.7 – 24.5)	17.7 (14.1 – 21.9)
Richer	20.3 (17.0 – 24.0)	21.9 (18.0 – 26.3)	19.1 (15.7 – 23.1)
Middle	20.1 (17.4 – 23.0)	20.2 (17.0 – 23.9)	19.9 (17.0 – 23.2)
Poor	21.4 (18.4 – 24.8)	20.2 (16.8 – 24.1)	22.3 (18.8 – 26.1)
Poorest	20.0 (16.5 – 24.0)	18.6 (15.1 – 22.5)	21.0 (16.8 – 25.8)

249

250 Table 2: Mean Child WHZ and sampling weight-adjusted sample size (Adj. N) of analyzed sample,
251 stratified by level of selected factors.

Variable	Adj. N	Mean child WHZ (95% CI)
All	3465	-0.3 (-0.36, -0.24)
Maternal BMI Category		
30+	545	-0.02 (-0.16, 0.11)
25 to < 30	956	-0.14 (-0.25, -0.03)
18.5 to <25	1602	-0.41 (-0.51, -0.31)
<18.5	361	-0.66 (-0.8, -0.51)
Mother currently employed		
No	3003	-0.29 (-0.36, -0.23)
Yes	462	-0.35 (-0.52, -0.19)
Youngest child		
No	1189	-0.21 (-0.29, -0.13)
Yes	2276	-0.35 (-0.42, -0.28)
Oldest child under-five		
No	1126	-0.4 (-0.51, -0.28)
Yes	2339	-0.26 (-0.32, -0.2)
Sex of child		
Male	1770	-0.32 (-0.39, -0.24)
Female	1695	-0.29 (-0.36, -0.21)
Area		
Urban	1134	-0.31 (-0.4, -0.21)
Rural	2331	-0.30 (-0.37, -0.22)
Wealth index (urban/ rural specific)		
Richest	633	0.04 (-0.09, 0.17)
Richer	702	-0.17 (-0.29, -0.06)
Middle	695	-0.36 (-0.5, -0.22)
Poor	743	-0.52 (-0.64, -0.4)
Poorest	693	-0.45 (-0.58, -0.31)

252
253

254 **Associations between mother and child weight status**

255
256 Maternal BMI was positively associated with child WHZ for all children among
257 children under-five. Children of normal weight, overweight, and obese women had WHZ
258 scores that were 0.22 [95% CI: 0.05, 0.38], 0.44 [95% CI: 0.26, 0.63], and 0.53 [95% CI:
259 0.33, 0.73] units higher than children of underweight women, respectively. Maternal BMI
260 was also positively associated with child WHZ among children ages 2 to 5, where children
261 of normal weight, overweight, and obese women had WHZ scores that were 0.27 [95%

262 CI: 0.09, 0.44), 0.51 [95% CI: 0.30, 0.72), and 0.61 [95% CI: 0.37, 0.85] units higher than
263 children of underweight women, respectively.

264 For all children under-five, household wealth index was positively associated with
265 child WHZ (Richest: 0.37 [95% CI: 0.18, 0.55]; Rich: 0.19 [95% CI: 0.02, 0.37]; Middle:
266 0.01 [95% CI: -0.17, 0.20]; Poor: -0.12 [95% CI: -0.30, 0.06]; Poorest [ref.]). Household
267 wealth was similarly positively associated with child WHZ for children under 2 (Richest:
268 0.44 [95% CI: 0.13, 0.74]; Rich: 0.21 [95% CI: -0.08, 0.49]; Middle -0.31 [95% CI: -0.58, -
269 0.44]; Poor: -0.15 [95% CI: -0.46, 0.17]; Poorest: [ref]) and for children aged two to five
270 (Richest: 0.31 [95% CI: 0.11, 0.51]; Rich: 0.17 [95% CI: -0.01, 0.35]; Middle 0.23 [95%
271 CI: 0.03, 0.43]; Poor -0.10 [95% CI: -0.28, 0.07]; Poorest: [ref.]).

272 There were no statistically significant interaction effects for any linear model. Table
273 3 summarizes information for each model.

274

275

276

277

278

279

280

281

282

283

284 Table 3: Linear regression results of the association between maternal BMI and child WHZ, along with
 285 selected covariates, among three population groups: all children under-five, children under-two, children 2
 286 to 5.

Variable (Category)	All children under-five			Children under-two			Children 2 to 5 years		
	Coefficient (95% CI)	t-value	p	Coefficient (95% CI)	t-value	p	Coefficient (95% CI)	t-value	p
Mother BMI category			(<0.001)			0.130			(<0.001)
30 or greater	0.53(0.33, 0.73)	5.201	<.001	0.32(-0.02, 0.67)	1.836	0.067	0.61(0.37, 0.85)	5.043	<.001
25 to <30	0.44(0.26, 0.63)	4.670	<.001	0.26(-0.05, 0.57)	1.642	0.101	0.51(0.30, 0.72)	4.857	<.001
18.5 to <25	0.22(0.05, 0.38)	2.587	0.010	0.06(-0.21, 0.34)	0.447	0.655	0.27(0.09, 0.44)	2.945	0.003
less than 18.5	*			*			*		
Mother currently working			0.363			0.812			0.168
No	-0.08(-0.25, 0.09)	-0.910	0.363	0.03(-0.21, 0.28)	0.265	0.791	-0.14(-0.33, 0.05)	-1.423	0.155
Yes	*			*			*		
Youngest child			0.020			0.024			0.065
No	0.15(0.02, 0.27)	2.335	0.020	0.36(0.05, 0.67)	2.285	0.023	0.11(-0.03, 0.25)	1.574	0.116
Yes	*			*			*		
Oldest child under-five			0.519			0.650			0.667
No	-0.05(-0.19, 0.10)	-0.646	0.519	-0.05(-0.24, 0.14)	-0.491	0.624	-0.05(-0.25, 0.16)	-0.445	0.657
Yes	*			*			*		
Sex of child			0.420			0.178			0.934
Male	-0.04(-0.14, 0.06)	-0.808	0.420	-0.12(-0.3, 0.06)	-1.358	0.175	0.00(-0.10, 0.11)	0.045	0.964
Female	*			*			*		
Area			0.474			0.768			0.455
Urban	-0.04(-0.15, 0.07)	-0.723	0.470	-0.03(-0.21, 0.16)	-0.287	0.774	-0.05(-0.19, 0.08)	-0.785	0.433
Rural	*			*			*		
Wealth index (urban/ rural specific)			(<.001)			(<.001)			(<.001)
Richest	0.37(0.18, 0.55)	3.942	<.001	0.44(0.13, 0.74)	2.825	0.005	0.31(0.11, 0.51)	3.088	0.002
Rich	0.19(0.02, 0.37)	2.155	0.032	0.21(-0.08, 0.49)	1.413	0.158	0.17(-0.01, 0.35)	1.812	0.071
Middle	0.01(-0.17, 0.2)	0.139	0.889	-0.31(-0.58, -0.04)	-2.271	0.024	0.23(0.03, 0.43)	2.227	0.026
Poor	-0.12(-0.3, 0.06)	-1.341	0.181	-0.15(-0.46, 0.17)	-0.916	0.360	-0.10(-0.28, 0.07)	-1.168	0.244
Poorest	*			*			*		
Mother age (yrs)	0.00(-0.01, 0.01)	-0.130	0.897	0.00(-0.02, 0.02)	0.229	0.819	-0.01(-0.02, 0.00)	-2.490	0.019
Child age (mos)	0.00(0.00, 0.01)	0.340	0.734	0.01(0.00, 0.02)	1.312	0.190	-0.01(-0.02, 0.00)	-2.490	0.528
R²	0.050			0.060			0.064		
Wald F	8.551			3.885			9.147		

287 *Overall variable p-values in parenthesis denote statistically significant associations (Holm-Bonferroni*
 288 *corrected)*

289 * denotes reference category

290 Discussion and Conclusion

291 We found that maternal BMI was positively associated with child WHZ in children
 292 under 5. This positive association was present for children ages 2 to 5, even after
 293 accounting for several socio-demographic factors, but not for children under-two. The
 294 findings suggest that there is intergenerational concordance in weight status in Pakistani
 295 mothers and young children which is unmasked with increased age.

296 The findings of parent-child concordance of BMI-WHZ for all children under-five
297 are consistent with previous literature which specifically explore parent-child concordance
298 of NCD risk [2,11–16] and of association studies in South Asia which have observed a
299 positive association between maternal and children under-five weight status [17,18].
300 However, previously described phenomenon of the double burden of malnutrition at
301 household level (DBMHL) suggests co-existence of undernourished children and
302 overweight mothers in the same household, but this phenomenon is primarily described
303 for child stunting and, in some cases, underweight [43–51]. Although a recent study by
304 Biswas et al., in which they analyzed the co-existence of overweight mothers with all types
305 of undernourished children under-five (stunted, wasted, and underweight) in South and
306 Southeast Asia from 2006 to 2017 data, described positive relationships between
307 maternal weight status and all three types of child undernutrition, with the greatest co-
308 existence in Pakistan. However, this study did not analyze the relationship between non-
309 overweight or obese mothers and their children, nor did it split the children under-five into
310 two age groups [52].

311 Furthermore, double burden of malnutrition (DBM) at the country or population
312 level is well-established in previous literature [53–56]. DBM prevalence has been
313 increasing in Pakistan as shown by PNNS 2018 results, with wasting noted in 17.7% of
314 children, the highest in Pakistan’s history, prevalence of overweight children under-five
315 almost doubling from 5% in 2011 to 9.5% in 2018, and women ages 15-49 experiencing
316 an increase in overweight and obesity increasing from 28% to 37.8% from 2011 to 2018.
317 Both, DBMHL and DBM, can co-exist with maternal-child BMI-WHZ concordance, as the
318 increasing burden of DBM, with an increasing percentage of overweight children, could

319 be leading to the more recent findings of positives association of maternal-child weight
320 status. Co-existence of these phenomena within the same society highlight the complexity
321 of interventions and approaches required to address malnutrition among LMICs, as
322 different regions and households are experiencing various challenges.

323 To our knowledge, there are no studies which specifically investigate the
324 relationship between maternal and child concordance under-five stratified by 2 years of
325 age, as we analyzed in this study. These findings, which show a positive association
326 between maternal BMI and child WHZ only in the 2 to 5 age group, potentially indicate
327 that concordance between child WHZ and maternal BMI is 'unmasked' after age 2, when
328 the majority of early childhood catch-up growth has occurred [37–39]. As in, children who
329 may be born SGA may not show concordance of WHZ with maternal BMI during the first
330 two years of life, but as they experience their expected catch up growth and have spent
331 more time in a shared environment with their mothers, their weight status starts to become
332 concordant with their mothers, as seen by our analysis which shows the difference in
333 association between children older than and younger than two. This suggests that
334 children under-two who do not initially appear to have WHZ concordant to their mother's
335 BMI may develop this positive relationship later in life. Furthermore, a targeted review on
336 maternal and child overweight and obesity in LMICs shows that junk food, such as potato
337 chips, sponge cakes, sugary biscuits, and sugary drinks are a significant part of child diet
338 by the first two years of life [57], potentially yielding an avenue of intervention and an
339 explanation for the development of mother-child concordance after two years of age. This
340 relationship could also be affected by other dietary aspects such as breastfeeding status
341 and weaning, as prolongation of breastfeeding has been shown to decrease excess

342 weight gain in children [58]. Overall, this may indicate an avenue for early interventions
343 for weight-related risks in children, where healthier nutritional sources during infancy may
344 lead to a decreased sharing of NCD risk found between very young children and their
345 mothers as children age.

346 Regarding other covariates which were analyzed, we showed wealth index as a
347 statistically significant positive factor across all GLM analyses. This association existing
348 in Pakistan is well established in the literature [59–64]. Increasing wealth quintile has
349 been suggested to correlate with access to nutrient rich foods and increased sedentary
350 behaviors [18,65–68].

351 Considering strengths of this study, the analysis was done on a nationally
352 representative dataset which involved a complex sampling process to obtain the best
353 possible demographic snapshot of Pakistan. The analysis also included a significant
354 number of covariates to unveil potential confounders and used rigid levels of correction
355 and validation regarding significance and multi-collinearity. Limitations of the study
356 included the use of cross-sectional data, whereas a cohort study could provide more in-
357 depth data on trends in specific households.

358 Overall, the findings of maternal-child concordance and, more specifically, the
359 ‘masking’ phenomenon could have important implications regarding interventions and
360 policy making, where these seemingly non-concordant groups may be ignored as an
361 avenue to improve NCD risk unless there is data suggesting that the concordance may
362 ‘unmask’ later. Future work should involve more detailed data collection of
363 cardiometabolic disease and, more broadly, NCD risk concordance and outcome

364 concordance on entire households, including young children, to develop family-wide
365 interventions to reduce long-term mortality from these diseases.

366 **Acknowledgements**

367 We thank the DHS Program supported by USAID, the National Institute of Population
368 Studies, and ICF for the data used in this study. We also thank the Emory Global
369 Diabetes Research Center and Global Health Equity Scholars NIH FIC TW010540.

370 **References**

- 371 1. Park HS, Park JY, Cho SI. Familial aggregation of the metabolic syndrome in Korean families
372 with adolescents. *Atherosclerosis*. 2006 May 1;186(1):215–21.
- 373 2. Dong F, Howard AG, Herring AH, Adair LS, Thompson AL, Popkin BM, et al. Concordance of
374 haemoglobin A1c, blood pressure and C-reactive protein between children and their
375 parents in Chinese households. *Pediatric Obesity*. 2017;12(5):422–30.
- 376 3. Feng Y, Zang T, Xu X, Xu X. Familial Aggregation of Metabolic Syndrome and Its Components
377 in a Large Chinese Population. *Obesity*. 2008;16(1):125–9.
- 378 4. Gupta S, Kapoor S. GENETIC AND ENVIRONMENTAL INFLUENCES ON BLOOD PRESSURE IN
379 AN URBAN INDIAN POPULATION. *Journal of Biosocial Science*. 2013 Jan;45(1):1–11.
- 380 5. Kim HC, Kang DR, Choi KS, Nam CM, Thomas GN, Suh I. Spousal Concordance of Metabolic
381 Syndrome in 3141 Korean Couples: A Nationwide Survey. *Annals of Epidemiology*. 2006 Apr
382 1;16(4):292–8.
- 383 6. Lee MH, Kim HC, Thomas GN, Ahn SV, Hur NW, Choi DP, et al. Familial concordance of
384 metabolic syndrome in Korean population—Korean National Health and Nutrition
385 Examination Survey 2005. *Diabetes Research and Clinical Practice*. 2011 Sep 1;93(3):430–6.
- 386 7. Patel SA, Dhillon PK, Kondal D, Jeemon P, Kahol K, Manimunda SP, et al. Chronic disease
387 concordance within Indian households: A cross-sectional study. *PLOS Medicine*. 2017 Sep
388 29;14(9):e1002395.
- 389 8. Smith KR, Zick CD. Linked Lives, Dependent Demise? Survival Analysis of Husbands and
390 Wives. *Demography*. 1994 Feb 1;31(1):81–93.

- 391 9. Umberson D. Family Status and Health Behaviors: Social Control as a Dimension of Social
392 Integration. *Journal of Health and Social Behavior*. 1987;28(3):306–19.
- 393 10. Meyler D, Stimpson JP, Peek MK. Health concordance within couples: A systematic review.
394 *Social Science & Medicine*. 2007 Jun 1;64(11):2297–310.
- 395 11. Hawkins SS, Cole TJ, Law C, Group and the MCSCHE. An ecological systems approach to
396 examining risk factors for early childhood overweight: findings from the UK Millennium
397 Cohort Study. *Journal of Epidemiology & Community Health*. 2009 Feb 1;63(2):147–55.
- 398 12. Reilly JJ, Armstrong J, Dorosty AR, Emmett PM, Ness A, Rogers I, et al. Early life risk factors
399 for obesity in childhood: cohort study. *BMJ*. 2005 Jun 9;330(7504):1357.
- 400 13. Salsberry PJ, Reagan PB. Dynamics of Early Childhood Overweight. *Pediatrics*. 2005 Dec
401 1;116(6):1329–38.
- 402 14. Robinson SM. Preventing childhood obesity: Early-life messages from epidemiology.
403 *Nutrition Bulletin*. 2017;42(3):219–25.
- 404 15. Clifford SA, Gillespie AN, Olds T, Grobler AC, Wake M. Body composition: population
405 epidemiology and concordance in Australian children aged 11–12 years and their parents.
406 *BMJ Open* [Internet]. 2019 Jul 1 [cited 2021 Nov 2];9(Suppl 3). Available from:
407 https://bmjopen.bmj.com/content/9/Suppl_3/95
- 408 16. Gibson LY, Byrne SM, Davis EA, Blair E, Jacoby P, Zubrick SR. The role of family and maternal
409 factors in childhood obesity. *Medical Journal of Australia*. 2007;186(11):591–5.
- 410 17. Cesare MD, Bhatti Z, Soofi SB, Fortunato L, Ezzati M, Bhutta ZA. Geographical and
411 socioeconomic inequalities in women and children’s nutritional status in Pakistan in 2011:
412 an analysis of data from a nationally representative survey. *The Lancet Global Health*. 2015
413 Apr 1;3(4):e229–39.
- 414 18. Bishwajit G, Yaya S. Overweight and obesity among under-five children in South Asia. *Child
415 and Adolescent Obesity*. 2020 Jan 1;3(1):105–21.
- 416 19. United Nations Population Division. *World Population Prospects: The 2022 Revision*
417 [Internet]. United Nations, Department of Economic and Social Affairs, Population Division
418 (2022). [cited 2022 Dec 4]. Available from: population.un.org/wpp
- 419 20. Government of Pakistan, UNICEF. *National Nutrition Survey 2018 - Full Report (3 Volumes)
420 & Key Findings Report* [Internet]. Islamabad, Pakistan: UNICEF; 2020 Oct. Available from:
421 [https://www.unicef.org/pakistan/reports/national-nutrition-survey-2018-full-report-3-
422 volumes-key-findings-report](https://www.unicef.org/pakistan/reports/national-nutrition-survey-2018-full-report-3-volumes-key-findings-report)

- 423 21. National Institute of Population Studies (NIPS) [Pakistan], ICF. Pakistan Demographic and
424 Health Survey 2017-18 [Internet]. Islamabad, Pakistan and Rockville, Maryland, USA: NIPS
425 and ICF; 2019. Available from: <http://dhsprogram.com/pubs/pdf/FR354/FR354.pdf>
- 426 22. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, et al. Maternal and child
427 undernutrition and overweight in low-income and middle-income countries. *The Lancet*.
428 2013 Aug 3;382(9890):427–51.
- 429 23. Nutrition and Food Safety WHO Team. WHO child growth standards:: Length/height-for-
430 age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age;
431 methods and development [Internet]. Geneva: WHO Press; 2006. 312 p. (WHO child growth
432 standards). Available from: <https://www.who.int/publications/i/item/924154693X>
- 433 24. ICF. Demographic and Health Surveys Standard Recode Manual for DHS7 [Internet].
434 Rockville, Maryland, USA: The Demographic and Health Surveys Program; [cited 2022 Dec
435 4]. Available from:
436 https://dhsprogram.com/pubs/pdf/DHSG4/Recode7_DHS_10Sep2018_DHSG4.pdf
- 437 25. World Health Organization. Training Course on Child Growth Assessment [Internet].
438 Geneva: WHO; 2008 [cited 2023 Jan 17] p. 58. Available from:
439 <https://apps.who.int/iris/handle/10665/43601>
- 440 26. Abera L, Dejene T, Laelago T. Prevalence of malnutrition and associated factors in children
441 aged 6–59 months among rural dwellers of damot gale district, south Ethiopia: community
442 based cross sectional study. *International Journal for Equity in Health*. 2017 Jun
443 26;16(1):111.
- 444 27. Mushtaq MU, Gull S, Shahid U, Shafique MM, Abdullah HM, Shad MA, et al. Family-based
445 factors associated with overweight and obesity among Pakistani primary school children.
446 *BMC Pediatr*. 2011 Dec;11(1):1–10.
- 447 28. Siddiqa M, Zubair A, Kamal A, Ijaz M, Abushal T. Prevalence and associated factors of
448 stunting, wasting and underweight of children below five using quintile regression analysis
449 (PDHS 2017–2018). *Sci Rep*. 2022 Nov 25;12(1):20326.
- 450 29. Mamabolo RL, Alberts M, Steyn NP, Waal HAD van de, Levitt NS. Prevalence and
451 determinants of stunting and overweight in 3-year-old black South African children residing
452 in the Central Region of Limpopo Province, South Africa. *Public Health Nutrition*. 2005
453 Aug;8(5):501–8.
- 454 30. Fotso JC. Child health inequities in developing countries: differences across urban and rural
455 areas. *International Journal for Equity in Health*. 2006 Jul 11;5(1):9.
- 456 31. Wieringa FT, Gauthier L, Greffeuille V, Som SV, Dijkhuizen MA, Lailou A, et al. Identification
457 of Acute Malnutrition in Children in Cambodia Requires Both Mid Upper Arm Circumference


- 458 and Weight-For-Height to Offset Gender Bias of Each Indicator. *Nutrients*. 2018
459 Jun;10(6):786.
- 460 32. Khan S, Zaheer S, Safdar NF. Determinants of stunting, underweight and wasting among
461 children < 5 years of age: evidence from 2012-2013 Pakistan demographic and health
462 survey. *BMC Public Health*. 2019 Dec;19(1):1–15.
- 463 33. Das S, Rahman RM. Application of ordinal logistic regression analysis in determining risk
464 factors of child malnutrition in Bangladesh. *Nutr J*. 2011 Nov 14;10:124.
- 465 34. Hajian-Tilaki K, Heidari B. Childhood Obesity, Overweight, Socio-Demographic and Life Style
466 Determinants among Preschool Children in Babol, Northern Iran. *Iran J Public Health*. 2013
467 Nov;42(11):1283–91.
- 468 35. Trevor N. Croft, Aileen M.J. Marshall, Courtney K. Allen. Guide to DHS Statistics [Internet].
469 Rockville, Maryland, USA: ICF; [cited 2022 Nov 22]. Available from:
470 https://dhsprogram.com/pubs/pdf/DHSG1/Guide_to_DHS_Statistics_DHS-7_v2.pdf
- 471 36. Lee AC, Katz J, Blencowe H, Cousens S, Kozuki N, Vogel JP, et al. National and regional
472 estimates of term and preterm babies born small for gestational age in 138 low-income and
473 middle-income countries in 2010. *Lancet Glob Health*. 2013 Jul;1(1):e26–36.
- 474 37. Cho WK, Suh BK. Catch-up growth and catch-up fat in children born small for gestational
475 age. *Korean J Pediatr*. 2016 Jan;59(1):1–7.
- 476 38. Hokken-Koelega AC, De Ridder MA, Lemmen RJ, Den Hartog H, De Muinck Keizer-Schrama
477 SM, Drop SL. Children born small for gestational age: do they catch up? *Pediatr Res*. 1995
478 Aug;38(2):267–71.
- 479 39. Karlberg J, Albertsson-Wikland K. Growth in Full- Term Small-for-Gestational-Age Infants:
480 From Birth to Final Height. *Pediatr Res*. 1995 Nov;38(5):733–9.
- 481 40. Thompson CG, Kim RS, Aloe AM, Becker BJ. Extracting the Variance Inflation Factor and
482 Other Multicollinearity Diagnostics from Typical Regression Results. *Basic and Applied Social
483 Psychology*. 2017 Mar 4;39(2):81–90.
- 484 41. Holm S. A Simple Sequentially Rejective Multiple Test Procedure. *Scandinavian Journal of
485 Statistics*. 1979;6(2):65–70.
- 486 42. Aickin M, Gensler H. Adjusting for multiple testing when reporting research results: the
487 Bonferroni vs Holm methods. *Am J Public Health*. 1996 May;86(5):726–8.
- 488 43. Anik AI, Rahman MM, Rahman MM, Tareque MI, Khan MN, Alam MM. Double burden of
489 malnutrition at household level: A comparative study among Bangladesh, Nepal, Pakistan,
490 and Myanmar. *PLOS ONE*. 2019 Aug 16;14(8):e0221274.

- 491 44. Jehn M, Brewis A. Paradoxical malnutrition in mother–child pairs: Untangling the
492 phenomenon of over- and under-nutrition in underdeveloped economies. *Economics &*
493 *Human Biology*. 2009 Mar 1;7(1):28–35.
- 494 45. Oddo VM, Rah JH, Semba RD, Sun K, Akhter N, Sari M, et al. Predictors of maternal and child
495 double burden of malnutrition in rural Indonesia and Bangladesh. *The American Journal of*
496 *Clinical Nutrition*. 2012 Apr 1;95(4):951–8.
- 497 46. Fernald LC, Neufeld LM. Overweight with concurrent stunting in very young children from
498 rural Mexico: prevalence and associated factors. *Eur J Clin Nutr*. 2007 May;61(5):623–32.
- 499 47. Garrett JL, Ruel MT. Stunted Child–Overweight Mother Pairs: Prevalence and Association
500 with Economic Development and Urbanization. *Food Nutr Bull*. 2005 Jan 1;26(2):209–21.
- 501 48. Doak CM, Adair LS, Bentley M, Monteiro C, Popkin BM. The dual burden household and the
502 nutrition transition paradox. *Int J Obes*. 2005 Jan;29(1):129–36.
- 503 49. Doak CM, Adair LS, Monteiro C, Popkin BM. Overweight and Underweight Coexist within
504 Households in Brazil, China and Russia. *The Journal of Nutrition*. 2000 Dec 1;130(12):2965–
505 71.
- 506 50. Kosaka S, Umezaki M. A systematic review of the prevalence and predictors of the double
507 burden of malnutrition within households. *British Journal of Nutrition*. 2017
508 Apr;117(8):1118–27.
- 509 51. Davis JN, Oaks BM, Engle-Stone R. The Double Burden of Malnutrition: A Systematic Review
510 of Operational Definitions. *Current Developments in Nutrition*. 2020 Sep 1;4(9):nzaa127.
- 511 52. Biswas T, Townsend N, Magalhaes RJS, Hasan M, Mamun A. Patterns and determinants of
512 the double burden of malnutrition at the household level in South and Southeast Asia. *Eur J*
513 *Clin Nutr*. 2021 Feb;75(2):385–91.
- 514 53. Popkin BM, Corvalan C, Grummer-Strawn LM. Dynamics of the double burden of
515 malnutrition and the changing nutrition reality. *The Lancet*. 2020 Jan 4;395(10217):65–74.
- 516 54. Wells JC, Sawaya AL, Wibaek R, Mwangome M, Poullas MS, Yajnik CS, et al. The double
517 burden of malnutrition: aetiological pathways and consequences for health. *The Lancet*.
518 2020 Jan 4;395(10217):75–88.
- 519 55. Shrimpton R, Rokx C. The Double Burden of Malnutrition [Internet]. World Bank; 2012
520 [cited 2022 Dec 7]. 1. (Health, Nutrition and Population Discussion Papers). Available from:
521 <http://elibrary.worldbank.org/doi/abs/10.1596/27417>
- 522 56. Double burden of malnutrition [Internet]. World Health Organization. World Health
523 Organization; [cited 2022 Dec 7]. Available from: [http://www.who.int/nutrition/double-](http://www.who.int/nutrition/double-burden-malnutrition/en/)
524 [burden-malnutrition/en/](http://www.who.int/nutrition/double-burden-malnutrition/en/)

- 525 57. Jaacks LM, Kavle J, Perry A, Nyaku A. Programming maternal and child overweight and
526 obesity in the context of undernutrition: current evidence and key considerations for low-
527 and middle-income countries. *Public Health Nutrition*. 2017 May;20(7):1286–96.
- 528 58. Griffiths LJ, Smeeth L, Hawkins SS, Cole TJ, Dezateux C. Effects of infant feeding practice on
529 weight gain from birth to 3 years. *Archives of Disease in Childhood*. 2009 Aug 1;94(8):577–
530 82.
- 531 59. Asim M, Nawaz Y. Child Malnutrition in Pakistan: Evidence from Literature. *Children*. 2018
532 May;5(5):60.
- 533 60. District Government Tharparkar. Analysis of Nutritional Data Community-Based
534 Management of Acute Malnutrition Project. Sindh, Pakistan: District Government
535 Tharparkar; 2014.
- 536 61. Achakzai P, Khan R. NUTRITIONAL STATUS AND ASSOCIATED FACTORS AMONG CHILDREN
537 LESS THAN FIVE YEARS OF AGE IN TEHSIL ZARGHOON TOWN, DISTRICT QUETTA,
538 BALUCHISTAN. *J Ayub Med Coll Abbottabad*. 2016 Mar;28(1):146–51.
- 539 62. Khan GN, Turab A, Khan MI, Rizvi A, Shaheen F, Ullah A, et al. Prevalence and associated
540 factors of malnutrition among children under-five years in Sindh, Pakistan: a cross-sectional
541 study. *BMC Nutrition*. 2016 Nov 24;2(1):69.
- 542 63. Ali W, Ayub A, Hussain H. PREVALENCE AND ASSOCIATED RISK FACTORS OF UNDER
543 NUTRITION AMONG CHILDREN AGED 6 TO 59 MONTHS IN INTERNALLY DISPLACED
544 PERSONS OF JALozAI CAMP, DISTRICT NOWSHERA, KHYBER PAKHTUNKHWA. *J Ayub Med
545 Coll Abbottabad*. 2015 Sep;27(3):556–9.
- 546 64. Gul R, Kibria Z. PREVALENCE AND PREDETERMINANTS OF MALNUTRITION IN CHILDREN
547 UNDER 3 YEARS OF AGE IN THE TWO RURAL COMMUNITIES OF PESHAWAR. *Khyber Medical
548 University Journal*. 2013;5(4).
- 549 65. Trang NHHD, Hong TK, van der Ploeg HP, Hardy LL, Kelly PJ, Dibley MJ. Longitudinal
550 Sedentary Behavior Changes in Adolescents in Ho Chi Minh City. *American Journal of
551 Preventive Medicine*. 2013 Mar 1;44(3):223–30.
- 552 66. Biadgilign S, Mgutshini T, Gebremichael B, Haile D, Berhanu L, Chitekwe S, et al. Correlates
553 of Sedentary Time Among Children and Adolescents in Ethiopia: A Cross-Sectional Study.
554 *Pediatric Exercise Science*. 2022 Jul 20;1(aop):1–8.
- 555 67. Carrillo-Larco RM, Miranda JJ, Bernabé-Ortiz A. Wealth index and risk of childhood
556 overweight and obesity: evidence from four prospective cohorts in Peru and Vietnam. *Int J
557 Public Health*. 2016 May 1;61(4):475–85.

558 68. Neupane S, K.C. P, Doku DT. Overweight and obesity among women: analysis of
559 demographic and health survey data from 32 Sub-Saharan African Countries. BMC Public
560 Health. 2016 Jan 13;16(1):30.

561



medRxiv preprint doi: <https://doi.org/10.1101/2023.05.04.23289509>; this version posted May 5, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

* If an EB was greater than 300 households, then the EB was segmented (determined by the field team before household listing) and only one segment was selected to form the cluster (probability of selection was proportional to the household size)

** No replacements or changes of pre-selected households were allowed during the implementing stage.

*** Survey successfully administered in 561 out of 580 EBs, 19 clusters dropped due to security issues during the fieldwork.

Fig 1