

The impact of remote consultations on the quality of primary care: A systematic review

Kate Campbell¹, Geva Greenfield¹, Edmond Li², Niki O'Brien², Benedict Hayhoe¹, Thomas Beaney¹, Azeem Majeed¹, Ana Luisa Neves¹

¹Department of Primary Care and Public Health, Imperial College London

²Patient Safety Translational Research Centre, Institute of Global Health Innovation, Imperial College London

Word Count: 3,300 words

Corresponding author:

Dr Ana Luisa Neves¹

Department of Primary Care and Public Health, Imperial College London

Abstract

Background: The adoption of remote consultations, catalysed by the COVID-19 pandemic, has transformed the delivery of primary care services. We evaluated the impact of remote consultations on the quality of primary care.

Methods: Six databases were searched. Studies evaluating the impact of remote consultations, for any disease, were included. Title and abstract screening, and full-text screening were performed by two pairs of investigators. Risk of bias was assessed using the Mixed Methods Appraisal Tool. A narrative synthesis of the results was performed.

Findings: Thirty studies (5,469,333 participants) were included in the review. Remote consultations generally had a positive or equivalent impact compared to face-to-face (F2F) consultations, particularly in reducing patient costs and improving time efficiency. The effectiveness of remote consultations was non-inferior to F2F care in six out of seven studies evaluating this aspect. Two studies found that remote consultations reduced wait times for appointments. Younger, female patients were more likely to use remote consultations and those of lower socioeconomic status were less likely to use video consultations than telephone appointments. The impacts on safety and patient-centeredness were largely inconclusive.

Interpretation: Remote consultations may be equally as effective as F2F care and have a potentially positive impact on the efficiency and timeliness of care. Those of lower socioeconomic status were more likely to use consultations delivered via telephone than videoconference. Developing a strong evidence-base capitalising on real-world data as well as clinical trials is crucial for the future development of remote consultations and tailoring them to patient needs and preferences.

Funding: National Institute for Health and Care Research Applied Research Collaboration Northwest London.

What is already known on this topic: Existing literature reviews exploring remote consultations have primarily been confined to assessing their impact on effectiveness, efficiency, or specific clinical conditions whilst utilising a broad definition regarding what constitutes remote services. Evidence was largely heterogeneous, often focussing on interventions delivered in secondary care facilities or by specialists only. There is a paucity of systematic reviews pertaining to primary care.

What this study adds: This systematic review investigates the impact of remote consultations on the quality of primary care. Our results show that remote consultations may be equally as effective as F2F care and have a potentially positive impact on efficiency, timeliness of care, and reduced rates of follow-up in secondary or tertiary care. Patients from lower socioeconomic backgrounds were more likely to use consultations delivered via telephone than video conference.

How this study might affect research, practice or policy: Our systematic review has demonstrated that remote consultations have the potential to be just as effective as F2F consultations by reducing waiting times, patient costs, and rates of follow-up in hospitals. However, there currently remains a lack of robust studies available exploring the effect of remote consultations on patient safety, equity, and patient-centredness, highlighting areas where future research efforts need to be devoted. Data collection methods more bespoke to the primary care context, better accounting for patient characteristics and needs, and inclusive of its intended end-users, are necessary to generate a stronger evidence base to inform future remote care policies.

Introduction

The onset of the COVID-19 pandemic in 2020 resulted in the rapid expansion of remote consultations in primary care.¹ The shift from a primarily face-to-face (F2F) model of healthcare provision led to approximately 70%² and 65%³ of primary care contacts being delivered remotely in the United Kingdom (UK) and the United States (US), respectively.

Remote consultations can be defined as real-time communication between patients and clinicians, through telephone or videoconferencing.⁴ It is argued that not only have remote consultations helped to minimise COVID-19 transmission, but they may also improve efficiency and access to care.⁵ This may be particularly important in rural areas with geographical disparities in service provision⁶ and resource-constrained settings with workforce shortages.⁴

However, concerns have been raised over the speed at which remote consultations were implemented, with both patients and clinicians reporting a lack of confidence in the underlying technology and potentially poorer clinical decision-making as key issues.⁷ Remote care limits clinicians' capacity to conduct physical examinations⁸ and increase reliance on patients' abilities to articulate their symptoms,⁵ posing potential safety risks. Moreover, those with limited access to technology or with lower digital literacy may be at risk of 'digital exclusion'.⁹

Previous reviews have investigated the impact of remote consultations on the effectiveness^{3,10-13} and efficiency^{14,15} of care, often focussing on specific clinical conditions or taking broad definitions of remote services. However, there is a notable lack of systematic reviews assessing the impacts on safety, patient-centredness, timeliness and equity, with much of the existing literature limited to scoping or rapid reviews of the evidence.¹⁶⁻¹⁹ Furthermore, while some systematic reviews investigate the impact of remote care on aspects of quality exclusively in primary care settings,^{11,14,15,20} others include heterogeneous evidence, including interventions delivered in secondary care facilities or by specialists.^{3,13} There is therefore a need to comprehensively evaluate the impact on all aspects of care quality, in this specific clinical setting.

The aim of this review is to systematically evaluate the impact of remote consultations on the quality of primary care. We chose to use the Institute of Medicine (IOM)'s theoretical framework to map the impact across six domains of quality, including efficiency, effectiveness, safety, patient-centredness, timeliness and equity.²¹

Methods

This systematic review followed the Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) checklist.²² The study protocol was registered with the International Prospective Register of Systematic Reviews (CRD42022362380).

Search Strategy

Six databases (MEDLINE, Embase, HMIC, PsychInfo, CINAHL and Cochrane) were searched on 20th June 2022. The search included studies published between January 2017 to June 2022, as the last 5 years have seen most of the shift toward remote care. A combination of free text and Medical Subject Headings (MeSH) terms was used (Appendix 1). The concepts of remote consultations and primary care were kept intentionally broad to address variations in language; search terms for the domains of quality were adapted from a previous review.²³ Reference lists of relevant systematic reviews and grey literature sources were also screened.

Study Selection Criteria

Studies were included if they focused on adult patients accessing primary care services; involved telephone or videoconference consultations delivered by healthcare professionals; compared outcomes with F2F consultations; and reported outcomes that fit under any of the IOM's quality of care domains. A detailed description of inclusion and exclusion criteria is provided in Appendix 2. Studies focussing on specific health conditions were not excluded, in order to characterise the general use of remote consultations in primary care.

Screening and Data Extraction

Following the removal of duplicates, citations were uploaded into an online screening tool (Covidence²⁴). Title and abstract screening, followed by full-text screening, were performed by two independent reviewers. Cohen's kappa was used to measure intercoder agreement in each screening phase (0.22 and 0.65, respectively). Disagreements were resolved by discussion with a third investigator. Data extraction was conducted using a standardised data extraction form (Appendix 3). Effect sizes such as mean differences, odds ratios (OR) and risk ratios (RR) were extracted. Where available, rates of intervention adherence, follow-up and survey response were also extracted.

Risk of Bias Assessment

Risk of bias was assessed using the Mixed Methods Appraisal Tool (MMAT v.18)²⁵ (Appendix 4). A study was considered high risk if it scored 'Yes' in two or fewer dimensions, moderate risk if it scored 'Yes' in three dimensions, and low risk if it scored 'Yes' in four or all dimensions.

Data Synthesis

A narrative synthesis of the findings was conducted for each domain of quality.

Role of the funding source

The study funders did not play a role in the study design; collection, analysis, or interpretation of data; manuscript writing; or in the decision to submit for publication. Researchers were independent of the funders. All authors had full access to all data included in this study and can take responsibility for its integrity and the accuracy of the data analysis.

Results

Search Results

Searches retrieved a total of 6,272 records (Figure 1). No relevant records were found from searching grey literature or from reference lists of relevant articles. Thirty papers met the inclusion criteria.

Description of Included Studies

The 30 included studies (Table 1) comprised a total of 5,469,333 subjects. Sample sizes ranged from 28²⁶ to 1,490,734²⁷ participants, and publication years ranged between 2017 and 2022. Study types included 14 retrospective cohorts,²⁷⁻⁴⁰ six cross-sectional,⁴¹⁻⁴⁶ four quasi-experimental,^{26,47-49} three randomised controlled trials (RCT),⁵⁰⁻⁵² two cohorts,^{53,54} and one cluster RCT.⁵⁵ Most studies took place in the US (n=20),^{27-31,33-41,43-46,50,55} with the rest conducted in Australia,^{32,42} Canada,^{47,49} Kenya,⁵¹ the UK,⁴⁸ Japan,⁵² Singapore,²⁶ New Zealand,⁵³ and Sweden.⁵⁴

Studies considered patients with a range of health conditions (i.e. mental illnesses,^{33,37,39,50} urgent and non-emergent conditions,^{30,41} overweight and obesity,^{28,55} low back pain,²⁹ alcohol use disorders,⁵¹ nicotine dependence,⁵² hard-to-heal ulcers,⁵⁴ and acute, non-urgent conditions.³⁴ The remaining studies (n=17) considered primary care users in general. All consultations were delivered in primary care settings. Three studies specified occurring in rural locations^{45,51,55} and one was conducted in an urban, socioeconomically deprived area.⁴⁸

Summary of Risk of Bias Assessment

Nineteen studies had a moderate risk,^{26,28-30,34-36,38-40,45-50,53-55} three had a low risk,^{32,37,52} and eight had a high risk of bias^{27,31,33,41-44,51} (Figure 2). For quantitative descriptive studies, the main sources of bias were poor sample representativeness and risk of non-response bias (i.e. potential lower engagement from those with lower digital literacy).^{42,43,47} Main sources of bias for RCTs included issues with blinding,^{51,52,55} low or unclear adherence to the intervention,^{50,51,55} lack of details on randomisation,⁵¹ and differences between groups at baseline.⁵⁰ Non-randomised studies had globally a high risk of bias, stemming from uncertain accuracy of measurements of exposure and outcome,^{27-32,34,35,38,41,44-46} confounders unaccounted for,^{27,29,30,40,41,44,45} overrepresentation of certain subgroups,⁴⁹ and selection bias.^{36,38,46}

Interventions

Four studies investigated telephone consultations only^{48,49,51,55} and ten assessed consultations delivered only over videoconference, using a range of platforms (i.e., Skype, Zoom, videophone or bespoke telehealth portals).^{26,28-30,41-43,50,52,54} The remaining sixteen studies considered both telephone and videoconference as the intervention. Almost all studies involved first consultations in an episode of care; other interventions consisted of consultations including behavioural therapy,^{39,50,55} motivational interviewing,⁵¹ and smoking cessation counselling.⁵²

Table 1. Characteristics of the included studies.

Author, year	Country	Time period	Study design	Participants (setting)	Sample size (N)	Study design	Consultation description	Risk of bias
Baughman, 2022a ²⁸	USA	April 2020 - September 2021	Retrospective cohort study	Primary care patients aged 26-70 with abnormal BMI scores (<18.5 or >25 (kg/m ²))	287,387(RC: 1,556; blended: 63,489; F2F: 222,333)	Comparison of outcomes between initial visit modalities: <ul style="list-style-type: none"> ▪ RC (Zoom video) ▪ blended RC (patients with RC and F2F visits within the timeframe) ▪ F2F 	Initial BMI screening visits	Moderate
Baughman, 2022b ²⁹	USA	July 2019 - June 2021	Retrospective cohort study	Primary care patients aged 18-50 with a lower back pain diagnosis	20,624 (RC: 5,334; F2F: 15,290)	Comparison of outcomes between initial visit modalities: <ul style="list-style-type: none"> ▪ RC (Zoom video) ▪ F2F 	Initial primary care consultation	Moderate
Befort, 2021 ⁵⁵	USA	February 2016 - October 2017	Cluster RCT	Primary care patients aged 20-75 (mean age 54.7) with BMI scores of 30 - 45, residing in rural locations	1,407 (group RC: 466; individual F2F: 473; group F2F: 468)	Practices (n=36) randomly assigned to: <ul style="list-style-type: none"> ▪ group RC (telephone conference call) ▪ group F2F ▪ individual F2F 	Individual F2F: 15-minute behavioural therapy; Group RC and F2F: 60-minute group behavioural therapy (14 patients per group)	Moderate

Author, year	Country	Time period	Study design	Participants (setting)	Sample size (N)	Study design	Consultation description	Risk of bias
Bernstein, 2021 ³⁰	USA	November 2015 - March 2019	Retrospective cohort study	Ambulatory care patients aged >60 with urgent and non-emergent conditions	1088(RC: 115; F2F: 973)	Comparison of outcomes between index visit modalities: <ul style="list-style-type: none"> ▪ RC (video) ▪ F2F 	Visit with physician including clinical assessment, prescriptions, or referrals if appropriate	Moderate
Chavez, 2022 ³¹	USA	April 2019 - March 2021	Retrospective cohort study	Patients (mean age 54.1) accessing care at an academic family medicine practice	57006 ^a (RC: 7,577; F2F: 49,429)	Comparison of outcomes between index visit modalities: <ul style="list-style-type: none"> ▪ RC (video or telephone) ▪ F2F Stratified by pre-pandemic and pandemic timeframes	Consultation with physician	High
Dai, 2022 ³²	Australia	March 2020 - August 2021	Retrospective cohort study	Geriatric primary care patients aged >65 in residential aged care facilities	27,980	Assessed associations between sociodemographic characteristics and visit modality: <ul style="list-style-type: none"> ▪ RC (video or telephone) ▪ F2F 	GP consultation	Low
Egede, 2017 ⁵⁰	USA	September 2006 - October 2012	RCT	Elderly veterans aged >58 with depression	241 (RC: 120; F2F: 121)	Comparison of outcomes between consultation modalities:	60-minute behavioural activation for depression	Moderate

Author, year	Country	Time period	Study design	Participants (setting)	Sample size (N)	Study design	Consultation description	Risk of bias
						<ul style="list-style-type: none"> ▪ RC (videophone) ▪ F2F 	delivered weekly for 8 weeks	
Frank, 2021 ³³	USA	March 2019 - December 2020	Retrospective cohort study	Mental health patients at an academic primary care practice aged 4 – 73 (mean age 28.32)	173	Comparison of outcomes between two waves of consultation modalities: <ul style="list-style-type: none"> ▪ RC (video or telephone): Mar - Dec 20 ▪ F2F: Mar - Dec 19 	30-minute mental health appointments	High
Gordon, 2017 ³⁴	USA	January 2014 - May 2015	Retrospective cohort study	Primary care patients aged <65 receiving care for acute, nonurgent conditions ^b	18,516 (RC: 4,635; F2F: 13,881)	Comparison of outcomes between consultation modalities: <ul style="list-style-type: none"> ▪ RC ▪ F2F 	Primary care consultations	Moderate
Govier, 2022 ³⁵	USA	March 2019 - July 2021	Retrospective cohort study	Primary care patients aged >18 diagnosed with COVID-19 between March - July 2020	11,326 (RC:1,360; F2F: 9,966)	Assessed associations between sociodemographic characteristics and visit modality: <ul style="list-style-type: none"> ▪ RC ▪ F2F 	Primary care consultations	Moderate
Graetz, 2022 ³⁶	USA	January 2016 - May 2018	Retrospective cohort study	Primary care patients of all ages (22.2% >65)	1,131,722	Comparison of outcomes between consultation modalities:	Primary care consultations requested by patients using an	Moderate

Author, year	Country	Time period	Study design	Participants (setting)	Sample size (N)	Study design	Consultation description	Risk of bias
						<ul style="list-style-type: none"> ▪ RC (video) ▪ RC (telephone) ▪ F2F 	online portal	
Haderlein, 2022³⁷	USA	March 2018 - October 2021	Retrospective cohort study	Veterans (mean age 48.7) seeking mental health care in an urban VA Primary Care-Mental Health Integration clinic	2,479	Comparison of outcomes between consultation modalities: <ul style="list-style-type: none"> ▪ RC (video or telephone) ▪ F2F 	Primary care mental health consultations	Low
Harder, 2020⁵¹	Kenya	September 2014 - December 2015	RCT	Primary care patients (mean age 38) with alcohol use disorders in rural primary health centre	300 (RC:104; F2F: 92; Waitlist control: 104)	Comparison of outcomes between consultation modalities: <ul style="list-style-type: none"> ▪ RC (mobile phone call) ▪ F2F ▪ Waitlist control 	One 30-minute motivational interviewing session	High
Li, 2022²⁷	USA	June 2019 - September 2020	Retrospective cohort study	Primary care patients (mean age 39.2)	1,490,734	Comparison of outcomes between practice level of RC use: <ul style="list-style-type: none"> ▪ High RC use ▪ Medium RC use ▪ Low RC use 	Primary care consultations	High

Author, year	Country	Time period	Study design	Participants (setting)	Sample size (N)	Study design	Consultation description	Risk of bias
Lovell, 2021 ⁴¹	USA	April 2016 - March 2017	Retrospective cohort study	Primary care patients aged <64 accessing care for low-acuity, urgent conditions ^c	5,919 (RC: 1,531; F2F: 4,388)	Comparison of outcomes between consultation modalities: <ul style="list-style-type: none"> ▪ RC (videocall) ▪ F2F 	Primary care consultations	High
Manski-Nankervis, 2022 ⁴²	Australia	October 2020 -May 2021	Descriptive cross-sectional study	Primary care patients (mean age 31.8) who completed a videoconference call with a health care professional	499	Online survey of experience with most recent RC visit (videoconference) with comparison to past experience of F2F visit	Primary care consultations	High
McGrail, 2017 ⁴⁷	Canada	2013 - 2014	Quasi-experimental (interrupted time series) with cross-sectional survey component	Primary care patients aged >18 living in British Columbia	29,267 (RC: 7,286; F2F: 21,981) Survey: 399	Comparison of outcomes between consultation modalities: <ul style="list-style-type: none"> ▪ RC ▪ F2F Online survey of experience with most recent RC visit with comparison to past experience of F2F visit	Primary care consultations	Moderate
Miller, 2019 ⁴⁸	UK	June 2014 – May 2017	Quasi-experimental (interrupted time series)	Primary care patients of all ages at an urban general practice in a socio-economically	27,589 (RC: 9,113; F2F: 18,476)	Comparison of outcomes between the two study phases: <ul style="list-style-type: none"> ▪ F2F (preintervention phase) 	Primary care consultations	Moderate

Author, year	Country	Time period	Study design	Participants (setting)	Sample size (N)	Study design	Consultation description	Risk of bias
				deprived area		<ul style="list-style-type: none"> RC (telephone-first phase) 		
Mohan, 2022⁴³	USA	April - December 2020	Descriptive cross-sectional study	Primary care patients (mean age 48.7) at academic medical centre	797	Online survey of experience with most recent RC visit with comparison to past experience of F2F visit	Primary care consultations	High
Neufeld, 2022⁴⁹	Canada	September 2020 - February 2021	Quasi-experimental study	Primary care patients aged 18-87)	66 (RC: 32; F2F: 34)	Comparison of outcomes between consultation modalities: <ul style="list-style-type: none"> RC (telephone) F2F Assessed by online survey	Family physician consultations	Moderate
Nomura, 2019⁵²	Japan	March - June 2018	RCT	Primary care patients (mean age 55) with nicotine dependence	115 (RC: 58; F2F: 57)	Comparison of outcomes between consultation modalities: <ul style="list-style-type: none"> RC (internet-based video call) F2F 	Smoking cessation counselling, 5 sessions over 24 weeks with access to smoking cessation mobile app and an exhaled CO checker	Low
Pierce, 2020⁴⁴	USA	March - April 2020	Analytical cross-sectional study	Primary care patients (mean age 45) at an academic family medicine centre	6,984	Assessed associations between sociodemographic characteristics and visit modality:	Family medicine consultations	High

Author, year	Country	Time period	Study design	Participants (setting)	Sample size (N)	Study design	Consultation description	Risk of bias
						<ul style="list-style-type: none"> RC (audio or video) F2F 		
Quinton, 2021⁴⁵	USA	March 2019 - March 2021	Analytical cross-sectional study	Patients aged >18 presenting for ambulatory visits at a rural healthcare provider	54,559	Assessed associations between patient characteristics and visit modality: <ul style="list-style-type: none"> RC (audio or video) F2F 	Ambulatory care visits	Moderate
Reed, 2020⁴⁶	USA	January 2016 - May 2018	Analytical cross-sectional study	Primary care patients of all ages at a large integrated health care delivery system	1,131,722	Assessed associations between patient characteristics and visit modality: <ul style="list-style-type: none"> RC (telephone or video) F2F 	Index primary care consultations (no visits within 7 days prior)	Moderate
Reed, 2021³⁸	USA	January 2016 - May 2018	Retrospective cohort study	Primary care patients of all ages (mean age 43) at a large integrated health care delivery system	1,131,722	Comparison of outcomes between consultation modalities: <ul style="list-style-type: none"> RC (telephone or video) F2F 	Index primary care consultations (no visits within 7 days prior)	Moderate
Rene, 2022³⁹	USA	October-2019 - May 2020	Retrospective cohort study	Primary care patients aged >18 with depression and/or anxiety	338 (RC: 181; F2F: 157)	Comparison of outcomes between consultation modalities: <ul style="list-style-type: none"> RC (pandemic 	Initial behavioural health consultation, followed by 30-	Moderate

Author, year	Country	Time period	Study design	Participants (setting)	Sample size (N)	Study design	Consultation description	Risk of bias
						cohort: Apr – May 20) <ul style="list-style-type: none"> F2F (pre-pandemic cohort: Oct– Nov 19) 	minute visits for behavioural activation, motivational interviewing, and psycho-education	
Ryskina, 2021⁴⁰	USA	March 2020 - May 2020	Retrospective cohort study	Primary care patients aged >65 (mean age 75.1)	17,103 (RC: 10,311; F2F: 6,792)	Comparison of outcomes between consultation modalities: <ul style="list-style-type: none"> RC (telephone or video) F2F 	Primary care consultations	Moderate
Tan, 2020²⁶	Singapore	April 2019 - May 2019	Quasi-experimental study (prospective, self-controlled)	Active military servicemen (93% aged 18-24) accessing primary care at a military medical centre	28	Comparison of outcomes between consultation modalities: <ul style="list-style-type: none"> RC (on-premises Zoom video) F2F (immediately following RC) 	Primary care consultations with assistant present to perform auscultations and physical examinations Symptom collection app used prior to RC and data available to the RC physician	Moderate
Ure, 2022⁵³	New Zealand	May - July 2021	Cohort study	Primary care patients of all ages (25% aged <5) at a general	454 (RC: 133; F2F: 321)	Comparison of outcomes between consultation modalities:	Primary care consultations following triage by telephone	Moderate

Author, year	Country	Time period	Study design	Participants (setting)	Sample size (N)	Study design	Consultation description	Risk of bias
				practice medical centre		<ul style="list-style-type: none"> ▪ RC ▪ F2F 		
Wickstrom, 2018⁵⁴	Sweden	October 2014 - September 2016	Cohort study	Primary care patients (RC mean age 77; F2F mean age 75) with hard-to-heal ulcers	Healing time study: 1,988 (RC: 100; F2F: 1,888) Waiting time study: 200 (RC: 100; F2F: 100)	Comparison of outcomes between consultation modalities: <ul style="list-style-type: none"> ▪ RC (Skype video) ▪ F2F 	Consultations involved ulcer diagnosis and discussion of treatment strategy	Moderate

BMI, body mass index; CO, carbon monoxide; F2F, face-to-face; GP, general practitioner; NA, not applicable; RCT, randomised controlled trial; RC, remote consultation; UK, United Kingdom; USA, United States of America

^a Number of index visits; number of participants not reported

^b Conditions included: sinusitis, upper respiratory infection, urinary tract infection, conjunctivitis, bronchitis, pharyngitis, influenza, cough, dermatitis, digestive symptom, or ear pain

^c Conditions included: sinusitis, conjunctivitis, urinary tract infection, upper respiratory infection, influenza/ pneumonia, bronchitis, dermatitis/ eczema, ear pain, digestive symptoms, and cough

Outcomes

Efficiency

Sixteen papers evaluated outcomes related to efficiency,^{26,27,29-31,33,34,38-42,47,48,50,53} including rates of follow-up visits and hospitalisations, patient costs, and appointment characteristics (i.e., length, attendance, cancellations, and no-shows) (Appendix 5). Eleven out of sixteen papers found a positive (n=6) or no impact (n=5) on efficiency in at least half of the outcomes extracted.^{26,29,33,34,38,40-42,47,48,50} Of the eight studies comparing rates of follow-up visits in primary care, five found that remote consultations resulted in a greater need for additional care,^{30,31,38,41,53} while three found no differences.^{34,47,48}

Out of seven studies evaluating the rates of follow-up consultations in secondary or tertiary care, three found no changes in rates of emergency department (ED) follow-up visits or hospitalisations.^{38,41,48} Another three studies^{34,40,47} found a significant reduction in follow-up visits after a remote consultation (including ED visits,³⁴ hospitalisations^{34,40} and ambulatory care sensitive conditions (ACSC) visits⁴⁰). Only one study found that high use of remote consultations was associated with more annual ACSC visits compared to practices with low use.²⁷

In terms of appointment characteristics, one study found that remote consultations led to a higher number of appointments attended and fewer cancellations.³³ However, another reported lower rates of attendance and increased cancellations in the context of mental health appointments that took place remotely during the pandemic.³⁹ One paper found that the use of remote consultation and a symptoms checker application resulted in a shorter appointment duration when compared with F2F care.²⁶ Lastly, four out of the five papers assessing the impact on patient costs demonstrated a reduction in the costs associated with remote consultations compared to F2F visits.^{34,41,42,47}

Effectiveness

Seven studies assessed the effectiveness of remote consultations (Appendix 6).^{28,33,39,51,52,54,55} Six found a non-inferior impact on effectiveness for at least half of the outcomes.^{33,39,51,52,54,55} Two studies investigating the effectiveness of remote mental health care in improving anxiety and depression symptoms also demonstrated its non-inferiority³⁹ or superiority³³ to F2F care. Remote consultations were found to be more effective for the care of hard-to-heal ulcers,⁵⁴ and equally as effective as F2F care in reducing alcohol consumption,⁵¹ abstinence from smoking,⁵² and for weight management.⁵⁵

Safety

Only a small study (n=28) considered outcomes related to the safety of care, finding an overall diagnostic agreement rate of 92% between remote and F2F assessments.²⁶ Agreement rate was 100% for headache, gastroenteritis, and conjunctivitis, but lower for dermatological conditions and upper respiratory tract infections (87.5% and 93.3%, respectively).²⁶

Patient-Centredness

Four of five studies assessing the impact on patient-centredness indicated that remote care had a positive or equivalent effect compared to F2F care (Appendix 7).^{26,42,43,47} One study found that those seen remotely reported lower perceived autonomy support.⁴⁹ Three studies asked patients to compare their recent teleconsultations with past experiences of F2F care,^{42,43,47} with most respondents agreeing that remote care was 'as good' (84%)⁴² and 'as thorough' (79%)⁴⁷ as F2F care; more convenient (91%) and of equal or better value (67%).⁴³ In another study, the majority of patients (39.9%) had no preference regarding consultation modality.²⁶

Timeliness

Two out of the three studies evaluating the impact on timeliness found an improvement when consultations were delivered remotely (Appendix 8).^{36,54} Graetz et al³⁶ reported that both video and

telephone consultations were more likely to occur within one day of scheduling. Similarly, a study at a wound healing clinic found that remote consultations took place significantly sooner after referral.⁵⁴ In contrast, a study at a Primary Care Mental Health clinic reported that patients who were initially assessed remotely were less likely to receive same-day mental health care.³⁷

Equity

Six studies assessed the impact on equity of care in terms of the utilisation of services by sex, age, ethnicity, socio-economic status (SES) and rural or urban residence (Appendix 9).^{32,35,40,44-46} Two studies found that women were more likely to use remote care than men.^{44,46} Regarding age, two papers reported that the likelihood of having remote consultations decreased with increasing age.^{32,46} Interestingly, one study performed in the first month of the pandemic found that those over 65 were more likely to have remote consultations compared to those aged 18 to 44.⁴⁴

Lower use of remote consultations was reported to be associated with both lower^{45,46} and higher SES.^{32,44} Notably, two studies found that those of lower SES were less likely to have video consultations than they were to have telephone consultations.^{44,46}

Findings regarding the impact of ethnicity on the utilisation of remote care were also mixed. Two studies suggested that black patients were less likely to use remote consultations than white patients,^{44,45} while another three found the opposite effect.^{35,40,46} Asian patients residing in the US were more likely to use video consultations than white patients but slightly less likely to use telephone consultations.⁴⁶

Lastly, a US study reported that patients living in rural areas had lower remote care use,⁴⁴ whereas an Australian study found the inverse effect.³²

Table 2. Summary of main findings

Domain of quality	Main findings
Efficiency	<ul style="list-style-type: none"> Remote consultations may reduce rates of hospitalisation and follow-up in secondary care, but may increase rate of follow-up in primary care compared to F2F consultations Remote care may lead to lower overall patient spending and be more time-efficient than F2F care
Effectiveness	<ul style="list-style-type: none"> Treatment delivered remotely is as effective in improving clinical outcomes as F2F care, particularly for psychological or behaviour-related conditions treated in primary care
Safety	<ul style="list-style-type: none"> Remote consultations delivered over videoconference may have similar diagnostic accuracy to F2F for most conditions There is a lack of studies investigating other aspects of safety, such as medication safety incidents, highlighting an important gap in knowledge
Patient-Centredness	<ul style="list-style-type: none"> Patients indicate that remote consultations are more convenient and are of similar value and quality to F2F consultations, though findings may be at risk of bias Remote consultations may reduce patients' perceived autonomy support compared to F2F care
Timeliness	<ul style="list-style-type: none"> Remote appointments may have lower wait times than F2F appointments
Equity	<ul style="list-style-type: none"> Women are more likely to use remote consultations than men

- The use of remote care declines with increasing age
- The impacts of SES, location of residence and ethnicity vary considerably between studies and are likely influenced by numerous contextual factors

Discussion

Summary of main findings

Our findings suggest that remote consultations are equally or more effective than F2F care for the management of conditions including mental illness, excessive smoking, and alcohol consumption. The evidence for the impact on clinical safety is extremely limited. Four studies indicated positive impacts on some aspects of patient-centredness, however, a negative impact was noted on patients' perceived autonomy support (i.e., the degree to which people perceive others in positions of authority to be autonomy supportive). Remote consultations may reduce waiting times, lower patient costs, and reduce rates of follow-up in secondary and tertiary care. However, there is evidence that remote consultations may increase the need for additional GP visits compared with those seen in person. Evidence regarding equity was considerably mixed. Overall, it appears that remote care is more likely to be used by younger, female patients, with disparities between other subgroups depending on contextual factors.

Interpretation of Findings in the Context of Previous Research

Efficiency

The indication that remote consultations may increase rates of follow-up visits in primary care is consistent with previous evidence.^{3,14} This negative impact on efficiency may to some extent be explained by the timeframes during which the studies took place. In those occurring shortly after the onset of the pandemic,^{31,53} when the rapid transition to remote care was necessary, the increased rates of follow-up may be a consequence of lower clinician or patient confidence in remote care due to its initial unfamiliarity.³⁴

Previous reviews have not specifically assessed the impact of remote consultations on follow-up at secondary or tertiary levels of care. This review's finding that they may reduce or have no impact on follow-up at these higher levels of care might be explained by retrospective study designs precluding adequate adjustment for confounders and by the heterogeneity of the interventions included.^{30,34,38,40}

The finding that remote care may be associated with lower patient costs is in line with previous research.^{3,15} Although this evidence is most relevant to countries in which patients pay for services out-of-pocket, it appears that patients accessing publicly funded health systems may also benefit financially from remote care, mainly due to reductions in travel expenses in time costs from loss of work.^{3,41,42} Furthermore, this review and wider evidence indicate that remote consultations are generally shorter than F2F visits.^{3,14,26} The decrease in consultation length reported by Tan et al²⁶ may be explained by the use of a symptom checker application prior to the teleconsultation, which could have improved efficiency for clinicians. However, it is so far unclear whether shorter appointments are indeed more cost-effective than longer appointments, and whether they allow enough time for discussion of more complex matters.

Effectiveness

Remote care seems to be as effective as F2F visits for certain clinical outcomes (i.e., depression and anxiety symptoms, alcohol use disorder scores, smoking abstinence rates, and ulcer healing times). Existing reviews have similarly found non-inferior outcomes when remote care was delivered by specialists,^{3,10} combined with remote patient monitoring,¹¹ or offered in addition to F2F care.¹³ This study suggests that remote consultations may be an effective substitute for F2F consultations in primary care settings.

Patient-centredness

The wider literature has reported similarly positive findings in terms of patient satisfaction across various types of secondary care and remote patient monitoring for diabetes.^{16,17} One possible explanation is that the use of remote consultations can reduce time pressure for physicians, allowing them to provide more patient-centric consultations. However, the notable positive impact on patient-centredness (i.e., convenience and preference) should be interpreted with caution, due to the significant risk of bias in these studies and the heterogeneity of measures used. Many of the included studies relied on non-validated surveys to assess satisfaction. In this review, the only study using validated questionnaires found that remote care led to lower perceived autonomy support, possibly due to the absence of non-verbal cues and decreased relational competence.⁴⁹

Timeliness

There is some evidence that opting for remote care may reduce wait times for initial consultations compared to F2F care,^{36,54} possibly facilitated by removing the barriers of job flexibility and travel times.³⁶ Shorter wait times is a key benefit of remote care perceived by patients.¹⁹ However, the finding that wait times for mental healthcare were increased following a remote appointment may reflect the possibility that patients with lower clinical need chose remote consultations to begin with, (and thus their concerns were assessed as less urgent) or indicate logistical issues in transitions of care.³⁷

Equity

Our findings regarding age and sex are in line with previous research,²⁰ and reflect older patients' lower average digital literacy and access to technology.⁴⁴ Evidence concerning the impacts of SES, ethnicity and location of residence on the use of remote care was inconsistent across studies, potentially due to differences in populations and study settings. To this end, access and utilisation of care is highly context-specific and will be shaped by both community and practice-level features. As engagement and participation in care is generally lowest for socially disadvantaged populations, disentangling the patient characteristics that may exclude them from remote care services, at the local level, is essential.

Safety

Limited conclusions can be drawn regarding the safety of remote consultations as only one study investigating outcomes in this domain was identified in this review.

Strengths and Limitations

This study provides insight into the recent changes in the delivery of primary care and the impact of remote consultations on care quality. The review uses the IOM's comprehensive quality framework and maps findings to this model, lending a structured approach to the evaluation of the impact of remote consultations.

This review did not consider paediatric populations, care delivered by non-healthcare professionals (e.g., community health workers), or outcomes related to medication prescribing. Many of the included studies consider patients who are part of specific subpopulations, such as elderly veterans⁵⁰ or young military servicemen,²⁶ which may limit the generalisability of the findings.

Eligible studies were restricted to those published within the last five years; while this timeframe was considered reasonable in line with the changing healthcare landscape, this might have potentially led to some earlier papers being missed.

All but one of the included studies were from high-income countries and most were from the US, highlighting the lack of research from low and middle-income countries. Many of the findings of this review will therefore have limited relevance outside of the US, and certainly outside of high-income countries with dissimilar health system financing structures or technological infrastructure. The lack of

papers evaluating safety aspects, such as medication safety incidents, demonstrates the gap in knowledge previously highlighted by Gleeson et al.¹⁸ Similarly, current evidence has common limitations of bias introduced through lack of adjustment for confounders, and selection bias towards inclusion of patients with greater digital literacy or engagement with healthcare.

Finally, the apparent lack of studies investigating the effectiveness of remote care for a wider range of morbidities in primary care highlights a key area for further research.

Implications for research, policy, and practice

While evidence of improved efficiency is likely to keep driving the implementation of remote care, it is critical to ensure that the transition to new service delivery models does not pose additional patient harm. The apparent lack of studies investigating the safety of remote consultations highlights a concerning gap in the literature, and future evaluations should focus on the evaluation of diagnostic error and medication safety in this context. Furthermore, health technology assessments investigating the impact on patient-centredness should capitalise on the use of validated patient-reported measures whenever possible, to allow a rigorous comparative approach.

Policy efforts to support improvements in data collection in primary care (i.e., consultation type, duration, and quality outcomes) will be critical to developing a strong evidence-base capitalising on real-world data. The mixed findings on the impact on equity highlight the need for investigations at the local level, which will be vital to develop context-specific strategies, tailored to community health needs and characteristics. Data collection should adopt an intersectional approach, considering a breadth of patient characteristics, to inform the design of locally appropriate interventions and ensure equitable access to care. Importantly, remote consultation interventions and access schemes must incorporate participatory approaches in their research and design, encouraging input from marginalised voices and including community knowledge, values, and preferences in decision-making processes.

Acknowledgments

This work is independent research supported by the National Institute for Health and Care (NIHR) Research Applied Research Collaboration Northwest London. EL and ALN are funded by NIHR Patient Safety Research Collaborative, with infrastructure support from Imperial NIHR Biomedical Research Centre. The views expressed in this publication are those of the author(s) and not necessarily those of the National Institute for Health Research or the Department of Health and Social Care.

Data availability statement

The search strategy and extracted data contributing to the narrative synthesis are available as Supplementary Data. Any additional data are available on request.

Contributor statement

ALN and KC contributed to the conception and design of the study. KC, EL, NO, and ALN contributed to the literature search and data extraction. KC, GG, TB, BH, and ALN contributed to data analysis and interpretation. All authors contributed to writing the manuscript and critical revision. All authors approved the manuscript. KC and ALN guarantee the integrity of the work.

Declaration of interests

KC, GG, EL, NO, TB, AM and ALN have nothing to disclose. BH is an employee of eConsult Health Ltd, a provider of electronic consultations for NHS primary, secondary and urgent / emergency care.

References

1. Dhaliwal JK, Hall TD, LaRue JL, Maynard SE, Pierre PE, Bransby KA. Expansion of telehealth in primary care during the COVID-19 pandemic: benefits and barriers. *Journal of the American Association of Nurse Practitioners*. 2022;34(2):224-9.
2. Royal College of General Practitioners. GP consultations post-COVID should be a combination of remote and face to face, depending on patient need, says College. Royal College of General Practitioners; 2021. Available from: <https://www.rcgp.org.uk/news/2021/may/gp-consultations-post-covid> [Accessed July 10, 2022].
3. Carrillo de Albornoz S, Sia K-L, Harris A. The effectiveness of teleconsultations in primary care: systematic review. *Family Practice*. 2022;39(1):168-82.
4. Kadir MA. Role of telemedicine in healthcare during COVID-19 pandemic in developing countries. *Telehealth and Medicine Today*. 2020;5(2).
5. Colbert GB, Venegas-Vera AV, Lerma EV. Utility of telemedicine in the COVID-19 era. *Reviews in cardiovascular medicine*. 2020;21(4):583-7.
6. Flumignan CDQ, Rocha APd, Pinto ACPN, Milby KMM, Batista MR, Atallah ÁN, et al. What do Cochrane systematic reviews say about telemedicine for healthcare? *Sao Paulo Medical Journal*. 2019;137:184-92.
7. Verma P, Kerrison R. Patients' and physicians' experiences with remote consultations in primary care during the COVID-19 pandemic: a multi-method rapid review of the literature. *BJGP open*. 2022;6(2).
8. Lopez AM, Lam K, Thota R. Barriers and facilitators to telemedicine: can you hear me now? *American Society of Clinical Oncology Educational Book*. 2021;41:25-36.
9. Pagliari C. Digital health and primary care: Past, pandemic and prospects. *Journal of global health*. 2021;11.
10. Scott AM, Bakht M, Greenwood H, Cardona M, Clark J, Krzyzaniak N, et al. Real-time telehealth versus face-to-face management for patients with PTSD in primary care: a systematic review and meta-analysis. *The Journal of Clinical Psychiatry*. 2022;83(4):41146.
11. Mabeza RMS, Maynard K, Tarn DM. Influence of synchronous primary care telemedicine versus in-person visits on diabetes, hypertension, and hyperlipidemia outcomes: a systematic review. *BMC primary care*. 2022;23(1):1-10.
12. Batsis JA, DiMilia PR, Seo LM, Fortuna KL, Kennedy MA, Blunt HB, et al. Effectiveness of ambulatory telemedicine care in older adults: a systematic review. *Journal of the American Geriatrics Society*. 2019;67(8):1737-49.
13. Flodgren G, Rachas A, Farmer AJ, Inzitari M, Shepperd S. Interactive telemedicine: effects on professional practice and health care outcomes. *Cochrane Database of Systematic Reviews*. 2015(9).
14. Downes MJ, Mervin MC, Byrnes JM, Scuffham PA. Telephone consultations for general practice: a systematic review. *Systematic reviews*. 2017;6(1):1-6.
15. De Guzman KR, Snoswell CL, Caffery LJ, Smith AC. Economic evaluations of videoconference and telephone consultations in primary care: A systematic review. *Journal of Telemedicine and Telecare*. 2021:1357633X211043380.

16. Agbali R, Balas AE, Beltrame F, De Leo G. A review of audiovisual telemedicine utilization and satisfaction assessment during the COVID-19 pandemic. *International Journal of Technology Assessment in Health Care*. 2022;38(1).
17. Sim R, Lee SWH. Patient preference and satisfaction with the use of telemedicine for glycemic control in patients with type 2 diabetes: a review. *Patient preference and adherence*. 2021;15:283.
18. Gleeson LL, Clyne B, Barlow JW, Ryan B, Murphy P, Wallace E, et al. Medication Safety Incidents Associated with the Remote Delivery of Primary Care: A Rapid Review. *medRxiv*. 2022.
19. Lindenfeld Z, Berry C, Albert S, Massar R, Shelley D, Kwok L, et al. Synchronous Home-Based Telemedicine for Primary Care: A Review. *Medical Care Research and Review*. 2022:10775587221093043.
20. Parker RF, Figures EL, Paddison CA, Matheson JI, Blane DN, Ford JA. Inequalities in general practice remote consultations: a systematic review. *BJGP open*. 2021;5(3).
21. Wolfe A. Institute of Medicine report: crossing the quality chasm: a new health care system for the 21st century. *Policy, Politics, & Nursing Practice*. 2001;2(3):233-5.
22. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Aki EA, Brennan SE, Chou R. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *International journal of surgery*. 2021 Apr 1;88:105906.
23. Neves AL, Freise L, Laranjo L, Carter AW, Darzi A, Mayer E. Impact of providing patients access to electronic health records on quality and safety of care: a systematic review and meta-analysis. *BMJ quality & safety*. 2020;29(12):1019-32.
24. Covidence systematic review software, Veritas Health Innovation, Melbourne, Australia; 2022. Available from: www.covidence.org [Accessed June 2022].
25. Hong QN, Pluye P, Fàbregues S, Bartlett G, Boardman F, Cargo M, et al. Mixed methods appraisal tool (MMAT), version 2018. *Registration of copyright*. 2018;1148552(10).
26. Tan N-G, Yang LW-Y, Tan MZ-W, Chng J, Tan MH-T, Tan C. Virtual care to increase military medical centre capacity in the primary health care setting: A prospective self-controlled pilot study of symptoms collection and telemedicine. *Journal of telemedicine and telecare*. 2020:1357633X20959579.
27. Li KY, Ng S, Zhu Z, McCullough JS, Kocher KE, Ellimoottil C. Association Between Primary Care Practice Telehealth Use and Acute Care Visits for Ambulatory Care–Sensitive Conditions During COVID-19. *JAMA Network Open*. 2022;5(3):e225484-e.
28. Baughman D, Baughman K, Jabbarpour Y, Waheed A. Comparable quality performance between telemedicine and office-based care for abnormal BMI screening and management. *Obesity Science and Practice*. 2022.
29. Baughman D, Ptasincki A, Baughman K, Buckwalter N, Jabbarpour Y, Waheed A. Comparable Quality Performance of Acute Low-Back Pain Care in Telemedicine and Office-Based Cohorts. *Telemedicine journal and e-health : the official journal of the American Telemedicine Association*. 2022.
30. Bernstein P, Ko KJ, Israni J, Cronin AO, Kurliand MM, Shi JM, et al. Urgent and non-emergent telehealth care for seniors: Findings from a multi-site impact study. *Journal of telemedicine and telecare*. 2021:1357633X211004321.

31. Chavez A, Pullins C, Yennie J, Singh C, Kosiorek H. Association of Telehealth with Short-Interval Follow-Up. *Journal of the American Board of Family Medicine : JABFM*. 2022;35(3):485-90.
32. Dai Z, Sezgin G, Li J, Franco GS, McGuire P, Datta S, et al. Telehealth utilisation in residential aged care facilities during the COVID-19 pandemic: A retrospective cohort study in Australian general practice. *Journal of telemedicine and telecare*. 2022:1357633X221094406.
33. Frank HE, Grumbach NM, Conrad SM, Wheeler J, Wolff J. Mental health services in primary care: Evidence for the feasibility of telehealth during the COVID-19 pandemic. *Journal of affective disorders reports*. 2021;5:100146.
34. Gordon AS, Adamson WC, DeVries AR. Virtual Visits for Acute, Nonurgent Care: A Claims Analysis of Episode-Level Utilization. *Journal of medical Internet research*. 2017;19(2):e35.
35. Govier DJ, Cohen-Cline H, Marsi K, Roth SE. Differences in access to virtual and in-person primary care by race/ethnicity and community social vulnerability among adults diagnosed with COVID-19 in a large, multi-state health system. *BMC health services research*. 2022;22(1):511.
36. Graetz I, Huang J, Muelly E, Gopalan A, Reed ME. Primary Care Visits Are Timelier When Patients Choose Telemedicine: A Cross-Sectional Observational Study. *Telemedicine journal and e-health : the official journal of the American Telemedicine Association*. 2022.
37. Haderlein TP, Dobalian A, Raja PV, Der-Martirosian C. Association Between Virtual Care Use and Same-Day Primary Care Access in VA Primary Care-Mental Health Integration. *Journal of primary care & community health*. 2022;13:21501319221091430.
38. Reed M, Huang J, Graetz I, Muelly E, Millman A, Lee C. Treatment and Follow-up Care Associated With Patient-Scheduled Primary Care Telemedicine and In-Person Visits in a Large Integrated Health System. *JAMA network open*. 2021;4(11):e2132793.
39. Rene R, Cherson M, Rannazzisi A, Felter J, Silverio A, Cunningham AT. Transitioning from In-Person to Telemedicine Within Primary Care Behavioral Health During COVID-19. *Population health management*. 2022.
40. Ryskina KL, Shultz K, Zhou Y, Lautenbach G, Brown RT. Older adults' access to primary care: Gender, racial, and ethnic disparities in telemedicine. *Journal of the American Geriatrics Society*. 2021;69(10):2732-40.
41. Lovell T, Albritton J, Dalto J, Ledward C, Daines W. Virtual vs traditional care settings for low-acuity urgent conditions: An economic analysis of cost and utilization using claims data. *Journal of telemedicine and telecare*. 2021;27(1):59-65.
42. Manski-Nankervis J-A, Davidson S, Hiscock H, Hallinan C, Ride J, Lingam V, et al. Primary care consumers' experiences and opinions of a telehealth consultation delivered via video during the COVID-19 pandemic. *Australian journal of primary health*. 2022;28(3):224-31.
43. Mohan S, Lin W, Orozco FR, Robinson J, Mahoney A. Patient Perceptions of Video Visits in a Fee-for-Service Model. *Journal of the American Board of Family Medicine : JABFM*. 2022;35(3):497-506.
44. Pierce RP, Stevermer JJ. Disparities in use of telehealth at the onset of the COVID-19 public health emergency. *Journal of telemedicine and telecare*. 2020:1357633X20963893.
45. Quinton JK, Ong MK, Vangala S, Tettele-Burns A, Webb A, Sarkisian C, et al. The Association of Broadband Internet Access and Telemedicine Utilization in rural Western Tennessee: an observational study. *BMC Health Services Research*. 2021;21(1):1-6.

46. Reed ME, Huang J, Graetz I, Lee C, Muelly E, Kennedy C, et al. Patient Characteristics Associated With Choosing a Telemedicine Visit vs Office Visit With the Same Primary Care Clinicians. *JAMA network open*. 2020;3(6):e205873.
47. McGrail KM, Ahuja MA, Leaver CA. Virtual Visits and Patient-Centered Care: Results of a Patient Survey and Observational Study. *Journal of medical Internet research*. 2017;19(5):e177.
48. Miller D, Loftus AM, O'Boyle PJ, McCloskey M, O'Kelly J, Mace D, et al. Impact of a telephone-first consultation system in general practice. *Postgraduate medical journal*. 2019;95(1129):590-5.
49. Neufeld A, Bhella V, Svrcek C. Pivoting to be Patient-Centered: A Study on Hidden Psychological Costs of Tele-Healthcare and the Patient-Provider Relationship. *Internet Journal of Allied Health Sciences & Practice*. 2022;20(1):1-10.
50. Egede LE, Gebregziabher M, Walker RJ, Payne EH, Acierno R, Frueh BC. Trajectory of cost overtime after psychotherapy for depression in older Veterans via telemedicine. *Journal of affective disorders*. 2017;207:157-62.
51. Harder VS, Musau AM, Musyimi CW, Ndetei DM, Mutiso VN. A randomized clinical trial of mobile phone motivational interviewing for alcohol use problems in Kenya. *Addiction*. 2020;115(6):1050-60.
52. Nomura A, Tanigawa T, Muto T, Oga T, Fukushima Y, Kiyosue A, et al. Clinical Efficacy of Telemedicine Compared to Face-to-Face Clinic Visits for Smoking Cessation: multicenter Open-Label Randomized Controlled Noninferiority Trial. *Journal of medical Internet research*. 2019;21(4):e13520.
53. Ure A. Investigating the effectiveness of virtual treatment via telephone triage in a New Zealand general practice. *Journal of primary health care*. 2022;14(1):21-8.
54. Wickstrom HL, Oien RF, Fagerstrom C, Anderberg P, Jakobsson U, Midlov PJ. Comparing video consultation with inperson assessment for Swedish patients with hard-to-heal ulcers: registry-based studies of healing time and of waiting time. *BMJ open*. 2018;8(2):e017623.
55. Befort CA, Vanwormer JJ, Desouza C, Ellerbeck EF, Gajewski B, Kimminau KS, et al. Effect of Behavioral Therapy with In-Clinic or Telephone Group Visits vs In-Clinic Individual Visits on Weight Loss among Patients with Obesity in Rural Clinical Practice: A Randomized Clinical Trial. *JAMA - Journal of the American Medical Association*. 2021;325(4):363-72.




Figure 1. PRISMA flow chart of included studies.




Figure 2. Risk of bias assessments. Cells were coloured coded green for 'yes', grey for 'can't tell', and red for 'no'. *McGrail is categorised as both quantitative descriptive and non-randomised due to the two distinct cross-sectional survey and quasi-experimental components of the study.