

1 **World Health Organization Danger Signs to predict bacterial sepsis in newborns: A pragmatic**
2 **prospective cohort study**

3
4 Omolabake Akinseye, MBBS^{1,#}, Constantin R. Popescu, MD, MSc^{2,3,#}, Msandeni Chiume-Kayuni
5 MMED Peads, MSc^{4,5}, Michael A. Irvine, PhD^{6,7}, Norman Lufesi DCM, MPhil⁸, Tisungane Mvalo,
6 MBBS, MMED^{9,10}, Niranjan Kissoon, MD, MCCM^{1,2}, Matthew O. Wiens, PharmD, PhD¹¹, Pascal M.
7 Lavoie, MDCM, PhD^{1,2,*}

8
9 # contributed equally as co-first authors

10
11 **Affiliations:**

12 ¹ Department of Pediatrics, Faculty of Medicine, University of British Columbia, Vancouver, Canada

13 ² British Columbia Children's Hospital Research Institute, Vancouver, Canada

14 ³ Department of Pediatrics, Université Laval, Québec, Canada

15 ⁴ Department of Pediatrics, Kamuzu Central Hospital, Lilongwe, Malawi

16 ⁵ Kamuzu University of Health Sciences, Lilongwe, Malawi

17 ⁶ British Columbia Centre for Disease Control, Vancouver, Canada

18 ⁷ Faculty of Health Sciences, Simon Fraser University, Burnaby, Canada

19 ⁸ Department of Curative and Medical Rehabilitation, Ministry of Health, Lilongwe, Malawi

20 ⁹ University of North Carolina Project Malawi, Lilongwe, Malawi

21 ¹⁰ Department of Pediatrics, University of North Carolina at Chapel Hill School of Medicine, Chapel
22 Hill, NC, USA

23 ¹¹ Department of Anesthesiology, Pharmacology & Therapeutics, Faculty of Medicine, University of
24 British Columbia, Vancouver, Canada

25
26 ***Corresponding author:** Dr. Pascal M. Lavoie
27 BC Children's Hospital Research Institute
28 4th Floor, Translational Research Building
29 950 West 28th Avenue
30 Vancouver, British Columbia V5Z 4H4, Canada
31 Email: plavoie@bcchr.ca
32 Fax: +1-604-875-3106

33
34 **Short title:** Prediction of bacterial sepsis in infants.

35
36 **Conflict of Interest Statement:** The authors declare no conflicts of interest.

37
38 **Funding Support:** This study was supported by a grant from Grand Challenges Canada (R-ST-POC-
39 1707-04630). PML is supported by the BC Children's Hospital Research Foundation through the
40 Investigator Grant Award Program. CRP was funded by a Sue Carruthers Graduate Studentship from
41 the BC Children's Hospital Foundation. The other authors received no additional funding.

42
43 **Role of Funder/Sponsor:** The funder/sponsor did not participate in the work.

44
45 **Abbreviations:** WHO: World Health Organization; DS: danger signs; LMICs: low and middle-income
46 countries; CSF: cerebrospinal fluid; AUC-ROC: Area Under Curve – Receiver Operator
47 Characteristics; bpm: breaths per minute.

48

49 **Abstract**

50

51 Background: The World Health Organization (WHO) has developed danger signs (DS) to help front-
52 line health workers triage interventions in children with severe illnesses. Our objective was to evaluate
53 the extent to which DS predict bacterial sepsis in young infants presenting with acute illness.

54

55 Methodology/Principal Findings: This prospective study evaluated nine DS in infants younger than 3
56 months with suspected sepsis in a large regional hospital in Lilongwe, Malawi, between June 2018 and
57 April 2020. The main outcomes were positive blood or cerebrospinal fluid (CSF) cultures and
58 mortality. Blood (n=85/401) and CSF (n=2/204) cultures were positive in 21.2% and 1% of infants,
59 respectively (N=401; gestational age mean \pm SD: 37.1 \pm 3.3 weeks, birth weight 2865 \pm 785 grams). In-
60 hospital deaths occurred in 9.7% (N=39/401) of infants (61.5% within 48h of admission). In univariate
61 analyses, all DS were associated with mortality except for temperature instability and tachypnea,
62 whereas “infant was unable to feed” was the only DS significantly associated with bacterial sepsis.
63 After co-variable adjustments, number of DS predicted mortality (OR: 1.75; 95%CI: 1.43–2.16;
64 $p < 0.001$; AUC-ROC: 0.756) but not positive cultures (OR 1.08; 95%CI: 0.92–1.30; $p = 0.336$). Whether
65 potential bacterial contaminants were included or not did not change results meaningfully.

66

67 Conclusion/Significance: DS predicted fatal outcomes but not positive cultures in a large regional
68 hospital setting. These data imply that the incidence of bacterial sepsis and attributable mortality are
69 unlikely to be accurate based on clinical signs alone, in infants in LMIC settings.

70 **Introduction**

71 Three million children die each year worldwide in their first month of age, with a significant proportion
72 of these deaths occurring during the first week(1). Severe infections (including sepsis, pneumonia,
73 diarrhea and viral diseases) are common causes of serious illnesses in young infants (2). Up to 25% of
74 cases in low- and middle-income countries (LMICs) have identified bacterial causes (3). However, the
75 lack of diagnostic resources, particularly in LMICs makes it difficult to accurately quantify this burden.
76 This gap has enormous implications in terms of implementing policies and guidelines for the use of
77 antimicrobials to prevent these deaths (1). At the individual level, identifying bacterial causes for sepsis
78 is crucial to prevent fatalities and guide the correct empiric antibiotic therapy in the context of rising
79 antimicrobial resistance. Indeed, the misuse, or even the overuse, of antibiotics carries a risk of
80 subsequent treatment failure due to colonization by resistant bacteria (4, 5).

81

82 To assist front-line health workers, the World Health Organization (WHO) and United Nations
83 International Children's Emergency Fund developed algorithms in the early 1990s to help triage
84 interventions for sick children (6). This Integrated Management of Childhood Illness (IMCI) strategy
85 included the identification of clinical signs to alert the need for escalation of care, later termed Danger
86 Signs (DS) (6). Subsequently, the Young Infant Clinical Signs Study Group carried out a multi-
87 centered study in six LMICs and found that seven signs (i.e., history of difficulty feeding, grunting,
88 convulsions, movement only when stimulated, respiratory rate ≥ 60 breaths per minute (bpm), severe
89 chest indrawing, temperature below 35.5°C or above 37.5°C , prolonged capillary refill, cyanosis and
90 stiff limbs) best predicted the need for higher level of care, with a sensitivity of 74% to 85%, and
91 specificity of 75% to 79%, depending on infants' age (i.e., 0 to 6 days versus 7 to 59 days old) (7). A
92 systematic review of five studies on 17,506 infants found similar results (8).

93 Over time, the use of DS has been extended to provide a reasonable basis for the initiation of empiric
94 antibiotic treatment (9). However, these signs have neither been developed nor validated to diagnose
95 sepsis of bacterial etiology. Neonatal disorders such as transient tachypnea of the newborn,
96 hypoglycemia, hyaline membrane disease, birth asphyxia, and viral infections (to mention only a few),
97 present in a fashion similar to bacterial sepsis (10). Hence, a key question is whether DS can
98 reasonably identify newborns or young infants who truly benefit from antibiotic treatment. Only a few
99 studies have attempted to answer this question, but certainly none that have, to the best of our
100 knowledge, been conducted prospectively, in a pragmatic setting where clinicians, not research staff,
101 pragmatically reported DS (11).

102

103 The objective of this study was to prospectively determine the extent to which WHO DS predict
104 bacterial sepsis in young infants, in a LMIC hospital setting where front-line health workers were asked
105 to assess and record the presence of DS in infants who presented with suspected sepsis.

106

107 **Materials and Methods**

108 Study design, setting & population: This study was a pragmatic cohort study, and secondary analysis of
109 data from a parent study that aimed to identify molecular signatures of bacterial sepsis in young infants
110 with suspected sepsis (12). In the parent study, infants below 3 months of age were sequentially
111 enrolled “around the clock” at the time they presented with suspected sepsis before, to the neonatal or
112 pediatric units at Kamuzu Central Hospital in Lilongwe, Malawi between June 5th, 2018, and April 6th,
113 2020. To minimize bias towards false-negative blood culture, infants were excluded if they had
114 received antibiotics for >4 hours at the time of enrollment. In the original (parent) study, about 851
115 infants presented with suspected sepsis and the vast majority were not enrolled because staff were

116 unable to get consent - either parents declined or the conditions were not sufficiently permissive for
117 proper consent due to other clinical demands. Enrolled infants were followed during hospitalization.

118

119 The Kamuzu Central Hospital the largest referral hospital in central Malawi, where approximately
120 3,000 infants are admitted to its neonatal unit annually. “Suspected sepsis” was defined as per the
121 clinical staff, essentially in an infant that looked unwell where bacterial sepsis could not be excluded
122 clinically. Before the study, the staff at Kamuzu Central Hospital did not have routinely access to blood
123 culture or other ancillary tests, IV fluids and gavage feeds were also most often not available due to
124 limited resources and supplies. Available interventions include antibiotics, assistance with feeding and
125 routine surgeries when necessary. Oxygen and continuous positive pressure ventilation were only
126 available to an extremely limited number of infants, on a limited basis when the hospital had electrical
127 power. Invasive mechanical ventilation was not possible. Written informed consent was obtained from
128 all parents/guardian of each participant, in English or Chichewa (main local language). During the
129 study, the authors did not have access to information that could identify participants, with the exception
130 of the consent forms who were kept under the responsibility of the local study PI (MCK). The study
131 was approved by the University of British Columbia’s Children’s and Women’s Research Ethics Board
132 (#H16-02639) and the National Health Science Research Committee of Malawi (#17/8/1819).

133

134 Clinical data and blood samples collection: Each infant had an initial clinical evaluation by a healthcare
135 worker (or staff, either a nurse or clinical officer), and the presence of the following DS were
136 prospectively collected using electronic forms, on a tablet device: history of poor feeding (interviewing
137 the parent/caregiver), entered as “feeding well”, “unable to feed” or “uninterested in feeding”,
138 respiratory rate (visually counted by attended clinician), axillary body temperature (measured using an
139 electronic Welch Allyn SureTemp thermometer, model 692), drowsiness/difficulty to wake, grunting,

140 severe chest recession, central cyanosis and convulsions (defined pragmatically by attending clinician).
141 All infants also had a standardized aerobic blood culture (minimum 2 mL of blood) collected within 4
142 hours of presentation, using a rigorously implemented aseptic protocol (with in-training of all front-line
143 healthcare providers prior to study initiation), as detailed (13). Infants also had cerebrospinal fluid
144 (CSF) and urine cultures collected as clinically indicated. Fast breathing was defined as a respiratory
145 rate of ≥ 60 bpm. Temperature instability was defined as a body temperature $< 35.5^{\circ}\text{C}$ or $> 38^{\circ}\text{C}$.
146 Blood cultures were processed into BACTEC PEDS Plus culture bottles incubated in a BD BACTEC
147 9050 instrument. In keeping with a pragmatic design, the study did not attempt to enforce strict criteria
148 to define DS or illness status. Identification of organisms from blood or CSF cultures was completed
149 in-house using bioMerieux API, and BD Crystal kits at a local University of North Carolina-affiliated
150 research facility, as described (12). Culture results were made available to clinicians within 5 days.
151 Bacterial isolates that could be restored (~60% of isolates) were also analyzed for further confirmation
152 of identity (not shown) on a Microflex LT MALDI-TOF mass spectrometer, using the BioTyper
153 software version 3.1 (Bruker Daltonik, Germany).

154

155 Outcome measures: The primary outcomes of interest were *mortality*, and *culture-positive sepsis*
156 defined as the bacterial growth of a known neonatal pathogen in blood or CSF culture sample. Micro-
157 organisms not commonly considered as pathogens in term or near-term infants, such as coagulase-
158 negative Staphylococcus, Micrococcus or Bacillus were excluded (14), but these potential
159 contaminants were included in a sensitivity analysis using an alternate bacterial sepsis outcome
160 (**Supplemental Table 1**). In-hospital mortality was recorded until discharge home or transfer to
161 another hospital facility. Data were considered missing when infants were transferred within 72 hours
162 after initial admission to the hospital.

163

164 Statistical Analysis: In the parent study, the sample size was set conveniently at 500 infants over 2
165 years, but the study needed to be stopped in April 2020 due to the COVID-19 pandemic. Demographic
166 characteristics were reported descriptively using the mean/median with dispersion. The following nine
167 WHO DS were used as predictors: *not feeding well, convulsions, drowsy/difficult to wake, movements*
168 *only when stimulated or no movements, fast breathing, grunting, severe chest recessions, temperature*
169 *instability (hypo/hyperthermia), central cyanosis*. All quantitative variables were analyzed as such,
170 except for fast breathing and temperature instability which were categorized according to WHO
171 definitions (see above). Odds ratios were calculated for each DS and the main outcomes of bacterial
172 sepsis or mortality. Logistic regressions were conducted between the number of DS and the main
173 outcomes. In separate analyses, odds ratios were calculated using logistic regressions between mean
174 DS (i.e., the number of DS divided by the available DS data) to determine the effect of missing DS
175 data. Unadjusted logistic regression models, including demographic variables, were conducted to
176 determine the predictive value of the DS. The predictive performance of DS for each outcome was
177 assessed using the area under receiver operator characteristic curve (AUC-ROC). Sociodemographic
178 variables associated with mortality, as well as their independent effects were analyzed using regression
179 models. Only complete cases were used in the analysis. Statistical significance was considered at p-
180 value <0.05 with no adjustment made for multiple testing in the unadjusted models. Analyses were
181 performed using R (version 4.1.0) and SPSS (version 20; IBM, Armonk, NY, USA).

182

183 **Results**

184 Study population: All infants originally enrolled in the parent study were included, for a total of 401
185 infants enrolled in the original study and included in this analysis. The majority were term or late
186 preterm infants, born vaginally, presented from home in their first week of post-natal life, and had
187 roughly equal female/male distribution (**Table 1**). Most infants were reported by clinicians to be ill at

188 the time of initial presentation (69.3%; 278/401), 22.2% (89/401) were severely ill, and 8.5% appeared
 189 well (34/401). In total, 79.8% (320/401) presented at least one DS. Overall, 56.9% (228/401) of infants
 190 had been ill for at least 24 hours prior to initial presentation, and their parents had traveled a median
 191 distance of 17 kilometers, often by foot or taxi, to reach the nearest hospital. In total, 94/401 (23.4%)
 192 infants had received antibiotics prior to the blood culture, for a median of 2 hours (range 0.75 to 3.5
 193 hours). The duration of antibiotic use in infants ranged from 1 to 30 days (**Table 1**). Few infants with
 194 negative blood culture had antibiotics discontinued after 48 hours (21/401, 5.2%).

195
 196 **Table 1: Socio-demographic characteristics of the study population.**

Participant characteristics		N with data
Gestational age at birth (weeks), mean (SD)	37.1 (3.3)	397
Birth weight (gram), mean (SD)	2865 (785)	380
Female, no (%)	185 (46)	401
Admitted from, no (%)		401
Inborn	161 (40)	
Transferred from another facility	224 (56)	
Home	16 (4)	
Post-natal age at presentation (days), median (IQR)	4 (2-16)	401
Mode of delivery, no (%)		398
Cesarean section	101 (25.2)	
Vaginal delivery	297 (71.8)	
Duration of illness before presentation to hospital (hours), median (IQR)	24 (12-48)	341

Distance of travel to hospital (km), median (IQR)	17 (11-25)	342
Illness status, number on infants (%)		401
Well	34 (8.5)	
Ill	278 (69.3)	
Severely ill	89 (22.2)	
Duration of antibiotics (days), median (IQR)	6 (5-9)	401
Duration of hospital stay (days), median (IQR)	7 (5-9)	401
Danger Signs, number of infants (%)		
Not feeding well	171 (43.3)	395
Convulsions	28 (7.0)	401
Drowsy / Difficult to wake	33 (8.2)	401
Movements only when stimulated or no movements	26 (6.5)	401
Fast breathing ≥ 60 per min	104 (27.1)	384
Grunting	30 (7.5)	401
Severe chest recessions	51 (12.7)	401
Temperature instability $>38^{\circ}\text{C}$ or $<35.5^{\circ}\text{C}$	118 (32.7)	361
Central cyanosis	29 (7.2)	401

197

198

199 Outcome data: Blood (n = 85/401) and CSF (n = 2/204) cultures were positive in 21.2% and 1% of the
200 infants, respectively. However, about half (46 / 85) had bacterial types considered to be contaminants
201 (e.g., *bacillus*, *micrococcus*, *corynebacterium* and *coagulase-negative staphylococcus* species and non-
202 pathogenic *streptococcus* species such as *oralis*), so the actual proportion with bacterial sepsis,
203 considering only known pathogens in term and near-term infants, was effectively 9.7% (39/401).

204 Excluding contaminants, gram-positive bacteria (53.8%; 21/39) were slightly more prominent than
205 gram-negative bacteria (46.2%; 18/39) (**Supplemental Table 1**). Among all infants with presumed
206 sepsis, pathogen sepsis was more common in the inborn (17/161) than in the infants transferred from
207 another hospital facility or from home (22/240) ($p=0.029$). The proportion of gram-positive sepsis was
208 comparable between inborn infants (10/240) versus in infants transferred from another hospital or
209 home (11/161) ($p=0.240$). Three infants were transferred before 72 hours, so among infants in whom
210 data were available, the mortality was 9.7% (39/398), of which 61.5% (24/39) died within 48h.

211

212 Associations with mortality and culture-positive outcomes: The following DS were significantly
213 associated with mortality (**Table 2**): *not feeding well, convulsions, drowsy/difficult to wake, movements*
214 *only when stimulated or no movements, grunting, severe chest recessions, central cyanosis*. In contrast,
215 *fast breathing or temperature instability* were not significantly associated with mortality. The
216 cumulative number of DS was also significantly associated with mortality (OR 1.75 [95%CI: 1.43 –
217 2.16] per additional DS; $p<0.001$) (**Supplemental Figure 1**). Admission weight, but not age of the
218 mother, gestational age, birth weight and sex were significantly associated with mortality
219 (**Supplemental Table 2**). When adjusting for all of these co-variables, cumulative number of DS
220 remained significantly associated with mortality (**Supplemental Table 2**; $p<0.001$).

221 **Table 2: Associations between DS and mortality.**

Variable	Odds ratio (OR)	95%CI	P value
Not feeding well	7.21	(3.27 - 18.2)	<0.001
Convulsions	3.52	(1.31 - 8.59)	0.008
Drowsy / Difficult to wake	5.04	(2.12 - 11.4)	<0.001
Movements only when stimulated or no movements	4.89	(1.88 - 11.9)	<0.001
Fast breathing ≥ 60 per min	1.86	(0.87 - 3.85)	0.100
Grunting	3.95	(1.54 - 9.34)	0.002
Severe chest recessions	3.01	(1.34 - 6.37)	0.005
Temperature instability $>38^{\circ}\text{C}$ or $<35.5^{\circ}\text{C}$	1.70	(0.75 - 3.75)	0.191
Central cyanosis	4.15	(1.62 - 9.87)	0.002

222

223

224 Bacterial sepsis with a known pathogen was significantly associated with mortality (OR 3.29 [95%CI:
225 1.37 – 7.36]; $p = 0.005$). However, the only DS significantly associated with bacterial sepsis when
226 considering only known neonatal pathogens was *not feeding well* (OR 2.04 [95%CI: 1.05 – 4.06];
227 $p=0.038$) (**Table 3**). Associations between convulsions, drowsiness, lethargy (i.e., movements only
228 when stimulated or no movement), tachypnea, grunting, severe chest recession, temperature instability
229 and central cyanosis were all non-significant (**Table 3**). Association was between cumulative number
230 of DS and bacterial sepsis was also non-significant (OR 1.21 [95% CI: 0.98 – 1.47]; $p=0.072$) in
231 regression models (**Supplemental Figure 2**). At discharge, clinical staff selected non-bacterial sepsis
232 in 76.3% (306/401), bacterial sepsis in 15.5% (62/401) and “other non-sepsis syndromes” in 8.2%
233 (33/401) of cases, from these options, for study purposes.

234

235 **Table 3: Association between DS and bacterial sepsis when organism is a known pathogen.**

Variable	Odds ratio (OR)	95%CI	P value
Not feeding well	2.04	(1.05 - 4.06)	0.038
Convulsions	0.69	(0.11 - 2.45)	0.626
Drowsy / Difficult to wake	0.91	(0.21 - 2.73)	0.886
Movements only when stimulated or no movements	1.75	(0.49 - 4.90)	0.327
Fast breathing ≥ 60 per min	1.16	(0.53 - 2.38)	0.703
Grunting	2.54	(0.89 - 6.31)	0.058
Severe chest recessions	1.61	(0.62 - 3.71)	0.288
Temperature instability $>38^{\circ}\text{C}$ or $<35.5^{\circ}\text{C}$	1.80	(0.86 - 3.70)	0.113
Central cyanosis	0.67	(0.11 - 2.34)	0.588

236

237 When including all positive blood cultures (**Supplemental Table 3**), culture positivity was not
238 significantly associated with mortality (OR 1.77 [95% CI: 0.83 – 3.6]; $p=0.123$), or any DS. The
239 cumulative number of DS was also not significantly associated with bacterial sepsis when including
240 positive cultures with potential bacterial contaminants (OR 1.21 [95% CI: 0.98 – 1.47]; $p=0.0715$).

241

242 **Discussion**

243 This study found that most WHO DS were significantly associated with mortality, but only one DS
244 (i.e., not feeding well) as significantly associated with culture-proven bacterial sepsis. Additionally, the
245 majority of DS were not associated with culture-proven bacterial sepsis in this LMIC neonatal cohort.
246 The results held true whether only bacterial pathogens, or potential bacterial contaminants were
247 considered. To the best of our knowledge, the current study is the first to pragmatically examine
248 whether DS predict bacterial sepsis in infants. These findings were obtained prospectively in an LMIC

249 setting where DS were identified by front line healthcare workers as they would normally do
250 pragmatically outside a research protocol – we assumed that this study design would more closely
251 reflects the real-life utility of DS. Few studies have attempted to determine how DS can predict
252 bacterial sepsis caused by bacterial pathogens in newborns or young infants. A retrospective Brazilian
253 study reviewed data on 83 sick neonates over a 12 months period, and came to the same conclusion that
254 DS alone or in combination did not predict severe serious bacterial infection (15). The findings of the
255 current study are also consistent with those of another study carried out in a primary care, rural setting
256 in Bangladesh, India and Pakistan. In the latter study, clinical signs could not differentiate infants with
257 viral or bacterial illnesses (16). The current study findings have implications for a hospital setting, but
258 further research is needed to determine the generalizability of the results to other hospital contexts.
259
260 In addition to the main findings, the current study also found that only a minor proportion of infants
261 with suspected sepsis showed positive blood culture. Only few other studies have prospectively
262 estimated the incidence of bacterial sepsis in LMIC cohorts, and it is therefore useful to present the
263 current findings in the context of those other studies. A hospital centre study in Delhi, India, included
264 13,530 neonates (all comers; mean GA±SD: 36.0±3.4 weeks) monitored daily for clinical signs of
265 sepsis with per-protocol completion of a sepsis work-up (blood culture and lumbar puncture) prior to
266 initiating antibiotics, and found a 14.3% incidence of culture-positive sepsis (17), which is similar to
267 the findings of the current study. Another study in community settings in Madagascar, Senegal and
268 Cambodia reported 8.2% culture-confirmed sepsis among 514 infants assessed for “possible serious
269 bacterial infection” out of 3,688 neonates (18). One large study followed 12,622 live-births up to 60
270 days of age in rural India between April 2002 and March 2005, with daily home visits and clinical
271 assessment, and standardized blood and CSF cultures following enrollment of infants with suspected
272 sepsis. It reported a 6.3% incidence of culture-confirmed sepsis (53 with positive blood cultures out of

273 842 infants assessed, excluding potential contaminants), for a population-level incidence of culture-
274 confirmed sepsis of 6.7 per 1,000 births (19). Taken together, these findings suggest that bacterial
275 sepsis may be uncommon in this age group even in LMICs, when sepsis is clinically suspected.

276

277 Blood cultures are the gold standard for diagnosing bacterial sepsis in young infants (20). However,
278 these tests lack sensitivity in neonates (21). In the current study we were careful to prescribe a
279 minimum blood volume to maximize culture yield. Yet, a key question is whether we misestimate the
280 proportion of bacterial sepsis in infants in whom cultures are negative? The Aetiology of Neonatal
281 Infection in South Asia (ANISA) study (22) also attempted to identify the cause of sepsis for 6022
282 cases of possible serious bacterial infections in infants in the first 60 days after birth, in a population of
283 63,114 livebirth). Bacterial and viral (e.g., respiratory syncytial virus) causes were identified in 16%
284 and 12% of cases, respectively. For the remaining 72% cases, no pathogen could be identified despite
285 systematic blood and respiratory cultures, which is consistent with findings from the current study.

286

287 Evidently the use of antibiotics in the current study was high, with the vast majority of infants receiving
288 antibiotics for a minimum of five days during hospitalization, as per hospital practices. As in other
289 LMIC settings, blood cultures were not routinely available at the study site prior to the study. As such,
290 no approach had been developed to stop antibiotics after 48 hours given a negative blood culture. In
291 another study in Thailand, the duration of antibiotic use was also high with durations of 15 and 8 days
292 for culture-proven and culture-negative early-onset neonatal sepsis, respectively (23). These results
293 highlight the need to rely on DS, and potential coercive effect of the lack of laboratory resources
294 capable of providing basic laboratory test results to help “rule out” sepsis (24).

295

296 A strength of this study is its prospective design, with standardized blood culture procedures to ensure
297 that the chances of contamination during blood sampling were minimized, and robust microbiological
298 identification methods including clinical laboratory protocols on-site augmented by advanced MALDI-
299 TOF species determination in another, North American clinical microbiology laboratory. The strong
300 association between bacterial sepsis and mortality suggest that the infants with positive cultures were
301 correctly identified. However, a number of study limitations should be acknowledged. Although a
302 sizeable proportion of the recruited infants presented from home, data were collected in a single
303 hospital-type setting; as such, caution should be exercised when generalizing to other hospital settings,
304 or extrapolating to non-hospital primary care settings. For logistical reasons CSF was collected from
305 only approximately half of the infants, which may have underestimated the incidence of bacterial
306 sepsis. Moreover, due to the extremely limited diagnostic capabilities of the study LMIC settings, in
307 most cases, causes of deaths or alternate diagnoses could not be determined. Similarly, the study did
308 not include non-bacterial testing, which precluded identification of viral sepsis syndromes.

309

310 In conclusion, our study validates the use of WHO DS to predict fatal outcomes, but not bacterial
311 sepsis, in young infants. This study further emphasizes the need for better diagnostics for bacterial
312 sepsis, in order to ensure that health gains continue to improve, in areas of the world where sepsis
313 represents a huge burden and where antimicrobial resistance is also increasing at alarming rates (25).

314

315 **Acknowledgments**

316 We thank Dr. David Goldfarb for expert input into data interpretation and Amber Johnson from the
317 British Columbia Children's Hospital Microbiology Laboratory for confirming bacterial isolates by
318 mass spectrometry.

319 **REFERENCES**

- 320 1. GBD 2016 Causes of Death Collaborators. Global, regional, and national age-sex specific mortality
321 for 264 causes of death, 1980-2016: a systematic analysis for the Global Burden of Disease Study
322 2016. *Lancet*. 2017;390(10100):1151-210.
- 323 2. Rudd KE, Johnson SC, Agesa KM, Shackelford KA, Tsoi D, Kievlan DR, et al. Global, regional, and
324 national sepsis incidence and mortality, 1990-2017: analysis for the Global Burden of Disease Study.
325 *Lancet*. 2020;395(10219):200-11.
- 326 3. Fleischmann-Struzek C, Goldfarb DM, Schlattmann P, Schlapbach LJ, Reinhart K, Kissoon N. The
327 global burden of paediatric and neonatal sepsis: a systematic review. *The Lancet Respiratory*
328 *medicine*. 2018;6(3):223-30.
- 329 4. Costelloe C, Metcalfe C, Lovering A, Mant D, Hay AD. Effect of antibiotic prescribing in primary
330 care on antimicrobial resistance in individual patients: systematic review and meta-analysis. *BMJ*.
331 2010;340:c2096.
- 332 5. Klingenberg C, Kornelisse RF, Buonocore G, Maier RF, Stocker M. Culture-Negative Early-Onset
333 Neonatal Sepsis - At the Crossroad Between Efficient Sepsis Care and Antimicrobial Stewardship.
334 *Front Pediatr*. 2018;6:285.
- 335 6. Costello A. Integrated management of childhood illness. *Lancet*. 1997;350(9087):1266.
- 336 7. Young Infants Clinical Signs Study Group. Clinical signs that predict severe illness in children under
337 age 2 months: a multicentre study. *The Lancet*. 2008;371(9607):135-42.
- 338 8. Opiyo N, English M. What clinical signs best identify severe illness in young infants aged 0-59 days
339 in developing countries? A systematic review. *Archives of disease in childhood*. 2011;96(11):1052-
340 9.

- 341 9. World Health Organization. Pocket Book of Hospital Care for Children: Guidelines for the
342 Management of Common Childhood Illnesses. WHO Guidelines Approved by the Guidelines
343 Review Committee. 2nd ed. Geneva, 2013.
- 344 10. Wynn JL, Polin RA. Progress in the management of neonatal sepsis: the importance of a consensus
345 definition. *Pediatr Res*. 2018;83(1-1):13-5.
- 346 11. Clinical prediction of serious bacterial infections in young infants in developing countries. The WHO
347 Young Infants Study Group. *Pediatr Infect Dis J*. 1999;18(10 Suppl):S23-31.
- 348 12. Popescu CR, Tembo B, Chifisi R, Cavanagh MMM, Lee AH, Chiluzi B, et al. Whole blood genome-
349 wide transcriptome profiling and metagenomics next-generation sequencing in young infants with
350 suspected sepsis in a low-and middle-income country: A study protocol. *Gates Open Res*.
351 2020;4:139.
- 352 13. Popescu C, Tembo B, Chifisi R, Cavanagh M, Chiluzi B, Lufesi N, et al. Procedure policies, data
353 collection form and model consent form (supplemental material)~Improving the Early Diagnosis of
354 Neonatal Sepsis in Malawi. In: *Scholars Portal Dataverse*, editor. 2 ed 2020.
- 355 14. Downie L, Armiento R, Subhi R, Kelly J, Clifford V, Duke T. Community-acquired neonatal and
356 infant sepsis in developing countries: efficacy of WHO's currently recommended antibiotics--
357 systematic review and meta-analysis. *Archives of disease in childhood*. 2013;98(2):146-54.
- 358 15. Ronfani L, Vilarim JN, Dragovich D, Bacalhau AF, Cattaneo A. Signs of severe bacterial infection
359 in neonates. *J Trop Pediatr*. 1999;45(1):48-51.
- 360 16. Arvay ML, Shang N, Qazi SA, Darmstadt GL, Islam MS, Roth DE, et al. Infectious aetiologies of
361 neonatal illness in south Asia classified using WHO definitions: a primary analysis of the ANISA
362 study. *The Lancet Global health*. 2022;10(9):e1289-e97.
- 363 17. Study IotDNI. Characterisation and antimicrobial resistance of sepsis pathogens in neonates born in
364 tertiary care centres in Delhi, India: a cohort study. *The Lancet Global health*. 2016;4(10):e752-60.

- 365 18. Huynh BT, Kermorvant-Duchemin E, Chheang R, Randrianirina F, Seck A, Ratsima EH, et al. Severe
366 bacterial neonatal infections in Madagascar, Senegal, and Cambodia: A multicentric community-
367 based cohort study. *Plos Medicine*. 2021;18(9).
- 368 19. Panigrahi P, Chandel DS, Hansen NI, Sharma N, Kandefer S, Parida S, et al. Neonatal sepsis in rural
369 India: timing, microbiology and antibiotic resistance in a population-based prospective study in the
370 community setting. *J Perinatol*. 2017;37(8):911-21.
- 371 20. Polin RA. Management of neonates with suspected or proven early-onset bacterial sepsis. *Pediatrics*.
372 2012;129(5):1006-15.
- 373 21. Popescu CR, Cavanagh MMM, Tembo B, Chiume M, Lufesi N, Goldfarb DM, et al. Neonatal sepsis
374 in low-income countries: epidemiology, diagnosis and prevention. *Expert Rev Anti Infect Ther*.
375 2020;18(5):443-52.
- 376 22. Saha SK, Schrag SJ, El Arifeen S, Mullany LC, Shahidul Islam M, Shang N, et al. Causes and
377 incidence of community-acquired serious infections among young children in south Asia (ANISA):
378 an observational cohort study. *Lancet*. 2018;392(10142):145-59.
- 379 23. Kiatchoosakun P, Jirapradittha J, Sirikarn P, Laopaiboon M, Pattanittum P, Chandrakachorn W, et
380 al. Early-Onset Neonatal Sepsis and Antibiotic Use in Northeast Thailand. *Am J Perinatol*.
381 2019;36(12):1295-303.
- 382 24. Lavoie PM, Popescu CR, Molyneux EM, Wynn JL, Chiume M, Keitel K, et al. Rethinking
383 management of neonates at risk of sepsis. *Lancet*. 2019;394(10195):279-81.
- 384 25. Iregbu K, Dramowski A, Milton R, Nsutebu E, Howie SRC, Chakraborty M, et al. Global health
385 systems' data science approach for precision diagnosis of sepsis in early life. *Lancet Infect Dis*.
386 2022;22(5):e143-e52.
- 387