

1 **Factors associated with the near miss of pregnant and postpartum women hospitalized by**
2 **Covid-19**

3
4 **Maternal near miss by Covid-19**

5
6 Ana Carolina Schultz da Silva¹, Márcia Moroskoski², Natan Nascimento de Oliveira², Franciele Aline
7 Machado de Brito², Marcela de Andrade Pereira Silva³, Maria Aparecida Salci², Lígia Carreira², João
8 Ricardo Nickenig Vissoci⁴⁻⁵, Rosana Rosseto de Oliveira²

9
10 ¹Nursing, Centro Universitário Uningá.

11 ² Nursing Departament, Universidade Estadual de Maringá, Maringá, Brazil.

12 ³ Health Science Departament, Universidade Estadual de Maringá, Maringá, Brazil.

13 ⁴Division of Emergency Medicine, Duke University Medical Center, Durham, North Carolina, USA.

14 ⁵Duke Global Health Institute, Duke University, Durham, North Carolina, USA.

15

16 Corresponding author: João Ricardo Nickenig Vissoci. E-mail: joao.vissoci@duke.edu

17

18 **ABSTRACT**

19 **Objective:** To analyze the factors associated with maternal near miss due to Covid-19 in Brazil.

20 **Method:** cross-sectional study that assessed the cases of pregnant and postpartum women
21 hospitalized with SARS due to Covid-19, aged between 10 and 49 years and from March 2020 to
22 March 2022. Secondary data available in the Influenza Epidemiological Surveillance Information
23 System (SIVEP-Flu) were used. Data were analyzed using SPSS statistical software, with univariate
24 analysis followed by logistic regression. **Results:** the significant independent factors associated with
25 near miss in pregnant postpartum women due to Covid-19 were black/brown race, O2 saturation

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

26 <95%, dyspnea, comorbidities, need for invasive and non-invasive ventilatory support. **Conclusion:**
27 the factors associated with near misses in pregnant and postpartum women with covid-19 reveal
28 characteristics and risks inherent to the pathophysiology of the disease that contribute to worsening
29 the condition.

30

31 **Keywords:** Coronavirus Infections; Near Miss; Maternal Mortality; Epidemiology; Nursing.

32

33 INTRODUCTION

34

35 In December 2019, the world was surprised by the emergence of a new virus in China, SARS-
36 CoV-2. Already in March 2020, the World Health Organization (WHO) had declared it a pandemic
37 due to its high transmissibility and the alarming number of cases and deaths in several countries and
38 continents.¹

39 Studies show an increased mortality rate from Covid-19 to the vulnerability of different
40 population groups, with or without morbidity, and in different age groups.² Among the most
41 vulnerable groups are pregnant women, who are susceptible to severe infections due to physiological
42 changes during pregnancy.³

43 In addition, although research has not shown evidence of vertical transmission of the virus, there is
44 an increased prevalence of preterm birth, miscarriage, emergency cesarean section, intrauterine
45 growth restriction, maternal death, maternal near misses, and other complications during pregnancy
46 in women with Covid-19.⁴

47 In Brazil, the Covid-19 Observatory of the Oswaldo Cruz Foundation (Fiocruz) announced in
48 2021 that the country has the highest number of maternal deaths caused by the Covid-19 virus.⁵
49 Another study showed that about eight out of 10 pregnant and postpartum women who have become

50 victims of the new coronavirus in the world are Brazilian.⁶ This rate could be even higher due to
51 underreporting of cases and the fact that many women of reproductive age (10-49 years) have died.⁷

52 Reducing maternal mortality is a global challenge for health systems and has become even
53 more challenging with the advent of Covid-19.⁸ The United Nations (UN), through the Sustainable
54 Development Goals, agreed to the Brazilian reality of maternal mortality rate not exceeding 30 deaths
55 per 100,000 live births by 2030.⁹ However, in 2021, this rate was 107 deaths per 100,000 births.¹⁰

56 The target set by the United Nations will only be possible through measures to improve the
57 capacity to care for pregnant women before, during, and after childbirth.⁸ On the other hand, non-
58 compliance will indicate the ineffectiveness of public policies aimed at women, since most cases are
59 considered preventable deaths.¹¹

60 In this sense, one of the strategies to reduce maternal mortality is to study the Near Miss
61 Materno (NMM), which consists of analyzing cases in which the woman almost died due to a serious
62 complication that occurred during pregnancy, childbirth, or the puerperium.¹²

63 It is believed that hospitalization of a pregnant or postpartum woman in intensive care for
64 Covid-19 could be avoided through prevention and health promotion. Therefore, it is of paramount
65 importance to study and train professionals working in these areas to recognize the factors associated
66 with complications and thus avoid them.¹³

67 Therefore, this study aims to analyze the factors associated with maternal near misses due to
68 Covid-19 in Brazil, considering the importance of the topic and the need to deepen the discussion and
69 knowledge about this disease.

70

71 **METHOD**

72

73 This is an observational, cross-sectional, and analytical study of notifications of pregnant and
74 postpartum women hospitalized for Covid-19 in Brazil from March 2020 to March 2022. The study

75 is based on the STROBE (Strengthening the Reporting of Observational Studies in Epidemiology)
76 tool, which aims to strengthen the appropriate reporting of observational studies in epidemiology.¹⁴

77 The notifications analyzed correspond to the 27 states of the country, with an estimated
78 population of 214.3 million inhabitants in 2022.¹⁵ The age range of 10 to 49 years was used because
79 it is defined in Brazil as the group of women of childbearing age¹⁶, filtering also the cases of pregnant
80 and postpartum women positive to Covid-19.

81 The notifications used were recorded in the public records of the Influenza Epidemiological
82 Information System (SIVEP-Influenza) of the Ministry of Health (MS). SIVEP-Influenza is a
83 surveillance program for the influenza virus in the country, from the sentinel surveillance network of
84 the influenza syndrome. Due to the pandemic, the Health Surveillance Secretariat of the Ministry of
85 Health (SVS/MS) adapted the system to include Covid-19 data.¹⁷ The data used in the study were
86 extracted from the platform on April 15, 2021 using R software.

87 To achieve the objective of analyzing the factors associated with maternal near misses due to
88 Covid-19, the dependent variable was the hospitalization of the pregnant and/or postpartum woman
89 for Covid-19 in the ICU. Pregnant and postpartum women who died regardless of ICU or ward
90 admission were excluded. Independent variables were grouped into sociodemographic characteristics
91 (race, age, education, area of residence, and region); signs and symptoms (fever, sore throat, dyspnea,
92 respiratory distress, saturation <95%, vomiting, diarrhea, and other symptoms); pre-existing
93 conditions (comorbidity, cardiac, hematologic, neurologic, renal, hepatic, Down syndrome, asthma,
94 diabetes, obesity, pneumonia, immunodeficiency); and clinical characteristics (obstetric population,
95 ventilatory support, nosocomial infection, influenza vaccine, Covid-19 vaccine, antiviral).

96 The statistic used was logistic regression, which aims to construct a model from a set of
97 observations that allows the prediction of the values taken by a categorical variable, in order to
98 identify the factors that characterize each group of sick individuals compared to healthy individuals.¹⁸
99 Thus, in the univariate regression analysis, the variables with $p < 0.20$ were selected.

100 The forward stepwise logistic regression model was then used, and the measures of association
101 were the adjusted odds ratio (adjusted OR) with a 95% confidence interval (CI), taking into account
102 the absence of multicollinearity in the model. Data were analyzed using the Statistical Package for
103 the Social Sciences (SPSS, version 20.1, IBM, USA).

104 The study used secondary and public domain databases and did not require the approval of an
105 ethics committee, according to the recommendations of Resolution No. 510/2016 of the National
106 Health Council.¹⁹

107

108 RESULTS

109

110 We analyzed 19,908 hospitalizations of pregnant and postpartum women with Covid-19 that
111 occurred in Brazil between March 2020 and March 2022. Of these total hospitalizations, 16,358
112 (82.16%) were pregnant women and 3,550 (17.84%) were postpartum women. Considering the
113 obstetric population, 4,132 (20.75%) experienced the situation of a near miss due to Covid-19,
114 represented by ICU admission, and 15,776 (79.25%) were admitted to wards (Table 1).

115 The sociodemographic aspects of pregnant and postpartum women with Covid-19
116 hospitalized in Brazil are shown in Table 1. The odds of ICU admission were higher in women aged
117 35 years and older (OR=1.41; CI=1.309-1.525), illiterate (OR=1.81; CI=1.087-3.021) and living in
118 peri-urban areas (OR=1.67; CI=1.072-2.603). Among the Brazilian regions, the Southeast region had
119 the highest odds ratio for the outcome (OR=1.86; CI=1.63-2.11) (Table 1).

120

121 Table 1 – Univariate analysis of hospitalizations of pregnant and postpartum women with Covid-19,
122 according to sociodemographic characteristics. Brazil, 2020-2022

Region	UTI		Infirmery		Total	OR	IC	p-value
	n (4132)	%	n (15776)	%				
South	644	15,59	2796	17,72	3440	1,4	1,21-1,62	<0,001

Southeast	1969	47,65	6449	40,88	8418	1,86	1,63-2,11	<0,001
Midwest	397	9,61	1325	8,40	1722	1,82	1,55-2,15	<0,001
Northeast	808	19,55	3297	20,90	4105	1,49	1,29-1,72	<0,001
North	314	7,60	1909	12,10	2223	-	-	-
Age								
<20	259	6,27	1456	9,23	1715	0,73	0,634-0,838	<0,001
20 to 34	2578	62,39	10564	66,96	13142	-	-	-
≥35	1295	31,34	3756	23,81	5051	1,41	1,309-1,525	<0,001
Race/Color								
White	1566	37,90	5685	36,04	7251	-	-	-
Black/Brown	1885	45,62	7619	48,29	9504	0,90	0,833-0,968	0,005
Yellow	28	0,68	113	0,72	141	0,90	0,593-1,366	0,619
Indigenous	12	0,29	138	0,87	150	0,32	0,175-0,571	<0,001
Education								
Illiterate	24	0,58	44	0,28	68	1,81	1,087-3,021	0,023
Elementary	414	10,02	1930	12,23	2344	0,71	0,609-0,835	<0,001
High School	815	19,72	3761	23,84	4576	0,72	0,626-0,828	<0,001
College	366	8,86	1216	7,71	1582	-	-	-
Zone		0,00		0,00				
Urban	3391	82,07	13352	84,63	16743	-	-	-
Rural	216	5,23	1018	6,45	1234	0,83	0,718-0,972	0,02
Peri-urban	28	0,68	66	0,42	94	1,67	1,072-2,603	0,023

123

124 Regarding the signs and symptoms of hospitalized pregnant and postpartum women with
 125 Covid-19 (Table 2), the results show that having had a fever (OR=1.47; CI=1.362-1.591), cough
 126 (OR=1.71; CI=1. 562-1.869), dyspnea (OR=4.57; CI=4.177-5.012), respiratory distress (OR=3.51;
 127 CI=3.231-3.810) and O2 saturation <95% (OR=5.62; CI=5.174-6.100) were factors associated with
 128 ICU admission. The sore throat was a protective factor for ICU admission (OR=0.77; CI=0.700-
 129 0.843).

130

131 Table 2 – Univariate Analysis of Hospitalizations of Pregnant and Postpartum Women with Covid-
 132 19 by Signs and Symptoms Brazil, 2020-2022

	UTI		Infirmary		Total	OR	IC	p-value
	n (4132)	%	n (15776)	%				
Fever								
Yes	2333	56,46	7577	48,03	9910	1,47	1,362-1,591	<0,001
No	1190	28,80	5689	36,06	6879	-	-	-
Cough								
Yes	2967	71,81	9847	62,42	12814	1,71	1,562-1,869	<0,001
No	708	17,13	4014	25,44	4722	-	-	-
Sore throat								
Yes	704	17,04	3502	22,20	4206	0,77	0,700-0,843	<0,001
No	2248	54,40	8592	54,46	10840	-	-	-
Dyspnea								
Yes	3082	74,59	6558	41,57	9640	4,57	4,177-5,012	<0,001

No	654	15,83	6367	40,36	7021	-	-	-
Respiratory Distress								
Yes	2429	58,79	5171	32,78	7600	3,51	3,231-3,810	<0,001
No	972	23,52	7260	46,02	8232	-	-	-
O2 Saturation <95%								
Yes	2490	60,26	3610	22,88	6100	5,62	5,174-6,100	<0,001
No	1036	25,07	8438	53,49	9474	-	-	-
Puke								
Yes	354	8,57	1491	9,45	1845	0,94	0,831-1,064	0,331
No	2539	61,45	1057	6,70	3596	-	-	-
Diarrhea								
Yes	342	8,28	1429	9,06	1771	0,95	0,837-1,076	0,411
No	2550	61,71	10108	64,07	12658	-	-	-
Other Symptoms								
Yes	1652	39,98	6745	42,75	8397	0,98	0,906-1,064	0,653
No	1373	33,23	5504	34,89	6877	-	-	-

133

134 Among the pre-existing comorbidities in pregnant and postpartum women with Covid-19, the
 135 diagnosis of heart disease (OR=1.49; CI=1.282-1.721), asthma (OR=1.45; CI=1.205-1.735), diabetes
 136 (OR=1.20; CI=1.039-1.381), obesity (OR=2.37; CI=2.053-2.733), kidney disease (OR=1.69;
 137 CI=1.132-2.510) or the simple presence of comorbidity (OR=1.53; CI=1.423-1.639) were associated
 138 with a maternal near miss (Table 3).

139

140 Table 3 – Univariate analysis of hospitalizations of pregnant and postpartum women with Covid-19
 141 according to pre-existing comorbidities. Brazil, 2020-2022

	UTI		Infirmary		Total	OR	IC	p-value
	n (4132)	%	n (15776)	%				
Comorbidities								
Yes	2604	63,02	8320	52,74	10924	1,53	1,423-1,639	<0,001
No	1528	36,98	7456	47,26	8984	-	-	-
Cardiopathy								
Yes	304	7,36	729	4,62	1033	1,49	1,282-1,721	<0,001
No	1312	31,75	4674	29,63	5986	-	-	-
Haematological disease								
Yes	28	0,68	65	0,41	93	1,46	0,935-2,286	0,096
No	1512	36,59	5132	32,53	6644	-	-	-
Down syndrome								
Yes	7	0,17	23	0,15	30	1,03	0,440-2,397	0,952
No	1531	37,05	5164	32,73	6695	-	-	-
Liver disease								
Yes	10	0,24	32	0,20	42	1,06	0,520-2,161	0,873
No	1516	36,69	5141	32,59	6657	-	-	-
Asthma								
Yes	184	4,45	439	2,78	623	1,45	1,205-1,735	<0,001
No	1406	34,03	4851	30,75	6257	-	-	-
Diabetes								
Yes	314	7,60	895	5,67	1209	1,20	1,039-1,381	0,013

No	1332	32,24	4548	28,83	5880	-	-	-
Obesity								
Yes	380	9,20	592	3,75	972	2,37	2,053-2,733	<0,001
No	1268	30,69	4679	29,66	5947	-	-	-
Neurological disease								
Yes	32	0,77	75	0,48	107	1,45	0,954-2,201	0,082
No	1504	36,40	5109	32,38	6613	-	-	-
Pneumopatias								
Yes	29	0,70	75	0,48	104	1,31	0,852-2,024	0,217
No	1505	36,42	5112	32,40	6617	-	-	-
Immunodeficiency								
Yes	34	0,82	123	0,78	157	0,94	0,639-1,378	0,747
No	1491	36,08	5063	32,09	6554	-	-	-
Kidney disease								
Yes	37	0,90	75	0,48	112	1,69	1,132-2,510	0,010
No	1486	35,96	5078	32,19	6564	-	-	-

142

143

144

145

146

147

148

149

150

151

Regarding the analysis of clinical characteristics of pregnant and postpartum women hospitalized with Covid-19, it was associated with ICU admission being postpartum (OR=1.41; CI=1.299-1.538), having received invasive ventilatory support (OR=94.77; CI=79.039-113.644), non-invasive ventilation (OR=5.53; CI=5.044-6.068), not having been vaccinated against influenza (OR=1.40; CI=1.243-1.585) and/or against Covid-19 (OR=1.71; CI=1.520-1.924) and having received antiviral treatment (OR=1.14; CI=1.003-1.290) (Table 4).

Table 4 – Univariate Analysis of Hospitalizations of Pregnant and Postpartum Women with Covid-19 by Clinical Characteristics Brazil, 2020-2022

	UTI		Infirmity		Total	OR	IC	p-value
	n (4132)	%	n (15776)	%				
Obstetric Population								
Pregnant	3219	77,90	13139	83,28	16358	-	-	-
Postpartum	913	22,10	2637	16,72	3550	1,41	1,299-1,538	<0,001
Ventilatory support								
Invasive	1217	29,45	159	1,01	1376	94,77	79,039-113,644	<0,001
Non-invasive	1928	46,66	4315	27,35	6243	5,53	5,044-6,068	<0,001
No	735	17,79	9101	57,69	9836	-	-	-
Nosocomial Infection								
Yes	59	1,43	187	1,19	246	1,24	0,926-1,671	0,148
No	3203	77,52	12626	80,03	15829	-	-	-
Flu vaccine								
Yes	420	10,16	2620	16,61	3040	-	-	-
No	1119	27,08	4973	31,52	6092	1,4	1,243-1,585	<0,001

Covid* Vaccine

Yes	447	10,82	2537	16,08	2984	-	-	-
No	1318	31,90	4374	27,73	5692	1,71	1,520-1,924	<0,001

Antiviral

Yes	350	8,47	1242	7,87	1592	1,14	1,003-1,290	0,045
No	2750	66,55	11097	70,34	13847	-	-	-

152 * The vaccine against COVID-19 was not made available to pregnant and postpartum women in Brazil until April 2021,
 153 with a gradual introduction between regions of the country and between age groups, which means that many pregnant
 154 and postpartum women in the study population did not receive the vaccine until the time of hospitalization (MINISTRY
 155 OF HEALTH, 2021).
 156

157 In logistic regression analysis, the independent significant factors associated with a near miss
 158 in pregnant and postpartum women with Covid-19 were being black/brown (ORaj=0.83; CI=0.720-
 159 0.966), having O2 saturation <95% (ORaj=1.92; CI=1.609-2.299), presenting dyspnea (ORaj=1.47;
 160 CI=1.216-1.778), having comorbidities (ORaj=1.49; CI=1.282-1.737), requiring invasive
 161 mechanical ventilation (ORaj=44.99; CI=33.099-61.167), requiring non-invasive mechanical
 162 ventilation (ORaj=3.32; CI=2.754-3.994), and not having been vaccinated against influenza
 163 (ORaj=1.18; CI=1.006-1.390), with all these factors controlled by yellow and indigenous race/color
 164 (Table 5).

165

166 Table 5 – Final multiple logistic regression model of factors associated with outcome in hospitalized
 167 pregnant and postpartum women with Covid-19, according to ICU admission. Brazil, 2020-2022

	ORaj*	IC(95%)	p
Maternal characteristics			
Race/black/brown color	0,83	0,720-0,966	0,016
Signs and symptoms			
O2 Saturation <95%	1,92	1,609-2,299	<0,001
Dyspnea	1,47	1,216-1,778	<0,001
Pre-existing comorbidities			
Comorbidity present	1,49	1,282-1,737	<0,001
Clinical features			
Invasive ventilatory support	44,99	33,099-61,167	<0,001
Noninvasive ventilatory support	3,32	2,754-3,994	<0,001
Not having a flu vaccine	1,18	1,006-1,390	0,042

168 * Adjusted for yellow and indigenous race/color.

169

170 **DISCUSSION**

171

172 This study was able to identify the characteristics of women who developed a maternal near
173 miss due to Covid-19. Factors associated with maternal near misses were black/brown color, history
174 of comorbidities, O₂ saturation <95%, dyspnea, need for invasive and non-invasive ventilatory
175 support, and no influenza vaccination.

176 Regarding the socio-demographic profile, black and brown women were predominant,
177 demonstrating that they are more susceptible to Covid-19 infection. However, the proportion of
178 black and brown women admitted to intensive care was lower, which may be related to their lack of
179 access to health services and technologies. The smaller number of black and brown pregnant and
180 postpartum women admitted to ICU beds compared with those admitted to wards confounded the
181 odds ratio, and black/brown race/color could be confused with a protective factor. However, a large
182 body of literature shows the opposite, with whites being the most hospitalized population and
183 black/browns having higher mortality rates.²⁰

184 In this sense, the historical difficulties of this population group in accessing the health system
185 are notorious, as evidenced by unfavorable indicators such as maternal mortality, since they represent
186 approximately 65% of the total maternal deaths in the country.²¹

187 Another important aspect is the social inequality between whites and blacks/brown. A survey
188 conducted during the pandemic showed that the per capita household income of the white Brazilian
189 population is much higher. In July 2020, the per capita income of whites was R\$ 1,639.00, while that
190 of the black and brown population was R\$ 971.00.²² Therefore, social inequalities deserve attention
191 due to their strong relationship with situational health indicators.

192 In terms of signs and symptoms, there was a predominance of O₂ saturation <95% and
193 dyspnea. Physiologically and anatomically, pregnant women undergo pregnancy-related changes that

194 increase the risk of complications and death from Covid-19. Pulmonary physiology undergoes
195 hormonal and functional changes that make it less tolerant to hypoxia. In early pregnancy,
196 progesterone levels act on the brainstem to increase respiratory rate and tidal volume, and chest wall
197 compliance and airway resistance decrease. In the last trimester, the uterus constricts the diaphragm,
198 reducing total lung capacity.²³

199 In addition to these respiratory changes, there are important immunologic changes that place
200 pregnant and postpartum women at risk for developing more severe respiratory infections, which can
201 lead to hypoxemia, which in combination with pulmonary infection can lead to vasoconstriction and
202 worsening of the condition.²³

203 Dyspnea is very common in Covid-19, reported in 81% of hospitalized patients. The presence
204 of this symptom is related to the severity of the disease, being less common in the early phase.²⁴ When
205 present, dyspnea occurs because the disease uses angiotensin-converting enzyme 2 (ACE 2) as a
206 receptor, which is widely distributed in pulmonary, intestinal, renal, and vascular tissues. As a result,
207 cytokines are released that damage lung, kidney, and heart tissue and worsen organ system function.²⁵

208 Among pre-existing comorbidities, cardiac disease, hematologic disease, diabetes, asthma,
209 neurologic disease, renal disease, pulmonary disease, and obesity are considered risk factors for
210 adverse outcomes in coronavirus-infected patients.²⁶⁻²⁷

211 In this regard, the present study found that most pregnant and postpartum women had
212 comorbidities. A review of 77 studies concluded that advanced maternal age, high body mass index,
213 and previous comorbidities were the most relevant risk factors in pregnant women who developed
214 the most severe form of the disease and required intensive care. These findings suggest a risk profile
215 for adverse maternal outcomes such as ICU admission and clinical deterioration, which may progress
216 to death.²⁸

217 Among the most relevant clinical characteristics, the use of ventilatory support (invasive and
218 noninvasive) stands out. Studies conducted in Brazil showed that 20% of pregnant and postpartum

219 women with the severe acute respiratory syndrome (SARS) required ICU admission, 22% required
220 noninvasive ventilation, and 10 to 25% required invasive ventilation.²⁹⁻³⁰

221 In light of Covid-19 and SARS, ventilatory support has become a common requirement in the
222 management of affected patients. In several contexts, the use of invasive techniques for oxygenation
223 was higher than that of non-invasive techniques, with a significant increase in 2021, due to the
224 prolonged and fatal nature of the disease, indicating the intensification of care to reverse the severe
225 condition.³¹

226 Regarding vaccination against influenza (H1N1), an association has been observed with
227 hospitalizations for Covid-19. In this sense, unvaccinated patients had a higher risk of ICU admission,
228 confirming studies showing that vaccination reduces the number of positive tests for Covid-19 and
229 improves clinical outcomes.³²

230 Another study suggests that influenza vaccination is associated with a lower risk of
231 cardiovascular events and reduces the likelihood of overlapping secondary infections, as patients with
232 Covid-19 can acquire the influenza virus at the same time.³³

233 Although some international studies have shown high morbidity and mortality in pregnant and
234 postpartum women with the severe form of COVID-19, the clinical course of the disease in pregnant
235 women from other countries has not shown a significant increase in the rate of outcomes with worse
236 prognosis compared to the general population and the Brazilian reality.³⁴⁻³⁵

237 As a result, WHO, together with the American College of Gynecology and Obstetrics,
238 observed an increase in weekly deaths among pregnant and postpartum women and included this
239 group as a priority in the country's vaccination campaigns as an important preventive measure, given
240 that the effectiveness of the vaccine is comparable to that of nonpregnant women.³⁶ In light of these
241 findings and the pandemic scenario, it is essential that health services promote the vaccination of the
242 obstetric population, which has been shown to have an impact on the development of severe cases,
243 particularly respiratory disease.

244 Given these findings and the pandemic scenario, health services must promote the vaccination
245 of the obstetric population, which has been shown to have a positive impact on the development of
246 severe cases, particularly those with respiratory failure requiring ventilatory support and invasive
247 ventilation, reaffirming the importance of vaccination against Covid-19.

248 It is also necessary to implement new protocols and preventive measures aimed at combating
249 and preventing new transmissions of the virus, culminating in new configurations and adaptations of
250 assistance to pregnant and postpartum women. In addition, policies that address racial inequalities
251 are essential to contribute to the improvement of care, which must comply with the principles of the
252 Single Health System (SUS) in terms of universality, inclusiveness, and equity.

253 It is worth mentioning that the present study has some limitations, mainly related to the use of
254 secondary data, which are subject to underreporting or completion problems. However, the health
255 information systems of the Brazilian Ministry of Health are of great value, as they allow the
256 epidemiological study of diseases at the national level, which helps in the planning of health policies
257 and programs. In addition, during the Covid-19 pandemic, SIVEP-Influenza was the main source of
258 data on Covid-19 hospitalizations in the country.

259

260 **CONCLUSION**

261

262 ICU admissions of pregnant and postpartum women with Covid-19, considered in this study
263 as maternal near misses, were associated with black and brown race/color, with O₂ saturation <95%,
264 with dyspnea, with comorbidities, with a need for invasive or non-invasive ventilatory support, and
265 with lack of influenza vaccination.

266 In this sense, recognition of the factors involved in maternal near misses can help to direct
267 actions to qualify the care of women in pregnancy and the puerperium. In addition, it allows us to
268 identify which women are more likely to have a fatal outcome, thus allowing early intervention.

269

270 ACKNOWLEDGEMENTS

271

272 We would like to thank the Universidade Estadual de Maringá for their support in carrying out the
273 research and the Centro Universitário Uningá and Duke University for the partnership.

274

275 REFERENCES

276

277 1. Organização Mundial da Saúde (OMS). OMS declara pandemia do novo Coronavírus.

278 Universidade Aberta do Sus (UNA-SUS). 11 de março de 2020 [Cited 2022 jan 23].

279 Available from: [https://www.unasus.gov.br/noticia/organizacao-mundial-de-saude-declara-](https://www.unasus.gov.br/noticia/organizacao-mundial-de-saude-declara-pandemia-de-coronavirus)
280 [pandemia-de-coronavirus.](https://www.unasus.gov.br/noticia/organizacao-mundial-de-saude-declara-pandemia-de-coronavirus)

281

282 2. Zaigham M, Andersson O. (2020). Resultados maternos e perinatais com COVID-19: Uma

283 revisão sistemática de 108 gestações. Acta Obstet Gynecol Scand. Available from:

284 <https://doi.org/10.1111/aogs.13867>.

285

286 3. Yan J, Guo J, Fan C, et al. (2020). Coronavirus disease 2019 in pregnant women: a report

287 based on 116 cases. Am J Obstet Gynecol, Available from:

288 <https://pubmed.ncbi.nlm.nih.gov/32335053/>

289

290 4. Fernandes JN, Rezende MCD, Otsubo BYV, et al. (2022). Correlation between COVID-19
291 and gestational complications: a systematic review. Brazilian Journal of Health Review, v5,

292 n 2, 6405-6411 [Cited 2023 apr 11]. Available from: <https://doi.org/10.34119/bjhrv5n2-213>

293

- 294 5. Fundação Oswaldo Cruz (Fiocruz). (2021). Boletim Extraordinário, 25 de junho de 2021.
295 [Cited 2022 jan 24]. Available from:
296 [https://agencia.fiocruz.br/sites/agencia.fiocruz.br/files/u34/boletim_extraordinario_2021-](https://agencia.fiocruz.br/sites/agencia.fiocruz.br/files/u34/boletim_extraordinario_2021-junho-23-parte2-pags09-17.pdf)
297 [junho-23-parte2-pags09-17.pdf](https://agencia.fiocruz.br/sites/agencia.fiocruz.br/files/u34/boletim_extraordinario_2021-junho-23-parte2-pags09-17.pdf)
298
- 299 6. Bastos, LS, Niquini RP, Lana RM, et al. (2020) COVID-19 and hospitalizations for SARI in
300 Brazil: a comparison up to the 12th epidemiological week of 2020. Cad Saúde Pública.
301 Available from:
302 <https://www.scielo.br/j/csp/a/KQxzHZdFHcPx5CftPXZKwgs/?format=pdf&lang=en>
303
- 304 7. Almeida JP, Santana VS, Santos KM, et al. (2021). Internações por SRAG e óbitos por
305 Covid-19 em gestantes brasileiras: uma análise da triste realidade. Brazilian Journal of
306 Health Review. Available from:
307 <https://www.brazilianjournals.com/index.php/BJHR/article/view/31570>
308
- 309 8. Motta CT (2021). Subsidies for the compliance of SDG 3.1 of the 2030 Agenda: an analysis
310 of maternal mortality in Brazil from 1996 to 2018. M. Sc. Dissertação, Escola Nacional de
311 Saúde Pública Sérgio Arouca. Available from:
312 [https://www.arca.fiocruz.br/bitstream/handle/icict/48404/caio_tavares_motta_ensp_mest_20](https://www.arca.fiocruz.br/bitstream/handle/icict/48404/caio_tavares_motta_ensp_mest_2021.pdf;jsessionid=8396E5143894CDCB6684F3F1C8D35951?sequence=2)
313 [21.pdf;jsessionid=8396E5143894CDCB6684F3F1C8D35951?sequence=2](https://www.arca.fiocruz.br/bitstream/handle/icict/48404/caio_tavares_motta_ensp_mest_2021.pdf;jsessionid=8396E5143894CDCB6684F3F1C8D35951?sequence=2)
314
- 315 9. Instituto de Pesquisa Econômica Aplicada (IPEA). ODS – Metas Nacionais dos Objetivos
316 de Desenvolvimento Sustentável. 2018. [Cited 2022 jan 24]. Available from:
317 [https://repositorio.ipea.gov.br/bitstream/11058/8855/1/Agenda_2030_ods_metas_nac_dos_o](https://repositorio.ipea.gov.br/bitstream/11058/8855/1/Agenda_2030_ods_metas_nac_dos_obj_de_desenv_susten_propos_de_adequa.pdf)
318 [bj_de_desenv_susten_propos_de_adequa.pdf](https://repositorio.ipea.gov.br/bitstream/11058/8855/1/Agenda_2030_ods_metas_nac_dos_obj_de_desenv_susten_propos_de_adequa.pdf)

319

320 10. Observatório Obstétrico Brasileiro (OOBr). Óbitos de Gestantes e Puérperas. 2022. [Cited
321 2022 jan 24]. Available from: [https://observatorioobstetrico.shinyapps.io/obitos-grav-
322 puerp](https://observatorioobstetrico.shinyapps.io/obitos-grav-
322 puerp) .

323

324 11. Souza ASR, Amorim MMR. (2021). Maternal mortality by Covid-19 in Brazil. Rev Bras
325 Saúde Matern Infant. Available from:
326 <https://www.scielo.br/j/rbsmi/a/R7MkrmCgdmyMpBcL7x77QZd/?format=pdf&lang=en>

327

328 12. Organização Mundial da Saúde (OMS). Avaliação da qualidade do cuidado nas
329 complicações graves da gestação: A abordagem do near miss da OMS para a saúde materna.
330 2011. [Cited 2022 jan 27]. Available from: [https://www.paho.org/clap/dmdocuments/CLAP-
331 Trad05pt.pdf](https://www.paho.org/clap/dmdocuments/CLAP-
331 Trad05pt.pdf)

332

333 13. Lima R. (2021). Direito sexuais e direitos reprodutivos das mulheres. Escola de Assistência
334 Jurídica da Defensoria Pública. [Cited 2022 aug 18]. Available from: [Cartilha Direito
335 Sexuais e Reprodutivos Das Mulheres | PDF \(scribd.com\)](#)

336

337 14. Malta M, Cardoso LO, Bastos FI et al. (2010) STROBE initiative: guidelines on reporting
338 observational studies. Rev Saude Publica, v. 44, n. 3, p. 559–565. DOI: [\[OBJ\]](#)

339

340 15. Instituto Brasileiro de Geografia e Estatística (IBGE). (2022). Projeção da população do
341 Brasil e das Unidades da Federação. [Cited 2022 jun 21]. Available from:
342 <https://ibge.gov.br/apps/populacao/projecao/>

343

- 344 16. Brasil. (2008). Ministério da Saúde. Portaria nº 1,119, de 5 de junho de 2008. Regulamenta
345 a vigilância de óbitos maternos. [Cited 2022 jun 21]. Available from:
346 https://bvsmms.saude.gov.br/bvs/saudelegis/gm/2008/prt1119_05_06_2008.html
347
- 348 17. Bahia. Secretária da Saúde do Estado da Bahia. Guia Rápido SIVEP – Gripe. 2020. [Cited
349 2022 jun 14]. Available from: [https://docs.bvsalud.org/biblioref/2021/03/1147534/guia-
350 rapido-sivep-gripe_agosto-2020.pdf](https://docs.bvsalud.org/biblioref/2021/03/1147534/guia-rapido-sivep-gripe_agosto-2020.pdf)
351
- 352 18. Garson GD. (2014). Logistic Regression: Binary & Multinomial [Cited 2022 aug 21].
353 Available from:
354 [file:///C:/Users/marci/Downloads/Logistic%20Regression_%20Binary%20and%20Multino
355 mial%20\(%20PDFDrive%20\).pdf](file:///C:/Users/marci/Downloads/Logistic%20Regression_%20Binary%20and%20Multinomial%20(%20PDFDrive%20).pdf)
356
- 357 19. Brasil. (2016). Resolução nº 510, de 7 de abril de 2016. [Cited 2022 jun 21]. Available
358 from: <https://conselho.saude.gov.br/resolucoes/2016/Reso510.pdf>
359
- 360 20. Zeiser FA, Donida B, Costa CA, et al. (2022) First and second COVID-19 waves in Brazil:
361 A cross-sectional study of patients’ characteristics related to hospitalization and in-hospital
362 mortality. *The Lancet Regional Health - Americas*, v. 6, n. November, p. 1–16, 2022. DOI:
363 <https://doi.org/10.1016/j.lana.2021.100107>
364
- 365 21. Lima LCBB. (2021). A ocorrência de falhas assistenciais durante o trabalho de parto: uma
366 revisão integrativa. Trabalho de Conclusão de Curso. Pontifícia Universidade Católica de
367 Goiás. [Cited 2023 mar 07]. Available from:
368 <https://repositorio.pucgoias.edu.br/jspui/handle/123456789/2510?mode=full>

369

370 22. Instituto de Pesquisa Econômica e Aplicada (IPEA). (2021). A pandemia de Covid-19 e a
371 desigualdade racial de renda. Boletim de Análise Político-Institucional, n. 26, março 2021.
372 [Cited 2023 mar 08]. Available from:
373 https://repositorio.ipea.gov.br/bitstream/11058/10519/1/BAPI_26_DesRacial.pdf

374

375 23. Czeresnia RM, Trad ATA, Britto ISW, et al. (2020). Sars-Cov2 and Pregnancy: a review of
376 the facts. Rev Bras Ginecol Obstet, v42, n9, 562-568. [Cited 2022 aug 21]. Available from:
377 <https://doi.org/10.1055/s-0040-1715137>

378

379 24. Nascimento IJB, Groote TC, O’Mathúna DP, et al. (2020). Clinical, laboratory and
380 radiological characteristics and outcomes of nves coronavírus (SARS-CoV-2) infection in
381 humans: A systematic review and series of meta-analyses. PLoS One. [Cited 2022 aug 22].
382 Available from: [10.1371/journal.pone.0239235](https://doi.org/10.1371/journal.pone.0239235)

383

384 25. Hoffmann M, Kleine-Weber H, Schoeder S, et al. (2020) SARS-CoV-2 cell entry depends
385 on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. Cell, v
386 181, 271-280. [Cited 2022 Aug 22]. Available from:
387 <https://doi.org/10.1016/j.cell.2020.02.052>

388

389 26. Elshafeey F, Magdi R, Hindi N, et al. (2020). Asistematic scoping review of Covid-19
390 during pregnancy and childbirth. International Journal of Gynecology & Obstetrics, v1, n2.
391 [Cited 2022 Aug 21]. Available from: <https://doi.org/10.1002/ijgo.13182>

392

- 393 27. Jering KS, Claggett BL, Cunningham JW, et al. (2021). Clinical characteristics and
394 outcomes of hospitalized women giving birth and without Covid-19. *JAMA Intern Med*,
395 181, v 5, 714-717. [Cited 2022 aug 21]. Available from:
396 <https://doi.org/10.1001/jamainternmed.2020.9241>
397
- 398 28. Allotey J, Fernandez S, Bonet M, et al. (2020). Clinical manifestations, risk factors, and
399 maternal and perinatal outcomes of coronavirus disease 2019 in pregnancy: living
400 systematic review meta-analysis. *The BMJ*, 370, m3320. [Cited 2022 aug 22]. Available
401 from: Available from: <https://doi.org/10.1136/bmj.m3320>
402
- 403 29. Godoi APN, Bernardes GCS, Almeida NA, et al. (2021) Severe Acute Respiratory
404 Syndrome by COVID-19 in pregnant and postpartum women. *Rev Bras Saúde Matern Infant*,
405 v 21, n 2, 5461-5469. [Cited 2022 aug 22]. Available from: [https://doi.org/10.1590/1806-](https://doi.org/10.1590/1806-9304202100S200008)
406 [9304202100S200008](https://doi.org/10.1590/1806-9304202100S200008)
407
- 408 30. Leal LF, Merckx J, Fell DB, et al. (2021). Characteristics and outcomes of pregnant woman
409 with SARS-CoV-2 infection and other severe acute respiratory infections (SARI) in Brazil
410 from January to November 2020. *Braz J Infect Dis*. [Cited 2022 aug 24]. Available from:
411 [10.1016/j.bjid.2021.101620](https://doi.org/10.1016/j.bjid.2021.101620)
412
- 413 31. Takemoto MLS, Menezes MO, Andreucci CB, et al. (2020). The tragedy of COVID-19 in
414 Brazil: 124 maternal deaths and counting. *Int J Gynaecol Obstet*, v 151, n 1, 154-156. [Cited
415 2022 aug 22]. Available from: Available from: <https://doi.org/10.1002/ijgo.13300>
416

- 417 32. Conlon A, Ashur C, Washer L. et al. (2021). Impacto f the influenza vaccine on Covid-19
418 infection rates and severity. American Journal of Infection Control, v 49, 694-700. [Cited
419 2022 aug 22]. Available from: <https://doi.org/10.1016/j.ajic.2021.02.012>
420
- 421 33. Behrouzi B, Campoverde MVA, Liang K, et al. (2020). Influenza Vaccination to Reduce
422 Cardiovascular Morbidity and Mortality in Patients With COVID-19. Journal of the
423 American College of Cardiology, v. 76, n. 15, p. 1–19, 2020. DOI:
424 [10.1016/j.jacc.2020.08.028](https://doi.org/10.1016/j.jacc.2020.08.028)
425
- 426 34. Dashraath P, Wong JLJ, Lim MXK, et al. (2020). Coronavirus Disease 2019 (COVID-19)
427 pandemic and pregnancy. Am J Obstet Gynecol, v 222, n 6, 521-531. [Cited 2022 aug 18].
428 Available from: <https://doi.org/10.106/j.ajog.2020.03.021>
429
- 430 35. Pacagnella RCP, Nakamura-Pereira M, Gomes-Sponholz F, et al. (2018). Maternal mortality
431 in Brazil: Proposals and strategies for its reduction. Rev Bras Ginecol Obstet, v 40, 501-
432 506). [Cited 2022 aug 18]. Available from: <https://doi.org/10.1055/s-0038-1672181>
433
- 434 36. Rodrigues FO, Antunes Neto A, Silva RG, et al. (2021). Desfechos maternos da COVID-19
435 e atualizações sobe a vacinação em gestantes e puérperas. Revista Brazilian Journal of
436 Development, v 7, n 6, 57232-57247. [Cited 2021 jan 27]. Available from:
437 <https://doi.org/10.34117/bjdv7n6-227>
438
439
440