Title: Endovascular treatment of ruptured very small intracranial aneurysms: a

systematic review and meta-analysis

Authors: Hidetoshi Matsukawa, M.D., Ph.D. 1, Sameh Samir Elawady, M.D. 1,

Mohammad Mahdi Sowlat, M.D.¹, Sami Al Kasab, M.D.¹, Kazutaka Uchida, M.D.,

Ph.D.^{1,2}, Shinichi Yoshimura, M.D., Ph.D.^{1,2}, Alejandro M Spiotta, M.D.¹

Affiliation

1 Department of Neurosurgery, Medical University of South Carolina

2 Department of Neurosurgery, Hyogo Medical University

Corresponding author

Alejandro M Spiotta

Medical University of South Carolina

Department of Neurosurgery, 171 Ashley Avenue, Charleston, South Carolina 29425.

E-mail: spiotta@musc.edu.

Cover title: EVT of RVSIAs

Word count: 5144/6000 words

Number of tables and figures: 6

Abstract (298 of 300 words)

Background: Endovascular treatment (EVT) is widely accepted for intracranial

aneurysms due to its safety and efficacy. However, EVT of ruptured very small

intracranial aneurysms (RVSIA) (≤3 mm) is still challenging and the risk-benefit ratio

of EVT remains unclear. The aim of this study was to evaluate the safety and efficacy of

EVT of RVSIA.

Methods: We performed a systematic review and meta-analysis of the studies on EVT

of RVSIA. Pooled prevalence rates were calculated for initial and follow-up complete

occlusion rates (Raymond Roy Grade 1), recanalization, retreatment, long-term

favorable outcome (modified Rankins scale score 0 to 2 or Glasgow Outcome Scale 4 or

5), procedure-related complications (coil herniation, thromboembolism, and

intraprocedural re-rupture), and procedure-related mortality. Pooled odds ratios were

calculated to compare these outcomes between simple coiling and stent-assisted coiling

(SAC).

Results: Of the 600 studies screened, 24 studies with a total of 1355 RVSIAs treated

with EVT were included. The initial and follow-up complete aneurysm occlusion rates

were 64% (95% confidence interval [CI]: 52-74%) and 85% (95% CI: 74-92%). The

rates of recanalization and retreatment were 6% (95% CI: 3-10%) and 3% (95% CI: 2-

4%). The favorable long-term follow-up outcome was observed in 91% (95% CI: 89-

93%) of patients. The rates of coil herniation, thromboembolism, and intraprocedural

rupture were 2% (95% CI: 1–8%), 4% (95% CI: 3–6%), and 4% (95% CI: 2–7%),

respectively. Mortality was 3% (95% CI: 2-4%). Comparison of outcomes between

simple coiling and SAC revealed no significant difference, except for a higher

likelihood of recanalization in the coiling group (Odds ratio, 3.51 [95% CI, 1.31–9.45]).

Conclusions: Our meta-analysis demonstrates that EVT for RVSIA is a feasible,

effective, and safe approach that is associated with favorable clinical outcomes in both

the short and long term.

Abbreviations: EVT = endovascular treatment; RVSIA = ruptured very small

intracranial aneurysm; SAC = stent-assisted coiling

Key words: endovascular treatment; meta-analysis; ruptured very small intracranial

aneurysms; systematic review.

Introduction

The incidence of aneurysmal subarachnoid hemorrhage (aSAH) is approximately 6–11 per 100 000 persons per year. ^{1,2} In spite of the understanding of the pathophysiology, around 30% of patients who survive after aSAH will not regain full independence. ³ Very small intracranial aneurysm (\leq 3 mm) represents 13.2% - 15.1% of all intracranial aneurysms. ^{4,5} Although the risk of rupture is known to increase with the increasing size of an aneurysm^{6,7}, a large number of patients have subarachnoid hemorrhage as a result of rupture of intracranial aneurysms with diameters \leq 5 mm. ⁸⁻¹²

Very small aneurysms may be too small to accept a clip without narrowing or tearing the parent vessel. ^{13,14} Endovascular treatment (EVT) of such aneurysms is also considered to be challenging due to the thin fragile aneurysm wall, with limited space to obtain a stable microcatheter position for coil deployment, and with a high risk of intraprocedural rupture. ^{4,15-18} However, recent advances in endovascular techniques and devices have led to the rapid evolution of coil embolization techniques for intracranial aneurysms, which might allow for safer coil approach to ruptured very small intracranial aneurysm (RVSIA). In fact, some studies demonstrated successful EVT of RVSIA with stent-assisted coiling (SAC)¹⁹⁻²¹ although the number of patients was small

to reach a meaningful conclusion. Therefore, the risk-benefit ratio of EVT of RVSIA remains unclear. To examine the contemporary safety and efficacy of EVT of RVSIA and to compare outcomes between primary coiling and SAC, we aimed to

systematically review the literature and conduct a meta-analysis.

Material and Methods

Search strategy

This study was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. A comprehensive search of PubMed and Scorpus databases up to February 18, 2023, was conducted to identify relevant studies. We used the following keywords: "burst," "ruptured," "small," "tiny," "3mm," "aneurysm," "brain," "intracranial," "berry," "cerebral," "circle of Willis," "saccular," "fusiform," "embolize*," "coil," "stent," and "endovascular." These keywords were

Selection criteria

combined using "OR" or "AND."

Our study included original articles that reported the outcomes of EVT for ruptured very small intracranial aneurysms without any restrictions on age, sex, race, ethnicity,

country of origin, or publication date. The article was not eligible if it met any of the following exclusion criteria: (1) aneurysm size > 3mm; (2) blood blister and dissecting aneurysms; (3) unreliable, non-extractable, duplicated, and overlapped data sets; (4) no full text available; (5) conference/poster abstracts; (6) case reports/series of ≤ 10 cases; or (7) review articles. Studies were identified based on the abovementioned eligibility criteria by two independent reviewers. Any disagreement was resolved through cooperative discussion.

Data extractions

Two independent reviewers evaluated the studies and extracted the relevant the data. The extracted data included the first author, year of publication, study design, number of patients with coiling or SAC, age, gender, posterior circulation aneurysm, Hunt-Hess grade²², size of sac and neck, and angiographic follow-up period (month), technical success, initial and long-term complete occlusion rates, recanalization rate, retreatment long-term favorable neurologic outcome, intraoperative complications rate, (intraprocedural parent artery vasospasm, coil herniation, thromboembolic complications, and intraprocedural re-rupture), and mortality. Any discrepancies between reviewers were resolved through discussion. Complete aneurysm occlusion was defined as Raymond-Roy occlusion classification I.²³ Favorable outcome was set as

modified Rankin scale 0 to 2²⁴ or Glasgow Outcome Scale 4 or 5.²⁵

Quality assessment of the included studies

Risk of bias in each included study was assessed by two independent reviewers using the Risk of Bias In Non-randomized Studies of Interventions (ROBINS-I) tool. This tool evaluates the methodological quality of the studies in various domains, including confounding factors, selection of participants into the study, classification of interventions, deviations from intended interventions, missing data, measurement of outcomes, and selection of the reported results.²⁶

Statistical analysis

We analyzed all the data using R software version 4.2.3. A "meta" package was used to calculate pooled prevalence (%) or pooled odds ratios, and their corresponding 95% confidence intervals (CI) using either a fixed-effect or random-effect model, depending on the degree of heterogeneity present among the included studies. Heterogeneity was evaluated using Q statistics and I2 test. We considered I2 value > 50% or P-value < 0.05 to indicate substantial heterogeneity. Due to the non-comparative nature of most included studies and the small number of comparative studies (< 10 per analysis), we were unable to test for publication bias or perform meta-regression.

Results

Literature Search

600 studies were initially identified, 470 were excluded after title and abstract screening.

130 studies were assessed for full-text eligibility, and 110 were excluded. Additional 4

studies were identified through manual search of the references of included studies. 24

studies were included in the final analysis. A study flow diagram is shown in Fig 1.

Baseline Characteristics

24 included studies with 1355 patients underwent EVT for RVSIAs. The characteristics of the included studies are summarized in Table 1. Thirteen studies were published from China^{19,21,27-37}; 3 were from USA³⁸⁻⁴⁰; 2 were from Japan^{41,42}; and 1 each from France⁴³, Iran⁴⁴, Korea⁴⁵, Netherland⁴, Taiwan⁴⁶, and UK.⁴⁷ All studies were retrospective cohort except one study. The proportion of male patients ranged from 20% to 65.8%. The mean age ranged from 34.3 to 65.2 years old (standard deviation [SD]:±6.0) and Hunt and Hess grade ranged from 1.4 to 2.8 (SD:±0.35). The median size of aneurysm dome and neck were 2.5 mm (SD:±0.28) and 1.8 mm (SD:±0.27). The mean duration of angiographic follow-up ranged from 4.6 to 28.5 months (standard deviation:±6.7).

Risk of Bias

According to the ROBINS quality assessment tool, 10 of our included studies had a moderate risk of bias and the remaining 14 studies had a low risk (Supplement Fig. 1).

Outcome analysis

Angiographic and Clinical outcomes

Initial technical failure occurred in 2% (95% CI, 1–5%) of patients. Initial complete aneurysm occlusion rate was 64% (95% CI, 52–74%), while follow-up complete occlusion rate was 85% (95% CI, 74–92%). Recanalization and retreatment were observed in 6% (95% CI, 3–10%) and 3% (95% CI, 2–4%) of the coiled RVSIAs, respectively. Favorable outcome was observed in 91% (95% CI, 89–93%).

Adverse Events

Overall complications were seen in 9% (95% CI, 7–12%) of subjects. Parent artery vasospasm, coil herniation, and thromboembolism were observed in 6% (95% CI, 5–9%), 2% (95% CI, 1–8%), and 4% (95% CI, 3–6%), respectively. Intraprocedural re-rupture rate was observed in 4% of RVSIAs (95% CI, 2–7%). Mortality rate was 3% (95% CI, 2–4%).

Coiling and SAC

Proportion of initial and follow-up complete aneurysm occlusion rates were not significantly different between the coiling and SAC (initial, OR 1.11, [95% CI: 0.77–

1.61]; *P*=0.57; follow-up, OR 0.19, [95% CI: 0.01–4.36]; *P*=0.24). Recanalization rate was significantly higher in the coiling group (OR 3.51, [95% CI: 1.31–9.45]; *P*=0.013). Pooling of the 2 studies comparing favorable outcomes between the two groups showed no significant difference (OR 1.00, [95% CI: 0.47-2.15]; *P*=0.99). Additionally, there were no significance between two groups in procedure-related complications including; coil herniation (OR 1.05: 95% CI: 0.17–6.30, *P*=0.96) and intraprocedural rupture (OR 0.64, [95% CI: 0.11–1.70]; *P*=0.23).

Discussion

The results of this meta-analysis of the contemporary EVT of RVSIA demonstrate an acceptable low rate of complications, intraprocedural re-ruptures, and retreatment, and high rate of follow-up complete aneurysm occlusion and long-term follow-up favorable outcome. In addition, comparison of coiling and SAC revealed no significant difference in patient outcomes or complications, except for a higher recanalization rate in the coiling group.

Recently, EVT has been widely used in the treatment of ruptured intracranial aneurysms. The findings of the International Subarachnoid Aneurysm Trial (ISAT) suggested that for patients with SAH who were suitable for both EVT and neurosurgical

clipping, those with EVT were more likely to survive independently after 1 year than neurosurgical clipping. 49 On the other hand, the smallest subset of ruptured intracranial aneurysms, those less than 3mm, are generally considered difficult and high-risk to treat with EVT and therefore are often approached with microsurgical clipping. The main challenges of EVT for RVSIA are related to the geometrical challenges they present as they are often broad-necked with a shallow dome. This results in a smaller margin of error in accessing the aneurysm with a microcatheter, as well as difficulty in maintaining microcatheter stability during coil deployment and maintaining the coils within the aneurysm while preventing prolapse into the parent artery. These factors theoretically lead to difficulty in achieving satisfactory coil packing, and higher risk of intraprocedural re-rupture and early rebleeding. 4,50 Another previous study showed that in comparison with EVT of larger ruptured aneurysms, tiny aneurysm coiling was related to an increased absolute technical failure rate of 13.7% suggesting that this subset of aneurysms represented the most challenging of all to approach with EVT.⁵¹ However, as challenging as these aneurysms may be to treat, over time there have been considerable advances in EVT devices and techniques; ranging from improved guide catheters microwires and coils. This contemporary meta-analysis demonstrates high technical success rate (98%) and low intraprocedural rupture rate (4%) among RVSIA

treated with EVT. The initial and follow-up complete aneurysm occlusion rate was 64% and 85%. During follow-up, recanalization observed in only 6% and retreatment was performed in just 3%. Favorable outcome was achieved in 91% of subjects and mortality rate was 3%. In addition to re-rupture, other intraprocedural events such as thromboembolism, coil herniation, and parent artery vasospasm are also significant issues to be avoided. In this analysis, the incidence of thromboembolism, coil herniation, and parent artery vasospasm was 4%, 2%, and 6%, respectively. Compared to the results of previous meta-analysis, the rate of initial complete occlusion, intraprocedural rupture and retreatment was lower, the rate of long-term favorable outcome was higher, and follow-up complete occlusion rate was similar to our meta-analysis (Table 2). This could be in part due to a more conservative coiling approach by treating physicians attempting to avoid the most dreaded complication of all intraprocedural re-rupture. The efficacy and safety of EVT of RVSIAs may be due to not only the accumulated experience of neurointerventional physicians but also the introduction of new microcatheters and microguidewires with improved trackability, pushability, torque, and novel hydrophilic coatings, which could help neurointerventionalists safely access a very small aneurysm dome. Furthermore, the evolution of various kinds of soft and extra soft coils could also have contributed to the higher success rate of EVT of RVSIA.

While relatively safe and effective, 9% and 4% frequency of overall complications and retreatment are acceptable but not insignificant, therefore, neurointerventionalists should keep continue to improve clinical and radiological outcomes in the management of RVSIA with EVT.

Stent placement could help maintain coils inside the aneurysm, thus increasing the coil packing density and decreasing the stent porosity at the neck plane. Stent struts across the aneurysm neck also could provide the structural basis for endothelialization and moderate the hemodynamic condition.²¹ Since the majority of RVSIAs are wide-necked, SAC is often needed from a purely mechanical and geometric sense. However, in the acute stage of subarachnoid hemorrhage, the patient is in a hypercoagulable state and, therefore, SAC may increase the risk of thromboembolic events as well as hemorrhagic complications such as external ventricular drain site bleeds. In fact, Roh et al. reported that the rate of thromboembolic complications in stent-assisted coil embolization of ruptured intracranial aneurysms was approximately twice as high as that in non-stent coil embolization.⁵² This meta-analysis showed no significant difference between coiling and SAC of RVSIAs in the rate of initial and follow-up complete aneurysm occlusion, intraprocedural rupture, coil herniation, and favorable outcome.

Although both coiling and SAC could be selected in the treatment of RVSIAs,

Limitations

This meta-analysis has several limitations. First, moderate heterogeneity was found. Second, there is also bias inherent in the retrospective nature of most studies included. Third, another periprocedural complications, such as extraventricular drainage tract hemorrhage, intraventricular hemorrhage, and aSAH-induced delayed cerebral vasospasm, and antiplatelet managements are also important issues in the management of aSAH, though, these variables could not be sufficiently investigated due to lack of data. Fourth, whether balloon-assisted coiling was performed could not be collected due to missing data. In this regard, some studies showed that balloon use was related to thromboembolic complications and intraprocedural rupture^{4,41}, though the other indicated that it was not associated with intraprocedural rupture.⁵³ Fifth, thorough investigation of delayed post-treatment aneurysm rupture, a relevant concern, was not possible, due to lack of long term data. Finally, mortality, retreatment, thromboembolism, parent artery vasospasm could not be compared between coiling and SAC due to the lack of data. Nevertheless, this contemporary meta-analysis is currently

the largest study assessing the EVT of RVSIAs with more than 1300 subjects.

Conclusion

This meta-analysis demonstrates that EVT is a feasible, effective, and safe approach for the treatment of RVSIA and is associated with favorable clinical outcomes in both the short and long term. Further well- designed randomized controlled trials with large

Acknowledgement:

sample size are warranted to verify our findings.

NA

Source of funding:

NA

Disclosure:

H. Matsukawa received a non-related lecture fee from Daiichi-Sankyo and consulting services fee from B. Braun. K. Uchida received a non-related lecture fee from

Daiichi-Sankyo, Bristol-Myers Squibb, Stryker, and Medtronic. S. Yoshimura received

a non-related lecture fee from Stryker, Medtronic, Johnson & Johnson, Kaneka Medics.

A. Spiotta Research support (STAR) Avail, RapidAI, Penumbra, Microvention,

Medtronic, Stryker, Brain Aneurysm Foundation. Consulting RapidAI, Avail,

Penumbra, Medtronic.

Availability of data and materials

The data that support the findings of this study are available from the corresponding author, Alejandro M Spiotta, upon reasonable request.

References

1. Linn FH, Rinkel GJ, Algra A, van Gijn J. Incidence of subarachnoid hemorrhage:

- Epidemiology of aneurysmal subarachnoid hemorrhage in Australia and New Zealand: incidence and case fatality from the Australasian Cooperative Research on Subarachnoid Hemorrhage Study (ACROSS). Stroke. 2000;31:1843-1850.
- 3. Hop JW, Rinkel GJ, Algra A, van Gijn J. Case-fatality rates and functional outcome after subarachnoid hemorrhage: a systematic review. *Stroke.* 1997;28:660-664.
- 4. van Rooij WJ, Keeren GJ, Peluso JP, Sluzewski M. Clinical and angiographic results of coiling of 196 very small (< or = 3 mm) intracranial aneurysms. *AJNR Am J Neuroradiol*. 2009;30:835-839. doi: 10.3174/ajnr.A1429
- 5. Weir B, Disney L, Karrison T. Sizes of ruptured and unruptured aneurysms in relation to their sites and the ages of patients. *J Neurosurg.* 2002;96:64-70. doi: 10.3171/jns.2002.96.1.0064
- 6. Morita A, Kirino T, Hashi K, Aoki N, Fukuhara S, Hashimoto N, Nakayama T, Sakai M, Teramoto A, Tominari S, et al. The natural course of unruptured cerebral aneurysms in a Japanese cohort. *The New England Journal of Medicine*. 2012;366:2474-2482. doi: 10.1056/NEJMoa1113260
- 7. Wiebers D. Unruptured intracranial aneurysms: natural history, clinical outcome, and risks of surgical and endovascular treatment. *The Lancet*. 2003;362:103-110.
- 8. Beck J, Rohde S, Berkefeld J, Seifert V, Raabe A. Size and location of ruptured and unruptured intracranial aneurysms measured by 3-dimensional rotational angiography. *Surg Neurol*. 2006;65:18-25; discussion 25-17. doi: 10.1016/j.surneu.2005.05.019
- 9. Juvela S, Porras M, Heiskanen O. Natural history of unruptured intracranial aneurysms: a long-term follow-up study. *J Neurosurg*. 1993;79:174-182. doi: 10.3171/ins.1993.79.2.0174
- 10. Juvela S, Porras M, Poussa K. Natural history of unruptured intracranial aneurysms: probability and risk factors for aneurysm rupture. *Neurosurg Focus*. 2000;8:Preview 1.
- 11. Juvela S, Porras M, Poussa K. Natural history of unruptured intracranial aneurysms: probability of and risk factors for aneurysm rupture. *J Neurosurg*. 2008;108:1052-1060. doi: 10.3171/JNS/2008/108/5/1052
- 12. Sonobe M, Yamazaki T, Yonekura M, Kikuchi H. Small unruptured intracranial aneurysm verification study: SUAVe study, Japan. *Stroke*. 2010;41:1969-1977. doi: 10.1161/STROKEAHA.110.585059
- 13. Fujiwara S. Fujii K. Nishio S. Fukui M. Long-term results of wrapping of

- intracranial ruptured aneurysms. *Acta Neurochir (Wien)*. 1990;103:27-29. doi: 10.1007/BF01420188
- 14. Deshmukh VR, Kakarla UK, Figueiredo EG, Zabramski JM, Spetzler RF. Long-term clinical and angiographic follow-up of unclippable wrapped intracranial aneurysms.

 *Neurosurgery**. 2006:58:434-442; discussion 434-442. doi: 10.1227/01.NEU.0000199158.02619.99
- Nguyen TN, Raymond J, Guilbert F, Roy D, Berube MD, Mahmoud M, Weill A. Association of endovascular therapy of very small ruptured aneurysms with higher rates of procedure-related rupture. *J Neurosurg*. 2008;108:1088-1092. doi: 10.3171/JNS/2008/108/6/1088
- 16. Lim YC, Kim BM, Shin YS, Kim SY, Chung J. Structural limitations of currently available microcatheters and coils for endovascular coiling of very small aneurysms. *Neuroradiology*. 2008;50:423-427. doi: 10.1007/s00234-008-0365-y
- 17. Singh V, Gress DR, Higashida RT, Dowd CF, Halbach VV, Johnston SC. The learning curve for coil embolization of unruptured intracranial aneurysms. *AJNR Am J Neuroradiol*. 2002;23:768-771.
- 18. Elijovich L, Higashida RT, Lawton MT, Duckwiler G, Giannotta S, Johnston SC, Cerebral Aneurysm Rerupture After Treatment I. Predictors and outcomes of intraprocedural rupture in patients treated for ruptured intracranial aneurysms: the CARAT study. *Stroke*. 2008;39:1501-1506. doi: 10.1161/STROKEAHA.107.504670
- 19. Zhao R, Shen J, Huang QH, Nie JH, Xu Y, Hong B, Yang PF, Zhao WY, Liu JM. Endovascular treatment of ruptured tiny, wide-necked posterior communicating artery aneurysms using a modified stent-assisted coiling technique. *J Clin Neurosci*. 2013;20:1377-1381. doi: 10.1016/j.jocn.2012.12.012
- 20. Fang C, Li MH, Zhang PL, Wang W, Tan HQ, Xu HW, Zhou B. Endovascular Treatment for Very Small Supraclinoid Aneurysms with Stent-Assisted Coiling. Long-Term Follow-up in Six Cases. *Interv Neuroradiol*. 2009;15:37-44. doi: 10.1177/159101990901500106
- Zhang Y, Yang M, Zhang H, Zhang X, Li Y, Jiang C, Liu J, Yang X. Stent-Assisted Coiling May Prevent the Recurrence of Very Small Ruptured Intracranial Aneurysms:
 A Multicenter Study. World Neurosurg. 2017;100:22-29. doi: 10.1016/j.wneu.2016.12.107
- 22. Hunt WE, Hess RM. Surgical risk as related to time of intervention in the repair of intracranial aneurysms. *J Neurosurg*. 1968;28:14-20. doi: 10.3171/jns.1968.28.1.0014
- 23. Mascitelli JR, Moyle H, Oermann EK, Polykarpou MF, Patel AA, Doshi AH, Gologorsky

- Y, Bederson JB, Patel AB. An update to the Raymond-Roy Occlusion Classification of intracranial aneurysms treated with coil embolization. *Journal of neurointerventional surgery*. 2015;7:496-502. doi: 10.1136/neurintsurg-2014-011258
- van Swieten JC, Koudstaal PJ, Visser MC, Schouten HJ, van Gijn J. Interobserver agreement for the assessment of handicap in stroke patients. *Stroke*. 1988;19:604-607.
- 25. Jennett B, Bond M. Assessment of outcome after severe brain damage:: A practical scale. *The Lancet*. 1975;305:480-484.
- Sterne JA, Hernan MA, Reeves BC, Savovic J, Berkman ND, Viswanathan M, Henry D, Altman DG, Ansari MT, Boutron I, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*. 2016;355:i4919. doi: 10.1136/bmj.i4919
- 27. Fang C, Li MH, Zhu YQ, Tan HQ, Zhang PL, Xu HW, Wang W, Zhou B. The effectiveness and feasibility of endovascular coil embolization for very small cerebral aneurysms: mid- and long-term follow-up. *Ann Vasc Surg.* 2010;24:400-407. doi: 10.1016/j.avsg.2009.10.005
- 28. Hong B, Yang PF, Zhao R, Huang QH, Xu Y, Yang ZG, Liu J. Endovascular treatment of ruptured tiny intracranial aneurysms. *J Clin Neurosci*. 2011;18:655-660. doi: 10.1016/j.jocn.2010.09.013
- 29. Li CH, Su XH, Zhang B, Han YF, Zhang EW, Yang L, Zhang DL, Yang ST, Yan ZQ, Gao BL. The stent-assisted coil-jailing technique facilitates efficient embolization of tiny cerebral aneurysms. *Korean J Radiol*. 2014:15:850-857. doi: 10.3348/kjr.2014.15.6.850
- 30. Yu M, Liu F, Jiang S, Nie B. Stent-assisted coiling for the treatment of ruptured micro-intracranial wide-necked aneurysms. *Interv Neuroradiol*. 2015;21:40-43. doi: 10.15274/inr-2014-10097
- 31. Wu P, Ocak PE, Wang D, Ocak U, Xu S, Li Y, Zhang T, Shi H. Endovascular Treatment of Ruptured Tiny Intracranial Aneurysms with Low-Profile Visualized Intraluminal Support Device. *J Stroke Cerebrovasc Dis.* 2019:28:330-337. doi: 10.1016/j.jstrokecerebrovasdis.2018.09.052
- 32. Zhao B, Xing H, Fan L, Tan X, Zhong M, Pan Y, Wan J. Endovascular Coiling versus Surgical Clipping of Very Small Ruptured Anterior Communicating Artery Aneurysms. *World Neurosurg.* 2019;126:e1246-e1250. doi: 10.1016/j.wneu.2019.03.074
- 33. Wang JW, Li CH, Tian YY, Li XY, Liu JF, Li H, Gao BL. Safety and efficacy of endovascular treatment of ruptured tiny cerebral aneurysms compared with ruptured

- larger aneurysms. *Interv Neuroradiol*. 2020;26:283-290. doi: 10.1177/1591019919897446
- 34. Ni H, Zhao LB, Liu S, Jia ZY, Cao YZ, Shi HB. The Safety and Efficacy of Endovascular Treatment for Very Small Ruptured Anterior Communicating Artery Aneurysms: A Large Single-Center Experience With 81 Consecutive Cases. *World Neurosurg*. 2021;152:e576-e582. doi: 10.1016/j.wneu.2021.06.025
- 35. Qin F, Liu J, Zhao X, Wu D, Lai N, Zhang Z, Li Z. Endovascular Treatment of Ruptured Very Small Intracranial Aneurysms: Complications, Recurrence Rate, and Clinical Outcomes. *Front Neurol.* 2021;12:767649. doi: 10.3389/fneur.2021.767649
- 36. Zhang G, Wu Y, Wei Y, Xue G, Chen R, Lv N, Zhang X, Duan G, Yu Y, Li Q, et al. Stent-assisted coiling vs. coiling alone of ruptured tiny intracranial aneurysms: A contemporary cohort study in a high-volume center. Front Neurol. 2022;13:1076026. doi: 10.3389/fneur.2022.1076026
- 37. Jin Y, Guo X, Quan T, Chen Z, Liu C, Guan S. Safety and efficacy of endovascular treatment for tiny ruptured intracranial aneurysms with low-profile visualized intraluminal support stents. *Interv Neuroradiol*. 2023;29:141-147. doi: 10.1177/15910199221079967
- 38. Nguyen TN, Raymond J, Guilbert F, Roy D, Bérubé MD, Mahmoud M, Weill A. Association of endovascular therapy of very small ruptured aneurysms with higher rates of procedure-related rupture. *J Neurosurg*. 2008;108:1088-1092. doi: 10.3171/jns/2008/108/6/1088
- 39. Dalfino J, Nair AK, Drazin D, Gifford E, Moores N, Boulos AS. Strategies and outcomes for coiling very small aneurysms. *World Neurosurg*. 2014;81:765-772. doi: 10.1016/j.wneu.2013.01.090
- 40. Anokwute MC, Braca JA, Bohnstedt B, DeNardo A, Scott J, Cohen-Gadol A, Sahlein DH. Endovascular treatment of ruptured tiny (≤3mm) intracranial aneurysms in the setting of subarachnoid hemorrhage: A case series of 20 patients and literature review. *J Clin Neurosci*. 2017;40:52-56. doi: 10.1016/j.jocn.2017.01.011
- 41. Suzuki S, Kurata A, Ohmomo T, Sagiuchi T, Niki J, Yamada M, Oka H, Fujii K, Kan S. Endovascular surgery for very small ruptured intracranial aneurysms. Technical note. *J Neurosurg*. 2006;105:777-780. doi: 10.3171/jns.2006.105.5.777
- 42. Tsutsumi M, Aikawa H, Onizuka M, Kodama T, Nii K, Matsubara S, Iko M, Etou H, Sakamoto K, Kazekawa K. Endovascular treatment of tiny ruptured anterior communicating artery aneurysms. *Neuroradiology*. 2008;50:509-515. doi: 10.1007/s00234-008-0371-0
- 43. Pop R. Aloraini Z. Mihoc D. Burta H. Manisor M. Richter JS. Simu M. Chibbaro S.

- Proust F, Wolff V, et al. Embolization of very small (≤ 3 mm) unruptured intracranial aneurysms: A large single-center experience on treatment of unruptured versus ruptured cases. *World Neurosurg*. 2019;128:e1087-e1095. doi: 10.1016/j.wneu.2019.05.070
- 44. Mohammadian R, Asgari M, Sattarnezhad N, Mansourizadeh R, Mohammadian F, Shimia M, Talebi M, Meshkini A, Amirkolahy M. Endovascular treatment of very small and very large ruptured aneurysms of the anterior cerebral circulation: a single-center experience. *Cerebrovasc Dis.* 2013;35:235-240. doi: 10.1159/000347078
- 45. Kim JH, Choi CH, Lee JI, Lee TH, Ko JK. Endovascular treatment of ruptured tiny aneurysms. *J Cerebrovasc Endovasc Neurosurg.* 2019;21:67-76. doi: 10.7461/jcen.2019.21.2.67
- 46. Yang MS, Wong HF, Yang TH, Chen YL, Chan SW, Lee HJ, Tung YW, Tu CW, Chen TY. Alternative option in the treatment of very small ruptured intracranial aneurysms. Surg Neurol. 2009;72 Suppl 2:S41-46. doi: 10.1016/j.wneu.2009.09.003
- 47. Chung KH, Herwadkar A, Laitt R, Patel HC. Rate and clinical impact of intra-procedural complications during coil embolisation of ruptured small (3 mm or less) cerebral aneurysms. *Clin Neurol Neurosurg.* 2013;115:1356-1361. doi: 10.1016/j.clineuro.2012.12.020
- 48. Li J, Su L, Ma J, Kang P, Ma L, Ma L. Endovascular Coiling Versus Microsurgical Clipping for Patients With Ruptured Very Small Intracranial Aneurysms: Management Strategies and Clinical Outcomes of 162 Cases. *World Neurosurg.* 2017;99:763-769. doi: 10.1016/j.wneu.2015.11.079
- 49. Molyneux AJ, Kerr RS, Yu LM, Clarke M, Sneade M, Yarnold JA, Sandercock P, International Subarachnoid Aneurysm Trial Collaborative G. International subarachnoid aneurysm trial (ISAT) of neurosurgical clipping versus endovascular coiling in 2143 patients with ruptured intracranial aneurysms: a randomised comparison of effects on survival, dependency, seizures, rebleeding, subgroups, and aneurysm occlusion. Lancet. 2005;366:809-817. doi: 10.1016/S0140-6736(05)67214-5
- 50. Yamaki VN, Brinjikji W, Murad MH, Lanzino G. Endovascular Treatment of Very Small Intracranial Aneurysms: Meta-Analysis. *AJNR Am J Neuroradiol*. 2016;37:862-867. doi: 10.3174/ajnr.A4651
- 51. Pierot L, Barbe C, Spelle L, investigators A. Endovascular treatment of very small unruptured aneurysms: rate of procedural complications, clinical outcome, and anatomical results. *Stroke*. 2010;41:2855-2859. doi: 10.1161/STROKEAHA.110.588830

anatomical results. *Stroke*. 2010,41.2000-2009. doi: 10.1101/Strokeana.110.500050

53. Moret J, Cognard C, Weill A, Castaings L, Rey A. [Reconstruction technic in the treatment of wide-neck intracranial aneurysms. Long-term angiographic and clinical results. Aprenos of 56 cases] / Neuroradia/ 1997:24:30-44

clinical results. Apropos of 56 cases]. *J Neuroradiol*. 1997;24:30-44.

Table 1. Summary of studies evaluating the EVT of RVSIA

Author year	Location	Study design	EVT,	Coiling,	SAC,	Mean age,	Male, n	PC aneurysm, n	Mean Hunt-Hess	Mean size		Mean angiography follow up, months
Author year	Location	Study design	n	n (%)	n (%)	years, (SD)	(%)	(%)	grade, (SD)	Dome	Neck	(SD)
Suzuki et al. 2006^{41}	Japan	Retrospective	21	21 (100)		60.0 (14.6)	5 (23.8)	5 (23.8)		2.64 (0.35)		26
Nguyen et al. 2008 ¹⁵	USA	Retrospective	60	60 (100)				20 (33.3)				
Tsutsumi et al. 2008 ⁴²	Japan	Retrospective	19	19 (100)		65.2 (15.6)	8 (42.1)		2.42 (1.00)	2.81 (0.29)	2.47 (0.42)	28.5 (15.4)
Rooij et al. 2009^4	Netherlands	Retrospective	149	149 (100)			51 (34.2)		2.13 (1.3)			
Yang et al. 2009 ⁴⁶	Taiwan	Retrospective	12	12 (100)		55.8 (18.8)	4 (33.3)	0 (0)	2.58 (0.67)	2.60 (0.49)	2.12 (0.23)	13.4 (7.5)
Fang et al. 2010 ²⁷	China	Retrospective	19	10 (52.6)	9 (47.4)	59.1 (16.4)	5 (26.3)	3 (15.8)		2.82 (0.90)		
Hong et al. 2011 ²⁸	China	Retrospective	40	29 (72.5)	11 (27.5)	50.1 (10.8)	22 (55.0)	3 (7.50)		2.34 (0.56)	1.82 (0.58)	17.6
Chung et al. 2013 ⁴⁷	UK	Retrospective	72	72 (100)		51.6 (11.0)	24 (33.3)	10 (13.9)		2.80 (0.40)		10.8
Mohammadian et al. 2013 ⁴⁴	Iran	Prospective	21	21 (100)		34.4 (9.3)		0 (0)	2.33 (0.48)			12.5 (5.0)

Zhao et al. 2013 ¹⁹	China	Retrospective	17		17 (100)	55 (7.8)	11 (64.7)	0 (0)	2.00 (1.12)	2.26 (0.49)		12.5 (7.8)
Dalfino et al. 2014 ³⁹	USA	Retrospective	17	15 (88.2)	2 (11.8)	55.4 (15.7)	5 (25)	2 (11)	2.17 (0.52)	2.76 (0.43)		19.9 (11.4)
Li et al. 2014 ⁴⁸	China	Retrospective	16		16 (100)	45	7 (44)		2.00 (0.89)	1.59 (0.22)	1.51 (0.27)	7.7 (5.1)
Yu et al. 2015 ³⁰	China	Retrospective	35		35 (100)	53	23 (65.8)		1.40 (0.50)	2.60 (0.40)		
Anokwute et al. 2017 ⁴⁰	USA	Retrospective	20	20 (100)		57.2 (13.5)	4 (20)	7 (35)	2.15 (1.35)	2.41 (0.64)		4.6 (4.0)
Zhang et al. 2017 ²¹	China	Retrospective	93	42 (45.2)	51 (54.8)	51.9 (10.4)	33 (35.5)	6 (6.5)		2.60 (0.40)	2.20 (0.50)	13
Kim et al. 2019 ⁴⁵	Korea	Retrospective	35	35 (100)		56 (14.4)	8 (22.9)	1 (3)	2.60 (0.84)	2.53 (0.38)	1.73 (0.44)	18.6
Pop et al. 2019 ⁴³	France	Retrospective	30	30 (100)		54.2 (13.9)	8 (26.7)	4 (13)		2.51 (0.41)	1.81 (0.39)	
Wu et al. 2019 ³¹	China	Retrospective	32		32 (100)	53 (14.5)	11 (34.4)	4 (12.5)	2.19 (0.47)	2.28 (0.53)	2.40 (0.50)	9.3 (1.9)
Zhao et al. 2019 ³²	China	Retrospective	46	38 (82.6)	8 (17.4)	54.5 (11.2)	22 (47.8)	0 (0)		2.80 (0.40)	2.20 (0.60)	
Wang et al. 2020 ³³	China	Retrospective	33	17 (51.5)	16 (48.5)			5 (15.2)				

Ni et al.	China	D atus au anticas	01	52	29	56.5 (9.6)	20 (27)	0	2.29 (0.00)	2 (0 (0 40)	1.80		
2021 ³⁴	China	Retrospective	81	(64.2)	(35.8)	56.5 (8.6)	30 (37)	0	2.38 (0.90)	2.60 (0.40)	(0.80)		
Qin et al.	China	Retrospective	152	90	62	56.2 (11.2)	54 (35.5)	7 (4.6)		2.50 (0.40)	1.90		
2021 ³⁵	Cillia	Retrospective	132	(59.2)	(40.8)	30.2 (11.2)	34 (33.3)	7 (4.0)		2.30 (0.40)	(0.50)		
Zhang et al.	China	Retrospective	245	152	93	55.6 (11.7)	92 (37.6)	0 (0)	2.39 (0.82)	2.40 (0.48)	1.80	14.1 (12.5)	
2022^{36}	Cillia	Retrospective	243	(62.0)	(38.0)	55.6 (11.7)	92 (37.6)	0 (0)	2.39 (0.02)	2.40 (0.46)	(0.60)	14.1 (12.3)	
Jin et al.	China	Retrospective	90	62	28	56.1 (10.3)	38 (42)	12 (13)	1.71 (0.77)	2.53 (0.39)	1.99	7.3 (1.9)	
2023^{37}	Cnina	Renospective	70	(68.9)	(31.1)	30.1 (10.3)	38 (42)	12 (13)	1.71 (0.77)	2.33 (0.39)	(0.59)	7.3 (1.9)	

EVT indicates endovascular treatment; PC, posterior circulation; RVSIA, ruptured very small intracranial aneurysm; SAC, stent-assisted coiling.


Table 2. Comparison of the results of previous and present study about EVT of RVSIAs

0.4	Previou	s (n=844) ⁵⁰	Present (n= 1355)			
Outcome	%	95% CI	%	95% CI		
Angiographic and clinical outcomes						
Initial complete occlusion	88	79-93	64	52-74		
Follow-up complete occlusion	86	80-90	85	74-92		
Recanalization	9	5-17	6	3-10		
Retreatment	7	3-14	3	2-4		
Favorable outcome (mRS0-2)	74	59-85	91	89-93		
Mortality	4	2-7	3	2-4		
Adverse events						
Thromboembolism	4	2-9	4	3-6		
Intraprocedural rupture	9	6-12	4	2-7		

CI indicates confidence interval: EVT, endovascular treatment; mRS, modified Rankin Scale; RVSIA, ruptured very small intracranial aneurysm; SAC, stent-assisted coiling.

Figure legends:

- Figure 1. Flow diagram of the search strategy and study selection
- Figure 2. The pooled prevalence of radiological and clinical outcomes
- Figure 3. The pooled prevalence of adverse events
- Figure 4. Forest plot for outcomes between coiling and SAC



Initial complete occlusion

Study Events Total 95%-CI Proportion Fang 2010 0.26 [0.09; 0.51] 0.27 [0.16; 0.41] Lu 2011 0.35 [0.16; 0.57] Wang 2020 0.37 [0.21; 0.55]yu 2015 Wu 2019 [0.24; 0.59]Yang 2009 [0.42 [0.15; 0.72]0.56 [0.30; 0.80]Ui 2014 Zhang 2022 0.59 [0.52; 0.65]144 Dalfino 2014 [0.33; 0.82]Zhao 2013 0.59 [0.33; 0.82] 0.60 [0.36; 0.81] Anokwute 2017 Jin 2022 0.61 [0.48; 0.73] 62 Kim 2019 [0.54; 0.85]Hong 2011 0.72 [0.56; 0.85]40 0.73 [0.62; 0.82]Ni 2021 81 Qin 2021 0.76 [0.69; 0.83]152 116 Tsutsumi 2018 0.84 [0.60; 0.97] 0.95 [0.76; 1.00] Mohamnadian 2013 Suzuki 2006 1.00 [0.84; 1.00] 0.62 [0.50; 0.73] Random effects model Heterogeneity: $I^2 = 78\%$, $\tau^2 = 0.8567$, p < 0.01

Follow-up complete occlusion

Study	Events	Total				Proportion	95%-CI
Chung 2013	13	35				0.37	[0.21; 0.55]
Zhang 2022	123	245				0.50	[0.44; 0.57]
Qin 2021	73	105				0.70	[0.60; 0.78]
Anokwute 2017	13	18				0.72	[0.47:0.90]
Dalfino 2014	13	18				0.72	[0.47; 0.90]
Jin 2022	66	90					[0.63: 0.82]
Wu 2019	23	28			-		[0.63: 0.94]
Tsutsumi 2018	14	16				0.88	[0.62: 0.98]
Mohammadian 2013	22	25					[0.69; 0.97]
Kim 2019	20	22					[0.71:0.99]
Ni 2021	55	60				0.92	[0.82: 0.97]
LI 2014	1.5	16			_	0.94	[0.70; 1.00]
Hong 2011	31	33					[0.80:0.99]
Suzuki 2006	20	21					[0.76: 1.00]
yu 2015	34	35					[0.85: 1.00]
Fang 2010	20	20				1.00	[0.83; 1.00]
Random effects model		787				0.85	[0.74; 0.92]
Heterogeneity: $I^2 = 84\%$, τ	= 1.212	θ , $\rho < 0.0$					
			0.4	0.6	0.8		

Recanalization


0.6

Retreatment

Study	Events	Total	Proportion	95%-CI	Study	Events	Total	Proportion	95%-CI
NI: 2024				nn no n	lin 2022		90		[0.00; 0.04]
Ni 2021		81		[0.00; 0.07]	Jin 2022				
Chung 2013	1	72	0.01	[0,00;00]	Chung 2013	0	72	0.00	[0.00; 0.05]
Anokwute 2017	0	20	0.00	[0.00; 0.17]	Zhao 2019	O	46	0.00	[0.00; 0.08]
Mohammadian 2013	1	21	0.05	[0.00; 0.24]	Ni 2021	1	81	0.01	[0.00; 0.07]
Zhang 2017	5	93	0.05	[0.02; 0.12]	Yu 2015	O	35	0.00	[0.00; 0.10]
Wang 2020	2	31	0.06	[0.01; 0.21]	Anokwute 2017	O	20	0.00	[0.00; 0.17]
Zhang 2022	17	245	0.07	[0.04; 0.11]	Tautaumi 2018	0	16	0.00	[0.00; 0.21]
Yang 2009	1	12	0.08	[0.00; 0.38]	Zhang 2022	10	245	0.04	[0.02; 0.07]
Jin 2022	8	90	0.09	[0.04; 0.17]	Suzuki 2006	1	21	0.05	[0.00; 0.24]
Kim 2019	2	22	0.09	[0.01; 0.29]	Qin 2021	7	105	0.07	[0,03; 0.13]
Tsutsumi 2018	2	16	0.12	[0.02; 0.38]	Dalfino 2014	2	19	0.11	[0.01; 0.33]


Overall complications

Initial failure



Vasospasm

Coil herniation



Thromboembolisms

Intraprocedural rupture

Study Proportion 05% CI

Initial complete occlusion

Follow-up complete occlusion

		oilling		SAC							oiling		SAC				
Study		lotai	Events	lotai	Odds Ratio	OR		Weight	Study	Evente	Total		Iotal	Odda Ratio	OR	95%-CI	maign
Zhang 2022	85	152	50	93		0.73	0.43; 1.24	60.8%	Yu 2015	0	35	34	35 -		0.00	[0.02]	12.9%
Não 2021	39	52	20	29		1.35	0.49; 3.69	12.1%	Li 2014	0	18	15	16		0.00	[0.00; 0.08]	12.9%
Jim 2022	38	82	15	28					2022 2022	57	152	66	93		0.25	[0.14; 0.43]	16.3%
Dattino 2014	9	15	1	2			0.08: 28.89			43	62	23	28		0.49	[0.16: 1.49]	15.9%
Famo 2010	2	10	1	8		2.00	0.15; 26.73	1.6%	NI 2021	33	36	22	24		1.00	[0.15: 6.48]	15.1%
Hong 2011	22	29	6	11	-		0.61; 11.28		Oalfino 2014	11	16	2	3		1.10	[0.08: 15.15]	13.9%
Zhang 2017	33	42	42	51	+		0.70; 11.04		Hong 2011	16	16		7		15.00	[0.62; 362.88]	13.0%
Common effect model Heterogeneity: I ² = 0%, τ ²		382		223	0.1 0.5 1 2 10	1.11	0.77; 1.61	100.0%	Random effects mod Heterogeneity: J ² = 815		333	01	206	0.001 0.1 1 10 1000	0.19	[0.01; 4.36]	100.0%

Recanalization

Favorable outcome

Study	Events	Total	Events	SAAC	Odds Ratio	OR	85%-Cl Weight	Study	Events	oiling Total E	venta	BAC	Odds Ratio	OR	95%-CI Y	Meight
Zhang 2022 Zhang 2017 Jin 2022	13	152	4	33 51 28		41	[0.66; 6.58] 84.9% [0.58; 50.38] 15.1% [0.50; 159.63] 0.0%	Ni 2021 Zhang 2022		137	27	90			[0.11; 3.01] [0.50; 2.85]	
Common effect model Heterogeneity: /2 = 0%, τ2				172	1 0.1 1 10 100		[1.31; 9.45] 100.0%	Common effect mode Heterogeneity: /2 = 0%, x		189		119	0.2 0.5 1 2 5	1.00	[0.47; 2.15] 1	00.0%

Coil herniation

Intraprocedural rupture

	Cc	illing		SAC									
Study	Events	Total	Events	Total	Odds Ratio	OR 25%-C	1 Weight		C	oilling	8AC		
								Study	Events	Total Ev	rents Total	Odds Ratio	OR 95%-CI Weight
medRxiv preprint doi: https://doi.org/10.1101/2023.05.16.23290074; this version posted May 17, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. All rights reserved. No reuse allowed without permission.	2	9	3	9		0.57 [0.07; 4.64	100.0%						
Zhang 2017		41	0	51		3.81 (0.15; 96.15			2	152	5 93		0.23 [0.04; 1.24] 100.0%
								Jim 2022	2	62	0 28		2.36 [0.11; 50.68] 0.0%
Common effect mode	1	50		80		1.05 [0.17; 6.30	100.0%						
Heterogeneity: /* = 0%, x	- o, o - o	33						Common effect model		214	121		0.44 [0.11; 1.70] 100.0%
					0.1 0.51 2 10			Heterogeneity: $I^2 = 40\%$, s	c = 1.074	7. p = 0.20			
												0.1 0.51 2 10	