

## **Comparing the Demographic and Health Survey's timing-based measure of unintended pregnancy to the London Measure of Unplanned Pregnancy in Bangladesh**

Md Nuruzzaman Khan<sup>1,2</sup>, Ph.D, Shimlin Jahan Khanam<sup>1</sup>, MSS, Melissa L. Harris<sup>2</sup>, Ph.D.

<sup>1</sup>Department of Population Science, Jatiya Kabi Nazrul Islam University, Mymensingh, Bangladesh.

<sup>2</sup> Centre for Women's Health Research, University of Newcastle, Australia.

**Short running title:** Discordant response in unintended pregnancy

### **Corresponding author**

Md Nuruzzaman Khan, Ph.D.

Assistant Professor

Department of Population Science

Jatiya Kabi Kazi Nazrul Islam University, Trishal, Mymensingh, Bangladesh.

Email: [sumonrupop@gmail.com](mailto:sumonrupop@gmail.com)

[mdnuruzzaman.khan@uon.edu.au](mailto:mdnuruzzaman.khan@uon.edu.au)

### **Word Count:**

Abstract: 236

Manuscript: 3010

Table: 03

Figure: 01

Reference: 30

Supplementary file: 01

## **Abstract**

**Background:** Demographic and Health Survey's timing-based measure is commonly used in Low and Middle Income Countries to estimate unintended pregnancy despite its limitations, including ambivalent responses and failure to consider the partner's intention, while LMUP, which can address these limitations, is not widely used in LMICs and is yet to be administered in Bangladesh. This study compared unintended pregnancy rates measured by the DHS timing-based measure and LMUP, as well as explored the extent of discordance between the measures and their determinants.

**Methods:** A cross-sectional survey was conducted in four districts of Bangladesh using two-stage stratified random sampling. Data was collected from 1,200 postnatal women. The study focused on discordance in reporting pregnancy intention between DHS timing-based measure and LMUP. Multivariate logistic regression models were used to identify predictors of discordant responses in reporting pregnancy intention.

**Results:** The prevalence of unintended pregnancy was found to be 24.3% through the DHS timing-based measure and 31.0% through the LMUP. Discordance in responses to pregnancy intention in the two measures was around 28%. Key predictors of discordance included older age, female last child, more than two children, poorer wealth quintile, and rural residence.

**Conclusions:** Prevalence of unintended pregnancy in Bangladesh and other LMICs, measured by DHS timing-based measure, may grossly underestimated. This suggests that the negative effects of unintended pregnancy are even more significant than currently believed, further highlighting the need to strengthen the family planning program in Bangladesh.

**Keywords:** Demographic and Health Survey timing-based measure of unintended pregnancy, London Measure of Unintended Pregnancy, discordance in pregnancy intention measures, Bangladesh, contraception.

## **Background**

Globally, almost half (47%) of all pregnancies, equivalent to approximately 121 million, are unintended, with low- and middle-income countries (LMICs) experiencing the majority of these cases due to lower usage of long-acting modern contraception and higher unmet need for contraception (1, 2). In particular, unintended pregnancy remains a key concern in Bangladesh, with over 47% of pregnancies at conception and around 23% of live births reported as unintended (3). Unintended pregnancies have significant negative consequences for maternal and child health, especially in LMICs, such as Bangladesh, where safe abortion services are restricted by law (1, 4). Approximately two-thirds of total unintended pregnancies in LMICs and Bangladesh end by induced abortions, contributing to around 13% of the total maternal mortality and costing billions of dollars in the treatment of abortion-related complications (1, 5). Unintended pregnancies resulting in live births can lead to delayed or inaccessible intra-partum and post-partum care, increasing the risk of several maternal and child health outcomes, such as stillbirths, preterm birth, low birth weight, child undernutrition, neonatal, and maternal mortality (3-7). Studies conducted in LMICs and Bangladesh have also found intergenerational effects of unintended pregnancy, such as higher rates of school dropout (8-10). Improving access to modern contraception and comprehensive sexuality education is crucial to reducing unintended pregnancies in Bangladesh and other LMICs (3).

The accuracy of current estimates of unintended pregnancy is a global concern, especially in LMICs where prospective data on reproductive events (e.g., conception, birth) are scarce, and data of unintended pregnancy usually come from the nationally representative Demographic and Health Survey (5). The survey collects timing-based data on pregnancy intention using two-time dependent questions which have several limitations. Specific limitations to this timing-based data include recall bias, which arises from difficulties in accurately remembering pregnancy intentions at the time of conception. Additionally, it can lead to a higher chance of ambivalent responses,

particularly when the child's sex is opposite to parental desire or when pregnancy intentions change after having a child (5). The major reasons for such limitations in data include: (i) they only apply to women who have given birth (i.e., excluding women who have had induced abortions), (ii) they do not include partners' intentions, despite often being the dominant group in deciding pregnancy timing and number, and (iii) they do not consider contraceptive use/non-use status at the time of conception (11, 12). These factors can further result in ambivalence, denial, and confusion about pregnancy and misreporting of partner's intentions, as a significant portion of reported unintended pregnancies in LMICs is wanted from the husband's perspective (5, 11, 12).

To comprehensively measure pregnancy intention, it is necessary to consider multiple dimensions, such as contraception use, feelings about pregnancy at the time of conception and after giving birth, and agreement with the partner (12). The London Measure of Unplanned Pregnancy (LMUP) can cover these issues and can overcome the limitations of the DHS timing-based measure(13). However, the use of LMUP in LMICs is still limited and has not been included in national-level survey (11-14). Furthermore, there has been no previous investigation into the discordance between the DHS timing-based measure and LMUP or the factors contributing to this discordance. The aim of this study is to estimate the prevalence of unintended pregnancy in Bangladesh using two measures: DHS timing-based estimate and LMUP. The study also seeks to determine the degree of discordance between the prevalence estimates obtained from these two measures and identify the factors that contribute to such discordance.

## **Methods**

### **Study settings and data**

This study employed a cross-sectional survey design conducted in four randomly selected districts (Mymensingh, Pabna, Kurigram, Satkhira) using a two-stage stratified random sampling method. In the first stage, a total of 20 hospitals were selected. These comprised district level hospitals, one upazila health complex, one union health complex, and two private hospitals. Data were collected from 1200 women who met the inclusion criteria: having recently delivered (either by caesarean section or vaginal delivery) and were visiting the selected hospitals for post-natal healthcare services. The sample size required for this study was determined to be 378 using a sample determination formula (15). Data were collected through a face-to-face interview by using a pre-developed and pre-tested structured questionnaire. The questionnaire contained questions pertaining to both the DHS timing-based measure of unintended pregnancy and LMUP (16). In addition, other relevant questions from the Bangladesh Demographic and Health Survey, which is a part of the worldwide recognized Demographic and Health Survey program of the USA, were included. The questionnaire was initially developed and field-tested, with subsequent refinements made based on feedback received. The final version of the questionnaire and the data collection methodology were approved by the Institutional Ethical Board of the Institution of Biological Science at the University of Rajshahi, Bangladesh.

### **Outcome measure**

Discordant responses in pregnancy intention (yes, no) in DHS timing-based measure and LMUP was our outcome of interest. The variable was generated by comparing pregnancy intention estimates produced through the DHS timing-based measure and the LMUP. In the DHS-timing based measure, eligible women (those who had at least one live birth within five years of the survey) were asked two questions. The first question inquired whether they intended to become pregnant at the time of conception, with response options of "yes" or "no." If the answer was "no," a follow-up second question was asked to determine whether the unwantedness was for a short period (response option "later") or permanent (response option "not at all"). We categorized these responses into intended pregnancy (positive response to the first question) and

unintended pregnancy (negative response to the first question and "later" or "not at all" in the second question) (see Supplementary Table 1). On the other hand, total of six questions were asked to record pregnancy intention through LMUP. Based on the response to six questions and corresponding scores, the LMUP produces estimates for unplanned pregnancy (score: 0-3), ambivalence (score: 4-9) and planned pregnancy (score: 10-12) (Supplementary table 2). We considered the responses to be concordant in pregnancy intention if women consistently reported a similar type of pregnancy intention in both measures, such as intended pregnancy and planned pregnancy or unintended pregnancy or unplanned pregnancy. Any mismatch between these two measures, such as intended pregnancy and ambivalence or intended pregnancy and planned pregnancy, were considered discordant pregnancy intention responses.

### **Explanatory variables**

Explanatory variables were selected in two stages. Firstly, a list of potential explanatory variables were compiled by reviewing the relevant literature in LMICs, particularly Bangladesh. As this is the first study of its kind in LMICs, we considered similar research, such as studies investigating determinants of unintended pregnancy measured through the DHS timing-based measure or the LMUP (3, 5, 12-14, 17, 18). Next, we checked the availability of the selected variables in the survey and created a list of available variables. The selected variables included women's age (categorized as  $\leq 19$  years, 20-34 years, and  $\geq 35$  years), women's education (categorized as no formal education, primary education, secondary education, and higher education), women's formal work engagement (yes or no), number of children ever born (categorized as 1-2 children and  $>2$  children), partner's education (categorized as no formal education, primary education, secondary education, and higher education), partner's occupation (categorized as agriculture worker, physical worker, service, business, and others), sex of the last child (male or female), household wealth quintile (categorized as poorer, poorest, middle, richer, and richest), place of residence (urban or rural), and district of residence (Pabna, Mymensingh, Satkhira, and Kurigram).

## Statistical analysis

Descriptive statistics were used to depict the baseline characteristics of the study participants. To present the discordant responses in pregnancy intention between the DHS timing-based measure and the LMUP, we utilized cross-tabulation. Multivariate logistic regression model was used to explore the factors associated with pregnancy intention discordance. We estimated both crude and adjusted associations while ensuring that multicollinearity was addressed prior to running the model. The results were reported as odds ratios (ORs) along with their corresponding 95% confidence intervals (CIs). The statistical software package Stata (version 15.2) was used for all statistical analyses.

## Results

Data were collected from a total of 1200 respondents, and the background characteristics of the participants are summarized in Table 1. The majority of respondents were aged between 20-34 years (80.2%), had secondary level education (40.0%), and were not formally employed (69.0%). The majority of partners had a higher education level (36.17%) and reported business as their occupation (28.3%). The study population had an almost equal proportion of male and female children (49.0% and 51.0%, respectively). The majority of participants lived in rural areas (78.7%).

**Table 1: Background characteristics of the respondents, N=1200**

Characteristics	N (%)
<b>Mother's age at birth of the last child</b>	
≤19 years	123 (10.3)
20-34 years	962 (80.2)
≥35 years	115 (9.58)
<b>Mothers' education</b>	
No formal education	115 (9.6)
Primary	233 (19.4)
Secondary	480 (40.0)
Higher	372 (31.0)
<b>Mother's formal employment status</b>	
Yes	372 (31.0)
No	828 (69.0)
<b>Partner's education status</b>	
No formal education	180 (15.0)


Primary	211 (17.6)
Secondary	375 (31.25)
Higher	434 (36.17)
<b>Partner's occupation</b>	
Agriculture worker	183 (15.3)
Physical worker	262 (21.8)
Services	335 (27.9)
Business	340 (28.3)
Other	80 (6.7)
<b>Sex of the last child</b>	
Male	588 (49.0)
Female	612 (51.0)
<b>Number of children ever born</b>	
1-2 children	892 (74.3)
>2 children	308 (26.7)
<b>Household wealth status</b>	
Poorest	238 (19.8)
Poorer	240 (20.0)
Middle	243 (20.3)
Richer	237 (19.8)
Richest	242 (20.2)
<b>Place of residence</b>	
Urban	256 (21.3)
Rural	944 (78.7)
District	
Pabna	306 (25.5)
Mymensingh	296 (24.7)
Satkhira	310 (25.8)
Kurigram	288 (24.0)

The prevalence of unintended pregnancy, measured using the DHS timing-based measure and the LMUP, are presented in Supplementary file 1 and 2, respectively. A higher prevalence of unplanned pregnancy was found using the LMUP (31.0%) compared to the DHS timing-based measure (24.3%). Nearly 47.5% of women intended to become pregnant at the time of conception, and almost 53% reported that their pregnancy occurred at the right time. Additionally, approximately 29% of women reported that they had never discussed pregnancy with their partner, and around 43% reported that they did not take any health actions (such as, took folic acid, sought medical/health advice) before becoming pregnant (Figure 1).

A total of 27.8% of the responses to pregnancy intention were discordant between the DHS timing-based measure and the LMUP (see Table 2).







**Figure 1:** Women’s response to different questions on London Measure of Unintended Pregnancy

**Table 2: Comparison of pregnancy intention measured by Demographic and Health Survey’s timing-based measure and London Measure of Unplanned Pregnancy**

LMUP classifications*	DHS timing-based measure		Total
	Unintended	Intended	
Unplanned	194 (CR)	179 (DR)	373 (31.0)
Ambivalent	30 (DR)	58 (DR)	78 (6.5)
Planned	68 (DR)	681 (CR)	749 (62.4)
Total	292 (24.3)	908 (75.7)	
<b>Comparison in DHS timing-based measure and LUMP</b>			
Concordant responses		865 (72.2)	
Discordant responses		335 (27.8)	

**Note:** \*Classified based on scores presented in Supplementary table 2. Score 0-3: unplanned pregnancy; score 4-9: ambivalent; score 10-12: planned pregnancy. CR: Concordant response in both measures, DR: Discordant responses between two measures

Table 3 presents the factors associated with discordant responses in pregnancy intention, as determined by both crude and adjusted ORs. Increasing age of the respondent, female sex of the last child, having more than two children, being in the poorest wealth quintile, and residing in Kurigram district were associated with higher likelihood of discordant responses in the crude estimate. These findings remained consistent after adjusting for potential confounding factors in the multivariate model.

Mothers aged 20-34 and  $\geq 35$  years had 32% (95% CI, 1.06-1.53) and 38% (95% CI, 1.06-1.88) increased odds of having a discordant responses in pregnancy intention, respectively, compared to those aged  $\leq 19$  years. On the other hand, respondents with secondary (OR, 0.96, 95% CI, 0.80 - 0.95) and higher (OR, 0.84, 95% CI, 0.63 - 0.95) education had decreased odds of having a discordant responses compared to those with no formal education or primary education. A 15-

fold increase (ORs, 1.15, 95% CI, 1.01 - 1.38) in discordant responses was found among not formally employed mothers compared to employed mothers. Furthermore, mothers whose partners were higher educated (aOR, 0.89, 95% CI, 0.79 - 0.98) had lower odds of reporting discordant responses in pregnancy intention than mothers whose partners were not formally educated. Those having more than two children had a 16% (aOR, 1.16, 95% CI, 1.07-1.40) increased odds of having a discordance responses compared to those with 1-2 children. The poorest (aOR, 1.32, 95% CI, 1.06 – 1.82) and poorer (aOR, 1.34, 95% CI, 1.06 – 1.76) mothers were more likely to report discordance responses in pregnancy intention than middle wealth quintile mothers. Rural mothers were 30% (aOR, 1.30, 95%CI, 1.03 – 1.51) more likely to report discordance responses in pregnancy intention than urban mothers. Lastly, mothers in Kurigram district had an 8% (aOR, 1.08, 95% CI, 1.02-1.17) increased odds of discordant responses in pregnancy intention compared to those in Pabna district.

**Table 3:** Factors Associated with Discordance Responses in Pregnancy Intention Measured through Demography and Health Survey’s timing-based measure and London Measure of Unplanned Pregnancy Using Multivariate Logistic Regression Model.

	<b>Discordance response, Crude odds ratio (95% CI)</b>	<b>Discordance response, Adjusted odds ratio (95% CI)</b>
<b>Mother’s age at birth of the last child</b>		
≤19 years	1.00	1.00
20-34 years	1.28** (1.10 -1.49)	1.32** (1.06 -1.53)
≥35 years	1.45* (1.08 -1.92)	1.38* (1.06 -1.88)
<b>Mothers’ education</b>		
No formal education	1.00	1.00
Primary	0.78 (0.63-1.26)	0.82 (0.67 - 1.42)
Secondary	0.94* (0.82 - 0.99)	0.96* (0.80 - 0.95)
Higher	0.82** (0.68-0.96)	0.84* (0.63 - 0.95)
<b>Mother’s formal employment status</b>		
Yes	1.00	1.00
No	1.14 (0.96-1.32)	1.15** (1.01 - 1.38)
<b>Partner’s education status</b>		
No formal education	1.00	1.00
Primary	0.93 (0.83 - 1.03)	0.96 (0.85 - 1.07)
Secondary	0.91 (0.81 - 1.04)	0.95 (0.85 - 1.11)
Higher	0.86 (0.78 - 0.94)	0.89** (0.79 - 0.98)
<b>Partner’s occupation</b>		

Physical worker	1.00	1.00
Agriculture worker	1.23 (0.93-1.53)	1.26 (0.95-1.56)
Services	1.13 (0.80 -1.46)	1.18 (0.85 -1.48)
Business	1.15 (0.84 – 1.46)	1.18 (0.82 – 1.48)
Other	1.06 (0.92 – 1.20)	1.11 (0.88 – 1.26)
<b>Sex of the last child</b>		
Male	1.00	1.00
Female	1.38** (1.08 – 1.72)	1.34** (1.06 – 1.67)
<b>Number of children ever born</b>		
1-2 children	1.00	1.00
>2 children	1.24** (1.09 – 1.39)	1.16* (1.07 – 1.40)
<b>Household wealth status</b>		
Poorest	1.39** (1.05 – 1.78)	1.32** (1.06 – 1.82)
Poorer	1.31** (1.02 – 1.64)	1.34** (1.06 – 1.76)
Middle	1.00	1.00
Richer	0.95 (0.86 - 1.04)	0.98 (0.87 - 1.08)
Richest	0.91* (0.87 – 0.95)	0.94 (0.87 – 1.01)
<b>Place of residence</b>		
Urban	1.00	1.00
Rural	1.26 (1.04 – 1.48)	1.30** (1.03 – 1.51)
<b>Districts</b>		
Pabna	1.00	1.00
Mymensingh	0.98 (0.92 -1.06)	0.99 (0.93 -1.05)
Satkhira	1.00 (0.99-1.03)	1.03 (0.98-1.05)
Kurigram	1.06* (1.01 - 1.10)	1.08** (1.02 - 1.17)

## Discussion

This study investigated unintended pregnancy prevalence using two measures - DHS timing-based and LMUP - and explored the degree of discordance between these measures, as well as the determinants of discordant responses. The study reveals that the prevalence of unplanned pregnancy reported through the LMUP (31.0%) was higher than through the DHS timing-based measure (24.3%). Additionally, approximately 28% of women provided discordant responses to pregnancy intention between the two measures. Significant predictors of discordant responses were increasing age, female sex of the last child, women having more than two children, being in the poorest wealth quintile, and residing in Kurigram district. These findings suggest that the actual occurrence of unintended pregnancy in Bangladesh and other LMICs may be higher than currently estimated through the DHS timing-based measure. Therefore, the burden on maternal

and child health in LMICs related to unintended pregnancy may be even more significant than what is currently believed.

The reported prevalence of unintended pregnancy measured through the DHS-timing based measure in this study is consistent with reported national estimates in Bangladesh, where the same approach is adopted, as with other LMICs (1, 3, 17, 18). However, the use of a more comprehensive unintended pregnancy measure, the LMUP, revealed a much higher prevalence of unplanned pregnancy, with almost 38% of all pregnancies either unplanned (31%) or ambivalent (7%). In particular, individual dimensions of the LMUP revealed a more complex picture of unintended pregnancy, with only 54% of total pregnancies occurring with mutual agreement between the husband and wife. Furthermore, 43% of women reported not taking any prior health actions before getting pregnant, indicating a lack of adequate counselling on family planning and pre-pregnancy health check-ups. We could not validate these findings because of a lack of relevant literature in Bangladesh and LMICs. However, these findings highlight the urgent need for more comprehensive family planning services that address social norms and misconceptions around pregnancy, as well as provide counselling on family planning and preconception care (11, 12). Failure to address these issues can lead to poor maternal and child health outcomes in Bangladesh and other LMICs.

Our findings highlight that mixed feelings or uncertainty about pregnancy intention were common among women in Bangladesh, with one in four women reporting discordance. Young women and those with two or more children were more likely to report discordance in pregnancy intention, possibly due to their desire to delay or limit childbirth, respectively (3, 19-21). Additionally, women whose last child was female were more likely to report discordance, reflecting the persistent preference for male children in Bangladesh and other LMICs (22). This preference is especially prevalent among women from lower wealth quintiles and rural areas, who rely heavily on agriculture and view male children as primary workers in agriculture and primary

caregivers in old age. This is in addition to perceptions surrounding the continuation of the family lineage and name(22-26). However, we also found that education played an important role in reducing discordance in pregnancy intention, as educated parents were less likely to prefer a child of a particular sex and were more likely to have access to family planning and contraception. As such, they appeared to have greater control over pregnancy planning (26-28).

The higher prevalence of discordance towards pregnancy intention suggests that a significant proportion of women have conflicting feelings about their pregnancy. This situation can have detrimental effects on various aspects. For instance, women who have contradictory feelings about their pregnancy are more likely to delay or have reduced access to maternal healthcare services, as evidenced by previous studies (3, 6, 7, 29). Additionally, women with unintended pregnancies or pregnancy ambivalence are more prone to engaging in negative behaviours during pregnancy, such as smoking, and have a higher prevalence of depression and anxiety(30). These concerns often stem from worries about their career or education(4). Collectively, these factors contribute to an increased risk of adverse maternal and child health outcomes, including high rates of maternal and child mortality already observed in Bangladesh and other LMICs (5, 8). Therefore, there is an urgent need for comprehensive family planning services and contraception, with a focus on comparatively young, poor, and less educated women. This direction should be incorporated into existing family planning services along with strengthening of current services.

This study has several key strengths including a large sample size. We exceeded the required sample size and increased the statistical power of our findings. Additionally, using two measures of unintended pregnancy, the DHS timing-based measure and the LMUP, provides a comprehensive understanding of the phenomenon. The study's rigorous sampling method and

inclusion criteria also enhanced internal validity and generalizability to the study population. Despite this, some limitations exist. The cross-sectional study design precludes establishing causal relationships in the reported associations in this study. The study's inclusion of only women who had recently delivered and were receiving post-natal healthcare services at selected hospitals limited generalizability to the broader population of women in Bangladesh, particularly women who do not access delivery healthcare services and post-natal care. Reliance on self-reported interview data may also introduce reporting bias, as some women may have provided socially desirable responses. Lastly, the study did not collect data on other potential factors, such as social norms about child sex preference, that could influence discrepancies between the two measures of unintended pregnancy. Despite these limitations, this study is the first in LMICs to compare the commonly used DHS timing-based measure of unintended pregnancy with the more advanced LMUP, revealing a higher discordance rate. This study findings underscores the potential underestimation of the true prevalence of unintended pregnancy in Bangladesh when measured using the DHS-timing based measure. Consequently, the reported adverse outcomes associated with unintended pregnancy may be even higher than currently reported. This underscores the urgent need for effective family planning interventions and the promotion of modern contraception methods to ensure that all pregnancies are desired and planned. Considering that other low- and middle-income countries (LMICs) also employ similar measurement approaches for unintended pregnancy as Bangladesh, the findings of this study also emphasize the necessity for further research to accurately assess the extent of unintended pregnancy in LMICs and implement appropriate interventions accordingly.

## **Conclusion**

The prevalence of unplanned pregnancy estimated through the LMUP was 31%, which was almost 7% higher than the prevalence estimated through the DHS timing-based measure of 24.3%. About 28% of women provided discordant responses in pregnancy intention, with a



significantly higher likelihood among women who were comparatively older in age, those whose last child was female, those who already had more than two children, and those in the poorest wealth quintile. These findings suggest that the actual extent of unintended pregnancy in Bangladesh may be higher than the currently available estimates derived from the DHS timing-based measure. As a result, the negative effects of unintended pregnancy may be more significant than what is currently perceived. Comprehensive programs that ensure family planning and contraception are necessary to empower couples to plan their pregnancies and to ensure every pregnancy is intended and conceived at the right time. Further, a comprehensive national-level study employing advanced techniques, such as LMUP, should be conducted to accurately determine the prevalence of unintended pregnancy. The findings of such a study can provide valuable insights for the development of comprehensive programs aimed at reducing the occurrence of unintended pregnancy and its related adverse outcomes.

**Conflict of Interest:** None

**Ethics approval:** Ethical approval for this study was obtained from Human Research Ethics Committee of the Jatiya Kabi Kazi Nazrul Islam University with the approval number JKKNIU2023-07. The research protocol, including the study design, data collection procedures, and informed consent process, was thoroughly reviewed and deemed ethically sound and compliant with all applicable regulations and guidelines.

**Acknowledgment:** We acknowledge the support Department of Population Science at Jatiya Kabi Kazi Nazrul Islam University, Bangladesh where this study was conducted.

**Funding:** This research did not receive any specific funds.

**Authors' contribution:** MNK designed this study, conducted data analysis. SJK write the first draft of this manuscript along with MNK. MLH critically reviewed this manuscript. All authors approve the final version of this manuscript.



## References

1. Bearak J, Popinchalk A, Alkema L, Sedgh G. Global, regional, and subregional trends in unintended pregnancy and its outcomes from 1990 to 2014: estimates from a Bayesian hierarchical model. *The Lancet Global Health*. 2018;6(4):e380-e9.
2. Bearak J, Popinchalk A, Ganatra B, Moller AB, Tunçalp Ö, Beavin C, et al. Unintended pregnancy and abortion by income, region, and the legal status of abortion: estimates from a comprehensive model for 1990-2019. *Lancet Glob Health*. 2020;8(9):e1152-e61. Epub 20200722.
3. Khan MN, Harris M, Loxton D. Modern contraceptive use following an unplanned birth in bangladesh: an analysis of national survey data. *International perspectives on sexual and reproductive health*. 2020;46:77-87.
4. Khan MN, Harris ML, Shifti DM, Laar AS, Loxton D. Effects of unintended pregnancy on maternal healthcare services utilization in low-and lower-middle-income countries: systematic review and meta-analysis. *International journal of public health*. 2019;64(5):743-54.
5. Khan M. Effects of unintended pregnancy on maternal healthcare services use in Bangladesh. Faculty of Health and Medicine, School of Medicine and Public Health, The. 2020;1.
6. Khan MN, Harris ML, Loxton D. Does unintended pregnancy have an impact on skilled delivery care use in Bangladesh? A nationally representative cross-sectional study using Demography and Health Survey data. *Journal of Biosocial Science*. 2020:1-17.
7. Khan MN, Harris ML, Loxton D. Low utilisation of postnatal care among women with unwanted pregnancy: A challenge for Bangladesh to achieve Sustainable Development Goal targets to reduce maternal and newborn deaths. *Health & Social Care in the Community*. 2020.
8. Gipson JD, Koenig MA, Hindin MJ. The effects of unintended pregnancy on infant, child, and parental health: a review of the literature. *Studies in family planning*. 2008;39(1):18-38.
9. Rahman MM. Is Unwanted Birth Associated with Child Malnutrition in Bangladesh? *Int Perspect Sex Reprod Health*. 2015;41(2):80-8.

10. Alam MZ, Islam MS. Is there any association between undesired children and health status of under-five children? Analysis of a nationally representative sample from Bangladesh. *BMC Pediatr.* 2022;22(1):445. Epub 20220725.
11. Kaufmann RB, Morris L, Spitz AM. Comparison of two question sequences for assessing pregnancy intentions. *Am J Epidemiol.* 1997;145(9):810-6.
12. Aiken AR, Westhoff CL, Trussell J, Castaño PM. Comparison of a Timing-Based Measure of Unintended Pregnancy and the London Measure of Unplanned Pregnancy. *Perspect Sex Reprod Health.* 2016;48(3):139-46. Epub 20160830.
13. Hall JA, Barrett G, Copas A, Stephenson J. London Measure of Unplanned Pregnancy: guidance for its use as an outcome measure. *Patient Relat Outcome Meas.* 2017;8:43-56. Epub 20170406.
14. Das S, Hall J, Barrett G, Osrin D, Kapadia S, Jayaraman A. Evaluation of the Hindi version of the London Measure of Unplanned Pregnancy among pregnant and postnatal women in urban India. *BMC Pregnancy Childbirth.* 2021;21(1):602. Epub 20210904.
15. Althubaiti A. Sample size determination: A practical guide for health researchers. *Journal of General and Family Medicine.* 2023;24(2):72-8.
16. Santelli J, Rochat R, Hatfield-Timajchy K, Gilbert BC, Curtis K, Cabral R, et al. The measurement and meaning of unintended pregnancy. *Perspectives on sexual and reproductive health.* 2003:94-101.
17. Bakibinga P, Matanda DJ, Ayiko R, Rujumba J, Muiruri C, Amendah D, et al. Pregnancy history and current use of contraception among women of reproductive age in Burundi, Kenya, Rwanda, Tanzania and Uganda: analysis of demographic and health survey data. *BMJ open.* 2016;6(3):e009991.
18. Khan MN, Islam MM. Women's experience of unintended pregnancy and changes in contraceptive methods: evidence from a nationally representative survey. *Reprod Health.* 2022;19(1):187. Epub 20220901.

19. Achana FS, Bawah AA, Jackson EF, Welaga P, Awine T, Asuo-Mante E, et al. Spatial and socio-demographic determinants of contraceptive use in the Upper East region of Ghana. *Reproductive health*. 2015;12(1):29.
20. Lin L, Lu C, Chen W, Li C, Guo VY. Parity and the risks of adverse birth outcomes: a retrospective study among Chinese. *BMC pregnancy and childbirth*. 2021;21(1):1-11.
21. Meselu W, Habtamu A, Woyraw W, Birlew Tsegaye T. Trends and predictors of modern contraceptive use among married women: Analysis of 2000-2016 Ethiopian Demographic and Health Surveys. *Public Health Pract (Oxf)*. 2022;3:100243. Epub 20220313.
22. Khan MN KS, Khan MMA, Islam MM, . Sibling Sex Composition and Contraceptive Use among Women in Bangladesh. *Sexual and Reproductive Health Matters* (under-review). 2023.
23. Rajan S, Nanda P, Calhoun LM, Speizer IS. Sex composition and its impact on future childbearing: a longitudinal study from urban Uttar Pradesh. *Reprod Health*. 2018;15(1):35. Epub 20180227.
24. Robey B. Sex preference and fertility: what is the link? *Asia Pac Pop Policy*. 1987(2):1-4.
25. Calhoun LM, Nanda P, Speizer IS, Jain M. The effect of family sex composition on fertility desires and family planning behaviors in urban Uttar Pradesh, India. *Reproductive health*. 2013;10:1-11.
26. Asadullah MN, Mansoor N, Randazzo T, Wahhaj Z. Is son preference disappearing from Bangladesh? *World Development*. 2021;140:105353.
27. Eliason S, Baiden F, Tuoyire DA, Awusabo-Asare K. Sex composition of living children in a matrilineal inheritance system and its association with pregnancy intendedness and postpartum family planning intentions in rural Ghana. *Reproductive health*. 2018;15:1-10.
28. Khan MN, Akter S, Islam MM. Availability and readiness of healthcare facilities and their effects on long-acting modern contraceptive use in Bangladesh: analysis of linked data. *BMC Health Serv Res*. 2022;22(1):1180. Epub 20220921.

29. Guliani H, Sepehri A, Serieux J. Determinants of prenatal care use: evidence from 32 low-income countries across Asia, Sub-Saharan Africa and Latin America. *Health policy and planning*. 2014;29(5):589-602.
30. Qiu X, Zhang S, Sun X, Li H, Wang D. Unintended pregnancy and postpartum depression: a meta-analysis of cohort and case-control studies. *Journal of Psychosomatic Research*. 2020;138:110259.