

Multiple introductions and country-wide spread of DENV-2 genotype II (Cosmopolitan) in Brazil

Tiago Gräf^{1#}, Caroline do Nascimento Ferreira¹, Gustavo Barbosa de Lima², Raul Emídio de Lima², Lais Ceschini Machado^{3,4}, Tulio de Lima Campos⁴, Michelle Orane Schemberger⁵, Helisson Faoro⁵, Marcelo Henrique Santos Paiva⁶, Matheus Filgueira Bezerra⁷, Valdinete Nascimento⁸, Victor Souza⁸, Fernanda Nascimento⁸, Matilde Mejía⁸, Dejanane Silva⁸, Yasmin Silva de Oliveira⁸, Luciana Gonçalves^{8,9}, Tatyana Costa Amorim Ramos⁹, Daniel Barros de Castro⁹, Ana Ruth Arcanjo¹⁰, LACEN-AM team*, Herton Augusto Pinheiro Dantas¹¹, LAFRON-AM team*, Mayra Marinho Presibella¹², LACEN-PR team*, Sandra Bianchini Fernandes¹³, LACEN-SC team*, Tatiana Schaffer Gregianini¹⁴, CEVS-RS team*, Keilla Maria Paz e Silva¹⁵, LACEN-PE team*, Claudio Tavares Sacchi¹⁶, IAL team*, Ana Cecília Ribeiro Cruz^{17,18}, IEC team*, Claudia Nunes Duarte dos Santos¹, Ana Maria Bispo de Filippis¹⁹, Richard Steiner Salvato¹⁴, Gonzalo Bello¹⁹, Gabriel Luz Wallau^{3,4,20}, Felipe Naveca^{8,19}

¹ Laboratório de Virologia Molecular, Instituto Carlos Chagas, Fundação Oswaldo Cruz, Curitiba 81350-010, Brazil

² Núcleo de Plataformas Tecnológicas (NPT), Instituto Aggeu Magalhães (IAM), FIOCRUZ-Pernambuco, Recife, Pernambuco, Brazil

³ Departamento de Entomologia, Instituto Aggeu Magalhães (IAM)-Fundação Oswaldo Cruz-FIOCRUZ, Recife 50670-420, Brazil

⁴ Núcleo de Bioinformática (NBI), Instituto Aggeu Magalhães (IAM), FIOCRUZ-Pernambuco, Recife, Pernambuco, Brazil

⁵ Laboratório de Ciências e Tecnologias Aplicadas em Saúde, Instituto Carlos Chagas, Fundação Oswaldo Cruz, Curitiba 81350-010, Brazil

⁶ Núcleo de Ciências da Vida, Universidade Federal de Pernambuco (UFPE), Centro Acadêmico do Agreste-Rodovia BR-104, km 59-Nova Caruaru, Caruaru 55002-970, Brazil

⁷ Departamento de Microbiologia, Instituto Aggeu Magalhães (IAM), FIOCRUZ-Pernambuco, Recife, Pernambuco, Brazil

⁸ Laboratório de Ecologia de Doenças Transmissíveis na Amazônia, Instituto Leônidas e Maria Deane, FioCruz, Manaus, Brazil

⁹ Fundação de Vigilância em Saúde do Amazonas - Dra Rosemary Costa Pinto, Manaus, Amazonas, Brazil

¹⁰ Laboratório Central de Saúde Pública do Amazonas (LACEN-AM), Manaus, Amazonas, Brazil

¹¹ Laboratório de Fronteira de Tabatinga (LAFRON-AM), Tabatinga, Amazonas, Brazil

¹² Laboratório Central de Saúde Pública do Estado do Paraná (LACEN-PR), São José dos Pinhais, Paraná, Brazil

¹³ Laboratório Central de Saúde Pública do Estado de Santa Catarina (LACEN-SC), Florianópolis, Santa Catarina, Brazil

¹⁴ Laboratório Central de Saúde Pública do Rio Grande do Sul (LACEN-RS), Porto Alegre, Rio Grande do Sul, Brazil

¹⁵ Laboratório Central de Saúde Pública de Pernambuco (LACEN-PE), Recife, Pernambuco, Brazil

¹⁶ Laboratório Estratégico, Instituto Adolfo Lutz (IAL), Sao Paulo, 01246-902, Brazil

¹⁷ Department of Arbovirology and Hemorrhagic Fevers, Evandro Chagas Institute, Health and Environment Surveillance Secretariat, Ministry of Health, Ananindeua 670030-000, Brazil

¹⁸ Institute of Biological Sciences, Federal University of Pará, Belém 66075-110, PA, Brazil

¹⁹ Laboratório de Arbovírus e Vírus Hemorrágicos, Instituto Oswaldo Cruz, FioCruz, Rio de Janeiro, Rio de Janeiro, Brazil

²⁰ Department of Arbovirology, Bernhard Nocht Institute for Tropical Medicine, WHO Collaborating Center for Arbovirus and Hemorrhagic Fever Reference and Research. National Reference Center for Tropical Infectious Diseases. Bernhard-Nocht-Straße 74, 20359 Hamburg - Germany.

#Corresponding author: tiago.graf@fiocruz.br

* Lists of authors and their affiliations appear at the Supplementary Consortia List

Abstract

Dengue virus serotype 2, genotype Cosmopolitan (DENV-2-GII), is one of the most widespread DENV strains globally. In the Americas, DENV-2 epidemics has been dominated by DENV-2 genotype Asian-American (DENV-2-GIII) and the first cases of DENV-2-GII were only described in 2019, in Peru, and in 2021 in Brazil. To gain new information about the circulation of DENV-2-GII in Brazil, we sequenced 237 DENV-2 confirmed cases sampled between March 2021 and March 2023 and revealed that DENV-2-GII is already present in all geographic regions of Brazil. Phylogeographic analysis inferred that DENV-2-GII was introduced at least four times in Brazil, between May 2020 and August 2022, generating multiple clades that spread throughout the country with different success. Despite multiple introductions of DENV-2-GII, analysis of the country-wide laboratory surveillance data showed that the Brazilian dengue epidemic in 2022 was dominated by DENV-1 in most states. We suggest that massive circulation of DENV-2-GIII in previous years in Brazil might have created a population immune barrier for widespread transmission of DENV-2-GII, leading to sustained cryptic circulation and localized outbreaks of this new genotype. In summary, our study stresses the importance of arboviral genomic surveillance to close monitoring and better understand the potential impact of DENV-2-GII in the coming years.

Research letter

Dengue virus serotype 2 (DENV-2) can be divided into five endemic/epidemic (nonsylvatic) genotypes: American, Cosmopolitan, Asian-American, Asian-II, and Asian-I¹, also known as genotypes I-V, respectively. In the Americas, the DENV-2 American genotype (DENV-2-GI) was the first to be identified, in the 1940s. However, it was later replaced by the Asian-American genotype (DENV-2-GIII) in the 1980s, and cases of DENV-2-GI were not found since then². In Brazil, DENV-2-GIII was first introduced in the country in the 1990s and at least three other introductions happened between 1990 and 2014, all with origin in Caribbean or Northern South America countries^{3,4}. Besides DENV-2, in Brazil all four DENV serotypes were already identified, presenting a complex dynamics of introductions, co-circulation, and alternation of dominance, which is associated with an increasing number of severe dengue cases and deaths in recent years⁵.

In 2021, the first cases of DENV-2 Cosmopolitan genotype (DENV-2-GII) were identified in the Brazilian states of Goiás (GO)⁶ and Acre (AC)⁷, located in the Central-Western and Northern country regions, respectively. Phylogenetic analysis suggested that the Brazilian DENV-2-GII sequences derived from an outbreak of this genotype in Peru in 2019⁸, representing the first documented introduction of DENV-2-GII in the Americas. DENV-2-GII is one of the most widespread genotypes, circulating in Asia-Pacific, Middle East, Africa, and Oceania, substantially contributing to the global dengue burden⁹. Here we present new data on the DENV-2-GII spread in Brazil, showing that shortly after multiple introductions in the country, it is already present in all geographical regions.

Genomic surveillance for DENV was performed in the Brazilian states of Amazonas (AM), Pernambuco (PE), São Paulo (SP), Paraná (PR), Santa Catarina (SC), and Rio Grande do Sul (RS) by each State Central Laboratories (LACEN) and Fiocruz laboratories. A selection of DENV-2 positive samples in RT-qPCR test were submitted for whole genome amplification and sequencing using Illumina's Viral Surveillance Panel or COVIDseq Test adapted to DENV-2. Genomes were assembled in ViralFlow software¹⁰ or using Geneious Prime 2022 and consensus sequences were genotyped using online tools (methods detailed in **Supplementary text**).

Among 237 DENV-2 sequenced samples, we identified 60 DENV-2-GII (**Supplementary Table 1**) which were aligned with a DENV-2-GII representative global dataset. Maximum likelihood phylogenetic analysis revealed that all DENV-2-GII Brazilian genomes, from this and previous studies^{6,7}, clustered in a monophyletic clade with sequences from Peru and Bangladesh (**Supplementary Figure 1**). Our phylogeographic analysis estimated that DENV-2-GII was introduced from Bangladesh into Peru in April 2019 (November 2018 - August 2019, 95% High Posterior Density [HPD]), and from there, it was introduced in Brazil at least four times (**Figure 1A**). The first introduction occurred through the AC state (North) in May 2020 (June 2019 - November 2020, 95% HPD) and from AC, DENV-2-GII disseminated to all other four Brazilian regions represented by the states of GO (Central-West), PE (Northeast), SP (Southeast), and SC (South). The oldest DENV-2-GII Brazilian samples collected in 2021 in AC (February to March), PE (July) and GO (November) clustered within this clade. A second introduction of DENV-2-GII happened in the state of SP in March 2021 (September 2020 - July 2021, 95% HPD), and from there, DENV-2-GII spread southwards to the states of PR, SC, and RS. A third introduction of DENV-2-GII occurred in January 2022 (October 2021 - March 2022, 95% HPD) in the state of RS, without evidence of further dissemination to other Brazilian states. Finally, the most recent introduction happened in AM state in August 2022 (June 2022 - October 2022, 95% HPD), causing an outbreak in cities close to the border with Peru and Colombia¹¹ . It is interesting to note that the first introductions of DENV-2-GII into Brazil most probably occurred during the period (May 2020 - March 2021) in which the country was most heavily affected by the SARS-CoV-2 pandemic, and that despite the control measures implemented to restrict human mobility in that period, DENV-2-GII viruses introduced in AC and SP were able to disseminate to other Brazilian states.




Figure 1. Spatio-temporal evolution of DENV-2 genotype II (Cosmopolitan) in South America and circulation dynamics of DENV serotypes in Brazil in recent years. (a) Time-scaled phylogenetic tree with ancestral locations inferred by the phylogeographic model. Branches are colored according to the most likely ancestral location, and pie-charts show the posterior probability of each location for key nodes with high uncertainty. Error bars show the 95% high posterior density of node heights related to introductions to Brazil. Black dots on branches indicate a posterior probability higher than 0.9. (b) Relative frequency of each DENV serotype in all Brazilian states between 2018 and 2022. States acronyms are according to the ISO 3166-2:BR.

To assess the potential of DENV-2-GII to spread and trigger relevant outbreaks in Brazil, we analyzed the dynamics of DENV serotypes' circulation in the country as identified through PCR tests by the laboratory surveillance and informed in the national system of diseases of compulsory notification (SINAN). Between 2018 and 2022, the Brazilian dengue epidemic was dominated by DENV-1 and DENV-2, with significant regional differences (**Figure 1B**). A predominance of DENV-2 after 2020, which could indicate a genotype II outbreak, was observed in the Northern states of AC (where genomic surveillance detected DENV-2-GII circulation), RO and RR, in the Southeastern states of RJ and Espirito Santo (ES), and in the Northeastern states of Ceara (CE), Paraíba (PB), Rio Grande do Norte (RN), Sergipe (SE) and PE. In the latter, only one sequence of DENV-2-GII was found compared to 80 sequences of DENV-2-GIII in 2021, indicating that the dominance of DENV-2 in PE at that time was not caused by genotype II, which could also be the case for the neighboring Northeastern states. A high prevalence of DENV-2 was observed in several states between 2018 and 2020, but there was no evidence of circulation of DENV-2-GII in Brazil before 2021 (**Supplementary Figure 2**). Finally, in other states including AM and RS, where genomic surveillance detected recent introductions of DENV-2-GII in 2022, DENV-1 was responsible for most dengue cases in the whole period supporting that DENV-2-GII only caused localized outbreaks in those states. In fact, in 2022, DENV-1 was responsible for more than 80% of the dengue cases in 17 out of 25 Brazilian states with data available.

In summary, this study confirms that DENV-2-GII is circulating nationally in Brazil, expanding the findings of previous localized studies^{6,7}. Our phylogenetic analysis unveiled multiple introductions of this genotype into Brazil from Peru, generating multiple clades that spread in the country with different success. Despite the wide geographic dispersion of DENV-2-GII, dengue epidemics in 2021-2022 in most Brazilian states were dominated by DENV-1, supporting that circulation of DENV-2-GIII in previous years probably created a high population immunity barrier against transmission of DENV-2-GII in Brazil. However, the rapid spread of DENV-2-GII across the country's territory and the heterogeneous population immune landscape across regions, could create an epidemic scenario for sustained cryptic circulation and localized outbreaks. Indeed, DENV-2-GII was probably responsible for most DENV-2 infections in Brazil in 2022 and this warrants close monitoring to better understand the potential impact of this genotype in the coming years.

References

- 1 Twiddy SS, Farrar JJ, Vinh Chau N *et al.* Phylogenetic relationships and differential selection pressures among genotypes of dengue-2 virus. *Virology* 2002; **298**: 63–72.
- 2 Allicock OM, Lemey P, Tatem AJ *et al.* Phylogeography and population dynamics of dengue viruses in the Americas. *Mol Biol Evol* 2012; **29**: 1533–1543.
- 3 de Jesus JG, Dutra KR, Sales FC da S *et al.* Genomic detection of a virus lineage replacement event of dengue virus serotype 2 in Brazil, 2019. *Mem Inst Oswaldo Cruz* 2020; **115**: e190423.
- 4 Brito AF, Machado LC, Oidtmann RJ *et al.* Lying in wait: the resurgence of dengue virus after the Zika epidemic in Brazil. *Nat Commun* 2021 121 2021; **12**: 1–13.
- 5 Fares RCG, Souza KPR, Añez G, Rios M. Epidemiological Scenario of Dengue in Brazil. *BioMed Res Int* 2015; **2015**: 1–13.
- 6 Giovanetti M, Pereira LA, Santiago GA *et al.* Emergence of Dengue Virus Serotype 2 Cosmopolitan Genotype, Brazil. *Emerg Infect Dis* 2022; **28**: 1725.
- 7 Amorim MT, Hernández LHA, Naveca FG *et al.* Emergence of a New Strain of DENV-2 in South America: Introduction of the Cosmopolitan Genotype through the Brazilian-Peruvian Border. 2023. doi:10.20944/preprints202305.1014.v1.
- 8 García MP, Padilla C, Figueroa D, Manrique C, Cabezas C. Emergence of the Cosmopolitan genotype of dengue virus serotype 2 (DENV2) in Madre de Dios, Peru, 2019. *Rev Peru Med Exp Salud Pública* 2022; **39**: 126–128.
- 9 Yenamandra SP, Koo C, Chiang S *et al.* Evolution, heterogeneity and global dispersal of cosmopolitan genotype of Dengue virus type 2. *Sci Rep* 2021; **11**: 13496.
- 10 Dezordi FZ, Neto AM da S, Campos T de L *et al.* ViralFlow: A Versatile Automated Workflow for SARS-CoV-2 Genome Assembly, Lineage Assignment, Mutations and Intra-host Variant Detection. *Viruses* 2022; **14**: 217.
- 11 Naveca FG, Investigação sobre o surto de dengue no segundo semestre de 2022 nos municípios da região do Alto Solimões, região da tríplice-fronteira Brasil, Colômbia e Peru - Relatório preliminar. *Personal communication* 2023. Available at: https://www.genomahcov.fiocruz.br/wp-content/uploads/2023/05/Investigacao_sobre_o_surto_de_DENV2_cosmopolita_n_o_Amazonas_Relatorio_preliminar_v2_assinado.pdf