

The Mediterranean Diet Scale (MDS): translation and validation of the Arabic version

Authors: Raghdah Aljehani,¹ Ghaidaa Aljehani,¹ Hanaa Alharazi,² Paula M Horta,³ Camila Kümmel Duarte,³ Gabriela Lima de Melo Ghisi.^{4,5*}

Affiliations:

1. Rehabilitation Department, King Abdullah Medical City, Makkah, Saudi Arabia.
2. Cardiology Department, King Abdullah Medical City, Makkah, Saudi Arabia.
3. Department of Nutrition, Federal University of Minas Gerais, Belo Horizonte, Brazil.
4. Department of Physical Therapy, University of Toronto, Toronto, Canada.
5. KITE, Toronto Rehabilitation Institute, University Health Network, Toronto, Canada.

*Corresponding author:

Department of Physical Therapy, University of Toronto, Toronto, Canada.
160-500 University Ave, Toronto, Canada M5G 1V7
Tel: (416) 597-3422 x. 5261 Fax: (416) 425-0301
email: gabriela.meloghisi@uhn.ca

Author Contributions:

Conceptualization, RA and GLMG; Methodology, GLMG; Formal Analysis, GLMG;
Investigation, RA, GA, HA, PMH, CK; Resources, RA; Data Curation, RA, GLMG; Writing –

- 23 Original Draft Preparation, GLMG; Writing – Review & Editing RA, GA, HA, PMH, CK;
- 24 Project Administration, GLMG.

Abstract

Objectives: The self-administered version of the Mediterranean Diet Scale (MDS) has been developed to test the inherent characteristics of this dietary pattern in a quick and simple way, due to the need of this assessment in the clinical and research setting. This study aimed to translate and psychometrically validate the self-administered MDS in Arabic (CRBS-A).

Methods: The original (English) version was originally translated to Arabic, followed by back-translation. Next, 10 healthcare providers, followed by 10 cardiovascular disease (CVD) patients rated the face and content validity (CV) of materials, providing input to improve cross-cultural applicability. Then, 200 patients from Saudi Arabia completed the questionnaire, of which factor structure, internal consistency, criterion and construct validity were assessed.

Results: Content and face validity was supported based on experts and patients' reviews (ranges: CV scores 0.9-1.0/1.0 and clarity 3.5 to 4.5/5). Minor edits were made. Subsequent factor analysis revealed 4 factors consistent with the original version of the instrument, all internally consistent. Total CRBS-A α was 0.74. Criterion validity was confirmed by the significantly higher scores in patients who participated in CR. Construct validity was also established by significant associations between MDS scores and monthly family income, having the diagnosis of acute coronary syndrome or with a history of valve repair or replacement, being obese or having dyslipidemia.

Conclusions: Overall, these results confirm the validity and reliability of the MDS in Arabic-speaking patients.

Keywords: Questionnaires and Surveys; Psychometrics; Diet, Mediterranean; Cardiac Rehabilitation; Coronary Artery Disease/Prevention and Control; Diet Surveys.

48 Introduction

49 Cardiovascular diseases (CVDs) are the most notable causes of disease burden across
50 many nations [1], including Arabic-speaking countries [2,3]. In addition to health impacts, CVDs
51 also carry a substantial economic burden on patients, their families, and societies due to medical
52 costs, diminished productivity from disability and premature death [4-6]. Cardiovascular
53 associations - regionally and internationally – recommend secondary prevention interventions to
54 address this burden, which rely on successful health behaviour change [7-10]. These
55 recommendations are based on findings linking CVDs to modifiable risk factors, particularly
56 unhealthy behaviours such as poor-quality diet [10-12].

57 Cardiac rehabilitation (CR) is a comprehensive outpatient chronic disease management
58 program designed to facilitate *heart*-health behaviours [9]. CR typically includes core
59 components that address CVD risk factors, including exercise training, patient education,
60 psychosocial management risk factor modification and nutritional counseling [13]. Research has
61 shown that when all these multifactorial components are delivered, CR participation is associated
62 with *reductions of mortality up to 25%*, as well as beneficial effects on morbidity, symptoms,
63 exercise tolerance and capacity, lipid and blood pressure levels, blood pressure, psychosocial
64 functioning and behaviour change [14-16]. In order to tailor adequate CR strategies and
65 sustainable recommendations, it is crucial to fully understand patients' habits that determine
66 cardiovascular health [17]. One of those is an adequate nutrition [18].

67 Over the years, various dietary patterns have been studied with the goal to identify the
68 one that is mostly effective on heart health. The Mediterranean Dietary Pattern (MDP) has been
69 greatly studied, with evidence showing its effects on the prevention and treatment of CVDs, as
70 well as on the reduction of CVD mortality (MDP) [19-21]. While the MDP is widely

recommended to reduce CVD risk, there is not robust evidence for adherence in CR after the many changes that happened in programs due to the COVID-19 pandemic [22,23].

In this context, questionnaires are considered useful means to collect important information that can support choices related to the care of patients [24]; however, not always their application can be implemented into clinical practices. For example, full-length food frequency questionnaires (FFQ) or food records used in dietary assessments have limitations related to accuracy and time [25], leading to a significant burden on the respondents and healthcare team [26]. The 14-item Mediterranean Diet Score (MDS) was developed to assess adherence to the MDP [27], due to its effectiveness for primary and secondary prevention of CVDs [28-30]. In order to eliminate barriers for its use, a self-administered version of the MDS was developed and validated by Ghisi et al (2018) [31].

The prevalence and control of CVDs is considered the biggest public health challenge in Arabic-speaking countries [32], with risk factors such as obesity more than doubling in the last years [33]. Changes in food consumption patterns have been observed in these countries, including a diet with high content of fat, sugars, sodium and cholesterol [34,35]. In addition, availability of CR programs in Arab countries is extremely limited, with only 8 programs identified in a previous global survey of CR programs [36]. In this context, having a validated questionnaire to evaluate the MDP adherence in Arabic-speaking patients is timely, supporting practitioners to develop and deliver dietary recommendations that are aligned with patients' needs. Therefore, this study aimed to translate and psychometrically validate the Arabic version of the MDS.

Methods

Design and Procedures

The study was reviewed and approved by the King Abdullah Medical City, Saudi Arabia (22-944). Data was collected between September 2022 and January 2023. Permission to conduct this Arabic translation and validation was obtained, and the first author of the original MDS self-administered developmental study was invited to be part of the Arabic project. Authors had no access to information that could identify individual participants during or after data collection.

Following best practices [37], this study included the following steps: 1) translation from English to Arabic and cultural adaptation of the MDS, 2) review by an expert panel and Arabic-speaking patients, and 3) cross-sectional survey for evaluation of psychometric properties.

As the first step, the initial conceptual translation of the scale from English to the target language (Arabic) was performed by an independent translator. Then, a second translator (now blinded to the objectives of the study) back translated the scale. All versions – original, translated, and back translated – were reviewed and combined by the research team into one version, which was reviewed by an expert team comprised of 10 healthcare providers that care for CVD patients (step 2). This expert team assessed face and content validity; the later by rating each item on a 5-point Likert-type scale ranging from 1 = completely irrelevant to 5 = very relevant. Content validity index (CVI) for the items (I-CVI) and scale (S-CVI) were computed. For both, only values higher than 0.80 were considered appropriate [38]. The scale was then revised based on CVI scores and suggestions made by experts. As part of phase 2, a group of 10 individuals from the target population assessed clarity of the scale using a 5-point Likert-type scale ranging from 1 = not clear to 5 = very clear. Results were analyzed to further revise the scale.

Finally, psychometric properties were evaluated (step 3). The translated and culturally adapted tool was administered to a convenience sample of cardiac patients from a cardiology center in Saudi Arabia. Patients were invited to participate in the study while attending regular clinical consultations. Those that accepted, signed the consent form, and completed the Arabic MDS in-person. The following psychometric properties were assessed following the Consensus-based Standards for the selection of health Measurement Instruments (COSMIN) taxonomy [39]: factor structure, internal consistency, construct, and criterion validity.

The self-administered version of the MDS

The original 14-item MDS was developed in Spanish [27] and later translated to English [40]. The English version was used to develop the self-administered version [31]. This version included pictures to facilitate comprehension of questions and a simplified language. It is comprised of 13 items with yes/no options. The items are divided into 4 domains as follows: meats and processed foods; olive oil and sauce; fruits, vegetable, nuts, and legumes; and, fish and seafood [31]. Each item is scored as 0 or 1 in accordance with MDP adherence [41]. The final score ranges from 0 to 13 (< 5 = low adherence to MDP; > 10 = high adherence to MDP) [31]. The self-administered version of the MDS is also available in Brazilian [42] and Chinese [43] languages.

Participants

The inclusion criteria for patient participants were confirmed cardiac diagnosis or presence of cardiovascular risk factors, 18 years of age or older, and being able and willing to provide informed consent. The exclusion criteria for patient participants were inability to eat by mouth or having food and dietary restrictions (e.g., nut allergies) and having any impairment

which preclude the participant's ability to answer the questionnaire (e.g., visual). A minimum of 200 participants was required following the recommendation for factor analysis [39].

Patient participants self-reported their sociodemographic and clinical characteristics, including the following: age, sex, cardiovascular diagnosis and risk factors, living area, educational level, and family income. In addition, CR participation (yes/no) was self-reported to assess criterion validity.

Data Analysis

First, exploratory factor analysis was performed, using the main component method for factor extraction. Only factors with eigenvalues > 1.0 were considered. Item factor loadings > 0.3 were used in finalizing the items for each factor and interpreting the factors [44]. Next, internal consistency was determined by calculating Cronbach's *alpha* values of the scale and subscales. Cronbach's *alpha* higher than 0.70 was considered acceptable [45].

Criterion validity was assessed by differences in MDS total scores by CR participation using independent samples *t*-tests. Construct validity was assessed by exploring the associations between self-reported sociodemographic characteristics of participants and MDS scores using Pearson's correlation, independent samples *t*-tests, and analysis of variance.

Finally, a descriptive analysis of the Arabic MDS was performed. The Statistical Package for Social Sciences v. 28 (SPSS Inc., Chicago, IL, USA) was used for data analysis. The level of significance for all tests was set at 0.05.

Results

Translation, cultural adaptation and review by an expert panel and patients

Following translation and back translation, a revised Arabic MDS version was reviewed by an expert panel (n=10). The I-CVI ranged from 0.9 and 1.0 and the S-CVI was 0.9. These

results identify an acceptable content validity for the Arabic MDS. For patients' review (n=10), clarity scores ranged from 3.5 to 4.5/5 (mean 4.2±0.3). Given these results, two items were culturally adapted as follows: pork removed from items 10 and 11 due to religious and cultural reasons. In addition, items 6 and 13 were rephrased to increase clarity. Appendix 1 displays the Arabic MDS.

Cross-sectional survey for evaluation of psychometric properties

Overall, 200 patients completed the Arabic MDS. Their sociodemographic and clinical characteristics are presented in Table 1.

Results from the Kaiser-Meyer-Olkin index (KMO=0.78) and Bartlett's Sphericity tests ($X^2=403.921$, $p<0.001$) indicated that the data were suitable for exploratory factor analysis. Four factors were extracted, similar to the original validation [25]. These factors represented 62.3% of the total variance and were all internally consistent (Cronbach's alpha ranged from 0.70-0.79). Factor 1 reflected items related to meats/processed foods (5 items), factor 2 related to olive oil/sauce items (3 items), factor 3 related fruits/vegetables and nuts/legumes (4 items), and factor 4 related to fish/seafood (1 item). Table 2 presents the factor structure of the Arabic MDS SV, including item loadings. The total Cronbach's alpha of the scale was 0.74, confirming the internal consistency of the scale.

With regard to criterion validity, significant associations were observed between total scores and CR participation. Those who participated in CR had significantly higher adherence to the MDP (i.e., higher MDS scores) than those who did not participate ($p=0.002$; Table 1).

As also shown in Table 1, with regard to construct validity, there were significant associations between MDS scores and monthly family income ($p=0.02$), having the diagnosis of acute coronary syndrome ($p=0.02$), a history of valve repair or replacement ($p=0.03$), being

obese ($p=0.02$) or having dyslipidemia ($p=0.03$). In this context, those with a family monthly income under SAR\$10,000 presented a significantly lower MDP adherence than their counterparts. In addition, those who had a diagnosis of acute coronary syndrome, those that were obese, with dyslipidemia, or those who underwent a valve repair or replacement, presented MDP adherence than those without the diagnosis or a history of these procedures (Table 1).

Adherence to the MDP

The total score for the self-administered version of the MDS in this study was $11.0 \pm 8.0/13$. Overall, 57 (28.5%) patients were classified as having high adherence to MDP and 29 (14.5%) as having low adherence. Table 2 also displays the number of patients that adhere to MDP recommendations (i.e., responded “yes” to each item asked in the questionnaire). Items with the highest adherence included seasoning foods with a combination of tomato, garlic, onions, or leek ($n=171$; 85.5%), eating poultry more often than red meat ($n=164$; 82.0%), and eating three servings or more of legumes a week ($n=137$; 68.5%). One item - eating three servings or more of fish or seafood each week – had the percentage of yes lower than 50% ($n=87$; 43.5%).

Discussion

Research has proven the benefits of the MDP for both primary and secondary prevention of CVDs [46]. However, multi-level barriers affect adherence to dietary recommendations, including in participants of CR programs [47]. Having a validated and convenient tool to assess MDP adherence – and ultimately guide clinical and educational practices – is timely in countries like Saudi Arabia where a small percentage of population meet dietary recommendations and programmes to improve dietary behaviours are urgently needed to reduce the current and future burden of disease [48]. Therefore, this study established the Arabic version of the MDS to assess

adherence to the MDP through a multi-step process that included translation, cultural adaptation and assessment of psychometric properties. Content and face validity was supported based on experts and patients' reviews. Subsequent factor analysis revealed 4 factors consistent with the original version of the instrument [31], all internally consistent. Criterion validity was confirmed by the significantly higher scores in patients who participated in CR. Construct validity was also established by significant associations between MDS scores and monthly family income, having the diagnosis of acute coronary syndrome or with a history of valve repair or replacement, being obese or having dyslipidemia. Overall, these results confirm the validity and reliability of the MDS in Arabic-speaking patients.

The self-administered version of the MDS – originally developed in English [31] and later translated and validated to Portuguese [41] and Chinese [42] – was developed to simplify the assessment of MDP adherence in clinical and research settings. This is the 4th language available and other validation projects are underway (including Spanish and French). The Cronbach's alpha of the Arabic MDS was 0.74, which is higher than previous studies (range: 0.42-0.69). The total MDS score of this score was higher than the Canadian (10.2 ± 1.9) [31], Portuguese (6.94 ± 1.99) [41], and Chinese (7.7 ± 2.4) [42] versions. Items identified in the current study as the ones with the highest adherence included flavouring foods with a combination of tomato, garlic, onions, or leek, eating poultry more often than red meat, and eating three servings or more of legumes a week. These results are similar to responses identified in the previous studies using the MDS; in Canada [31], the items with highest adherence included eating poultry more often than meat and flavouring foods with a combination of tomato, garlic, onions or leek; in Brazil [41], eating three servings or more of legumes a week and flavoring foods with a combination of tomato, garlic, onions, or leeks; in China [42], flavouring foods with a

combination of tomato, garlic, onions or leek. On the other hand, the self-reported consumption vegetable in the Arab-speaking sample was lower than the other samples (50% vs. 85% in Canada, 79% in Brazil and 87% in China) [31,41,42], despite the fact that the Saudi dietary guidelines strongly recommend a higher intake of vegetables [49].

As previously mentioned, construct validity of the Arabic MDS was confirmed in this study, as well as in other versions of this scale [31,41,42]. Specifically, those with lower monthly family income had significantly lower MDP adherence than their counterparts. Despite its complexity, the impact of socioeconomic on dietary pattern is undeniably important [50-52]. In general, people with higher income tend to follow the MDP recommendations [53]; however, in any context, recommendations to follow healthy dietary habits should be given to CVD patients. Future research examining the changes in adherence to the MD over time and after educational interventions is warranted, including in groups of individuals with low socioeconomic status.

Those who participated in CR had significantly higher adherence to the MDP (i.e., higher MDS scores) than those who did not participate. These results support the effects of participation in CR – and delivery of nutrition counselling – on adherence to heart-healthy dietary patterns [54]. Furthermore, studies should describe the strategies, tools, and techniques employed within the interventions used to adhere to the MDP and barriers and facilitators to follow this behaviour.

There are some limitations in this study. First, findings are limited to one center from 1 Arabic-speaking country (Saudi Arabia); results cannot be generalizable to the other 21 countries in the world that speak Arabic. Second, there may be selection and retention bias, which also limits the generalizability of the findings. Third, because of the nature of this cross-sectional study, causal conclusions cannot be drawn; it is recommended to test if the scale is sensitive to change before and after a nutritional or educational intervention. The final limitation pertains to

252 the assessment of other psychometric properties such as test-retest reliability, which should be
253 explored in future studies.

254 **Conclusion**

255 In conclusion, the Arabic MDS is a valid and reliable tool that can be used to assess MDP
256 adherence of Arabic-speaking cardiac patients. It is hoped that this scale supports the adequate
257 assessment of food intake and changes in dietary habits, which plays a highly significant role in
258 the care of people living with a cardiovascular condition.

259 **Supplementary Materials**

260 Appendix 1: Arabic MDS

261

262 **Conflicts of Interest**

263 The authors declare no conflict of interest.

References

1. Roth GA, Mensah, GA, Johnson, CO, et al. Global Burden of Cardiovascular Diseases and Risk Factors, 1990–2019. J Am Coll Cardiol. 2020; 76 (25): 2982–3021.
<https://doi.org/10.1016/j.jacc.2020.11.010>.
2. Aljefree N, Ahmed F. Prevalence of Cardiovascular Disease and Associated Risk Factors among Adult Population in the Gulf Region: A Systematic Review. Adv Public Health. 2015; 2015:1–23. <https://doi.org/10.1155/2015/235101>.
3. Ahmed AM, Hersi A, Mashhoud W, et al. Cardiovascular Risk Factors Burden in Saudi Arabia: The Africa Middle East Cardiovascular Epidemiological (ACE) Study. J Saudi Heart Assoc; 2017;29 (4): 235–43. <https://doi.org/10.1016/j.jsha.2017.03.004>.
4. Carter HE, Schofield D, Shrestha R. Productivity Costs of Cardiovascular Disease Mortality across Disease Types and Socioeconomic Groups. Open Heart. 2019; 6 (1): e000939.
<https://doi.org/10.1136/openhrt-2018-000939>.
5. Oldridge NB. Economic Burden of Physical Inactivity: Healthcare Costs Associated with Cardiovascular Disease. Eur J Cardiovasc Prev Rehabil. 2008; 15 (2): 130–9.
<https://doi.org/10.1097/HJR.0b013e3282f19d42>.
6. Salvatore FP, Spada A, Fortunato F, et al. Identification of Health Expenditures Determinants: A Model to Manage the Economic Burden of Cardiovascular Disease. IJERPH. 2021; 18 (9): 4652. <https://doi.org/10.3390/ijerph18094652>.
7. Piepol MF, Hoes AW, Agewall S, et al. 2016 European Guidelines on Cardiovascular Disease Prevention in Clinical Practice: The Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (Constituted by Representatives of 10 Societies and by Invited Experts) Developed with the

287 Special Contribution of the European Association for Cardiovascular Prevention &
 288 Rehabilitation (EACPR). Eur Heart J. 2016; 37 (29): 2315–81.
 289 <https://doi.org/10.1093/eurheartj/ehw106>.

290 8. Thomas RJ, King M, Lui K, et al. AACVPR/ACCF/AHA 2010 Update: Performance
 291 Measures on Cardiac Rehabilitation for Referral to Cardiac Rehabilitation/Secondary Prevention
 292 Services: A Report of the American Association of Cardiovascular and Pulmonary Rehabilitation
 293 and the American College of Cardiology Foundation/American Heart Association Task Force on
 294 Performance Measures (Writing Committee to Develop Clinical Performance Measures for
 295 Cardiac Rehabilitation). Circulation. 2010; 122 (13): 1342–50.
 296 <https://doi.org/10.1161/CIR.0b013e3181f5185b>.

297 9. Grace SL, Turk-Adawi KI, Contractor A, et al. Cardiac Rehabilitation Delivery Model for
 298 Low-Resource Settings. Heart. 2016; 102 (18): 1449–55. [https://doi.org/10.1136/heartjnl-2015-](https://doi.org/10.1136/heartjnl-2015-309209)
 299 [309209](https://doi.org/10.1136/heartjnl-2015-309209).

300 10. Smith SC, Benjamin EJ, Bonow RO, et al. AHA/ACCF Secondary Prevention and Risk
 301 Reduction Therapy for Patients With Coronary and Other Atherosclerotic Vascular Disease:
 302 2011 Update: A Guideline From the American Heart Association and American College of
 303 Cardiology Foundation. Circulation. 2011; 124 (22): 2458–73.
 304 <https://doi.org/10.1161/CIR.0b013e318235eb4d>.

305 11. Yusuf S, Joseph P, Rangarajan S, et al. Modifiable Risk Factors, Cardiovascular Disease, and
 306 Mortality in 155 722 Individuals from 21 High-Income, Middle-Income, and Low-Income
 307 Countries (PURE): A Prospective Cohort Study. Lancet. 2020; 395 (10226): 795–808.
 308 [https://doi.org/10.1016/S0140-6736\(19\)32008-2](https://doi.org/10.1016/S0140-6736(19)32008-2).

309 12. Yusuf S, Hawken S, Ôunpuu S, et al. Effect of Potentially Modifiable Risk Factors
310 Associated with Myocardial Infarction in 52 Countries (the INTERHEART Study): Case-Control
311 Study. *Lancet*. 2004; 364 (9438): 937–52. [https://doi.org/10.1016/S0140-6736\(04\)17018-9](https://doi.org/10.1016/S0140-6736(04)17018-9)
312 13. Mehra VM, Gaalema DE, Pakosh M, et al. Systematic Review of Cardiac Rehabilitation
313 Guidelines: Quality and Scope. *Eur J Prev Cardiol*. 2020; 27 (9): 912–28.
314 <https://doi.org/10.1177/2047487319878958>.
315 14. Taylor RS, Dalal HM, McDonagh STJ. The Role of Cardiac Rehabilitation in Improving
316 Cardiovascular Outcomes. *Nat Rev Cardiol*. 2022; 19 (3): 180–94.
317 <https://doi.org/10.1038/s41569-021-00611-7>
318 15. Kabboul N, Tomlinson G, Francis T, et al. Comparative Effectiveness of the Core
319 Components of Cardiac Rehabilitation on Mortality and Morbidity: A Systematic Review and
320 Network Meta-Analysis. *JCM*. 2018; 7 (12): 514. <https://doi.org/10.3390/jcm7120514>
321 16. Shi W, Ghisi GLM, Zhang L, et al. Systematic Review, Meta-analysis and Meta-regression
322 to Determine the Effects of Patient Education on Health Behaviour Change in Adults Diagnosed
323 with Coronary Heart Disease. *J Clin Nurs*. 2022; jocn.16519. <https://doi.org/10.1111/jocn.16519>
324 17. Artinian NT, Fletcher GF, Mozaffarian D, et al. Interventions to Promote Physical Activity
325 and Dietary Lifestyle Changes for Cardiovascular Risk Factor Reduction in Adults: A Scientific
326 Statement From the American Heart Association. *Circulation*. 2010; 122 (4): 406–41.
327 <https://doi.org/10.1161/CIR.0b013e3181e8edf1>
328 18. Ravera A, Carubelli V, Sciatti E, et al. Nutrition and Cardiovascular Disease: Finding the
329 Perfect Recipe for Cardiovascular Health. *Nutrients*. 2016; 8(6): 363.
330 <https://doi.org/10.3390/nu8060363>

- 331 19. Lichtenstein AH, Appel LJ, Vadiveloo M, et al. 2021 Dietary Guidance to Improve
332 Cardiovascular Health: A Scientific Statement From the American Heart
333 Association. *Circulation*. 2021; 144(23): e472–87.
334 <https://doi.org/10.1161/CIR.0000000000001031>
- 335 20. Galbete C, Schwingshackl L, Schwedhelm C, et al. Evaluating Mediterranean diet and risk of
336 chronic disease in cohort studies: an umbrella review of meta-analyses. *Eur J Epidemiol*.
337 2018; 33(10): 909–31. <https://doi.org/10.1007/s10654-018-0427-3>
- 338 21. Dinu M, Pagliai G, Casini A, et al. Mediterranean diet and multiple health outcomes: an
339 umbrella review of meta-analyses of observational studies and randomised trials. *Eur J Clin*
340 *Nutr*. 2018; 72(1): 30–43. <https://doi.org/10.1038/ejcn.2017.58>
- 341 22. Linan Pinto M, Pinto R, Charneca S, et al. Body composition, lipid profile and mediterranean
342 diet adherence in cardiovascular disease patients attending a long-term exercise-based cardiac
343 rehabilitation program during COVID-19 pandemic. *Eur J Prev Cardiol*. 2021; 28(Suppl 1):
344 zwab061.187. <https://doi.org/10.1093/eurjpc/zwab061.187>
- 345 23. Della Valle PG, Mosconi G, Nucci D, et al. Adherence to the Mediterranean Diet during the
346 COVID-19 national lockdowns: a systematic review of observational studies. *Acta bio-medica*
347 *Atenei Parmensis*. 2021; 92(S6): e2021440. <https://doi.org/10.23750/abm.v92iS6.12233>
- 348 24. Testa MA, Simonson DC. Chapter 12 - The Use of Questionnaires and Surveys. In *Clinical*
349 *and Translational Science (Second Edition)*; Robertson D, Williams GH. Eds.; Academic Press,
350 2017; 207–26. <https://doi.org/10.1016/B978-0-12-802101-9.00012-0>
- 351 25. Willett W. Future Directions in the Development of Food-Frequency Questionnaires. *Am J*
352 *Clin Nutr*. 1994; 59 (1): S171–4. <https://doi.org/10.1093/ajcn/59.1.171S>

353 26. Kowalkowska J, Slowinska M, Slowinski D, et al. Comparison of a Full Food-Frequency
354 Questionnaire with the Three-Day Unweighted Food Records in Young Polish Adult Women:
355 Implications for Dietary Assessment. *Nutrients*. 2013; 5 (7): 2747–76.
356 <https://doi.org/10.3390/nu5072747>

357 27. Martínez-González MA, Salas-Salvadó J, Estruch R, et al. Benefits of the Mediterranean
358 Diet: Insights From the PREDIMED Study. *Prog Cardiovasc Dis*. 2015; 58 (1): 50–60.
359 <https://doi.org/10.1016/j.pcad.2015.04.003>

360 28. Martinez-Gonzalez MA, Bes-Rastrollo M. Dietary Patterns, Mediterranean Diet, and
361 Cardiovascular Disease. *Curr Opin Lipidol*. 2014; 25 (1): 20–6.
362 <https://doi.org/10.1097/MOL.0000000000000044>

363 29. Jimenez-Torres J, Alcalá-Díaz JF, Torres-Peña JD, et al. Mediterranean Diet Reduces
364 Atherosclerosis Progression in Coronary Heart Disease: An Analysis of the CORDIOPREV
365 Randomized Controlled Trial. *Stroke*. 2021; 52 (11): 3440–9.
366 <https://doi.org/10.1161/STROKEAHA.120.033214>

367 30. Guasch-Ferré M, Willett WC. The Mediterranean Diet and Health: A Comprehensive
368 Overview. *J Intern Med*. 2021; 290 (3): 549–66. <https://doi.org/10.1111/joim.13333>

369 31. Ghisi GLDM, Mahajan A, Da Silva Chaves GS, et al. Validation of a Self-Administered
370 Version of the Mediterranean Diet Scale (MDS) for Cardiac Rehabilitation Patients in Canada.
371 *Int J Food Sci Nutr*. 2019; 70 (2): 202–11. <https://doi.org/10.1080/09637486.2018.1486392>

372 32. Elmusharaf K, Grafton D, Jung JS, et al. The Case for Investing in the Prevention and
373 Control of Non-Communicable Diseases in the Six Countries of the Gulf Cooperation Council:
374 An Economic Evaluation. *BMJ Global Health*. 2022; 7 (6): e008670.
375 <https://doi.org/10.1136/bmjgh-2022-008670>

33. World Heart Organization. WHO Global Health Observatory: Prevalence of obesity among adults, BMI \geq 30, age-standardized estimates by country. [internet]. 2016. Accessed 3 May 2023. Available from: <https://www.who.int/data/gho/data/themes/topics/indicator-groups/indicator-group-details/GHO/bmi-among-adults>

34. Musaiger AO. Diet and Prevention of Coronary Heart Disease in the Arab Middle East Countries. Med Princ Pract. 2002; 11 (Suppl. 2): 9–16. <https://doi.org/10.1159/000066415>

35. Al Moraie N, Lietz G, Seal CJ. Dietary Patterns and Risk of Heart Disease in Populations from Different Geographical Locations in Saudi Arabia. Proc Nutr Soc. 2012; 71 (OCE2): E50. <https://doi.org/10.1017/S0029665112001073>

36. Turk-Adawi KI, Terzic C, Bjarnason-Wehrens B, et al. Cardiac Rehabilitation in Canada and Arab Countries: Comparing Availability and Program Characteristics. BMC Health Serv Res. 2015; 15 (1): 521. <https://doi.org/10.1186/s12913-015-1183-7>

37. Wild D, Grove A, Martin M, et al. Principles of Good Practice for the Translation and Cultural Adaptation Process for Patient-Reported Outcomes (PRO) Measures: Report of the ISPOR Task Force for Translation and Cultural Adaptation. Value Health. 2005; 8 (2): 94–104. <https://doi.org/10.1111/j.1524-4733.2005.04054.x>

38. Mastaglia B, Toye C, Kristjanson LJ. Ensuring Content Validity in Instrument Development: Challenges and Innovative Approaches. Contemporary Nurse. 2003; 14 (3): 281–91. <https://doi.org/10.5172/conu.14.3.281>

39. Terwee CB, Bot SDM, De Boer MR, et al. Quality Criteria Were Proposed for Measurement Properties of Health Status Questionnaires. J Clin Epidemiol. 2007; 60 (1): 34–42. <https://doi.org/10.1016/j.jclinepi.2006.03.012>

- 398 40. BACPR. The BACPR Standards and Core Components for Cardiovascular Disease
399 Prevention and Rehabilitation [internet]. 2023. Accessed 3 May 2023. Available from:
400 [https://www.bacpr.org/_data/assets/pdf_file/0021/64236/BACPR-Standards-and-Core-](https://www.bacpr.org/_data/assets/pdf_file/0021/64236/BACPR-Standards-and-Core-Components-2023.pdf)
401 [Components-2023.pdf](https://www.bacpr.org/_data/assets/pdf_file/0021/64236/BACPR-Standards-and-Core-Components-2023.pdf)
- 402 41. Schröder H, Fitó M, Estruch R, et al. A Short Screener Is Valid for Assessing Mediterranean
403 Diet Adherence among Older Spanish Men and Women. Journal Nutr. 2011; 141 (6): 1140–5.
404 <https://doi.org/10.3945/jn.110.135566>
- 405 42. Teixeira R, Jansen AK, Pereira DA, et al. Brazilian Portuguese Version of the Mediterranean
406 Diet Scale: Translation Procedures and Measurement Properties. Diabetes Metabol Syndr: Clin
407 Res Rev. 2021; 15 (4): 102165. <https://doi.org/10.1016/j.dsx.2021.06.002>
- 408 43. Li J, Ding H, Wang Z, et al. Translation, Cultural Adaptation, Reliability, and Validity
409 Testing of a Chinese Version of the Self-Administered Mediterranean Diet Scale. Front Nutr.
410 2022; 9: 831109. <https://doi.org/10.3389/fnut.2022.831109>
- 411 44. Hair JF, Anderson RE. Multivariate data analysis. New Jersey: Prentice Hall, 1998.
- 412 45. Nunnally JC. Psychometric Theory. McGraw-Hill: New York, United States, 1978.
- 413 46. Widmer RJ, Flammer AJ, Lerman LO, et al. The Mediterranean Diet, Its Components, and
414 Cardiovascular Disease. Am J Med. 2015; 128 (3): 229–38.
415 <https://doi.org/10.1016/j.amjmed.2014.10.014>
- 416 47. Vanzella LM, Rouse V, Ajwani F, et al. Barriers and Facilitators to Participant Adherence of
417 Dietary Recommendations within Comprehensive Cardiac Rehabilitation Programmes: A
418 Systematic Review. Public Health Nutr. 2021; 24 (15): 4823–39.
419 <https://doi.org/10.1017/S1368980021002962>

420 48. Moradi-Lakeh M, El Bcheraoui C, Afshin A, et al. Diet in Saudi Arabia: Findings from a
 421 Nationally Representative Survey. *Public Health Nutr.* 2017; 20 (6): 1075–81.
 422 <https://doi.org/10.1017/S1368980016003141>

423 49. Ministry of Health. The Dietary Guidelines for Saudis. [internet] 2023. Accessed 3 May
 424 2023. Available from:
 425 <https://www.moh.gov.sa/en/HealthAwareness/Pages/SaudihealthFoodGuide.aspx>

426 50. Mayén AL, Marques-Vidal P, Paccaud F, et al. Socioeconomic Determinants of Dietary
 427 Patterns in Low- and Middle-Income Countries: A Systematic Review. *Am J Clin Nutr.* 2014;
 428 100 (6): 1520–31. <https://doi.org/10.3945/ajcn.114.089029>

429 51. D’Innocenzo S, Biagi C, Lanari M. Obesity and the Mediterranean Diet: A Review of
 430 Evidence of the Role and Sustainability of the Mediterranean Diet. *Nutrients.* 2019; 11 (6): 1306.
 431 <https://doi.org/10.3390/nu11061306>

432 52. Vlismas K, Stavrinou V, Panagiotakos DB. Socio-Economic Status, Dietary Habits and
 433 Health-Related Outcomes in Various Parts of the World: A Review. *Cent Eur J Public Health.*
 434 2009; 17 (2): 55–63. <https://doi.org/10.21101/cejph.a3475>

435 53. Cavaliere A, Elisa De Marchi A, Banterle A. Exploring the Adherence to the Mediterranean
 436 Diet and Its Relationship with Individual Lifestyle: The Role of Healthy Behaviors, Pro-
 437 Environmental Behaviors, Income, and Education. *Nutrients.* 2018; 10 (2): 141.
 438 <https://doi.org/10.3390/nu10020141>

439 54. Kocanda L, Schumacher TL, Plotnikoff RC, et al. Effectiveness and reporting of nutrition
 440 interventions in cardiac rehabilitation programmes: a systematic review. *Eur J Cardiovasc Nurs.*
 441 2023; 22(1): 1–12. <https://doi.org/10.1093/eurjcn/zvac033>

442

443 **Tables**

Table 1: Characteristics of patient participants and corresponding mean total MDS scores (N=200)

Characteristic		MDS Total Score mean±SD	p*
<i>Age</i> , mean±SD	46.5±13.6	-	-
<i>Sex</i> , n (%)			0.11
Male	92 (46.0)	8.3±2.3	
Female	108 (54.0)	7.7±2.4	
<i>Area of Living</i> , n (%)			0.51
Urban	180 (90.0)	7.9±2.4	
Rural	19 (9.5)	8.5±1.7	
<i>Highest educational level</i> , n (%)			0.91
Illiterate	9 (4.5)	7.7±2.7	
Less than High School	5 (2.5)	8.2±1.9	
High School	62 (31.0)	7.8±2.5	
College certificate or diploma	109 (54.5)	8.0±2.3	
Higher education	15 (7.5)	8.3±2.0	
<i>Monthly family income</i> , n (%)			0.02
Under SAR\$10,000	78 (39.0)	6.6±2.5	
Between SAR\$10,001-20,000	76 (38.0)	7.6±2.2	

Between SAR\$20,001-\$30,000	38 (19.0)	7.5±2.4	
Higher than SAR\$30,001	8 (4.0)	9.1±2.2	
<i>Cardiac Diagnosis/Procedures, n (%)</i>			
Percutaneous Coronary Intervention	24 (12.0)	8.1±2.2	0.72
Arrhythmia	22 (11.0)	7.4±2.6	0.25
Acute Coronary Syndrome	21 (10.5)	9.0±2.2	0.02
Acute Myocardial Infarction	16 (8.0)	8.0±2.3	0.90
Valve disorders	16 (8.0)	8.5±3.3	0.40
Cardiac Device	11 (5.5)	8.1±2.2	0.85
Valve Repair/Replacement	9 (4.5)	9.4±3.2	0.03
Heart Failure	8 (4.0)	7.4±1.6	0.50
Coronary Artery Bypass Graft	2 (1.0)	7.5±2.1	0.78
<i>Risk factors, n (%)</i>			
Family History of CVD	79 (39.5)	7.7±2.4	0.30
Sedentary/inactivity	72 (36.0)	7.8±2.5	0.50
Smoking	64 (32.0)	8.1±2.1	0.70
Diabetes Type II	60 (30.0)	8.3±2.3	0.23
Hypertension	46 (23.0)	7.8±2.4	0.52
Obesity	44 (22.0)	5.7±2.3	0.02
Depression	43 (21.5)	8.2±2.5	0.44

Dyslipidemia	34 (17.0)	6.5±2.3	0.03
<i>CR participation, yes (%)</i>			0.002
Yes	22 (11.0)	9.5±2.2	
No	178 (89.0)	7.6±2.3	

MDS, Mediterranean Diet Scale; SD standard deviation; CVD cardiovascular diseases; CR cardiac rehabilitation.

*p is used for association of characteristics and total mean score, tested using t-tests, analysis of variance or Pearson's correlation, as applicable.

Note: Income shown in Saudi Riyal (SAR). 1 SAR corresponds to USD\$0.47 (currency: March 1, 2023)

Table 2: Factor loadings from exploratory factor analysis, reliability of factors and number of ‘yes’ responses per item, N=202

Item	Factors				Yes, n (%)
	1. Meats and Processed Foods	2. Olive Oil and Sauces	3. Fruits, Vegetables, Legumes and Nuts	4. Fish and Seafood	
1. Do you use olive oil as the main source of fat when you cook?		0.721			123 (61.5)
2. Do you use at least four tablespoons or more of olive oil when cooking your food each day?		0.733			118 (59.0)
3. Do you eat two servings or more of vegetables each day?			0.689		100 (50.0)
4. Do you eat three servings or more of fruit each day?			0.629		101 (50.5)
5. Do you eat less than one tablespoon of butter, hydrogenated margarine or cream each day?	0.466				111 (55.5)
6. Do you drink less than one serving of sweet or sweetened drinks each day?	0.408				100 (50.0)
7. Do you eat three servings or more of legumes a week?			0.469		137 (68.5)
8. Do you eat three servings or more of fish or seafood each week?				0.474	87 (43.5)
9. Do you eat one serving or more of nuts each week?			0.453		129 (64.5)
10. Do you eat poultry (chicken or turkey) more often than meat (beef, veal, hamburger, or sausage)?	0.453				164 (82.0)
11. Do you limit red meat and processed meats to one serving or less one or two times a week?	0.452				132 (66.0)
12. Do you eat less than three servings of sweets or pastries each week?	0.405				119 (59.5)
13. Do you add some flavorings to your food, such as a mixture of tomatoes, garlic, onions and leeks, two or more times a week?		0.652			171 (85.5)

Variance explained	27.4%	16.8%	9.6%	8.5%	-
Eigenvalues	2.9	1.5	1.3	1.1	-
Reliability	0.79	0.78	0.75	0.70	-

* Item culturally adapted.