

Increasing HIV treatment literacy among people living with HIV using a novel health communication aid: Evidence from KwaZulu Natal province, South Africa

Authors: Caroline Govathson¹, Neo Ndlovu¹, Letitia Rambally-Greener², Laura Schmucker³, Candice Chetty-Makkan¹, Jacqui Miot¹, Harsha Thirumurthy³, Sophie Pascoe¹, Shawn Malone², Alison Bутtenheim⁴

Affiliations:

¹ Health Economics and Epidemiology Research Office, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

² Population Services International (PSI)

³ Department of Medical Ethics and Health Policy, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

⁴ Department of Family and Community Health, School of Nursing, University of Pennsylvania, Philadelphia, PA, USA

§ Corresponding author: Caroline Govathson; cgovathson@heroza.org

1 **Abstract**

2 **Introduction:** Effective health communication is important for promoting adherence to
3 antiretroviral therapy (ART). During counselling sessions with people living with HIV (PLHIV)
4 who are initiating or re-initiating ART, we assessed whether a simple visual aid using bead bottles
5 to explain the concept of viral suppression resulted in changes in HIV treatment literacy.
6

7 **Methods:** At three public sector clinics in KwaZulu-Natal (KZN) province, South Africa, we
8 enrolled adults who tested HIV-positive and were newly initiating ART or re-engaging in HIV
9 care. Trained HIV counsellors used bottles with coloured beads (“B-OK bottles”) to explain
10 concepts related to viral load, viral suppression, and undetectable=Untransmittable (U=U). We
11 assessed participants’ knowledge, attitudes, and perceptions about ART before and after
12 counselling.
13

14 **Results:** Between November 2022 and January 2023, we enrolled 80 PLHIV. Participants’ median
15 age was 32 years (IQR: 24–41) and 58% were male. After receiving counselling with the B-OK
16 bottles, understanding of U=U increased from 6% to 99% and understanding of ‘viral suppression’
17 increased from 20% to 99%. Confidence in the protective effects of ART increased (64% to 100%
18 for one’s own health; 58% to 94% for transmission to partners) and was observed among
19 participants both initiating ART and re-engaging in care. The number of participants agreeing that
20 viral suppression means their sexual partners are safe from HIV even without condoms increased
21 from 14% to 93% p-value =0.0. However, 65% still expressed worry that ART does not completely
22 eliminate the risk of sexually transmitting HIV.
23

24 **Conclusions:** Use of B-OK bottles during ART counselling was acceptable and increased HIV
25 treatment literacy. While there was no significant effect on level of confidence in complete
26 elimination of transmission risk when virally suppressed, we observed a significant reduction in
27 concern about transmitting HIV and an increase in confidence in the protective effects of ART.
28

29 Clinical Trial Number (SANCTR): DOH-27-092022-8067
30

31 **Introduction**

32 Despite scientific evidence about the therapeutic and prevention benefits of antiretroviral therapy
33 (ART), awareness of these benefits remains low in much of sub-Saharan Africa (SSA)(1,2). The
34 prevention benefits of ART in particular have been substantially underestimated in multiple
35 segments of the population, including people living with HIV (PLHIV) and healthcare providers
36 (3,4). Several studies conducted in South Africa, which has the world's largest HIV epidemic,
37 show that individuals have limited comprehension of the meaning of viral suppression and
38 substantially underestimate the effects of ART on viral load and transmission risk (5,6).

39
40 While increased treatment literacy has the potential to improve ART uptake and retention in care
41 among PLHIV, few evaluations have been conducted on interventions that disseminate
42 information about ART, including the message of "undetectable=untransmittable" (U=U). The
43 most commonly evaluated interventions, including motivational interviewing, health promotion
44 programs, group or individual recipient of care counselling, and active visualisation, have had
45 varying results (7,8,9).

46
47 Visual aids have the potential to convey health information in an accessible way to individuals
48 with low numeracy or literacy; this can be particularly effective for facilitating discussions of
49 complex numerical information, such as the percentage reduction in transmission risk due to ART.
50 Visual aids allow for the use of narrative, metaphor, and analogy in HIV treatment adherence
51 counselling (10,11) and can also help recalibrate flawed or outdated mental models about the
52 effects of ART on HIV viral loads and transmission risk (12,13).

53
54 The B-OK bead bottles (**Figure 1**) are a simple visual aid that is designed to assist healthcare
55 workers improve HIV treatment literacy among care recipients. Developed by Population Services
56 International (PSI) in collaboration with Matchboxology in South Africa in 2021, using a human-
57 centred design process, the bottles are a conversation starter for nurses and counsellors, assisting
58 them with communicating the importance of treatment adherence, the benefits of viral suppression,
59 and the "good news" of U=U and treatment as prevention. This study examines changes in
60 treatment literacy among PLHIV in South Africa who were introduced to the B-OK bottles during
61 ART initiation or re-engagement in care.

62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92

Methods

At three public sector clinics in KwaZulu-Natal, South Africa, we enrolled adults who tested HIV-positive and were newly initiating ART or were re-engaging in HIV care after a treatment interruption of at least 90 days. In addition to the standard post-test counselling that all clients receive at the clinics, trained HIV counsellors introduced the three bottles with coloured beads (B-OK bottles, Fig. 1) and used the bottles to explain the meaning of viral load, the effect of ART on viral suppression, and the reduction in transmission risk as a result of viral suppression (i.e., U=U).

Before and immediately after introducing the B-OK bottles, counsellors administered a brief survey that assessed participants' demographic characteristics and level of treatment literacy. Treatment literacy was measured using nine questions about the need for timely ART initiation, the effect of ART on viral load and the effect of viral suppression on transmission risk (Appendix 1). Participants' responses to the questions were assigned a score of 1 for correct responses, 0 for "not sure" responses, and -1 for incorrect responses. We then calculated a treatment literacy index score that was the sum of the scores from the 9 items, with a maximum value of 9. The survey also assessed participants' overall understanding of HIV treatment and their awareness of the concepts of "viral suppression" and "U=U".

We compared differences in treatment literacy index scores measured before and after exposure to the B-OK bottles using Wilcoxon sign-rank test. This statistical method was chosen due to the paired nature of the data and that the data was non-parametric as determined by a Shapiro-Wilk test. The study was approved by the research ethics committees at the University of the Witwatersrand # M220707 and University of Pennsylvania # 852087.



93

94 **Figure 1:** B-OK bead bottles

95

96 **Results**

97 Between November 14, 2022 and January 18, 2023, we screened 90 adult PLHIV at study clinics.
98 Ten PLHIV were excluded due to age, ill health or refusal to participate. Among the 80
99 participants who were enrolled, 23 were newly initiating ART and 55 were re-engaging in care
100 after a treatment interruption of at least 90 days, 2 preferred not to tell us whether they were newly
101 initiating ART or re-engaging with care after a treatment interruption. Median age was 32 years
102 (IQR: 24–41), 46 (58%) were male, 20 (25%) were either married or in a relationship, and 63
103 (79%) had completed primary school (**Table 1**).

104

105

106

107

108

109

110

111

112 **Table 1. Participant characteristics**

Variable	Categories	All participants	Newly initiating ART	Unknown ART status**	Re-engagement*
Age		32 (24.0-41.5)	27 (23.0-32.0)	36 (32.0-40.0)	33 (28-42)
Gender	Male	46 (57.5%)	8 (34.8%)	0 (0.0%)	36 (65.5%)
	Female	34 (42.5%)	15 (65.2%)	0 (0.0%)	19 (34.5%)
Race	African	79 (78.8%)	23 (29.1%)	1 (50.0%)	55 (69.6%)
	Indian	1 (1.3%)	0 (0.0%)	1 (50.0%)	0 (0.00%)
Relationship status	Single	60 (75.0%)	17 (73.9%)	1 (50.0%)	42 (76.4%)
	Relationship	17 (21.2%)	6 (26.1%)	0 (0.0%)	11 (20.0%)
	Married	3 (3.8%)	0 (0.0%)	1 (50.0%)	2 (3.6%)
Education completed	Primary school	17 (21.3%)	4 (17.4%)	0 (0.0%)	13 (23.6%)
	High school	61 (76.3%)	19 (82.6%)	1 (50.0%)	41 (74.6%)
	Diploma/degree	2 (2.5%)	0 (0.0%)	1 (50.0%)	1 (1.8%)
Employment	Employed	24 (30.0%)	9 (39.1%)	1 (50.0%)	14 (25.5%)
	Self employed	6 (7.5%)	1 (4.4%)	0 (0.0%)	5 (9.1%)
	In school	5 (6.3%)	1 (4.4%)	0 (0.0%)	4 (7.3%)
	Unemployed	45 (56.3%)	15 (52.2%)	1 (50.0%)	32 (58.2%)
Mobile phone	No	7 (8.8%)	3 (13.0%)	0 (0.0%)	4 (7.3%)
	Yes	73 (91.3%)	20 (87.0%)	2 (100.0%)	51 (92.7%)
Time since diagnosis	< 6 months	25 (31.3%)	22 (95.7%)	0 (0.0%)	3 (5.5%)
	>= 6 months	55 (68.8%)	1 (4.4%)	2 (100.0%)	52 (94.6%)


113 **Unknown ART status: was not determined whether they were new initiates or had been on ART before

114 *Re-engagement: re-engaging in care after a treatment interruption of at least 90 days

115

116 Exposure to the B-OK bottles led to a significant increase in treatment literacy among participants
 117 **(Figure 2)**. Prior to being introduced to the B-OK bottles, the median treatment literacy index
 118 score was 2 (IQR 0-3) out of 9. After being exposed to the B-OK bottles, treatment literacy was
 119 significantly higher (median score 7, IQR 5-7, $p < 0.001$). The largest increases in treatment literacy
 120 were observed for questions about the effect of ART on improving one's own health (16% to 95%)
 121 and on reducing transmission risk (14% to 94%). The only component of the literacy index that
 122 did not change significantly was knowledge about the need for immediate ART initiation following
 123 diagnosis (73% to 81%, $p = 0.13$). The increase in treatment literacy following exposure to the B-
 124 OK bottles was significant both among participants newly initiating ART and those re-engaging
 125 in care; post-test scores in new initiates were higher compared to post-test scores for those re-
 126 engaging with care although the median scores were the same (7: IQR 7-7 among new initiates vs

127 7: IQR 5 -7 among re-engaging; $p < 0.001$) (Figure 2). Prior to the B-OK bottle discussion, 70% of
128 participants agreed that they “worried that HIV treatment did not completely eliminate the risk of
129 transmission”, 14% disagreed, and 17% were not sure. After the bead bottle discussion, 65% of
130 participants still agreed with this statement, while the percentage of those who disagreed increased
131 to 33% and only 3% remained unsure. Exposure to the B-OK bottles also led to a significant
132 increase in whether participants had ever heard of the terms “viral suppression” and “U=U”, from
133 20.0% to 98.8% and 8.75% to 98.8%, respectively (**Table A1**)
134



135
136 **Figure 2.** HIV treatment literacy scores before and after B-OK bead bottle discussion, by treatment
137 status (re-engaging in care after a treatment interruption of at least 90 days or initiating treatment)
138

139 Discussion

140 Use of visual aids in the form of coloured bead bottles to provide education about HIV and ART
141 led to a large increase in treatment literacy among PLHIV at selected clinics in KwaZulu-Natal,
142 South Africa. Data collected prior to use of the B-OK bottles confirmed findings from other studies
143 (1,6) with participants having low treatment literacy, including limited awareness of the

144 therapeutic and prevention benefits of ART. In contrast, following the short, 10-minute
145 introduction to the B-OK bottles, care recipients reported much higher knowledge about the
146 benefits of ART.

147
148 This study contributes to a growing body of research on strategies for increasing treatment literacy
149 among PLHIV and the general population. While relatively few studies have been conducted in
150 Sub-Saharan Africa (8,9), the existing evidence shows that provision of information about U=U in
151 the form of flyers and interpersonal communication as well as community meetings were
152 successful in increasing HIV testing uptake and reducing stigma (3). To our knowledge, this is one
153 of the first studies to examine changes in treatment literacy through the use of simple visual aids
154 that explain the benefits of ART.

155
156 Several studies have highlighted the difficulties in conveying the U=U message to PLHIV in a
157 way that is easily understood (2,3,14). The B-OK bottles have the potential to remove some of
158 these challenges. These bottles use visual aids, which have long been recognised as an effective
159 way of communicating complex health information to, care recipients to explain concepts
160 involving HIV in the body and the way that ART can reduce viral load and transmission risk. The
161 bead bottles are also useful for explaining concepts involving probability. Enabling care recipients
162 to engage with the bead bottles may have also facilitated their absorption of key health information
163 relating to ART. Some advantages of the B-OK bottles include the low cost and ease with which
164 they can be made and the short amount of time needed to use the bottles to explain HIV, ART and
165 VLS. The bottles can also be used effectively by lay providers.

166
167 This study has several limitations. First, because there was no control group, changes in treatment
168 literacy following the use of B-OK bead bottles may be confounded by other factors. Additionally,
169 we measured the change in treatment literacy immediately after explaining the B-OK bead bottles,
170 which limits our ability to assess whether care recipients' increased treatment literacy was
171 sustained over time and whether this impacted treatment uptake and adherence.

172
173 **Conclusions**

174 This study shows that a simple visual aid that explains the effect of ART on HIV viral loads has
175 the potential to improve awareness of the therapeutic and prevention benefits of ART among
176 PLHIV. Future research should include rigorous, larger scale evaluations that determine the effect
177 of the B-OK bottles and similar visual aids on not only treatment literacy, but also health
178 behaviours and clinical outcomes among PLHIV. Implementation research that assesses strategies
179 for introducing and adapting the B-OK bottles for large-scale use should also be a priority.

180

181

182 **Competing interests**

183 The authors declare that they have no competing interests.

184

185 **Authors' contributions**

186 HT, SP, AB, CCM and SM conceptualized and designed the study. CG conducted the analysis.

187 CG prepared the original draft. All authors reviewed and edited the draft.

188 **Author information**

189 We would like to thank the all the participants and staff including data capturers and nurses at the
190 facilities we conducted the research for their support and contribution to the success of the study.

191

192 **Acknowledgements**

193 The authors gratefully acknowledge the contributions of the study participants and dedication of
194 the staff at the study sites.

195

196 **Funding** [Optional]:

197 This analysis was funded by the Bill & Melinda Gates Foundation (INV-008318). The contents
198 are the responsibility of the authors and do not necessarily reflect the views of the sponsor. The
199 funders had no role in the study design, collection, analysis and interpretation of the data, in
200 manuscript preparation or the decision to publish.

201

202 **Data Availability Statement** [Mandatory]

203 The dataset analysed for the manuscript is available upon reasonable request.

204 **Supporting Information** [Optional]

205 Supporting Information file 1:

1

2 **Table A1: Comparison of HIV Treatment Knowledge Pre and Post Use of BOK Bead Bottles**

3	Pre-test			Post-test			Exact Chi square
	Agree	Disagree	Not sure	Agree	Disagree	Not sure	P-value
Someone who tests positive for HIV does not need to start HIV treatment right away if they are still feeling fine.	17 (21.3%)	59 (73.8%)	4 (5.0%)	15 (18.8%)	65 (81.3%)	0 (0.0%)	0.13
If I am virally suppressed, my sexual partners are safe from HIV even if we don't use a condom.	11 (13.8%)	46 (57.5%)	23 (28.8%)	75 (93.8%)	5 (6.3%)	0 (0.0%)	<0.01
Even if I am virally suppressed, I worry that there is still some risk of giving HIV to a sexual partner.	53 (66.3%)	9 (11.3%)	18 (22.5%)	52 (65.0%)	27 (33.8%)	1 (1.3%)	<0.01
If someone with HIV is virally suppressed, the HIV is under control and cannot harm them.	13 (16.3%)	37 (46.3%)	30 (37.5%)	76 (95.0%)	(3.8%)	1 (1.3%)	<0.01
Someone with HIV should take treatment to protect their sexual partners from getting it.	51 (63.8%)	17 (21.3%)	12 (15.0%)	80 (100.0%)	0 (0.0%)	0 (0.0%)	<0.01
I worry that HIV treatment does not completely eliminate the risk of passing HIV through sex.	56 (70.0%)	11 (13.8%)	13 (16.3%)	52 (65.0%)	26 (32.5%)	3 (2.5%)	<0.01
Having HIV doesn't mean I will eventually die from it	69 (86.3%)	8 (10.0%)	3 (3.8%)	76 (95.0%)	4 (5.0%)	0 (0.0%)	0.08
Once I'm virally suppressed, it means the HIV is gone and I can stop taking treatment.	3 (3.8%)	69 (86.3%)	8 (10.0%)	3 (3.8%)	77 (96.3%)	0 (0.0%)	0.01
HIV treatment can get rid of almost all the HIV in the body.	5 (6.3%)	58 (72.5%)	17 (21.3%)	6 (7.5%)	73 (91.3%)	1 (1.3%)	<0.01
It's fine to miss your HIV treatment some days, as long as you're taking it most of the time.	13 (16.3%)	62 (77.5%)	5 (6.3%)	4 (5.0%)	76 (95.0%)	0 (0.0%)	<0.01

1 References

- 2 1. Raberahona M, Lidamahasolo Z, Andriamamonjisoa J, Andriananja V, Andrianasolo RL,
3 Rakotoarivelo RA, et al. Knowledge, attitudes, perception and practices regarding
4 antiretroviral therapy among HIV-infected adults in Antananarivo, Madagascar: A cross-
5 sectional survey. *BMC Health Serv Res* [Internet]. 2019 May 28 [cited 2023 Mar
6 29];19(1):1–9. Available from:
7 <https://bmchealthservres.biomedcentral.com/articles/10.1186/s12913-019-4173-3>
- 8 2. Gaps in U=U awareness globally: healthcare providers need to do more to get the message
9 out | *aidsmap* [Internet]. [cited 2023 Mar 29]. Available from:
10 [https://www.aidsmap.com/news/jul-2022/gaps-uu-awareness-globally-healthcare-](https://www.aidsmap.com/news/jul-2022/gaps-uu-awareness-globally-healthcare-providers-need-do-more-get-message-out)
11 [providers-need-do-more-get-message-out](https://www.aidsmap.com/news/jul-2022/gaps-uu-awareness-globally-healthcare-providers-need-do-more-get-message-out)
- 12 3. Bor J, Fischer C, Modi M, Richman · Bruce, Kinker · Cameron, King R, et al. Changing
13 Knowledge and Attitudes Towards HIV Treatment-as-Prevention and “Undetectable =
14 Untransmittable”: A Systematic Review. *AIDS Behav* [Internet]. 2021 [cited 2023 Mar
15 29];25:4209–24. Available from: <https://doi.org/10.1007/s10461-021-03296-8>
- 16 4. Ndyabakira A, Chamie G, Emperador D, Marson K, Kamya MR, Havlir D V., et al.
17 Men’s Beliefs About the Likelihood of Serodiscordance in Couples with an HIV-Positive
18 Partner: Survey Evidence from Rural Uganda. *AIDS Behav* [Internet]. 2020 Mar 1 [cited
19 2023 Mar 29];24(3):967–74. Available from:
20 <https://link.springer.com/article/10.1007/s10461-019-02531-7>
- 21 5. Mooney AC, Gottert A, Khoza N, Rebombo D, Hove J, Suárez AJ, et al. Men’s
22 Perceptions of Treatment as Prevention in South Africa: Implications for Engagement in
23 HIV Care and Treatment. <https://doi.org/10.1521/aeap2017293274> [Internet]. 2017 Jun 26
24 [cited 2023 Mar 29];29(3):274–87. Available from:
25 <https://guilfordjournals.com/doi/10.1521/aeap.2017.29.3.274>
- 26 6. Bor J, Musakwa N, Onoya D, Evans D. Perceived efficacy of HIV treatment-as-
27 prevention among university students in Johannesburg, South Africa. *Sex Transm Infect.*
28 2021 Dec 1;97(8):596–600.
- 29 7. Ridgeway K, Dulli LS, Murray KR, Silverstein H, Santo LD, Olsen P, et al. Interventions
30 to improve antiretroviral therapy adherence among adolescents in low- and middle-income
31 countries: A systematic review of the literature. *PLoS One* [Internet]. 2018 Jan 1 [cited
32 2023 Mar 29];13(1):e0189770. Available from:
33 <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0189770>

- 34 8. Damulak PP, Ismail S, Manaf RA, Said SM, Agbaji O. Interventions to improve
35 adherence to antiretroviral therapy (Art) in sub-saharan africa: An updated systematic
36 review. *Int J Environ Res Public Health*. 2021 Mar 1;18(5):1–18.
- 37 9. Jones ASK, Coetzee B, Kagee A, Fernandez J, Cleveland E, Thomas M, et al. The Use of
38 a Brief, Active Visualisation Intervention to Improve Adherence to Antiretroviral Therapy
39 in Non-adherent Patients in South Africa. *AIDS Behav* [Internet]. 2019 Aug 15 [cited
40 2023 Mar 29];23(8):2121–9. Available from:
41 <https://link.springer.com/article/10.1007/s10461-018-2292-1>
- 42 10. Newman S, Longworth DL, Stoller JK. Metaphorical medicine: using metaphors to
43 enhance communication with patients who have pulmonary disease. *Ann Intern Med*
44 [Internet]. 2002 Sep 3 [cited 2023 Apr 24];137(5 Part 1):376–9. Available from:
45 <https://pubmed.ncbi.nlm.nih.gov/12204044/>
- 46 11. Casarett D, Pickard A, Fishman JM, Alexander SC, Arnold RM, Pollak KI, et al. Can
47 metaphors and analogies improve communication with seriously ill patients? *J Palliat Med*
48 [Internet]. 2010 Mar 1 [cited 2023 Apr 24];13(3):255–60. Available from:
49 <https://pubmed.ncbi.nlm.nih.gov/19922170/>
- 50 12. Audet CM, Gutin SA, Blevins M, Chiau E, Alvim F, Jose E, et al. The Impact of Visual
51 Aids and Enhanced Training on the Delivery of Positive Health, Dignity, and Prevention
52 Messages to Adult Patients Living with HIV in Rural North Central Mozambique. *PLoS*
53 *One* [Internet]. 2015 Jul 6 [cited 2023 Apr 24];10(7). Available from:
54 </pmc/articles/PMC4493138/>
- 55 13. Mbanda N, Dada S, Bastable K, Ingalill GB, Ralf W. S. A scoping review of the use of
56 visual aids in health education materials for persons with low-literacy levels. *Patient Educ*
57 *Couns*. 2021 May 1;104(5):998–1017.
- 58 14. Grace D, Stewart M, Blaque E, Ryu H, Anand P, Gaspar M, et al. Challenges to
59 communicating the Undetectable equals Untransmittable (U=U) HIV prevention message:
60 Healthcare provider perspectives. *PLoS One* [Internet]. 2022 Jul 1 [cited 2023 Mar
61 29];17(7 July). Available from: [https://www.aidsmap.com/news/jul-2022/gaps-uu-](https://www.aidsmap.com/news/jul-2022/gaps-uu-awareness-globally-healthcare-providers-need-do-more-get-message-out)
62 [awareness-globally-healthcare-providers-need-do-more-get-message-out](https://www.aidsmap.com/news/jul-2022/gaps-uu-awareness-globally-healthcare-providers-need-do-more-get-message-out)

63

64