Association between child marriage and intimate partner violence.

A comparative analysis of Uganda Demographic Health Surveys

2011 and 2016

4 Authors

1

2

3

12

13

- 5 Theopista Fokukora^{1*}, Deda Ogum Alangea¹, Emefa Modey Amoah¹, Anthony Godi²
- 6 ¹Department of Population, Family and Reproductive Health, School of Public Health,
- 7 University of Ghana, P. O. Box, LG 13, Legon, Ghana
- ²Department of Biostatistics, School of Public Health, University of Ghana, P. O. Box, LG 13,
- 9 Legon, Ghana
- 10 Corresponding author
- * preciatheo@gmail.com _ ORCID: https://orcid.org/0000-0002-3323-2807

Abstract

- The incidence of child marriage (CM) and intimate partner violence (IPV) are high in Uganda.
- 15 The study sought to assess the association between child marriage and IPV across two waves of
- the Ugandan Demographic and Health Survey (UDHS). The evidence is important to inform
- 17 policy implementation strategies aimed at curbing child marriages and IPV.
- We analyzed data from UDHS 2011 and 2016. The analysis was restricted to women who have
- 19 ever cohabited. Simple and multiple logistic regression models were used to assess the
- association between child marriage and IPV.
- Following the 2011 and 2016 UDHS findings, child marriage was 54.7% in 2011 and 47.4% in
- 22 2016. Child marriage among the current adolescents increased from 9.4% in 2011 to 14.2% in
- 23 2016. According to UDHS 2011, 40.0% and 39.0% experienced physical and sexual violence
- respectively. In 2011 and 2016, 23.1% and 17.4% experienced both physical and sexual
- violence. There are higher odds of physical violence among child marriages compared to adult
- 26 marriages in 2011 and 2016 (AOR = 1.44; 95% CI: 1.12-1.84 in 2011 and AOR = 1.26; 95% CI:
- 27 1.12-1.42). A wealthy or educated woman has lower odds (AOR = 0.61; 95% CI: 0.41-0.91 and

AOR = 0.41; 95% CI: 0.25-0.66) to experience both physical violence and sexual violence than a

poor or uneducated one. Alcohol is the most common cause of IPV. A woman married to an

husband who consumes alcohol has higher odds of experiencing physical, sexual or both

physical and sexual violence as compared to someone whose husband did not consume alcohol

(AOR = 2.23; 95% CI:1.61-3.09 in 2011 and AOR = 2.42; 95% CI: 2.15-2.73 in 2016)

33 Intimate Partner Violence is associated with child marriage, level of education, wealth quintile,

residence, and partner's alcohol consumption. Much emphasis needs to be directed to cultural, as

well as social economic factors but more specifically on the contribution of women

emancipation/empowerment to prevent IPV.

Keywords

28

29

30

31

32

34

35

36

37

38

39

41

42

43

44

45

46

47

48

49

Child Marriage, Intimate Partner Violence, Uganda Demographic and Health survey

Introduction

40 Child marriage (CM) is one of the concerns of public health since below 18 years, the child's

physical, psychomotor, and cognitive developments are still developing. CM is common in both

developed and developing countries. Each year globally, 15 million marry below age 18 [1, 2].

Further, studies have found out that worldwide, 5% marry below 15, and 25% marry between 15-

17 years [3]. The projection is likely that up to 150 million will be married in the next decade

[1]. Recent research shows that 2.2 million adolescents marry in Europe, particularly in the

Central and Eastern part [4]. Child marriage is highly prevalent in sub-Saharan Africa. It is

evidenced that 55% of women in Africa are married before the age of 18 [5]. The recent study

carried out in Sub Saharan Africa found out that child marriage varies, it ranged from 13.5% in

Rwanda to 77% in Chad [1]. A round of DHS analysis from most countries in West Africa found

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

out that child marriage is 28% in Nigeria, 25% Burkina Faso, and 60% Niger [6]. In Ghana, 20% of girls marry by 18 years [7]. On the other hand, In Ethiopia, 40% of them were married before 18 years [8, 9]. Intimate Partner Violence (IPV) on the other hand is common in communities. Globally IPV increased from 1.1 million in 2010 to 1.4 in 2011[10]. Tia in his research found that 40% of the women who had faced IPV, only 7% had been reported [10, 11]. It is important to note that 15%-71% of the women worldwide experience a form of violence [11]. Importantly, 37% of the ever partnered partners face violence in Sub-Saharan Africa [12]. IPV is presented in different categories. Evidence has shown that physical violence accounts to 39%, psychological violence accounts to 45%, while almost a quarter (20%) face sexual violence in low income countries [13]. Systematic reviews indicate that IPV is high in Sub Saharan Africa, ranging from 43.4% in Nigeria to 76.5% in Ethiopia [14, 15]. Theoretical and empirical studies have evidenced that perpetuation of violence begins in the early childhood development [16]. Researchers have shown that CM and IPV are closely linked. Globally, 35% of young people below 18 years suffered IPV, though it's most prevalent in East Africa 45.1% [1, 17]. One out of every four (29%) of IPV happened to those married before 18 years [4]. The WHO multi-country survey about violence against women found out that IPV ranged from 19% to 66% among 15-24 years, but mostly among the adolescent marrieds than the adults [12]. Factors other than CM have been found to be associated with IPV. These include socialeconomic norms and beliefs, social-economic status, employment, income stress, poor communication, alcohol consumption of the husband[18], in-law violence and gender inequitable norms [19, 20].

Uganda's CM currently has been decreasing overtime, from 56% in 2006 [38], to 49% in 2010 [39] and then 43% in 2016 [21][38]. IPV according to UDHS 2006 increased from 48% to 56% in 2011, and then reduced to 39.6% in 2016 [18, 22]. The culture in Uganda as far as IPV is concerned, is profoundly embedded in disciplining the wives so that husbands can be seen as the superiors and the controllers [23]. More than half of men and nearly a quarter of the women had attitude supportive of wife beating [21]. Recent research still confirms that IPV is socialculturally justified by both the perpetuators and the sufferer [24]. Generally, CM and IPV usually end up in complications including but not limited to iniuries. depression, emotional distress often resulting in suicide [25-27]. Studies have shown that 38% of murdered women were a result of IPV [27, 28]. Sexual violence results in complications including death of child and/or mother [27], in addition to sexually transmitted diseases like HIV [29]. There is limited research on the association between CM and IPV in Uganda, which this study intends to find out. The findings of this research will be relevant for triggering policy implementation thereby improving the general health of the adolescent and thus achieving the SDG goal no 5.

Materials and Methods

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

Data from the Ugandan DHS (2011 and 2016) were used in this analysis. Using a 2-stage cluster sampling, the census enumeration areas were primary sampling units while households were secondary sampling units. The 2014 population and housing census included women aged 15-49 years, permanently residing in Uganda, and these were eligible to participate. The sample weights were derived from the domestic violence sub-sample which was nationally representative. The weighted sample of 3,461 and 6,701 women was selected for the study for

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

2011 and 2016 respectively. The analysis included only women who had ever cohabited and had responded to the domestic violence module in survey. Women who never answered intimate partner violence questions were excluded from the study. Ever married women and violence from an intimate partner were the variables of interest. Spousal violence was measured by modified Conflict Tactic Scale which was developed by Murray Straus in 1970s. This tool included specific questions on violence. Concerns about physical violence included; a) Push you, shake you or thrown something at you, b) Slap you, c) Twist your arm or pull your hair, d) Punch you with his fist or with something that could hurt you, e) Kick, drag, or beat you up, f)Try to choke you or burn you on purpose, g)Threatened to attack you with a knife, gun or any other weapon. A yes to one or more for the next 7 items constituted physical violence. Concerns about sexual violence included; a) Physically force you to have sexual intercourse with him even when you did not want to? b) Force you to perform any sexual acts you did not want to. A yes to one or all the two last statements constitutes sexual violence. Age was at first cohabitation was binary grouped (below 18 years as child marriage and above 18 as adult marriage). Current married children variable was also binary grouped (below 18 years as child and above 18 as an adult). This provided a better understanding of the association between child unions and intimate partner violence. The socio-demographic variables included the level of education, wealth quintile, residence, ethnicity, religion, and alcohol intake. **Data analysis**

Data was declared as a complex survey by applying sampling weights included in the dataset.

The demographic variables were analyzed using descriptive statistics to estimate the weighted

percentages. The patterns of association between type of IPV, age at first cohabitation/union and other socio-demographic variables were analyzed using the complex survey design adjusted Pearson Chi-square tests. These socio-demographic variables were also analyzed against the type of IPV to determine if there were any associations. The statistical significance was set at an alpha of 0.05 (5%). The data was analyzed using Stata/SE version 16 (StataCorp, College Station, Texas, USA).

Ethics approval and consent to participate.

A written request to use the 2011 and 2016 UDHS was submitted and approved by Measure DHS. The analysis was performed in accordance with relevant regulations and guidelines. Data used in this study was anonymized before use. Since the participants in this study were below 18 years old, written consents were sought from parents/guardians while emancipated minors signed their informed consent.

Results

The results indicated that CM was 54.7% in 2011 and 47.4% in 2016. CM among the current adolescents increased from 9.4% in 2011 to 14.2% in 2016. Most (59.4% and 57.4%) of the participants had attained primary level of education in 2011 and 2016 respectively. Majority (80.2% in 2011 and 73.3% in 2016) of the women come from rural settings. A quarter (25%) of the women belonged to the richest quartile. Uganda is predominantly a Christian country with majority of the women being Catholics and Protestants (table 1 below).

Table 1: Socio-demographic characteristics of women of childbearing age Uganda (UDHS 2011&2016)

	UDHS 2011	UDHS 2016	
Category	Weighted Percentage	Weighted	
8 v	(%)	Percentage (%)	
Age at cohabitation		8 (
Below 18 years	54.7	47.4	
18 years and above	45.3	52.6	
Current marital status among <18 years			
Ever married or cohabited.	9.4	14.2	
Never married or cohabited	90.6	85.8	
Level of education	70.0	02.0	
None	12.9	9.6	
Primary	59.4	57.4	
Secondary	22.5	25.1	
Higher	5.2	7.9	
Level of income		, , ,	
Poorest	17.5	17.5	
Poorer	18.2	18.4	
Middle	18.5	18.7	
Richer	19.9	19.9	
Richest	25.9	25.5	
Place of residence			
Urban	19.8	26.7	
Rural	80.2	73.3	
Region			
Kampala	9.7	5.5	
Central 1	11.0	13.5	
Central 2	10.4	10.6	
East Central	10.0	9.1	
Eastern	14.6	17.2	
North	8.5	10.5	
Karamoja	3.3	2.0	
West-nile	5.8	6.7	
Western	14.1	12.8	
Southwestern	12.7	12.1	
Religion			
Catholic	40.6	39.7	
Protestant	30.0	31.2	
Muslim	13.0	12.9	
Pentecostal	13.3	13.3	
SDA	1.9	1.7	
Others	1.2	12.4	
TOTAL	100	100	

Bivariate analysis (in table 2 below) indicated that 40.3% in 2011 and 39.0% in 2016 of the women experienced physical violence. A quarter (27.5% in 2011 and 22.5% in 2016) of the women experienced sexual violence, while 19.2% 16.0% of the women experienced both physical and sexual violence in 2011 and in 2016 respectively. Child marriage was significantly associated with IPV (p<0.05). Physical violence among child marriages was higher as compared

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

with adult marriages (45.4% vs 34.2% in 2011 and 44.2% vs 34.3% in 2016; p<0.05). Similarly, child marriage experienced a both physical and sexual violence as compared to adult marriages (23.1% vs 14.4% in 2011 and 17.4% vs 14.7% in 2016; p<0.01). For the section of currently married teenagers (below 18 years), a substantial proportion (19.1%) in 2011 and 25.1% in 2016, p<0.001) suffer physical violence. Exactly 17.0% in 2011 and 13.6% in 2016 suffer sexual violence (p<0.05). The proportion of those among this group who suffered both physical and sexual violence reduced from 13% in 2011 to 7.8% in 2016 (p<0.05). Almost half of women who experienced physical violence had no formal education (45.8% in 2011 to 47.2% in 2016). Experience of physical violence reduced with higher levels of education (p<0.05). Rural women experience twice as much physical violence more than the urban settings (p<0.001). Women from Eastern, North and Karamoja region experience more physical violence compared to those form other regions (p<0.001). While the physical violence reduced among women with non-alcoholic husbands (30.3% in 2011 to 28.3% in 2016), it increased among those whose spouses drunk alcohol (51.6% in 2011 to 53.2% in 2016; p<0.001). The results also indicated that sexual violence reduced among other levels of education except in no formal education category (p<0.01). The higher wealth quintile was observed to reduce sexual violence among the women (p<0.01). Rural women experience more sexual violence compared to women urban settings (p<0.01). Sexual violence increased among women from Eastern, North and Karamoja region compared to those form other regions (p<0.001). Whereas the sexual violence greatly reduced among women with non-alcoholic spouses (24.7% in 2011 to 17.9% in 2016 p<0.001), it slightly reduced among those with alcoholic husbands (30.8% in 2011 to 28.6% in 2016; p<0.001).

Taking both physical and sexual violence into account, the results indicated that higher wealth and education levels were associated with the lower risk of the combined type of violence (p<0.01). Furthermore, both physical and sexual violence more than doubled among the women whose husbands take alcohol as compared to those whose husbands were not alcoholics (25.1% vs 14.0% in 2011 and 22.8% vs 10.8% in 2016; p<0.0001).

Table 2: Bivariate analysis of the experience of IPV across different socio-demographic groups

			UDHS 2011 2011		UDHS 2016			
Variables		Physical violence % (x²)	Sexual violence % (x²)	Physical & Sexual Violence % (x ²)	Physical violence % (x²)	Sexual violence % (x²)	Physical and Sexua violence % (x²)	
Age at	18 and above	34.2 (22.27) ***	22.9 (15.31) *	14.4 (20.56) ***	34.3 (77.89) ***	22.2 (0.60)	14.7 (9.67) **	
cohabitation (years)	Below 18	45.4	31.3	23.1	44.2	22.9	17.4	
Current age	18 and above	40.8 (6.99)	27.6 (2.26)	19.3 (0.92)	39.3 (11.07) **	22.7 (6.12) *	16.1 (6.82) *	
(years)	Below 18	19.1	17.0	13.0	25.1	13.6	7.8	
Level of		45.8 (49.09) *	22.1 (21.52) *	19.1 (17.69) **	47.2 (221.56) ***	21.3 (73.38) ***	17.8 (66.70) **	
ducation	Primary	44.2	31.1	21.8	43.1	25.1	18.0	
	Secondary	29.3	23.5	13.0	29.5	20.0	12.4	
	Post-secondary	13.8	14.9	8.9	17.5	9.8	6.4	
Level of	Poorest	53.5 (80.58) *	31.4 (20.11) *	27.4 (32.16) ***	50.2 (217.94) ***	23.2(59.58) ***	18.8 (53.01) ***	
ncome	Poorer	48.5	28.5	21.8	43.8	25.9	18.8	
	Middle	41.5	31.0	20.1	39.6	24.6	15.8	
	Richer	35.2	25.6	15.8	36.2	23.9	16.3	
	Richest	24.5	22.0	11.7	25.8	15.6	10.6	
Place of	Urban	29.9 (15.61) **	24.2 (2.00)	13.2 (7.61) *	32.0 (49.31) ***	18.9 (18.56) **	13.8 (8.79) **	
residence	Rural	42.4	28.2	20.3	41.2	23.7	16.7	
Region	Kampala	28.8 (87.63) *	21.5 (45.93)	10.9 (27.07) *	24.8 (169.62) ***	14.4 (264.24) ***	10.1 (155.65) ***	
O	Central	28.4	25.1	15.7	32.7	19.4	14.4	
	Central	30.6	33.6	16.4	31.0	18.0	11.3	
	East Central	35.1	32.8	18.1	31.3	24.6	14.8	
	Eastern	57.2	34.4	28.8	49.3	44.8	16.1	
	North	55.3	27.8	22.3	41.7	22.6	18.1	
	Karamoja	36.0	16.4	15.6	41.1	16.9	11.9	
	West-Nile	46.3	27.7	23.9	45.4	21.7	17.5	
	Western	40.5	25.4	18.1	37.5	30.3	18.8	
	Southwestern	37.9	20.6	16.9	43.2	26.1	19.5	
Religion	Catholic	44.2 (17.35)	27.4 (8.54)	21.1 (8.02)	43.5 (70.72) *	22.1 (5.03)	16.6 (11.40)	
8	Protestant	36.6	25.9	16.6	38.7	23.4	16.7	
	Muslim	33.4	31.7	18.1	31.1	21.6	13.4	
	Pentecostal	44.7	28.4	20.3	36.4	23.5	16.1	
	SDA	35.7	13.9	10.6	22.0	20.1	8.8	
	Others	41.1	32.8	32.8	31.2	16.0	13.0	
Husband'	No	30.3 (301.53) ***	24.7 (214.55) ***	14.0 (33.73) ***	28.2 (484.37) ***	17.9 (119.95) ***	10.8 (198.70) ***	
takes Alcohol	Yes	51.6	30.8	25.1	53.2	28.6	22.8	
Overall Percent	2000 00000	40.3	27.5	19.2	39.0	22.5	16.0	

*, **, *** corresponds with p<0.05, p<0.01, and p<0.001 respectively

Table 3: Logistic regression models showing adjusted OR between demographic characteristics and Different types of violence in Uganda (UDHS 2011 & 2016)

UDHS 2010 UDHS 2016

	UDIIS 2010			UDIIS 2010			
	UDHS 2011			UDHS 2016			
Variables	Physical Violence	Sexual violence	Physical and Sexual Violence	Physical violence	Sexual violence	Physical and Sexual Violence	
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	
Age at cohabitation (years)							
18 and above	1(ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	
Below 18	1.44 (1.12-1.84) *	1.43 (1.11-1.85)	1.64 (1.22-2.21) **	1.26 (1.12-1.42) ***	0.91 (0.79-1.04)	1.04 (0.89-1.21)	
Current age (years)							
18 and above	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	
Below 18	0.23 (0.08-0.63)	0.41 (0.41-1.18)	0.48 (0.14-1.58)	0.45 (0.28-0.71) **	0.54 (0.31-0.92)	0.43 (0.23-0.79) *	
Level of education							
None	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	
Primary	1.12 (0.81-1.56)	1.76 (1.18-2.63)	1.44 (0.92-2.25)	0.93 (0.79-1.10)	1.24 (1.03-1.50) *	1.03 (0.84-1.26)	
Secondary	1.02 (0.64-1.64)	1.60 (0.94-2.71)	1.30 (0.67-2.51)	0.71 (0.57-0.88) **	1.04 (0.82-1.32)	0.80 (0.60-1.06)	
Post-Secondary	0.42 (0.17-1.06)	1.02 (0.44-2.45)	0.98 (0.33-2.96)	0.37 (0.27-0.52) ***	0.49 (0.32-0.75) **	0.41 (0.25-0.66) ***	
Level of income							
Poorest	1(ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	
Poorer	0.82 (0.56-1.20)	0.76 (0.52-1.13)	0.71 (0.46-1.12)	0.88 (0.73-1.05)	0.95 (0.77-1.17)	0.89 (0.76-1.11)	
Middle	0.78 (0.52-1.19)	0.88 (0.57-1.34)	0.72 (0.44-1.17)	0.79 (0.65-0.95) *	0.88 (0.71-1.09)	0.73 (0.57-0.92) *	
Richer	0.62 (0.39-0.97) *	0.65 (0.40-1.06)	0.55 (0.31-0.97) *	0.76 (0.61-0.94) *	0.89 (0.71-1.12)	0.80 (0.62-1.04)	
Richest	0.46 (0.25-0.83) *	0.59 (0.32-1.11)	0.49 (0.23-1.03)	0.61 (0.47-0.80) ***	0.67 (0.48-0.95)	0.61 (0.41-0.91) *	
Place of residence							
Urban	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	
Rural	0.93 (0.57-1.52)	0.86 (0.50-1.46)	0.93 (0.50-1.70)	0.93 (1.76-1.15)	0.98 (0.78-1.24)	0.87 (0.67-1.13)	
4	T C C C C C C C C C C C C C C C C C C C			The state of the s			

Region						
Kampala	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)
Central	0.69 (0.30-1.56)	1.18 (0.52-2.68)	1.30 (0.50-3.38)	1.20 (0.81-1.75)	1.19 (0.72-1.95)	1.28 (0.75-2.17)
Central	0.72 (0.32-1.65)	1.67 (0.81-3.47)	1.28 (0.50-3.32)	1.00 (0.67-1.51)	0.99 (0.58 - 1.67)	0.89 (0.51-1.54)
East Central	0.97 (0.42-2.24)	1.57 (0.69-3.56)	1.45 (0.61-3.45)	1.05 (0.70-1.58)	1.47 (0.81-2.65)	1.26 (0.69-2.30)
Eastern	1.95 (0.88-4.30)	1.62 (0.72-3.63)	2.18 (0.90-5.26)	1.89 (1.31-2.73) **	3.39 (2.01-5.71) ***	2.85 (1.70-4.76) ***
North	1.30 (0.56-2.99)	1.02 (0.46-2.25)	1.16 (0.47-2.84)	1.12 (0.76-1.65)	0.34 (0.19-0.60) ***	0.35 (0.19-0.64) **
Karamoja	0.54 (0.21-1.27)	0.59 (0.22-1.61)	0.78 (0.24-2.49)	0.67 (0.41-1.08)	0.62 (0.33-1.17)	0.47 (0.24-0.92) *
West-Nile	1.09 (0.47-2.55)	1.14 (0.50-2.62)	1.53 (0.60-3.86)	1.27 (0.87-1.85)	1.00 (0.58-1.71)	1.07 (0.62-1.85)
Western	1.07 (0.43-2.68)	1.16 (0.52-2.58)	1.39 (0.60-3.22)	1.44 (1.00-2.10) *	1.69 (1.01-2.82) *	1.46 (1.87-2.46)
Southwestern	0.87 (0.39-1.97)	0.83 (0.36-1.91)	1.23 (0.50-3.04)	1.30 (0.89-1.90)	1.28 (0.77-2.14)	1.31 (0.77-2.24)
Religion						
Catholic	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)
Protestant	0.75 (0.55-1.01)	0.92 (0.67-1.25)	0.79 (0.56-1.11)	1.11 (0.96-1.27)	0.97 (0.83-1.13)	0.93 (0.82-1.04)
Muslim	1.01 (0.66-1.55)	1.30 (0.82-2.05)	1.22 (0.69-2.15)	1.13 (0.92-1.38)	1.20 (0.94-1.53)	0.77 (0.64-0.94)
Pentecostal	1.37 (0.93-2.03)	1.13 (0.77-1.66)	1.23 (0.78-1.94)	0.74 (0.42-1.28)	1.19 (0.94-1.49)	0.88 (0.75-1.03)
SDA	1.21 (0.41-3.53)	0.50 (0.16-1.56)	0.70 (0.21-2.36)	1.15 (0.92-1.42)	1.16 (0.63-2.16)	0.58 (0.37-0.91)
Others	0.96 (0.33-2.82)	1.29 (0.41-4.07)	2.11 (0.65-6.87)	0.88 (0.52-1.48)	0.75 (0.39-1.45)	0.49 (0.25-0.96)
Husband takes alcohol						
No	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)
Yes	2.48 (1.83-3.35) ***	1.63 (1.23-2.16) **	2.23 (1.61-3.09) **	2.67 (2.35-3.02) ***	2.04 (1.77-2.36) ***	2.42 (2.15-2.73) **

^{*, **, ***} corresponds with p<0.05, p<0.01, and p<0.001 respectively

CM recorded significantly higher odds of physical violence as compared to adult marriages in both 2011 (AOR = 1.44; 95% CI: 1.12-1.84) and 2016 (AOR = 1.26; 95% CI: 1.12-1.42). The odds of experiencing physical violence reduces with increasing level of education and wealth index. An educated woman had lower odds of experiencing physical violence as compared to uneducated woman in 2016 as compared to 2011 (AOR = 0.42; 95% CI: 0.17-1.06 in 2011 to AOR = 0.37; 95% CI: 0.27-0.52; p<0.05 in 2016). Conversely, the odds of experiencing physical violence among wealthy women were lower compared to a poor woman, however, there was an increased in IPV among wealthy women in 2016 when compared to 2011 (AOR = 0.46; 95% CI: 0.25-0.83 in 2011 to AOR = 0.61; 95% CI: 0.47-0.80; p<0.05 in 2016). The women in the Eastern regions had higher odds of experiencing both physical

and sexual violence as compared to other regions (AOR=2.85; 95% CI:1.70-4.76). Women whose husbands drink alcohol had higher odds of experiencing both physical and sexual violence as compared to those of a woman whose husbands were non-alcoholics (AOR = 2.23; 95% CI:1.61-3.09 in 2011 and AOR = 2.42; 95% CI: 2.15-2.73 in 2016) (table 3).

Discussion

The findings from this analysis indicated that CM reduced from 54.7% in 2011 to 47.4% in 2016. It is true and evident that CM still exist in Uganda. The Uganda Demographic and Health Survey (UDHS) findings indicate that CM currently has been decreasing overtime, from 56% in 2006 [22]. Uganda also contributes to the overall global findings whereby, 15 million marry below age 18 [1, 3]. As reflected, Sub-Saharan Africa has the highest proportion [3, 5, 30]. Evidence is derived from UNFPA and UNICEF database where, East and Southern Africa had 36% of child marriage [1]. The rate of child marriage in Uganda's is much higher than that of Ghana, where 1 in every 5 girls marry by 18 years [10]. Recent research reveals that child marriage in Rwanda was lower (13.5%), while Chad's proportion was high at 77% compared to Uganda's situation [31].

The findings from this study further indicated that both physical and sexual violence reduced among child marriages from 23.1% in 2011 to 17.4% in 2016. This could have been possible because of human rights implementation programs about adolescents. Elsewhere, similar results revealed that IPV ranges from 15% -71% in different countries, but most prevalent among CM [10, 11]. In agreement with the results from this study, different kinds of violence include physical violence, psychological/emotional violence, and sexual violence accounting 39%, 45% and 20% respectively [13, 17], of which married children are most vulnerable. There is a high prevalence (76.5% and 43.4%) of IPV in Ethiopia and Nigeria respectively [18–20], which is proportional to child marriage in those countries. Recent estimates show that IPV ranged from 17.5% in Mozambique to 42% in Uganda [31]. Though such combined violence reduced among the CM, more strategies targeting communities still need to be employed to eliminate such suffering among this vulnerable population.

This study found that the odds of experiencing IPV among married children was higher as compared to married adults. The communities in Uganda discipline children because of the belief that they are not yet mature to make their own decisions. In this respect, husbands to these children use threats and beatings to justify their dominance over them. The results agree with a multi-country analysis of SSA DHS data [31]. Peterman and his colleagues (2015) while identifying the global trends, found that 1/3 of the women who experienced physical or sexual abuse were initiated to sex at a teenage age [15]. At such a tender age, the adolescents do not reveal their grievances because they can't fight for their rights. Evidence has been found in a comparative study of 34 countries, where 29% of IPV happened to those married before 18, but 15-17 years were more at risk [3]. The results are in agreement with systematic reviews carried out in most low and middle income countries, revealing that 19% to 66% among 15-24 years were at a high risk of IPV as compared to adults [12]. Rural communities especially in Uganda need to be sensitized about helping such children to mature into responsible adults without necessarily inflicting pain to them.

The findings indicated that the level of education was associated to IPV. The higher level of education exposes the women to understanding their rights and gives them empowerment. It is likely that the individual who goes through higher levels of education, and attains employment, sustains her economic income, which reduces violence in the households. This is not the case with married children. DHS 2016 shows that women who had a low level of education were more prone to IPV [26, 32]. The results further agree with findings from a meta-analysis in Sub-Saharan Africa, which identified that the women who were not educated were more likely to face IPV [33]. Jesmin (YEAR??) identified however, that lower education was not necessarily associated with IPV [34].

Apart from education, but closely related, the level of income plays a vital role in women's risk for IPV. Our findings indicated that the odds of IPV reduced among wealthier as compared to poor women in both 2011 and 2016. The married children are dependent on their husbands, and because of the economic breakdown especially among the rural men, this induces wife beating and other forms of violence to them. This is supported by evidence from Kenya, Zimbabwe, Nigeria [14, 35, 36], and Malawi [37]. However, other research have found that high social economic class and women empowerment, if not well managed, results in misunderstandings in the couples ending up in IPV [19, 20].

Apart from the socio-economic situation, the study indicated that residence was significantly associated with IPV and CM. Most of the children who were married were predominantly from rural settings where culturally wife beating is justified as a disciplinary behaviour. Findings indicated that rural women experienced twice as much physical violence than those in urban settings. In villages and humanitarian settings especially Somalia and South Sudan which still adore and embrace culture, IPV among such rural communities was most prevalent there [38]. This could also be related to religiosity's fear to divorce as they would be seen as failures in churches and communities. Evidence from diverse researchers found that individual, relationship, community and societal factors were all associated with IPV [13, 39]. A recent study in Ghana identified that IPV is justified by beating, dominance, and social cultural justification [40]. The drivers to IPV were poverty, low education levels, and limited implementation of policy frameworks to promote human rights [21, 41]. These are similar scenarios identified in Uganda.

Partner alcohol intake was significantly associated with IPV in this study. This finding supports earlier findings in Uganda. A drunkard is believed to have less insight of the outcome of

violence, and, men drink to supplement their status of superiority, manhood and dominance [18,

42]. The results agree with findings in Nepal - that alcohol consumption by the husband

predisposed a partner to IPV [43]. The consistent drinking of alcohol is often associated with

poverty. Alcohol intake further predisposes to sexual violence and uncontrolled birth rates

resulting in high fertility rate.

Some limitations were identified in this study. The data was retrospective in nature and thus

subject to some level of recall bias. Further, the women who report violence are often socially

excluded for exposing the family affairs to the public. This could have introduced the desirability

bias because of its sensitivity. Further, data used for this study was quantitatively collected,

therefore some deeper expressions and experiences of women who face violence could have

been missed. The qualitative approach could also reveal household and community factors.

Conclusion and implication

The study found out that child marriage and IPV are prevalent in Uganda. There was a

significant association between CM and IPV. Married children are characterized by low level of

education, poor wealth quintile, and dependability. All these were significantly associated with

IPV. Because IPV among adolescents has detrimental complications, policy makers and

implementors need to put into consideration children rights to protect those who get married

from facing this violence in developing countries like Uganda. The implementation of these

rights needs to begin from the grassroot level (communities) up to the national and international

agencies with an aim to improve children's health and strengthen the awareness of children's

rights in the communities.

Acknowledgments

I acknowledge the support by the HRP Alliance, part of the UNDP-UNFPA-WHO-World Bank

Special programme of Research Development and Research training in Human Reproduction

(HRP), a co-sponsored programme executed by the World Health Organization (WHO) who

facilitated my PhD studies. The University of Ghana, School of Public Health for their academic

support in the manuscript development and submission process.

Authors' contributions

Conceptualization: Theopista Fokukora, Deda Alangea, Emefa Modey Amoah

Formal analysis: Theopista Fokukora, Anthony Godi

Methodology: Theopista Fokukora

Supervision: Deda Alangea, Emefa Modey Amoah, Anthony Godi

Writing – original draft: Theopista Fokukora

Writing – review & editing: Deda Alangea, Emefa Moday, Anthony Godi

Declarations

Consent for publication: Not applicable

Availability of data and materials: The data is available on request at

http://www.dhsprogram.com/data/dataset. The authors confirm that they followed general rules

of requesting for the data from the above website, therefore did not have any special access or

privileges to the data.

Competing interests: The authors declare that they have no competing interests.

Funding: The author did not receive any funding to develop this manuscript.

Abbreviations

CM Child marriage

IPV Intimate Partner Violence

OR Odds Ratio

UDHS Uganda Demographic and Health Survey

WHO World Health Organization

References

1. UNICEF. Global programme to accelerate action to end child marriage. 2019.

- 2. Petroni S, Steinhaus M, Fenn NS, Stoebenau K, Gregowski A. New findings on child marriage in sub-Saharan Africa. Annals of global health. 2017;83(5-6):781-90.
- 3. Kidman R. Child marriage and intimate partner violence: a comparative study of 34 countries. International journal of epidemiology. 2017;46(2):662-75.
- 4. Sibanda M. Married too soon: Child marriage in Zimbabwe. The Research Advocate Unit. 2011:1-22.
- 5. Efevbera Y, Bhabha J, Farmer P, Fink G. Girl child marriage, socioeconomic status, and undernutrition: evidence from 35 countries in Sub-Saharan Africa. BMC medicine. 2019;17(1):1-12.
- 6. Avogo WA, Somefun OD. Early marriage, cohabitation, and childbearing in West Africa. Journal of environmental and public health. 2019;2019.
- 7. de Groot R, Kuunyem MY, Palermo T. Child marriage and associated outcomes in northern Ghana: a cross-sectional study. BMC public health. 2018;18(1):1-12.
- 8. Mekonnen Y, Telake DS, Wolde E. Adolescent childbearing trends and sub-national variations in Ethiopia: a pooled analysis of data from six surveys. BMC pregnancy and childbirth. 2018;18(1):1-13.
- 9. Erulkar A. Early marriage, marital relations and intimate partner violence in Ethiopia. International perspectives on sexual and reproductive health. 2013:6-13.
- 10. Palermo T, Bleck J, Peterman A. Tip of the iceberg: reporting and gender-based violence in developing countries. American journal of epidemiology. 2014;179(5):602-12.
- 11. Devries KM, Knight L, Child JC, Kyegombe N, Hossain M, Lees S, et al. Witnessing intimate partner violence and child maltreatment in Ugandan children: a cross-sectional survey. BMJ open. 2017;7(2):e013583.
- 12. Stöckl H, March L, Pallitto C, Garcia-Moreno C. Intimate partner violence among adolescents and young women: prevalence and associated factors in nine countries: a cross-sectional study. BMC public health. 2014;14(1):1-14.
- 13. Semahegn A, Torpey K, Manu A, Assefa N, Tesfaye G, Ankomah A. Are interventions focused on gender-norms effective in preventing domestic violence against women in low and lower-middle income countries? A systematic review and meta-analysis. Reproductive health. 2019;16(1):1-31.

- 14. Bamiwuye SO, Odimegwu C. Spousal violence in sub-Saharan Africa: does household poverty-wealth matter? Reproductive health. 2014;11(1):1-10.
- 15. Peterman A, Bleck J, Palermo T. Age and intimate partner violence: an analysis of global trends among women experiencing victimization in 30 developing countries. Journal of Adolescent Health. 2015;57(6):624-30.
- 16. Herrenkohl TI, Fedina L, Roberto KA, Raquet KL, Hu RX, Rousson AN, et al. Child maltreatment, youth violence, intimate partner violence, and elder mistreatment: A review and theoretical analysis of research on violence across the life course. Trauma, Violence, & Abuse. 2022;23(1):314-28.
- 17. Memiah P, Cook C, Kingori C, Munala L, Howard K, Ayivor S, et al. Correlates of intimate partner violence among adolescents in East Africa: a multi-country analysis. The Pan African Medical Journal. 2021;40.
- 18. Tumwesigye NM, Kyomuhendo GB, Greenfield TK, Wanyenze RK. Problem drinking and physical intimate partner violence against women: evidence from a national survey in Uganda. BMC public health. 2012;12(1):1-11.
- 19. Peterman A, Pereira A, Bleck J, Palermo TM, Yount KM. Women's individual asset ownership and experience of intimate partner violence: evidence from 28 international surveys. American journal of public health. 2017;107(5):747-55.
- 20. Jewkes R, Flood M, Lang J. From work with men and boys to changes of social norms and reduction of inequities in gender relations: a conceptual shift in prevention of violence against women and girls. The Lancet. 2015;385(9977):1580-9.
- 21. Wandera SO, Kwagala B, Ndugga P, Kabagenyi A. Partners' controlling behaviors and intimate partner sexual violence among married women in Uganda. BMC Public Health. 2015;15(1):1-9.
- 22. Kawuki J, Sserwanja Q, Mukunya D, Sepenu AS, Musaba MW. Prevalence and factors associated with sexual violence among women aged 15–49 years in rural Uganda: evidence from the Uganda Demographic and Health Survey 2016. Public health. 2021;196:35-42.
- 23. Gardsbane D. Violence against Women Policy and Practice in Uganda-Promoting Justice within the Context of Patriarchy 2016.
- 24. Pöllänen K, de Vries H, Mathews C, Schneider F, de Vries PJ. Beliefs about sexual intimate partner violence perpetration among adolescents in South Africa. Journal of interpersonal violence. 2021;36(3-4):NP2056-78NP.
- 25. John NA, Edmeades J, Murithi L. Child marriage and psychological well-being in Niger and Ethiopia. BMC public health. 2019;19(1):1-12.
- 26. Amegbor PM, Pascoe L. Variations in emotional, sexual, and physical intimate partner violence among women in Uganda: a multilevel analysis. Journal of interpersonal violence. 2021;36(15-16):NP7868-NP98.
- 27. Adedokun O, Adeyemi O, Dauda C. Child marriage and maternal health risks among young mothers in Gombi, Adamawa State, Nigeria: implications for mortality, entitlements and freedoms. African health sciences. 2016;16(4):986-99.
- 28. WHO. Global Strategy on Women's, Children's and Adolescents' Health (2016-2030). 2016.
- 29. Irani M, Latifnejad Roudsari R. Reproductive and sexual health consequences of child marriage: A review of literature. Journal of Midwifery and Reproductive Health. 2019;7(1):1584-90.
- 30. Rumble L, Peterman A, Irdiana N, Triyana M, Minnick E. An empirical exploration of female child marriage determinants in Indonesia. BMC public health. 2018;18(1):1-13.
- 31. Ahinkorah BO, Onayemi OM, Seidu A-A, Awopegba OE, Ajayi AI. Association between girl-child marriage and intimate partner violence in Sub-Saharan Africa: insights from a multicountry analysis of demographic and health surveys. Journal of interpersonal violence. 2022;37(15-16):NP13560-NP80.
- 32. UDHS. Uganda Demographic and

Health Survey 2016. 2016.

- 33. Nabaggala MS, Reddy T, Manda S. Effects of rural—urban residence and education on intimate partner violence among women in Sub-Saharan Africa: a meta-analysis of health survey data. BMC women's health. 2021;21(1):1-23.
- 34. Jesmin SS. Social determinants of married women's attitudinal acceptance of intimate partner violence. Journal of Interpersonal Violence. 2017;32(21):3226-44.
- 35. Reed E, Donta B, Dasgupta A, Ghule M, Battala M, Nair S, et al. Household debt and relation to intimate partner violence and Husbands' attitudes toward gender norms: A study among young married couples in rural Maharashtra, India. Public Health Reports. 2015;130(6):664-71.
- 36. Ibrahim M, Bashir S, Umar A, Gobir A, Idris S. Men's perspectives on intimate partner abuse in an urban community in North-Western Nigeria. Annals of Nigerian Medicine. 2014;8(1):37-.
- 37. Conroy AA. Gender, power, and intimate partner violence: a study on couples from rural Malawi. Journal of interpersonal violence. 2014;29(5):866-88.
- 38. Perrin N, Marsh M, Clough A, Desgroppes A, Yope Phanuel C, Abdi A, et al. Social norms and beliefs about gender based violence scale: a measure for use with gender based violence prevention programs in low-resource and humanitarian settings. Conflict and health. 2019;13(1):1-12.
- 39. Simister J, Kowalewska G. Gender-Based Violence and Christianity: Catholic prevention of divorce traps women in an abusive marriage. Psychology. 2016;7(13):1624-44.
- 40. Dery I. "Give her a slap or two... she might change": Negotiating masculinities through intimate partner violence among rural Ghanaian men. Journal of interpersonal violence. 2021;36(19-20):9670-90.
- 41. Psaki SR, Melnikas AJ, Haque E, Saul G, Misunas C, Patel SK, et al. What are the drivers of child marriage? A conceptual framework to guide policies and programs. Journal of Adolescent Health. 2021;69(6):S13-S22.
- 42. Zegenhagen S, Ranganathan M, Buller AM. Household decision-making and its association with intimate partner violence: Examining differences in men's and women's perceptions in Uganda. SSM-population health. 2019;8:100442.
- 43. Clark CJ, Ferguson G, Shrestha B, Shrestha PN, Batayeh B, Bergenfeld I, et al. Mixed methods assessment of women's risk of intimate partner violence in Nepal. BMC women's health. 2019;19(1):1-8.