

1 **A Novel Artificial Intelligence Platform to Automate Clinical Consultation Notes and**
2 **Enhance Diagnostic Efficiency in the Outpatient Clinic: Proposal of a Protocol for a Multi-**
3 **Center, Multi-Disciplinary, Prospective Randomized Controlled Trial**

4
5
6
7 Karanvir Gill*¹, Giovanni Cacciamani¹, Jamal Nabhani*¹, Joshua Corb*², Tom Buchanan¹, Daniel
8 Park¹, Virinder Bhardwaj³, Onkarjit Marwah⁴, Moses Kim⁵, Deepak Kapoor⁶, Alexander Kutikov⁷,
9 Robert Uzzo⁷, Inderbir Gill*^{#1}

10
11
12
13
14 1- Catherine & Joseph Aresty Department of Urology, USC Institute of Urology, University of
15 Southern California, Los Angeles, CA; 2 - Basis Worldwide; 3- Department of Urology, Sierra
16 View Medical Center, Porterville, CA; 4- Heartbeat Cardiovascular Medical Group, Glendale,
17 CA, 5- Orange Coast Urology Associates, Newport Beach, CA; 6- Solaris Urology Associates,
18 New York, NY; 7- Department of Urology, Fox Chase Cancer Center, Philadelphia, PA
19
20
21
22
23
24
25
26

27 Words: Abstract 434

28 Text: 4699

29 Tables: 1

30 References: 24

31
32
33
34 *Conflict of Interest: OneLine Health
35
36
37

38 #Correspondence:

39 Inderbir S. Gill, MD

40 Catherine & Joseph Aresty Department of Urology

41 Keck School of Medicine, University of Southern California 1441 Eastlake Ave, Suite 7416

42 Los Angeles, CA 90089

43 Email: igill@med.usc.edu; gillindy@gmail.com

44 Cell: 216-312-6700
45
46
47
48
49
50
51

52 **ABSTRACT**

53 Presented herein is a proposal for a protocol for a multi-center, multi-disciplinary randomized
54 controlled trial (RCT) to evaluate a novel artificial intelligence (AI)-based technology that
55 automates the construction of the clinical consultation note (CCN) and enhances diagnostic
56 assessments in the outpatient clinic setting. This innovative tech-platform automatically
57 generates the CCN and presents it to the provider in advance of the patient consultation,
58 without any work done by the provider. The constructed CCN is presented either in the native
59 electronic health record (EHR) or in a secure web-based application, in a HIPAA-compliant
60 manner. The proposed prospective trial will compare this novel AI/ML technology
61 (NAMT) versus the current standard-of-care (SOC) in the outpatient setting. Outpatient clinic-
62 days will be randomized to either “SOC clinic-day” or the “NAMT clinic-day” based on whether
63 the SOC or the NAMT was used to construct the CCN for all patients seen on that particular
64 clinic-day. Randomized cross-over of each provider between “SOC clinic-day” and “NAMT clinic-
65 day” will result in each provider serving as her/his own internal control. Objective data will be
66 used to compare study endpoints between the SOC and the NAMT. Co-primary endpoints
67 include a) CCN diagnostic accuracy/quality (based on standardized QNOTE metrics); and b)
68 Work-outside-work (WOW) time required by providers to complete clinic-related documentation
69 tasks outside clinic hours (based on EHR meta-data). Secondary endpoints include a) Provider
70 productivity (based on provider “walk-in, walk-out” time from the consultation room); b) Provider
71 satisfaction (based on the standardized AHRQ EHR End User Survey); and c) Patient
72 satisfaction (based on the standardized Press Ganey/CG-CAHPS survey). To assess
73 generalizability across the health-care spectrum, the study will be conducted in four different
74 types of health-care settings (large academic medical center; non-academic hospital; rural
75 hospital; community private practice); in four different disciplines (cardiology; infectious disease;
76 urology; emergency medicine); using four different EHR systems (Cerner; Epic; AllScripts;
77 MediTech/UroChart). We estimate an aggregate RCT sample size of 150 clinic-days (involving

78 3,000 total patients; 15-30 providers). This will randomize 75 clinic-days (1,500 patients) to the
79 control SOC arm, and 75 clinic-days (1,500 patients) to the intervention NAMT arm. We will use
80 a two-sided Z-test of difference between proportions with 90% power and two-sided 5%
81 significance level. This RCT is the first to evaluate the efficiency and diagnostic accuracy of pre-
82 constructing CCNs in an automated manner using AI/ML technology, deployed at a large-scale,
83 multi-institutional, multi-disciplinary, multi-EHR level. Results from this study will provide
84 definitive level 1 evidence about the desirability and generalizability of AI-generated
85 automatically constructed CCNs, assessing its potential benefits for providers, patients, and
86 healthcare systems.

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104 **INTRODUCTION**

105 Electronic Health Records (EHR) provide secure, immediate access to patient records, store
106 patient's clinical information and coding/billing data, and allow patients secure access to their
107 own records, thereby providing comprehensive information at the point-of-care. Following the
108 2009 Health Information Technology for Economic and Clinical Health (HITECH) Act, EHRs
109 have gained widespread penetration into the U.S. healthcare system, with implementation
110 reaching 96% of U.S. hospitals by 2017 [1, 2].

111 However, for the providers, EHR usage is labor-intensive by its very nature, with its repetitive
112 data entry now the major contributor to provider burnout nationally [3-8]. In outpatient clinics,
113 providers typically devote ~50% of their working hours to the EHR, equating to nearly two hours
114 of EHR use for every hour of patient consultation [9, 10]. This substantial time and effort
115 required for EHR data entry detracts from the actual time providers spend interacting with the
116 patient (versus interacting with the computer) during the consultation. This emphasis on EHR
117 data entry results in suboptimal direct communication between providers and patients, detracts
118 from the quality of the consultation, and decreases patient/provider satisfaction, contributing to
119 the current high rates of physician burnout [11]. Additionally, providers often spend considerable
120 amount of personal time after-hours and during weekends to complete EHR-related tasks,
121 termed 'work-outside-work' (WOW). Small wonder then that currently up to 70% of U.S.
122 physicians report burnout, directly attributed to EHR usage [11-14]

123 Provider burnout is linked to various aspects of EHR use, including time spent on EHR, level of
124 organizational support, and system usability [15]. Burnout can decrease the provider's quality of
125 EHR usage, contributing to diagnostic errors, with implications for patient care [16].

126 Misalignment of providers' cognitive processes with the technical aspects of EHR usability and
127 workflow can increase the risk for diagnostic errors [17]. Collectively, these findings highlight the
128 multi-dimensional burdens confronting providers when incorporating EHR into their practice.

129 During outpatient consultations, providers are required to create a Clinical Consultation Note
130 (CCN) for each patient, which consolidates the patient's medical history, prior test data, and
131 treatment plans in a concise and structured format. Constructing the CCN in the outpatient clinic
132 takes on average 16 minutes of EHR usage per patient [12]. Thoroughness of the CCN drives
133 ICD clinical coding [18], and high CCN quality is also associated with improved physician
134 performance and patient outcomes [19]. Objective, standardized tools to assess CCN diagnostic
135 efficiency and quality are available [20, 21], and EHR metadata can help evaluate the individual
136 physician effort invested into creating this document [9, 10].

137 Emerging artificial intelligence (AI) and machine learning (ML) technologies present robust new
138 opportunities for healthcare innovation [22]. Given that a significant portion of CCN creation
139 involves repetitive, scribe-like work, it can be mitigated with technology. Efforts are underway to
140 use speech-to-text technology and AI to automate CCN creation during clinic visits [9, 10]. More
141 recently, a novel AI/ML and logic-based technology platform has been developed that automates
142 the construction of a detailed preliminary CCN before the actual consultation, without any work
143 on the part of the provider.

144 Herein, we intend to perform a prospective randomized controlled trial (RCT) to compare this
145 novel technology platform with current standard-of-care practices in the outpatient clinic setting.
146 We seek to evaluate the benefits and efficiencies that may accrue to providers and patients
147 alike from using AI to automate the creation of the CCN. Using a large multi-institutional, multi-
148 disciplinary, multi-EHR cohort, our RCT will compare the novel technology (wherein the CCN is
149 constructed automatically using AI before the actual consultation) vis-s-vis the current standard-
150 of-care practice (wherein the CCN is constructed manually during the actual consultation). The
151 generated data will assess CCN diagnostic efficiency and quality, provider 'work-outside-work'
152 effort, provider productivity, provider satisfaction, and patient satisfaction.

153

154

155 **NOVEL TECHNOLOGY**

156 The novel AI technology is comprised of a proprietary logic algorithm-based, symptom-tree
157 Active Sheet that extracts subjective patient data directly from the patient and analyzes and
158 reformats it in a structured manner. In parallel, the AI/ML algorithm extracts relevant,
159 unstructured, objective data from outside records/documents. Proprietary, embedded AI/ML
160 technology generates national guidelines-based recommendations for diagnostic testing in a
161 patient-specific manner. This preliminary CCN is automatically presented to the provider, in
162 advance of the patient's visit, either within the institution's EHR or in a secure web-based App, in
163 a HIPAA-compliant manner. During the actual consultation, the provider re-confirms the clinical
164 history with the patient, performs the physical exam, and finalizes the CCN, completing the
165 patient consultation. The platform's combination of AI/ML algorithms, Active Sheet logic
166 algorithms, and Optical Character Recognition (OCR) technology makes it adaptable to various
167 types of data inputs. Its capability of EHR-integrated and web-based deployment, as well as
168 usage via laptops and cell phones, also ensures high scalability, allowing implementation in an
169 institution with or without an existing EHR system, and for patients with or without laptop
170 computers.

171

172 **METHODS**

173 Our study protocol follows the Standard Protocol Items Recommendations for Interventional
174 Trials-Artificial Intelligence (SPIRIT-AI) Guidelines [23]. The current manuscript is only a
175 proposal for the protocol for this study; IRB approval and clinicaltrials.gov notification will
176 precede the actual trial initiation and consenting/enrollment of any clinical patients.

177

178 **Study Settings**

179 This is a prospective multi-institutional, multi-disciplinary, multi-EHR randomized trial. To assess
180 generalizability across the health-care landscape, the RCT will be conducted concomitantly

181 across four different types of healthcare settings: large academic medical center, non-academic
182 hospital, rural hospital, and community private practices. Participating centers include the
183 University of Southern California and Fox Chase Cancer Center (large academic medical
184 centers); Sierra View Medical Center, Verdugo Hills Hospital (non-academic, rural hospitals);
185 and Solaris Urology and Glendale Cardiology (community private practices). Four specialties
186 will be assessed, including two medical specialties (cardiology, infectious disease), a surgical
187 specialty (urology), and an urgent care specialty (emergency medicine). Four different types of
188 EHRs will be assessed, including Cerner, Epic, AllScripts, and MediTech.

189 This prospective trial will randomize the novel AI/ML technology (NAMT) versus the current
190 standard-of-care (SOC) in the outpatient setting. Randomization will be done per outpatient
191 clinic-days based on method of constructing the CCN. As such, randomization to either the
192 “SOC clinic-day” or the “NAMT clinic-day”, will be based on whether the SOC or the NAMT was
193 used to construct the CCN for all patients seen by that provider in that clinic on that day.

194 Randomized cross-over of each provider between “SOC clinic-day” and “NAMT clinic-day” will
195 allow each provider to serve as his/her own internal control. To provide a balanced and valid
196 comparison, we will use similar clinic workflow methodology for collecting outside records/data
197 in each arm. Thus, medical records clerks and/or nurses would collect any available outside
198 patient records/data in both arms per existing, legacy data collection methods at each
199 participating site. In the “SOC clinic-day” arm, the medical records clerk/nurse would upload
200 patient records to the native EHR platform, and then the CCN would be constructed manually
201 per current standard practice. In the “NAMT clinic-day” arm, the medical records clerk/nurse
202 would upload these records to the NAMT platform (typically housed within the native EHR, thus
203 maintaining similarity in data/records collection methods), but the CCN would be constructed
204 automatically by the NAMT platform. Thus, we will use the same data entry mechanism in each
205 arm to ensure consistency and enhance validity of results. Similar objective endpoint data will

206 be obtained for each provider and patient in the control “SOC clinic-day” arm and the
207 experimental “NAMT clinic-day” arm (Figure 1 – CONSORT-AI Flowchart [24]).
208 Study endpoints will compare the SOC vs the NAMT based on objective data. Co-primary
209 endpoints are: a) CCN quality and diagnostic accuracy (based on standardized QNOTE metrics
210 [14]); and b) Work-outside-work (WOW) time required by providers to complete clinic-related
211 documentation tasks outside normal clinic hours (based on EHR meta-data). Secondary
212 endpoints include: a) Provider satisfaction (based on standardized AHRQ EHR End User
213 Survey); b) Patient satisfaction (based on standardized Press Ganey/CG-CAHPS survey
214 (www.pressganey.com)); and c) Provider productivity (based on provider “walk-in, walk-out’ time
215 from the individual consultation room) and imputed financial data. In total, 15-30 providers will
216 participate. Provider is defined as a licensed medical provider who is constructing the CCN in
217 their daily practice currently, including practicing physicians, mid-level practitioners (physician
218 assistant, nurse practitioner), or trainees (post-graduate fellow or resident). Scribes are not
219 designated as a provider. At the time of scheduling the clinic appointment, patients will be
220 informed that, on their specific clinic-day, their provider is doing all consultations using the
221 specific methodology (SOC or NAMT), as dictated by the randomization. At study onset, all
222 participating providers will attest that they have equipoise between the SOC and NAMT
223 approaches.

224

225 **Randomization: Rationale**

226 Our objective is to evaluate the merits/demerits of the novel AI/ML technology for the automated
227 construction of CCN vis-a-vis the standard-of-care (SOC). Elements that could potentially be
228 randomized include patients, providers, or the intervention itself. However, practically speaking,
229 all of these would impact daily clinic workflow. For instance, if we were to randomize patients,
230 then using both the SOC and NAMT within the same clinic day would be highly disruptive to
231 clinic workflow, and challenging, if not prohibitive, to implement from a practical perspective. If

232 we randomized providers, it would be problematic to randomize the AI/ML technology, and vice-
233 versa. Also, providers with different practice styles conducting one or the other intervention
234 would not control for inter-personal disparities. However, randomization of the entire clinic-day
235 to either “SOC clinic-day” or “NAMT clinic-day” for all patients seen that day, and then equally
236 alternating each provider between “SOC clinic-day” and “NAMT clinic-day” would satisfy study
237 objectives and allow each provider to serve as her/his own internal control. Providers would be
238 a heterogeneous group as regards gender, race, and age. This study design is pragmatic and
239 would be minimally disruptive to daily clinic workflow or provider productivity. To minimize
240 provider bias, at enrollment, all consented providers would attest to equipoise between SOC
241 and NAMT.

242

243 **Eligibility Criteria**

244 Inclusion and exclusion criteria are reported separately for providers and patients.

245

246 a) Providers:

247 i. Inclusion criteria: Licensed medical providers currently engaged in patient
248 consultations in the outpatient clinic, including practicing physicians; mid-level
249 practitioners (physician assistant, nurse practitioner); or trainee (post-graduate fellow
250 or resident). Use of scribes by providers is allowed.

251 ii. Exclusion Criteria: Licensed medical providers who are currently not engaged in
252 patient consultations in the outpatient clinic.

253

254 b) Patients:

255 i. Inclusion criteria: Outpatient clinic patients at least 18 years of age or older being
256 seen in the cardiology, infectious disease, urology, and emergency medicine
257 outpatient settings.

258 ii. Exclusion criteria: Patients younger than 18 years of age.

259

260 **Intervention**

261 Licensed, actively practicing providers who satisfy inclusion criteria are recruited after written
262 consent. Their upcoming clinic days are randomized to either “SOC clinic-day” or “NAMT clinic-
263 day”, wherein CCNs for all patients seen on those respective days are constructed using either
264 the standard-of-care practice or the novel AI/ML technology (Figure 1). Clinic days will be
265 randomized 1-2 weeks prior to the actual date of that clinic. As such, all patients seen on the
266 clinic day randomized to the “SOC clinic-day” will have their CCN constructed per that provider’s
267 customary, standard-of-care practice. Patients seen on the clinic day randomized to the “NAMT
268 clinic-day” will have their CCN constructed 2 weeks to 1 day prior to the clinic visit using the
269 novel AI/ML technology. The automatically constructed CCN will be made available to the
270 provider in advance of the clinic day, either integrated within their native EHR or in the secure,
271 free-standing web-based App. After pre-viewing the CCN, the provider will proceed to perform
272 the clinical consultation. Provider “walk-in, walk-out” time from the consultation room will be
273 recorded by a third-party observer using a stopwatch. After completing the consultation, the
274 patient will be invited to complete the adapted Press Ganey/CG-CAHPS patient satisfaction
275 questionnaire; the patient will provide oral or written informed consent before filling out the de-
276 identified survey instrument. This survey questionnaire has been adapted to fit our study-related
277 objectives, including questions on patient age, gender, level of education, race/ethnicity,
278 provider attentiveness to the patient, provider spending time doing computer work versus
279 interacting with the patient, and provider’s prior knowledge about the patient’s clinical condition
280 before entering the patient’s consultation room. The questionnaire then provides the patient a
281 final opportunity to record an overarching ‘thumbs-up’ or ‘thumbs-down’ about their clinic
282 experience. Interventions for each group are provided below with sufficient details to allow
283 replication, including method and timing of administration.

284

285 **a) Study Intervention for providers**

286 Providers will be invited to participate in the study and informed consent obtained. Providers
287 and their teams will be trained on the novel technology and related clinical workflow over a
288 period of 1-3 clinic-days (“Wash-in” period) to make them knowledgeable about the novel
289 technology platform.

290 Outpatient clinic days will be randomized into one of 2 arms, “SOC clinic-day” and “NAMT clinic-
291 day”. (Details in Figure 1)

292

- 293 i. On “SOC clinic-day”: CCNs for all patients seen on this day will be constructed per
294 the usual, standard-of-care practice of that provider. Prior to the clinic day, the
295 medical records clerk/nurse would upload any outside patient records/data to the
296 native EHR platform per current standard practice, and then the CCN would be
297 constructed manually.
- 298 ii. On “NAMT clinic-day”: CCNs for all patients seen on this day will be pre-constructed
299 using the novel AI/ML technology. At the time of being given the clinic appointment,
300 patients will be sent the electronic link and advised to do two things from their home -
301 ‘the earlier, the better’ - prior to coming to the clinic: a) complete the Active Sheet
302 (AS): when completing the AS, the patient may solicit help from family/friends, or
303 from remote licensed provider extender (PA/NP) or referring provider staff; and b)
304 upload any lab/radiology test results: the medical records clerk/nurse would help the
305 patient upload outside records/data to the NAMT platform (typically housed within the
306 native EHR platform). After these two actions – completing the AS; uploading the
307 lab/radiology test results - are done, the novel AI software automatically constructs
308 the CCN and delivers it to the provider, either in their native EHR or in the secure,
309 free-standing web-app. This action occurs automatically in the background invisible

310 to the provider/staff, in advance of the patient's visit. The patient then comes to the
311 clinic on the scheduled appointment date, the provider reviews the pre-constructed
312 CCN and proceeds with the consultation.

313

314 **b) Study Intervention for patients**

315 Immediately upon completion of their consultation, patients will be invited to fill out the adapted
316 Press-Ganey/CG-CAHPS survey in a de-identified fashion to assess patient satisfaction.
317 Patients who agree to fill out the survey will be asked to first provide written informed consent
318 for inclusion in the study. Upon completion of the survey, the patient departs the clinic area.
319 Patient participation will be deemed completed upon survey completion.

320

321 **Outcomes and Measurements**

322 Table 1 summarizes the a) co-primary and b) secondary outcomes of interest. Details regarding
323 the outcome measurements are reported in the supplementary materials.

324

325 **a) Co-Primary Outcomes**

326 Our co-primary objectives are (a) delivering a CCN of high-quality and diagnostic accuracy
327 without any effort on the part of the provider, and (b) reducing the provider time for doing work-
328 outside-work (WOW) to complete clinic-related documentation activities performed outside of
329 the typical 8 am to 4 pm clinic schedule. Corresponding co-primary endpoints are the objective
330 data metrics to assess CCN quality and diagnostic accuracy (using the standardized QNOTE
331 methodology), and provider WOW time (using EHR meta-data). The rationale behind these
332 endpoints is rooted in the central tenet of this automated technology, which is to create an
333 automated CCN that matches or exceeds the current, manually created, standard-of-care CCN,
334 while minimizing workload for the healthcare provider. Furthermore, we aspire to decrease the
335 time spent by providers working on clinic notes outside regular work hours. To ascertain these

336 endpoints, we utilize QNOTE to measure CCN quality and diagnostic accuracy using objective
337 metrics; provider's WOW time is determined by analyzing the objective EHR metadata for that
338 clinic-day, provided by the institutional IT team. The putative mechanisms of action underpinning
339 these objectives lie in streamlining clinical processes, thus enhancing the quality of patient care
340 and reducing diagnostic errors.

341

342 **b) Secondary Outcomes**

343 The secondary objectives of the study assess provider satisfaction, patient satisfaction, and
344 physician productivity. Provider satisfaction will be assessed using an adapted AHRQ Electronic
345 Health Record End User Survey. We hypothesize that automating the CCN will reduce the
346 workload of physicians, and consequently improve satisfaction and decrease burnout rates. At a
347 minimum at the end of each clinic day, providers will fill out the AHRQ EHR End User Survey to
348 capture their clinic-day experiences and responses.

349 Patient satisfaction, another critical component of healthcare delivery, will be evaluated using an
350 adapted Press Ganey/CG-CAHPS survey. We hypothesize that if providers spend more time
351 directly interacting with patients, instead of interacting with and entering data into the EHR,
352 patients will experience higher satisfaction. As such, consented patients will be requested to
353 complete the adapted Press Ganey/CG-CAHPS survey immediately after their consultation to
354 measure their level of satisfaction. Finally, physician productivity, particularly in terms of time
355 savings, which can lead to increased patient throughput and financial return-on-investment
356 (ROI), will be documented. We believe that automating the CCN might lead to decreased, yet
357 higher quality, consultation times, thereby improving physician productivity. To measure this, the
358 physician's "walk-in, walk-out" time from each consult room will be recorded with a stopwatch.
359 This objective, real-time measurement will assess the impact of the intervention on physician
360 productivity. Financial ROI will be imputed.

361

362 **Sample Size**

363 Given the novelty of our AI algorithm, prior randomized data do not exist. Our sample size will
364 be designed to detect a minimum of 30% improvement in the CCN quality/diagnostic accuracy,
365 and a 30% reduction in the duration of WOW, our 2 co-primary endpoints. We estimate the total
366 sample size to be 150 clinic-days, i.e., 75 “SOC-days” and 75 “NAMT-days”. Conservatively
367 assuming 20 patients seen per clinic-day, we will accrue at least 3,000 total patients - 1,500 in
368 the control “SOC-day” arm, and 1,500 in the intervention “NAMT-day” arm. Clinic-days will be
369 divided between the 4 specialties, leading to 30-40 clinic-days per specialty. We estimate a total
370 of 15-30 study providers, resulting in 5-10 clinic-days per provider. No single specialty will
371 accrue more than 30% of the aggregate clinic-days. We will use a two-sided Z-test of the
372 difference between proportions with 90% power and two-sided 5% significance level. This
373 sample size exceeds minimum requirements and will provide robust data. Study data will be
374 collected by trained personnel, physically located at each clinical site. The validated QNOTE
375 scoring system will assesses quality and diagnostic accuracy of de-identified CCNs uploaded to
376 an online repository for scoring by independent, blinded raters. Data collection process will be
377 integrated into clinic workflows, preserving the user experience while introducing the new AI/ML
378 tool.

379

380 **Recruitment**

381 Providers will be recruited from the participating institutions. We anticipate between 15-30
382 providers to participate in this RCT. This would provide a heterogeneous group of providers as
383 regards gender, race, and age, with each provider participating in 5-10 study clinic-days.
384 Randomized alternating cross-over of providers will result in each provider serving as her/his
385 own internal control. Patients will be identified exclusively based on the participating providers'
386 clinic schedules. Patients will be recruited and consented to the study immediately after they
387 finish their clinical consultation with the provider, and before they are administered the Press

388 Ganey/CG-CAHPS survey instrument. We intend to recruit over 3000 patients over a one-year
389 period. No incentives will be provided to either physicians or patients for participating in this
390 RCT.

391

392 **Screening Failure**

393 In case a provider has been already enrolled but wishes to drop out of the study within the first 5
394 clinic-days, he/she will be replaced with another physician from that specialty. Any provider who
395 has already completed the first 5 clinic-days in the study will not be replaced by another
396 provider; their data will be used as is with censoring at study drop-out. If a patient chooses to
397 not fill out the Press Ganey/CG-CAHPS satisfaction survey, it will be recorded as such, and their
398 survey data will be excluded from study calculations.

399

400 **Discontinuation of intervention**

401 After initially agreeing to participate, if a provider decides to discontinue in the study, she/he
402 must express this desire prior to 5 clinic-days after enrolling in the study; in this case their data
403 over those 5 clinic-days will be used, and they will be replaced by another physician. After 5
404 clinic-days, they will be considered to have been part of the trial and while their data will be
405 included for analyses, they will not be replaced by another physician. Participation in this study
406 is voluntary. If a patient decides to not provide informed consent at the time of answering the
407 Press Ganey/CG-CAHPS questionnaire, their survey data will not be included in the study. If a
408 patient decides to not participate in the study after initially agreeing, they must express this
409 desire prior to completing their Press Ganey/CG-CAHPS questionnaire and leaving the clinic.
410 The sole criterion for patient participation in the study is patient willingness. Participation in this
411 study is voluntary.

412

413 **Statistical Hypothesis and Analysis**

414

415 a) Statistical Hypothesis

416 i. Primary Efficacy Endpoint(s): We hypothesize that the CCN generated by the novel
417 technology will be superior to the SOC as regards CCN quality and diagnostic
418 accuracy (as measured by QNOTE methodology), and CCN efficiency (WOW time
419 as measured by EHR meta-data). Our null hypothesis is that there will be no
420 difference in the quality, diagnostic accuracy, and efficiency of the CCN derived from
421 the novel technology compared to standard-of-care.

422 ii. Secondary Efficacy Endpoint(s): We hypothesize that the CCN generated by the
423 novel AI/ML technology will be superior to the SOC as regards provider satisfaction
424 (adapted AHRQ Electronic Health Record End User Survey); patient satisfaction
425 (adapted Press Ganey/CG-CAHPS survey); and physician productivity (“walk-in,
426 walk-out” consult time).

427

428 Our null hypothesis is that outcomes will be no difference in the above-mentioned three metrics
429 for CCNs generated by the novel AI/ML technology versus standard-of-care.

430

431 b) Statistical analysis

432 Analysis of the primary endpoint will assess mean data for CCN quality/accuracy and CCN
433 efficiency to demonstrate superiority of the NT compared to SOC.

434

435 i. Quality of CCN will be assessed by the adapted QNOTE instrument, a validated
436 methodology that utilizes 15 clinical elements and 7 evaluative components (as
437 detailed in supplementary materials). Each CCN will be de-identified by redacting
438 patient/physician identifiers prior to assessment and uploaded to an online repository
439 for QNOTE scoring by three blinded, third-party raters. Each patient will be assigned

440 a unique identifier number to ensure confidentiality and privacy. Depending on quality
441 of each CCN, a score between 0 (minimum) to 1500 (maximum) will be assigned by
442 independent raters. The average of the ratings from the 3 independent raters for
443 each CCN will denote the composite QNOTE score for that CCN, which will be the
444 basis for statistical analysis comparing CCNs created on “SOC clinic-day” versus
445 “NT clinic-day”.

446 ii. Work-outside-Work (WOW) time will be objectively assessed from EHR meta-data
447 per provider (as detailed in supplementary materials).

448

449 Analysis of the secondary endpoints will calculate mean data to assess outcomes of the NT
450 compared to SOC for the following three aspects:

451

452 i. Provider satisfaction (adapted AHRQ Electronic Health Record End User Survey):
453 dichotomous positive versus negative Likert scale answers will be compared for each
454 question and in the aggregate.

455 ii. Patient satisfaction (adapted Press Ganey/CG-CAHPS survey): dichotomous
456 positive versus negative Likert scale answers will be compared for each question
457 and in the aggregate.

458 iii. Provider productivity (Total consultation time): physician “walk in-walk out” time from
459 the consult room at each patient consultation will be measured by the study
460 coordinator using a stopwatch.

461

462 The sample size will be designed to detect a minimum of 30% improvement in the CCN
463 quality/diagnostic accuracy, and a 30% reduction in the duration of WOW, our 2 co-primary
464 endpoints. Analysis of primary and secondary objectives will proceed as follows. Missing data
465 will be dealt with by imputation, and potential outliers will be scrutinized by the Principal

466 Investigator. For each question, Likert scale responses will be analyzed, allowing comparisons
467 between the novel technology and the standard-of-care. Answers will be divided into positive
468 and negative categories per the "top box method," which is standard for assessing CG-CAHPS
469 data. Likert scale responses will be dichotomized, with scores of 4 (agree) and 5 (strongly
470 agree) representing agreement (positive) and scores of 1 (strongly disagree), 2 (disagree), or 3
471 (neutral) indicating disagreement (negative). Screening for outliers will be conducted based on
472 the absolute agreement of each individual question and distribution of responses. Continuous
473 variables will be represented as mean and standard deviation (SD), and categorical variables
474 will be presented as median and interquartile range (IQR). For the univariate analysis, Kruskal-
475 Wallis, chi-squared (χ^2), and Fisher's exact tests will be utilized to compare continuous and
476 categorical variables as appropriate. A 95% confidence interval and a two-tailed test with $p < .05$
477 will be taken as statistically significant.

478 Upon completion of data collection, each question will be analyzed using Cochran-Mantel-
479 Haenszel tests, stratified by provider to eliminate potential individual provider bias from the
480 results. P-values will be determined using Chi-Square tests and will be deemed significant at
481 less than 0.05. For the assessment of patient satisfaction (utilizing the adapted Press
482 Ganey/CG-CAHPS survey with additional queries), each question will be stratified by provider to
483 remove potential individual bias and analyzed using Cochran-Mantel-Haenszel tests.

484 Considering the study's novelty and the absence of prior randomized data, an interim analysis
485 will be conducted. Calculations will be performed using an α level of 0.05 and a power of 90%.
486 Based on the trends in the data from the first 50 patients, sample sizes will be calculated for
487 each question, assuming the variables to be dichotomous. After the complete data collection,
488 each question will be analyzed using Cochran-Mantel-Haenszel tests, stratified by providers to
489 remove any individual provider bias. P-values will be determined using Chi-Square tests, and a
490 value of less than 0.05 will be considered significant.

491

492 **CONCLUSION**

493 Herein we propose the protocol of the first prospective randomized controlled trial of a large-
494 scale, multi-institutional, multi-disciplinary cohort to evaluate the practicality and efficiency of
495 using AI/ML technology to pre-construct the CCN automatically before the actual patient-
496 provider consultation. The study will compare this AI/ML framework vis-a-vis the standard-of-
497 care, wherein the CCN is generated manually during the consultation encounter. Data
498 generated by this study will provide insights into the desirability and generalizability of AI-
499 generated automated clinical consultation notes across the healthcare spectrum, and its
500 potential benefits to providers, patients, and healthcare institutions.

501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533

534 **REFERENCES**

- 535
- 536 1. Goldstein, M.M. and H. Thorpe Jane, The First Anniversary of the Health Information
537 Technology for Economic and Clinical Health (HITECH) Act: the regulatory outlook for
538 implementation. *Perspectives in health information management*, 2010. 7(Summer): p. 1c.
539
- 540 2. HealthIT.gov, [https://www.healthit.gov/data/quickstats/office-based-physician-electronic-](https://www.healthit.gov/data/quickstats/office-based-physician-electronic-health-record-adoption)
541 [health-record-adoption](https://www.healthit.gov/data/quickstats/office-based-physician-electronic-health-record-adoption). 2020.
542
- 543 3. McPeck-Hinz, E., et al., Clinician Burnout Associated With Sex, Clinician Type, Work
544 Culture, and Use of Electronic Health Records. *JAMA Netw Open*, 2021. 4(4): p. e215686.
545
- 546 4. Melnick, E.R., et al., Analysis of Electronic Health Record Use and Clinical Productivity
547 and Their Association With Physician Turnover. *JAMA Netw Open*, 2021. 4(10): p. e2128790.
548
- 549 5. Rotenstein, L.S., C. Sinsky, and C.K. Cassel, How to Measure Progress in Addressing
550 Physician Well-being: Beyond Burnout. *Jama*, 2021.
551
- 552 6. Kroth, P.J., et al., Association of Electronic Health Record Design and Use Factors With
553 Clinician Stress and Burnout. *JAMA Netw Open*, 2019. 2(8): p. e199609.
554
- 555 7. Melnick, E.R., et al., The Association Between Perceived Electronic Health Record
556 Usability and Professional Burnout Among US Physicians. *Mayo Clin Proc*, 2020. 95(3): p. 476-
557 487.
- 558 8. Nguyen, O.T., et al., A systematic review of contributing factors of and solutions to
559 electronic health record-related impacts on physician well-being. *J Am Med Inform Assoc*, 2021.
560 28(5): p. 974-984.
561
- 562 9. Kumah-Crystal, Y.A., et al., Electronic Health Record Interactions through Voice: A
563
564 Review. *Applied clinical informatics*, 2018. 9(3): p. 541-552.
565
- 566 10. Kataria, S. and V. Ravindran, Electronic health records: a critical appraisal of strengths
567 and limitations. *J R Coll Physicians Edinb*, 2020. 50(3): p. 262-268.
568
- 569 11. Naidu, A., Factors affecting patient satisfaction and healthcare quality. *Int J Health Care*
570 *Qual Assur*, 2009. 22(4): p. 366-81.
571
- 572 12. Overhage, J.M. and D. McCallie, Physician Time Spent Using the Electronic Health
573 Record During Outpatient Encounters. *Ann Intern Med*, 2020. 173(7): p. 594-595.
574
- 575 13. Yan, Q., et al., Exploring the relationship between electronic health records and provider
576 burnout: A systematic review. *J Am Med Inform Assoc*, 2021. 28(5): p. 1009-1021.
577
- 578 14. Burke, H.B., et al., QNOTE: an instrument for measuring the quality of EHR clinical
579 notes.
580
581 *J Am Med Inform Assoc*, 2014. 21(5): p. 910-6.
582
- 583 15. Li, C., et al., Electronic medical record-related burnout in healthcare providers: a scoping
584

585
586 review of outcomes and interventions. *BMJ Open*, 2022. 12(8): p. e060865.
587
588 16. Graber, M.L., C. Byrne, and D. Johnston, The impact of electronic health records on
589 diagnosis. *Diagnosis (Berl)*, 2017. 4(4): p. 211-223.
590
591 17. Dixit, R.A., et al., Electronic Health Record Use Issues and Diagnostic Error: A Scoping
592 Review and Framework. *J Patient Saf*, 2023. 19(1): p. e25-e30.
593
594
595 18. (2008)., A.C.f.L.A.a.t.N.H.S.i.E.a.W., PbR Data Assurance Framework 2007/08: findings
596 from the first year of the national clinical coding audit programme. Audit Commission. 2008.
597
598 19. Mann, R. and J. Williams, Standards in medical record keeping. *Clin Med (Lond)*, 2003.
599
600 3(4): p. 329-32.
601
602 20. Logan, J.R., P.N. Gorman, and B. Middleton, Measuring the quality of medical records: a
603 method for comparing completeness and correctness of clinical encounter data. *Proc AMIA*
604 *Symp*, 2001: p. 408-12.
605
606 21. Hong, C.J., et al., Accuracy and completeness of electronic medical records obtained
607 from referring physicians in a Hamilton, Ontario, plastic surgery practice: A prospective
608 feasibility study. *Plast Surg (Oakv)*, 2015. 23(1): p. 48-50.
609
610 22. Lee, P., S. Bubeck, and J. Petro, Benefits, Limits, and Risks of GPT-4 as an AI Chatbot
611 for Medicine. *N Engl J Med*, 2023. 388(13): p. 1233-1239.
612
613 23. Cruz Rivera, S., et al., Guidelines for clinical trial protocols for interventions involving
614 artificial intelligence: the SPIRIT-AI extension. *Lancet Digit Health*, 2020. 2(10): p. e549-e560.
615
616 24. Liu, X., et al., Reporting guidelines for clinical trial reports for interventions involving
617 artificial intelligence: the CONSORT-AI extension. *Lancet Digit Health*, 2020. 2(10): p. e537-
618 e548.
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635

636 **Table 1**
637

OBJECTIVES	ENDPOINTS	JUSTIFICATION FOR ENDPOINTS	PUTATIVE MECHANISMS OF ACTION
CO-PRIMARY			
<p>1. Quality/Diagnostic Accuracy of Clinical Consultation Note (CCN)</p> <p>2. Efficiency of CCN construction</p>	<p>1. QNOTE: to assess CCN quality/diagnostic accuracy (<i>Objective data</i>)</p> <p>2. Objective Meta-data: from EMR (<i>Objective data</i>)</p>	<p>1. The quality/accuracy of the automated CCN should be, at a minimum, equivalent to the standard-of-care</p> <p>2. Extra time (beyond typical clinic hours) spent by provider to complete CCN should decrease</p>	<p>1. QNOTE: Blinded, objective third-party assessment of CCN quality/accuracy</p> <p>2. Work-outside-Work (WOW) time/provider: Objective per-patient EHR meta-data from IT team</p>
SECONDARY			
<p>3. Provider satisfaction</p> <p>4. Patient satisfaction</p> <p>5. Provider productivity</p>	<p>3. AHRQ EHR End User Survey (<i>Objective data</i>)</p> <p>4. Press Ganey/CG-CAHPS survey (<i>Objective data</i>)</p> <p>5. Consultation time savings (<i>Objective data</i>)</p>	<p>3. Automating the CCN should decrease provider workload & improve satisfaction</p> <p>4. Automating the CCN should allow provider to spend more time interacting with the patient (rather than with the EHR), improving patient satisfaction</p> <p>5. Consultation times should, at a minimum, be similar to standard-of-care, upon automating the CCN</p>	<p>3. Providers to fill out adapted AHRQ EHR End User Survey per new patient consult or per clinic-day</p> <p>4. Patients to fill out adapted Press Ganey/CG-CAHPS survey at end of the consult</p> <p>5. Physician “walk-in walk-out” time from consult room, measured by stopwatch</p>

638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658

CONSORT Flow Diagram (Ideal Scenario)

