1	Late reproduction is associated with extended female survival but not with familial				
2	longevity				
3					
4					
5	Authors:				
6	Annelien C de Kat ¹ , Femke Roelofs ¹ , P. Eline Slagboom ² , Frank JM Broekmans ¹ , Marian				
7	Beekman ² , Niels van den Berg ²				
8					
9	Affiliations:				
10	1) Department of Reproductive Medicine and Gynecology, University Medical Center				
11	Utrecht, Utrecht, The Netherlands.				
12	2) Department of Biomedical Data Sciences, section of Molecular Epidemiology, Leiden				
13	University Medical Center, Leiden, The Netherlands				
14					
15					

16 Abstract (350 words): 17 Objective 18 Female reproductive health comprises adequate oocyte quality and quantity, normal 19 fecundability, a normal course of pregnancy, and the delivery of healthy offspring. General 20 aging patterns and the maintenance of somatic health have been associated with female 21 reproductive health. However, it is unknown whether better long-term somatic 22 maintenance is directly related to reproductive outcomes, and whether there is a shared 23 genetic predisposition underlying both somatic and reproductive aging. Here we investigate 24 whether reproductive health is related to female lifespan or familial longevity. 25 26 Design 27 Observational study 28 29 Subjects 30 10,255 female members of a multigenerational historical cohort (1812-1910), the LINKing 31 System for historical family reconstruction (LINKS) and 1,258 females from 651 long-lived 32 families in the Leiden Longevity Study. 33 34 Main outcome measures 35 The association between reproductive characteristics and longevity was studied both on an 36 individual and familial level. Individual survival was studied in relation to age at last 37 childbirth and total number of children. Familial longevity was studied through parental 38 survival and related to age at last childbirth, total number of children and a polygenic risk 39 score (PRS) for age at menopause. 40 41 Results 42 Females giving birth to their last child at a higher age lived longer: for each year increase in 43 the age of the birth of the last child, a woman's lifespan increased with 0.06 years (22 days) 44 (p<0.005). Females who lived to be in the top 10% survivors of their birth cohort (n=2,241,

21.9%) on average gave birth to their last child at a 1% later age than the remaining cohort

45

46

(IRR=1.01, p<0.005).

47 Females with 1 or 2 long-lived parents did not have a higher mean age of last childbirth. 48 There was neither a significant association between an increasing number of long-lived 49 ancestral family members (familial longevity), nor the PRS. 50 51 Conclusion 52 Female reproductive health associates with a longer lifespan and with survival to more 53 extreme ages (longevity). The heritable component in familial longevity, however, does not 54 associate to extended reproductive health and the PRS underlying age at menopause does 55 not explain familial longevity. Other factors in somatic maintenance that support a longer 56 lifespan are likely to have an impact on reproductive health. 57 58 Keywords 59

60

Longevity - fertility - aging

Introduction

Female reproductive health encompasses the evolution of being born with a complete set of oocytes, fertility and pregnancy, to the deterioration of ovarian quality and quantity, and ultimately postmenopausal health. It is widely accepted that these milestones and transitions do not stand alone but may be subject to the same processes that govern overall somatic aging and health ¹. This relationship has not yet been fully clarified and it thus remains unknown to what extent the maintenance of somatic health is primarily essential to reproductive health or vice versa, and whether there is a genetic predisposition underlying both healthier somatic and reproductive aging.

Over the past decades a plethora of studies has sought to determine and explain the relationship between ovarian and overall somatic aging. Though there remains some dispute, several studies have observed that mothers who give birth to a child at an advanced age have a longer post-reproductive survival ^{2,3}. Studies also suggested a familial or genetic component underlying both an increasing somatic lifespan and a longer reproductive period or reproductive lifespan^{4,5}, although others have proposed a trade-off mechanism for an increasing lifespan and childbearing⁶. The latter results originate from studies with varying sample sizes and potential biases in the selection of their study population and await confirmation from well-defined, large-scale cohorts. If longevity and late reproductive aging coincide in families, the study of both traits in longevity families may reveal shared genetic loci predisposing to better maintenance of somatic and reproductive cell functions.

Genetic studies into menopause generated polygenic risk scores (PRS) that indicate risk of early menopause and involved loci in DNA repair processes, known as hallmark

mechanisms of ageing ⁷. It is unknown whether the genetic component in age at menopause associates with that in familial longevity.

The current study addresses the relationship between somatic and reproductive aging and health in both a large multigenerational historical cohort and a cohort of long-living families. We aim to test whether reproductive health is associated with longevity, as well as whether members of exceptionally long-living families exhibit more favorable reproductive outcomes, compared to families from the general population. Firstly, we investigate whether lifespan (age at death) and longevity (survival to extreme ages) associate with age at last childbirth and total number of children. Secondly, we investigate whether age at last childbirth and number of children are associated with the number of long-lived parents (0, 1 or 2 long-lived parents), as an indicator of familial longevity 12. Thirdly, we investigate whether an increasing number of long-lived ancestors associates with a PRS for age at menopause, capturing alleles associated with age at menopause, as a proxy for the total reproductive lifespan.

Methods

100 101 102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

LINKS Study population

We used data from the LINKing System for historical family reconstruction (LINKS) which is a historical cohort of inhabitants of the province of Zeeland, the Netherlands, from the early 19th century. The LINKS database contains demographic and genealogical information derived from the Netherlands linked vital event registration. In the Netherlands, birth, marriage and death certificates were registered from the year 1812 onward. Currently LINKS Zeeland contains 739,453 birth, 387,102 marriage, and 641,216 death certificates that were linked together to reconstruct intergenerational pedigrees and individual life courses⁸. Two generations were identified in the dataset (Suppl Figure 1); FO and F1, of which the F1 generation is the index generation comprising the study participants. The F0 generation was selected by identifying couples who were married between 1812 and 1850 and had at least two children, ensuring that the F1 persons had at least one sibling. The families were mutually exclusive, meaning that a parent in the FO generation could only contribute data for a single family. From the F1 generation, the LINKS research persons (RP) were selected using the following criteria: members of the female sex, with an age of death above 50 years and a single spouse who lived until the RP was at least 50 years, and who delivered at least one child, ensuring high data quality. This selection made it possible to study the reproductive outcomes in the study population throughout the entire fertile lifespan. In both generations, a distinction was made between persons who belonged to the top 10% of survivors in their birth cohort, and those who did not. This calculation was based on Dutch lifetables, nationally collected data of survival of historic cohorts from Statistics

Netherlands (CBS)⁸. Reproductive characteristics of the RPs are derived using information of

their children (F3 generation). The reproductive characteristics that could reliably be extracted from the historic data were age at last childbirth and total number of children.

Leiden Longevity Study population

124

125

126

127128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

The Leiden Longevity Study (LLS) was initiated in 2002 to study the mechanisms that lead to exceptional survival in good health. The LLS currently consists of 651 three-generational families, defined by siblings who have the same parents. Inclusion took place between 2002 and 2006 and initially started with the recruitment of living nonagenarian sibling pairs of European descent (F2 generation). Within a sibling pair, males were invited to participate if they were 89 years or older and females if they were 91 years or older (N=944, mean age=93 years), representing < 0.5% of the Dutch population in $2001^{9,10}$. Inclusion was subsequently extended to the offspring of the sibling pairs and the partners of these offspring (F3 generation). This study focuses on all F3 generation females, which are henceforth denoted as LLS RPs (offspring and partners combined). The LLS DNA samples were genotyped using Illumina Infinium HD Human660W-Quad and OmniExpress BeadChips (Illumina, San Diego, CA, USA). DNA genotyping for LLS was performed at baseline as described in detail in Beekman et al., 2006 11 with the Illumina Human660W and Illumina OmniExpress arrays (Illumina, San Diego, CA, USA). Genotype imputation was performed using 288,635 SNPS with SNP-wise call rate (>95%), minor allele frequency (>1%) and no derivation from the Hardy-Weinberg equilibrium (p-value > 1×10^{-4}) at the Michigan Imputation Server (https://imputationserver.sph.umich.edu/index.html) with Haplotype Reference Consortium (HRC) reference panels (HRC1.1). Mortality information was verified by birth or marriage certificates and passports whenever possible. Additionally, verification took place via personal cards which were obtained from the Dutch Central Bureau of Genealogy. In January 2021 all mortality information was updated through the Personal Records Database (PRD) which is managed by Dutch governmental service for identity information. https://www.government.nl/topics/personal-data/personal-records-database-brp. The combination of officially documented information provides very reliable and complete ancestral as well as current mortality information.

Construction of LRC score in LLS data

Familial longevity was quantified with the Longevity Relatives Count (LRC) score. The LRC score can be interpreted as a weighted proportion (ranging between 0 and 10) ¹². For example, an LRC score of 5 for an RP indicates 50% long-lived family members, weighted by the genetic distance between RPs and their family members.

Construction of PRS of age at menopause in LLS data

The genome-wide association analysis for age at menopause resulted in a polygenic risk score (PRS) 7 that could be constructed from 290 SNPs. After we removed all T/A SNPs, SNPSs with MAF<0.01 and HWE_{pvalue}<10⁻⁴, and an imputation quality <0.8 13 , we were able to use 195 SNPs to construct the PRS for age at menopause in the LLS data set.

Statistical analysis

First, in LINKS, lifespan (age at death) was regressed on 1) age at last child birth (quantitative and categorical) and 2) number of children (quantitative and categorical), while adjusting for maternal birth cohort and age at marriage using a linear mixed-model with a random effect for the unique sibship ID to account for within sibship correlation.

Second, in LINKS, 1) age at last child birth and 2) number of children were regressed on a) the number of long-lived parents (0, 1 or 2 parents that belonged to the top 10% survivors of their birth cohort), as an indicator of familial longevity ¹⁴, and b) whether the RP survived to the top 10% of her birth cohort, while adjusting for maternal birth cohort and age at marriage using a Poisson mixed-model with a random effect for the unique sibship ID to account for within sibship correlation.

Third, in the LLS, the standardized (mean=0, SD=1) PRS of age at menopause of RPs was regressed on the number of long-lived family members, as measured with the LRC score, while adjusting for standardized birth year using a linear mixed-model with a random effect for the family ID to account for within family correlation.

All analyses were performed with R version 4.0.2.

Results

Study populations

In the LINKS data, 10,255 female Research Persons (RPs; the F1 generation) were identified. Collectively, they had 7,664 mothers and 7,636 fathers (the F0 generation) and 72,895 children (the F2 generation). In total, there were 7,721 unique families, taking into account that the RP generation included siblings. Mean age at death of the RPs was 73.9 (± 10.4) years and their mean number of children was 7.1 (± 3.9). Further descriptive characteristics of the RP group are described in detail in Table 1.

The LLS cohort included 1,258 females with a mean age of 59 years.

Females giving birth to their last child at a higher age live longer

To confirm in the LINKS dataset that mothers who give birth to a child at an advanced age have a longer post-reproductive survival, we investigated the relation between age at last child and lifespan using linear mixed-model regression analysis. We observed that for each year increase in the age of the birth of the last child, a woman has a 0.06 years (22 days) longer lifespan (p-value= $2.16 \cdot 10^{-3}$). When we compared the lifespan of females with a low (≤ 40 years) vs high (≥ 45 years) age at last child birth, we observed that females who delivered their last child beyond the age of 45 lived 1.41 years (17 months) longer than females who had their last child before 40 years (p-value= $9.07 \cdot 10^{-5}$), while adjusting for age at marriage and the mother's birth year (Table 2).

The number of children may relate both to age at latest childbirth and to survival of the mother, so we investigated whether the number of children associates with female lifespan. We performed a linear mixed-model regression analysis with lifespan as outcome and the number of children as the independent variable of interest, while adjusting for age at marriage and maternal birth year. We observed that with each additional child the

lifespan of females increased with 0.05 years (p-value= $7.63\cdot10^{-2}$). Moreover, females who had 10 or more children (N=2,771) in comparison to females who had 4 and fewer children (N=2,990) lived on average 0.61 years (18 days) longer, but this did not reach statistical significance (Table 2; p-value= $6.25\cdot10^{-2}$).

Next, we studied whether females who lived as long as the top 10% survivors of their birth cohort (long-lived), gave birth to their last child at higher ages than other females using mixed-model Poisson regression analysis. While adjusting for age at marriage and the RP's maternal birth year, we observed that the top 10% survivor (N = 2,241, 21.9%)), on average gave birth to their last child at a 1% later age than the remaining cohort (N= 8,014; 78,1%) (IRR=1.01; p-value=2.75·10⁻³). Thus, with an increasing age of last child, an RP has a very small increasing chance to become long-lived. Moreover, females who lived as long as the top 10% of their birth cohort, had on average 2% more children than other females (IRR=1.02, p-value=9.19·10⁻²), although this effect was not statistically significant. We conclude from the LINKS cohort that women with more offspring and a high age at last reproduction have a longer lifespan and that top 10% female survivors of their birth cohort are capable to reproduce longer than other women.

Females with long-lived parents do not differ from females from the general population with respect to their age at last child and number of children

High quality somatic maintenance in individuals may beneficially affect both reproductive health and longevity. It has also been suggested that familial longevity and especially heritable mechanisms drive such joint beneficial effects. To study whether familial longevity associates with reproductive health, we investigated whether females from long-lived families have more offspring and higher age at last reproduction. We observed that RPs with 1 or 2 long-lived parents do not have a significantly higher number of children (IRR-

237

238

239

240

241242

243

244

245

246

247

248

249

250

251

252

253

254

255

1 LL parent=0.98, p-value=1.62·10⁻¹; IRR-2 LL parents=0.98, p-value=6.44·10⁻¹) nor a higher mean age at last child birth (IRR-1 LL parent=0.99, p-value=5.28·10⁻²; IRR2 LL parents=1.00. p-value=9.99·10⁻¹) compared to RPs without any long-lived parents (Table 3). Females with long-lived parents have a similar genetic predisposition for a late menopause as other females To further study the genetic component underlying both longevity and reproductive health we focused on the Leiden Longevity Study. We investigated whether an increasing number of long-lived ancestors of female RPs, as measured with the LRC score, associates with an increasing number of alleles marking a later age at menopause. In the Leiden Longevity Study we included a generation of RP women (F3 generation; N=1,258) with a quantified family history of longevity, as well as their genotypes through single nucleotide polymorphism (SNP) data. We calculated a PRS for age at menopause based on the most recent Genome Wide Association Study (GWAS) 7. Using mixed-model linear regression, no statistically significant association between an increasing number of long-lived family members (familial longevity) and our polygenic score for age at menopause was observed (Beta=0.014 (95% CI=-0.01-0.04), Pvalue=2.63·10⁻¹). Still, the effect size was in the expected direction: with every 10 percent

increase in long-lived family members, the age at menopause, as expressed by our polygenic

score, was 0.014 standard deviation lower.

Discussion

256257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

In the historic (1812-1910) multigenerational LINKS database of Zeeland (NL), we observed that females who gave birth to their last child at a higher age, and with increasing numbers of offspring had a longer lifespan. Moreover, females who were among the top 10% survivors of their birth cohort delivered their last child at a slightly higher age and overall had slightly more children. This could be explained by a better maintenance of both reproductive health and overall health that supports longevity. Next, we studied whether the genetic component in familial longevity might associate with that of reproductive health, possibly pointing to common mechanisms in of both traits. However, females descending from such long-lived families do not have a different number of offspring or age at last child than females from non-long-lived families. Moreover, in the LLS, there was no evidence for an association between the genetic predisposition for a delayed age at menopause and familial longevity, as measured by a score indicating the proportion of long-lived ancestors. Hence, we conclude that a high age of last childbirth and number of offspring are markers of good reproductive health and overall health supporting longevity. They are not, however, explained by the genetic component in longevity, neither in the LINKS population nor in the Leiden Longevity study. Finally, the heritable component underlying the clinical extremes of age at menopause as represented by a PRS ⁷ does not appear to coincide with that in familial longevity. Our study affirms previous research that supports a relationship between a later age

Our study affirms previous research that supports a relationship between a later age at last childbirth and increased post-reproductive survival^{2,3,15-19} and the notion that the functioning of the reproductive system can be representative of females' health, not necessarily driving it as our data imply. The ability for late reproduction and a larger number of offspring could be facilitated by a longer reproductive period, i.e. later age at menopause.

Genes of DNA repair strongly relate to age at menopause⁷, suggesting that the latter is the result of overall somatic aging²⁰. Although the genetic loci associated with age at menopause have not yet been directly correlated to human longevity or familial longevity, a meta-analysis relating SNPs to exceptional human longevity (in single cases) reported a correlation with several of the same SNPs that related to age at menopause²¹. Because the genetic predisposition for a late onset of menopause is not significantly associated to the familial component of human longevity in our study, the late reproducing females' health might also be affected by other factors that influence lifespan in singletons (long lived persons without long lived family members) such as good environmental circumstances, healthy lifestyles, or favorable social factors.

Besides oocyte quantity, oocyte quality is a necessary factor for reproductive success and is thought to be a causal factor of age-related fertility decline²². As females age, oocyte competence decreases, leading to an increased risk of aneuploidy and miscarriage, in turn leading to decreased fecundity. Suggested pathways of oocyte quality decline include deterioration of the maintenance of mitochondrial function²³ and the intrafollicular processes of DNA translation²⁴. These processes are in turn thought to be subject to oxidative stress, to which the oocyte becomes more vulnerable with increasing age, as shown in animal models²⁵. Fertility treatments such as in vitro fertilization (IVF) have previously, though not consistently, been linked to adverse cardiovascular outcomes in the short and long term²⁶⁻²⁸, which could suggest an adverse aging profile for the sub-fertile population, but it is not clear whether this can be attributed to effects of the treatment or population risk. While it is possible that the influence of the DNA damage repair genes associated with age at menopause additionally extends to oocyte quality, this remains to be further determined. Perhaps in oocytes, a group of cells that spend most of their life in

senescence, the pathways for cell maintenance are regulated somewhat differently than in somatic cells. This could be another explanation for the lack of association between familial longevity and the polygenic risk score for menopause as well as reproductive outcomes in our study. It is also possible that fecundity, and thus oocyte quality, cannot be adequately measured through the proxies of age at childbirth and number of offspring. In addition, because we study a by-definition relatively healthy group of females who lived to age 50 and underwent at least one successful pregnancy and delivery, our study precludes an indepth inquiry into the association between infertility or involuntary childlessness and longevity.

The method of linkage of families in the historical cohort makes our study uniquely suited to study the familial effects of reproduction and longevity. The methodological selection of the study population as well as the population size add to its strengths. Due to the historical nature of the data, the results are not influenced by the use of hormonal contraception or assisted reproductive techniques, therefore allowing for a reasonable assumption of unrestricted natural fertility.

Our results are limited by the obligatory use of proxy variables for fertility and reproductive success, as we were limited to the data stored in governmental registries. Furthermore, as mentioned we included a relatively healthy group of individuals who lived to be at least 50 years old. It is possible that this selection excludes individuals with accelerated aging genotypes and therefore attenuates any associations of reproduction with longevity.

In conclusion, we affirm that a late age at last childbirth is associated with a longer lifespan, and that traits of reproductive success seem to be markers of females' health in middle age, likely acquired by good environmental circumstances. Furthermore, we

- 328 conclude that neither parental nor more extended ancestral familial longevity are
- 329 characterized by reproductive success.

Table 1. LINKS study population selected for females that gave birth to at least one child.

	Total number	Mean ± SD	Range
Number F1 RPs (N, % female)	10,255 (100)	-	_
Top 10% survivors of their birth cohort	2,241 (21.8)		
Number of unique sibships (N)	7,721	-	-
Birth year (Mean)	-	1839	1812-1873
Age at death in years (Mean ± SD)	-	73.9 ± 10.4	50-104
Number of children (Mean ± SD)	72,985	7.1 ± 3.9	1-24
Number of children ≤ 4 (N, %)	2,990 (29.2)	2.6 ± 1.1	1-4
Number of children ≥ 10 (N, %)	2,771 (27.1)	12.2 ± 2.1	10-24
Age at first child in years (Mean ± SD)	-	26.9 ± 4.9	15-49
Age at last child in years (Mean ± SD)	-	39.2 ± 5.0	18-51
Last childbirth ≤ 40 years (N, %)	5,190 (50.6)	35.5 ± 4.4	18-40
Last childbirth ≥ 45 years (N, %)	981 (9.6)	45.8 ± 1.0	45-51
Age at marriage lpha in years (Mean \pm SD)	-	25.9 ± 4.6	16-46
Number of identified FO parents (N, %)	15,300 (99.1)	-	-
RPs with 0 long-lived# parents (N, %)	8,293 (80.9)	-	-
RPs with 1 long-lived [#] parent (N, %)	1,849 (18.0)	-	-
RPs with 2 long-lived [#] parents (N, %)	113 (1.1)	-	-

 $^{^{\}alpha}$ In the case of multiple marriages, age at first marriage was considered. *Belonging to the top 10% survivors of their birth cohort. 15.3% (2,344) of the identified parents have a missing age at death.

Table 2. Age at last child birth and number of offspring associates with female lifespan.

	N (mean/proportion)	Beta (CI)	P-value
Age at last child (quantitative) Age at last child (qualitative)	10,255 (39.23)	0.06 (0.02- 0.10)	2.16 x·10 ⁻³
Group 0: ≤ 40 years Group 1: ≥ 45 years	5,190 (50.6) 981 (9.6)	REF 1.41 (0.71-2012)	REF 9.07 x·10 ⁻⁵
Number of children (quantitative) Number of children (qualitative)	10,255 (7.12)	0.05 (-0.01-0.11)	7.63 x·10 ⁻²
Group 0: ≤ 4	2,990 (29.2)	REF	REF
Group 1: ≥ 10	2,771 (27.1)	0.61 (-0.03-1.25)	6.25 x 10 ⁻²

All analyses are adjusted for the maternal birth year and RPs age at marriage. In addition, our research design accounts for survival of RPs and their partner up to the age of 50 years and the number of marriages. Analyses are done using mixed-model linear regression using the lme4 and ImerTest package in R. Confidence intervals have been calculated in R with the "confint" function using the Wald method. 4 separate analyses have been done with age at death as outcome; 1 where age at last child (quantitative definition) was independent variable of interest, 1 where age at last child (qualitative definition) was independent variable of interest, 1 where number of children (qualitative definition) was independent variable of interest.

Table 3. Association of age at last child birth and number of children with familial longevity.

	N (mean/proportion)	IRR (CI)	P-value
Age at last child			
Number of long-lived parents			
0 long-lived parents	8,293 (80.9)	REF	REF
1 long-lived parent	1,849 (18.0)	0.99 (0.98-1.00)	5.28 x·10 ⁻²
2 long-lived parents	113 (1.1)	1.00 (0.97-1.03)	9.99 x·10 ⁻¹
RP long-lived			
No	8,014 (78.2)	REF	REF
Yes	2,241 (21.8)	1.01 (1.00-1.02)	2.75 x 10 ⁻³
Number of children			
Number of long-lived parents			
0 long-lived parents	8,293 (80.9)	REF	REF
1 long-lived parent	1,849 (18.0)	0.98 (0.96-1.01)	1.62 x·10 ⁻¹
2 long-lived parents	113 (1.1)	0.98 (0.89-1.08)	6.44 x 10 ⁻¹
RP long-lived	·	·	
No	8,014 (78.2)	REF	REF
Yes	2,241 (21.8)	1.02 (1.00-1.04)	9 19 x 10 ⁻²

All analyses are adjusted for the maternal birth year and RPs age at marriage. In addition, our research design accounts for survival of RPs and their partner up to the age of 50 years and the number of marriages. Analyses are done using mixed-model poissons regression using the lme4 and lmerTest package in R. Confidence intervals have been calculated in R with the "confint" function using the Wald method. 4 separate analyses have been done; 1 where age at last child was the outcome and the number of long lived parents was independent variable of interest, 1 where age at last child was the outcome and RP long-lived yes/no was independent variable of interest, and 1 where number of children was the outcome and RP long-lived yes/no was independent variable of interest, and 1 where number of children was the outcome and RP long-lived yes/no was independent variable of interest.

References 357 358 359 360 361 References 362 1. Laven JSE, Visser JA, Uitterlinden AG, Vermeij WP, Hoeijmakers JHJ. Menopause: Genome 363 stability as new paradigm. Maturitas. 2016; 92: 15-23. 364 2. Gagnon A, Smith KR, Tremblay M, Vézina H, Paré PP, Desjardins B. Is there a trade-off 365 between fertility and longevity? A comparative study of women from three large historical 366 databases accounting for mortality selection. Am J Hum Biol. 2009; 21: 533-540. 367 3. Gagnon A. Natural fertility and longevity. Fertil Steril. 2015; 103: 1109-1116. 368 4. Perls TT, Alpert L, Fretts RC. Middle-aged mothers live longer. Nature. 1997; 389: 133. 369 5. Smith KR, Gagnon A, Cawthon RM, Mineau GP, Mazan R, Desjardins B. Familial 370 aggregation of survival and late female reproduction. J Gerontol A Biol Sci Med Sci. 2009; 64: 371 740-744. 372 6. Westendorp RG, Kirkwood TB. Human longevity at the cost of reproductive success. 373 Nature. 1998; 396: 743-746. 374 7. Ruth KS, Day FR, Hussain J, Martínez-Marchal A, Aiken CE, Azad A, Thompson DJ, 375 Knoblochova L, Abe H, Tarry-Adkins JL, Gonzalez JM, Fontanillas P, Claringbould A, Bakker 376 OB, Sulem P, Walters RG, Terao C, Turon S, Horikoshi M, Lin K, Onland-Moret NC, Sankar A,

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

Hertz EPT, Timshel PN, Shukla V, Borup R, Olsen KW, Aguilera P, Ferrer-Roda M, Huang Y, Stankovic S, Timmers PRHJ, Ahearn TU, Alizadeh BZ, Naderi E, Andrulis IL, Arnold AM, Aronson KJ, Augustinsson A, Bandinelli S, Barbieri CM, Beaumont RN, Becher H, Beckmann MW, Benonisdottir S, Bergmann S, Bochud M, Boerwinkle E, Bojesen SE, Bolla MK, Boomsma DI, Bowker N, Brody JA, Broer L, Buring JE, Campbell A, Campbell H, Castelao JE, Catamo E, Chanock SJ, Chenevix-Trench G, Ciullo M, Corre T, Couch FJ, Cox A, Crisponi L, Cross SS, Cucca F, Czene K, Smith GD, de Geus EJCN, de Mutsert R, De Vivo I, Demerath EW, Dennis J, Dunning AM, Dwek M, Eriksson M, Esko T, Fasching PA, Faul JD, Ferrucci L, Franceschini N, Frayling TM, Gago-Dominguez M, Mezzavilla M, García-Closas M, Gieger C, Giles GG, Grallert H, Gudbjartsson DF, Gudnason V, Guénel P, Haiman CA, Håkansson N, Hall P, Hayward C, He C, He W, Heiss G, Høffding MK, Hopper JL, Hottenga JJ, Hu F, Hunter D, Ikram MA, Jackson RD, Joaquim MDR, John EM, Joshi PK, Karasik D, Kardia SLR, Kartsonaki C, Karlsson R, Kitahara CM, Kolcic I, Kooperberg C, Kraft P, Kurian AW, Kutalik Z, La Bianca M, LaChance G, Langenberg C, Launer LJ, Laven JSE, Lawlor DA, Le Marchand L, Li J, Lindblom A, Lindstrom S, Lindstrom T, Linet M, Liu Y, Liu S, Luan J, Mägi R, Magnusson PKE, Mangino M, Mannermaa A, Marco B, Marten J, Martin NG, Mbarek H, McKnight B, Medland SE, Meisinger C, Meitinger T, Menni C, Metspalu A, Milani L, Milne RL, Montgomery GW, Mook-Kanamori DO, Mulas A, Mulligan AM, Murray A, Nalls MA, Newman A, Noordam R, Nutile T, Nyholt DR, Olshan AF, Olsson H, Painter JN, Patel AV, Pedersen NL, Perjakova N, Peters A, Peters U, Pharoah PDP, Polasek O, Porcu E, Psaty BM, Rahman I, Rennert G, Rennert HS, Ridker PM, Ring SM, Robino A, Rose LM, Rosendaal FR, Rossouw J, Rudan I, Rueedi R, Ruggiero D, Sala CF, Saloustros E, Sandler DP, Sanna S, Sawyer EJ, Sarnowski C, Schlessinger D, Schmidt MK, Schoemaker MJ, Schraut KE, Scott C, Shekari S, Shrikhande A, Smith AV, Smith BH, Smith JA, Sorice R, Southey MC, Spector TD, Spinelli JJ, Stampfer M, Stöckl D, van

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

Meurs JBJ, Strauch K, Styrkarsdottir U, Swerdlow AJ, Tanaka T, Teras LR, Teumer A, Porsteinsdottir U, Timpson NJ, Toniolo D, Traglia M, Troester MA, Truong T, Tyrrell J, Uitterlinden AG, Ulivi S, Vachon CM, Vitart V, Völker U, Vollenweider P, Völzke H, Wang Q, Wareham NJ, Weinberg CR, Weir DR, Wilcox AN, van Dijk KW, Willemsen G, Wilson JF, Wolffenbuttel BHR, Wolk A, Wood AR, Zhao W, Zygmunt M, Biobank-based Integrative Omics Study (BIOS) Consortium, eQTLGen Consortium, Biobank Japan Project, China Kadoorie Biobank Collaborative Group, kConFab Investigators, LifeLines Cohort Study, InterAct consortium, 23andMe Research Team, Chen Z, Li L, Franke L, Burgess S, Deelen P, Pers TH, Grøndahl ML, Andersen CY, Pujol A, Lopez-Contreras AJ, Daniel JA, Stefansson K, Chang-Claude J, van der Schouw YT, Lunetta KL, Chasman DI, Easton DF, Visser JA, Ozanne SE, Namekawa SH, Solc P, Murabito JM, Ong KK, Hoffmann ER, Murray A, Roig I, Perry JRB. Genetic insights into biological mechanisms governing human ovarian ageing. Nature. 2021; 8. van den Berg N, van Dijk IK, Mourits RJ, Slagboom PE, Janssens AAPO, Mandemakers K. Families in comparison: An individual-level comparison of life-course and family reconstructions between population and vital event registers. Popul Stud (Camb). 2021; 75: 91-110. 9. Schoenmaker M, de Craen AJM, de Meijer PHEM, Beekman M, Blauw GJ, Slagboom PE, Westendorp RGJ. Evidence of genetic enrichment for exceptional survival using a family approach: the Leiden Longevity Study. Eur J Hum Genet. 2006; 14: 79-84.

10. Schoenmaker M, de Craen AJM, de Meijer PHEM, Beekman M, Blauw GJ, Slagboom PE, Westendorp RGJ. Evidence of genetic enrichment for exceptional survival using a family approach: the Leiden Longevity Study. Eur J Hum Genet. 2006; 14: 79-84. 11. Beekman M, Blauw GJ, Houwing-Duistermaat JJ, Brandt BW, Westendorp RGJ, Slagboom PE. Chromosome 4q25, microsomal transfer protein gene, and human longevity: novel data and a meta-analysis of association studies. J Gerontol A Biol Sci Med Sci. 2006; 61: 355-362. 12. van den Berg N, Rodríguez-Girondo M, Mandemakers K, Janssens AAPO, Beekman M, Slagboom PE. Longevity Relatives Count score identifies heritable longevity carriers and suggests case improvement in genetic studies. Aging Cell. 2020; 19: e13139. 13. Choi SW, Mak TS, O'Reilly PF. Tutorial: a guide to performing polygenic risk score analyses. Nat Protoc. 2020; 15: 2759-2772. 14. van den Berg N, Rodríguez-Girondo M, van Dijk IK, Mourits RJ, Mandemakers K, Janssens AAPO, Beekman M, Smith KR, Slagboom PE. Longevity defined as top 10% survivors and beyond is transmitted as a quantitative genetic trait. Nat Commun. 2019; 10: 35-0. 15. Helle S, Lummaa V, Jokela J. Are reproductive and somatic senescence coupled in humans? Late, but not early, reproduction correlated with longevity in historical Sami women. Proc Biol Sci. 2005; 272: 29-37. 16. Sun F, Sebastiani P, Schupf N, Bae H, Andersen SL, McIntosh A, Abel H, Elo IT, Perls TT. Extended maternal age at birth of last child and women's longevity in the Long Life Family

Study. *Menopause*. 2015; 22: 26-31.

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

- 17. Müller HG, Chiou JM, Carey JR, Wang JL. Fertility and life span: late children enhance
- female longevity. J Gerontol A Biol Sci Med Sci. 2002; 57: 202.
- 18. McArdle PF, Pollin TI, O'Connell JR, Sorkin JD, Agarwala R, Schäffer AA, Streeten EA, King
- 444 TM, Shuldiner AR, Mitchell BD. Does having children extend life span? A genealogical study
- of parity and longevity in the Amish. J Gerontol A Biol Sci Med Sci. 2006; 61: 190-195.
- 19. Jaffe D, Kogan L, Manor O, Gielchinsky Y, Dior U, Laufer N. Influence of late-age births on
- 447 maternal longevity. *Ann Epidemiol*. 2015; 25: 387-391.
- 448 20. Laven JSE. Early menopause results from instead of causes premature general ageing.
- 449 Reprod Biomed Online. 2022; 45: 421-424.
- 450 21. Sebastiani P, Bae H, Sun FX, Andersen SL, Daw EW, Malovini A, Kojima T, Hirose N,
- 451 Schupf N, Puca A, Perls TT. Meta-analysis of genetic variants associated with human
- 452 exceptional longevity. Aging (Albany NY). 2013; 5: 653-661.
- 453 22. Silber SJ, Kato K, Aoyama N, Yabuuchi A, Skaletsky H, Fan Y, Shinohara K, Yatabe N,
- Kobayashi T. Intrinsic fertility of human oocytes. Fertil Steril. 2017; 107: 1232-1237.
- 455 23. Chiang JL, Shukla P, Pagidas K, Ahmed NS, Karri S, Gunn DD, Hurd WW, Singh KK.
- 456 Mitochondria in Ovarian Aging and Reproductive Longevity. Ageing Res Rev. 2020; 63:
- 457 101168.
- 458 24. Llarena N, Hine C. Reproductive Longevity and Aging: Geroscience Approaches to
- 459 Maintain Long-Term Ovarian Fitness. *J Gerontol A Biol Sci Med Sci.* 2021; 76: 1551-1560.

460 25. Sasaki H, Hamatani T, Kamijo S, Iwai M, Kobanawa M, Ogawa S, Miyado K, Tanaka M. 461 Impact of Oxidative Stress on Age-Associated Decline in Oocyte Developmental 462 Competence. Front Endocrinol (Lausanne). 2019; 10: 811. 463 26. Henriksson P. Cardiovascular problems associated with IVF therapy. J Intern Med. 2021; 464 289: 2-11. 465 27. Yiallourou SR, Magliano D, Haregu TN, Carrington MJ, Rolnik DL, Rombauts L, Rodrigues 466 A, Ball J, Bruinsma FJ, Da Silva Costa F. Long term all-cause and cardiovascular disease 467 mortality among women who undergo fertility treatment. Med J Aust. 2022; 217: 532-537. 468 28. Bungum AB, Glazer CH, Arendt LH, Schmidt L, Pinborg A, Bonde JP, Tøttenborg SS. Risk of 469 hospitalization for early onset of cardiovascular disease among infertile women: a register-470 based cohort study. Hum Reprod. 2019; 34: 2274-2281. 471 472